

Rook I Project

Environmental Impact Statement

TSD XXI: Environmental Risk Assessment

Ecometrix Environmental INTELLIGENCE

ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT

TECHNICAL SUPPORT DOCUMENT

REPORT PREPARED FOR:

NexGen Energy Ltd. 3150-1021 W Hastings St Vancouver, BC V6E 0C3

REPORT PREPARED BY:

Ecometrix Incorporated www.ecometrix.ca
Mississauga, ON

Ref. 19-2574 April 2024



ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT

TECHNICAL SUPPORT DOCUMENT

Langping Wu, Ph.D.
Environmental Modeller

Rina Parker, M.A.Sc., P.Eng

Rima Packer

Project Manager and Technical Lead

Don Hart, Ph.D.

Project Principal and Reviewer

EXECUTIVE SUMMARY

NexGen Energy Ltd. (NexGen) is proposing to develop a new uranium mining and milling operation in northwestern Saskatchewan, called the Rook I Project (Project). The proposed Project is subject to both federal and provincial Environmental Assessment (EA) processes, and the Environmental Impact Statement (EIS) was prepared to support the EA. The environmental risk assessment (ERA) is used to inform the EA. The ERA encompasses a human health risk assessment (HHRA) and an ecological risk assessment (EcoRA), which have been prepared to be compliant with Canadian Standards Association Group (CSA) N288.6-22 Environmental Risk Assessments for Class I Nuclear Facilities and Uranium Mines and Mills (CSA 2022), and Canadian Nuclear Safety Commission Regulatory Document 2.9.1, Environmental Principles, Assessments and Protection Measures (CNSC 2020).

The ERA focused on constituents of potential concern (COPCs) that exceeded screening values in air and water based on predicted atmospheric releases and aqueous releases (e.g., mine effluent, sewage effluent, site runoff, and groundwater solute releases) from the Project. Based on the screening of atmospheric releases, no COPCs in air were advanced for further quantitative assessment in the ERA. Based on the screening of aqueous releases, arsenic, cobalt, copper, molybdenum, uranium, sulphate, and chloride were advanced for further quantitative assessment in the ERA. Radionuclides, including the uranium-238 series and radon, were included as COPCs because these constituents are of public interest.

An environmental transport and pathways model (IMPACT) was used to evaluate the effects of COPCs on the local environment including human and ecological receptors.

The ERA estimated dose and risk to human and ecological receptors for the Application Case, reasonable upper bound sensitivity scenario, and the Reasonably Foreseeable Development Case (RFD) for all phases of the Project (i.e., the Project lifespan) and the far-future projection.

The selection of human and ecological receptors was informed by Indigenous Knowledge and Traditional Land Use (IKTLU) Studies, information obtained during community information sessions, and Joint Working Group (JWG) meetings.

The HHRA focused on members of the public potentially exposed to low levels of airborne or waterborne constituents. The selected human health receptor groups included:

- camp worker at Patterson Lake camp residence;
- subsistence harvesters;
- seasonal residents/lodge operators; and
- future permanent resident.



The ecological receptors selected for the EcoRA were a subset of valued components identified for the EA so that the results from the ERA could be used in the effects assessments for fish, vegetation, wildlife, human health, Indigenous land and resource use, and other land and resource uses.

Non-radiological Human Health Risk Assessment

The potential effects on human receptors were evaluated by comparing the non-radiological exposures of receptors to recognized benchmarks, a dose-based toxicity reference value in the same units, for each COPC. For assessment of non-carcinogens, risk was estimated based on a Project hazard quotient (HQ). Project HQs were compared to a benchmark HQ value of 0.2 because exposures to background sources were not included. This approach is consistent with Health Canada's guidance on human health preliminary quantitative risk assessment (Health Canada 2021a).

No significant adverse effect on any human receptors, as a result of releases from the Project, was considered likely during all Project phases (Operations was considered the bounding Project phase) for the Application Case, reasonable upper bound sensitivity scenario, and RFD Case. The predicted Project HQs for all non-carcinogenic COPCs (cobalt, copper, molybdenum, and uranium) would remain below the acceptable risk level of 0.2 per pathway for all human receptors.

For assessment of risk for carcinogens (arsenic), the incremental risk of developing cancer over a lifetime (ILCR) was estimated and compared against the cancer risk level of 1 in 100,000 recommended by Health Canada (2021a). Incremental cancer risk was predicted to exceed the negligible cancer risk level of 1 in 100,000 for the subsistence harvester at Patterson Lake South Arm just outside the Project footprint but is not expected to exceed the negligible cancer risk within the RSA farther from the Project. These findings are based on the conservative assumption of high consumption of Traditional Foods including fish and terrestrial animals in the Project footprint and LSA.

Radiological Human Health Risk Assessment

The incremental radiation dose to all human receptors during all Project phases and the far-future projection were predicted to be below the regulatory public dose limit of 1 millisieverts per year (mSv/yr) for the Application Case, the reasonable upper bound sensitivity scenario, and the RFD Case.

If a dose constraint of 0.3 mSv/yr is applied, the dose to the subsistence harvester (one-year-old) would be less than the dose constraint for the Application Case, the reasonable upper bound sensitivity scenario, and RFD Case, and well below the regulatory public dose limit.

In the far-future projection, a future permanent resident, living at the decommissioned and reclaimed Project site following Closure would receive a dose up to 0.07 mSv/yr, well below the regulatory public dose limit and the dose constraint.



Overall, since the radiation dose estimates were predicted to be below the public dose limit, no discernable health effects would be anticipated due to exposure of human receptors to radioactive releases from the Project.

Non-radiological Ecological Risk Assessment

The potential for ecological effects was assessed by comparing exposure levels to toxicological benchmarks and characterized quantitatively in terms of HQs. An HQ greater than 1 indicates adverse effects may be possible for a given ecological receptor and further investigation would be warranted.

No significant adverse effect on either aquatic or terrestrial populations or communities, as a result of releases from the Project, would be likely during all Project phases for the Application Case, reasonable upper bound sensitivity scenario, and RFD Case. All estimated HQs for all COPCs (chloride, sulphate, arsenic, cobalt, copper, molybdenum, and uranium) for all ecological receptors were predicted to remain below the HQ benchmark of 1.

For the far-future projection, once groundwater solutes reach Patterson Lake North Arm – West Basin, HQs would remain below the benchmark of 1 for all COPCs with the exception of copper for the Application Case, reasonable upper bound sensitivity scenario, and RFD Case. Exceedances of the HQ benchmark value of 1 for the far-future projection would be limited spatially to Patterson Lake and would be limited in magnitude to just above the benchmark of 1. Exceedances of the HQ benchmark were predicted for lake whitefish, benthic invertebrates, and zooplankton.

The toxicity reference value (TRV) for lake whitefish, benthic invertebrates, and zooplankton corresponds to the Canadian Council of Ministers of the Environment's (CCME) default water quality guideline for copper. The CCME water quality guideline is a conservative benchmark for copper below which adverse effects for most aquatic biota are not expected to occur. The most sensitive species include fresh water snails (*Lymnaea stagnalis* and *Pyrgulopsis robusta*), bivalves (*Villosa iris*), and water flea (*Daphnia magna*; ECCC 2021b). The most sensitive endpoints for chronic exposure to copper include growth for benthic invertebrates, reproduction for zooplankton, and growth and reproduction for fish (ECCC 2021b). The CCME water quality guideline is based on no or low effect levels for the most sensitive species; thus, the marginal TRV exceedances predicted by HQs just above 1 would indicate a possible slight reduction in growth or reproduction for exposed individuals of sensitive species in Patterson Lake North Arm – West Basin. However, since the HQ exceedance would be localized to Patterson Lake North Arm – West Basin, population level effects would not be expected.

Species at risk were assessed using surrogate species. The rusty blackbird and little brown myotis were identified as surrogate species at risk. Since no HQs above 1 were identified for these ecological receptors, individual species at risk would also be considered protected.



Radiological Ecological Risk Assessment

Radiation dose benchmarks of 9.6 milligrays per day (mGy/d) and 2.4 mGy/d (UNSCEAR 2008) were selected for the assessment of effects on aquatic biota and terrestrial biota, respectively, as recommended in CSA N288.6-22.

There were no predicted exceedances of the 9.6 mGy/d radiation dose benchmark for aquatic biota and 2.4 mGy/d radiation dose benchmark for terrestrial and riparian biota for the Application Case, the reasonable upper bound sensitivity scenario, and the RFD Case, during the Project phases and the far-future projection.

Since there were no predicted exceedances of the respective dose benchmarks for any of the aquatic or terrestrial receptors, individual species at risk would also be considered protected.

Monitoring and Follow-up

The HHRA and EcoRA were developed based on the best available information for the Rook I Project, including baseline monitoring data, assumptions on estimates of source terms, Traditional Food diet (consumption rates and food types), among others.

Monitoring would focus on collecting data to verify ERA model predictions as well as provide data to improve model predictions as the Project begins. Recommended monitoring would support NexGen's adaptive management framework with the goal of reducing uncertainty over time through an iterative process.



TABLE OF CONTENTS

EXEC	UIIVI	E SUMMARY	l
TABI	E OF	CONTENTS	V
LIST	OF TA	ABLES	VIII
LIST	OF FIG	GURES	XI
LIST	OF AF	PPENDICES	XI
ABBI	REVIA	TIONS AND UNITS OF MEASURE	XII
1.0	INTE	RODUCTION	1.1
	1.1	Regulatory Context	1.5
	1.2	Objectives	1.5
		Scope Spatial Boundaries Temporal Boundaries Assessment Cases	1.6 1.9
	1.4	Statement of Limitations	1.11
2.0	SITE	DESCRIPTION	2.1
	2.1	Project Description	2.1
	2.2	Description of the Natural and Physical Environment	2.1
	2.3	Uncertainty in Site Characterization	2.2
3.0	INTE	GRATION OF INDIGENOUS AND LOCAL KNOWLEDGE	3.1
	3.1	Community Information Sessions	3.2
	3.2	Joint Working Groups	3.3
	3.3	Indigenous Knowledge and Traditional Land Use Studies	3.5
4.0	МОГ	DEL INTEGRATION AND EVALUATION OF SOURCES	4.1
	4.1	Model Integration	4.1
	4.2 4.2.1	Aqueous Sources Project-Related Aqueous Releases	

	4.2.2	Aqueous Sources	4.3
	4.2.3	Screening for Constituents of Potential Concern in Aquatic Environment	4.3
	4.2.4	Evaluation of Selenium in Fish Tissue	4.15
	4.2.5	Water Quality Constituents for Further Evaluation in the Environmental Risk Ass	essment 4.16
	4.3	Atmospheric Sources	
	4.3.1	Project-Related Atmospheric Releases	
		Atmospheric Sources	
		Screening for Atmospheric Constituents of Potential Concern	
	4.3.4	Air Quality Constituents for Further Evaluation in the Environmental Risk Assess	
	4.4 4.4	Final List of Constituents of Potential Concern for the Environmental Risk A	Assessment
5.0	HUM	1AN HEALTH RISK ASSESSMENT	5.1
	5.1	Problem Formulation	5.1
	5.1.1	Human Receptor Selection and Characterization	
	5.1.2 Asses	Constituents of Potential Concern Retained for Further Evaluation in the Human sement 5.8	n Health Risk
		Selection of Exposure Pathways	
		Human Health Conceptual Site Model	
	5.1.5	Uncertainty in Problem Formulation	
	5.2	Exposure Assessment	
	5.2.1	Exposure Locations, Duration, and Frequency	
		Exposure and Dose Calculations	
		Exposure Point Consentration and Doses	
		Exposure Point Concentration and Doses Uncertainty in Exposure Assessment	
	5.3	Toxicity Assessment	
	5.3.1	,	
		Radiation Dose Limits and Targets Uncertainty in the Toxicity Assessment	
		•	
		Risk Characterization	
	5.4.1	Risk Estimation	
	5.4.2	Uncertainty in the Risk Characterization	5.97
6.0	ECO	LOGICAL RISK ASSESSMENT	6.1
	6.1	Problem Formulation	
		Selected Ecological Receptors	
		Constituents of Potential Concern Retained for Further Evaluation in the Ecolog	ical Risk
		Assessment and Massurement Fodge into	C 17
		Assessment and Measurement Endpoints	
		Selection of Exposure Pathways Ecological Conceptual Site Model	
	6.2	Exposure Assessment	
	6.2.1	Exposure Locations	6.26

	6.2.2	Exposure Averaging	6.28
	6.2.3	Exposure and Dose Calculations	6.29
		Exposure Factors	
		Exposure Point Concentrations and Doses	
	6.2.6	Uncertainty in Exposure Assessment	6.48
	6.3	Effects Assessment	
	6.3.1	Toxicological Benchmarks	
		Radiation Benchmarks	
	6.3.3	Uncertainty in the Effects Assessment	6.64
	6.4	Risk Characterization	
		Risk Estimation	
	6.4.2	Uncertainty in the Risk Characterization	6.78
7.0	QUA	LITY ASSURANCE AND SENSITIVITY ANALYSIS	7.1
	7.1	Quality Assurance and Quality Control	7.1
	7.2	Sensitivity Analysis	7.1
	7.2.1	Effects of Annual Weather Patterns	7.2
	7.2.2	Deposition of Constituents of Potential Concern on Soil and Plants	7.6
		Traditional Food Assumptions for Subsistence Harvesters	
	7.2.4	Climate Change	7.9
8.0	CON	CLUSIONS AND MONITORING	8.1
	8.1	Human Health Risk Assessment	8.1
	8.1.1	Non-radiological Human Health Risk Assessment	8.1
	8.1.2	Radiological Human Health Risk Assessment	8.2
	8.2	Ecological Risk Assessment	8.2
	8.2.1	Non-radiological Ecological Risk Assessment	8.2
	8.2.2	Radiological Ecological Risk Assessment	8.3
	8.3	Monitoring and Follow-up	8.4
9.0	REFE	RENCES	9.1
	9.1	Acts and Regulations	9.1
	0.2	Literature Cited	0.1

LIST OF TABLES

Table 1-1: Spatial Boundaries for the ERA	1.7
Table 1-2: Project Phases Assessed in the Environmental Risk Assessment	1.9
Table 3-1: Dates and Locations of June 2019 Community Information Sessions	
Table 3-2: Participation at the Community Information Sessions	
Table 4-1: Screening Values for the Selection of Surface Water Quality Constituents of	
Potential Concern for the Environmental Risk Assessment	4.7
Table 4-2: Surface Water Screening of Constituents of Potential Concern for the	
Environmental Risk Assessment	4.10
Table 4-3: Sediment Quality Screening for the Rook I Project	4.14
Table 4-4: Background Concentrations of Air Quality Constituents Used in The Air Quality Model	4.19
Table 4-5: Concordance between Air Quality Model and Receptor Locations	4.20
Table 4-6: Screening Values for the Selection of Air Quality Contaminants of Potential	
Concern for the Environmental Risk Assessment	4.23
Table 4-7: Air Quality Screening for Short-term and Long-term Exposures to Constituents	
in Air at Human and Ecological Receptor Locations	4.29
Table 4-8: Air Quality Screening for Short-term Exposures to Constituents in Air for	
Hypothetical Human and Ecological Receptors at the Fence Line	4.32
Table 4-9: Summary of Air Quality Constituents that Exceed a Screening Value	4.35
Table 4-10: Soil Quality Screening for the Rook I Project	4.41
Table 4-11: Final List of Constituents of Potential Concern for the Project Environmental	
Risk Assessment	4.43
Table 5-1: Rationale for Selection of Human Health Receptor Groups	5.2
Table 5-2: Human Health Exposure Pathways Considered for the Rook I Project	5.9
Table 5-3: Traditional Food Diets for Each Human Receptor Group	5.10
Table 5-4: Summary of Residency Assumptions for Human Health Receptor Groups	5.19
Table 5-5: Annual Ingestion Rates for Traditional Food Items of the Total Food Diet for	
Non-consumers, and Average and High Consumers of Traditional Foods	5.25
Table 5-6: Representative Ecological Receptors used in the Traditional Foods Diet	5.27
Table 5-7: Estimated Non-carcinogen Doses to Human Receptors – Operations –	
Application Case and Upper Bound Scenario	5.31
Table 5-8: Estimated Carcinogen Doses for Arsenic to Human Receptors – Operations –	
Application Case and Upper Bound Scenario	5.38
Table 5-9: Estimated Radiation Doses to Human Receptors – Operations – Application	
Case, Upper Bound Scenario	5.40
Table 5-10: Estimated Radiation Doses to Human Receptors – Far-Future – Application	
Case, Upper Bound Scenario	
Table 5-11: Summary of Input Parameters for Radon Dose Calculation	5.50
Table 5-12: Estimated Non-carcinogen Doses to Human Receptors – Reasonably	
Foreseeable Development Case	5.51



Table 5-13:	Estimated Carcinogen Doses for Arsenic to Human Receptors – Reasonably	
	Foreseeable Development Case – Operations and Far-Future Projection	5.60
Table 5-14:	Estimated Radiation Doses to Human Receptors – Reasonably Foreseeable	
	Development Case	5.62
Table 5-15:	How Uncertainties in the Human Health Exposure Assessment Are Addressed	5.68
Table 5-16:	Human Health Oral Exposure Toxicity Reference Values	5.70
Table 5-17:	Radiation Dose Limits for Human Health	5.72
Table 5-18:	Estimated Non-carcinogen Risk to Human Receptors – Project Lifespan and	
	Far-Future – Application Case and Upper Bound Scenario	5.76
Table 5-19:	Estimated Incremental Lifetime Cancer Risk from Arsenic to Human Receptors	
	- Application Case and Upper Bound Scenario	5.82
Table 5-20:	Estimated Carcinogen Risks for Arsenic to Seasonal Residents – Operations –	
	Application Case and Upper Bound Scenario	5.83
Table 5-21:	Estimated Carcinogen Risks for Arsenic to Subsistence Harvesters – Operations	
	- Application Case and Upper Bound Scenario	5.84
Table 5-22:	Summary of All Radiation Doses to Human Receptors – Application Case and	- 0-
T	Upper Bound Scenario	5.85
Table 5-23:	Total Radiation Dose to Camp Worker from all Radionuclides including Radon	F 0.0
Т-Ы-Г 24.	Progeny – Project Lifespan	5.86
Table 5-24:	Estimated Non-carcinogen Risk to Human Receptors – Operations –	г 07
Tabla E 2E.	Reasonably Foreseeable Development Case Estimated Incremental Lifetime Cancer Risk from Arsenic to Human Receptors	.5.87
1 able 5-25.	Reasonably Foreseeable Development Case	E 02
Tabla 5 26.	Estimated Carcinogen Risks for Arsenic to Subsistence Harvesters and Seasonal	. 5.95
Table 3-20.	Resident – Operations – Reasonably Foreseeable Development	5 01
Tahla 5-27·	Summary of All Radiation Doses to Human Receptors – Reasonably	. 3.34
Table 5-21.	Foreseeable Development Case	5 95
Table 5-28 [.]	Summary of Radiation Doses to Limiting Human Receptors	
	Total Radiation Dose to Camp Worker from all Radionuclides including Radon	. 5.50
Tuble 5 E5.	Progeny – Project Lifespan (RFD Case)	5.97
Table 6-1: F	Rationale for Selection of Ecological Receptors for the Rook I Project Ecological	. 5.5 .
	Risk Assessment	6.6
Table 6-2: 9	Selected Ecological Receptors for the Rook I Project Ecological Risk Assessment	
	ist of Identified Species with Conservation Status in the Vicinity of the Project	
	Surrogate Species for Identified Species at Risk	
	Assessment Endpoints, Measurement Endpoints, and Lines of Evidence	
	Complete Exposure Pathways for All Selected Ecological Receptors to be	
	Assessed using the IMPACT Model	6.22
Table 6-7: F	Residency Factors for Terrestrial Ecological Receptors with Large Home Ranges	6.28
	Body Weights and Intake Rates for Bird and Mammal Ecological Receptors	
	Exposure Factors Used in the Rook I Project IMPACT Model	
	Estimated Non-radiological Doses to Ecological Receptors – Operations –	
	Application Case and Upper Bound Scenario	6 35



Table 6-11: Estimated Non-radiological Doses to Ecological Receptors – Far-Future	
Projection – Application Case and Upper Bound Scenario	6.37
Table 6-13: Estimated Non-radiological Doses to Ecological Receptors – Reasonably	
Foreseeable Development Case	
Table 6-14: Procedure for Adjusting Test Endpoints to Chronic 20% Effect Concentration	
Table 6-15: Selected Toxicity Reference Values for Aquatic Biota	
Table 6-16: Selected Toxicity Reference Values for Terrestrial Biota	6.58
Table 6-17: Summary of Selected Toxicity Reference Values for Mammal and Bird Test	
Species – Arsenic	6.59
Table 6-18: Summary of Selected Toxicity Reference Values for Mammal and Bird Test	
Species – Cobalt	6.60
Table 6-19: Summary of Selected Toxicity Reference Values for Mammal and Bird Test	
Species – Copper	6.61
Table 6-20: Summary of Selected Toxicity Reference Values for Mammal and Bird Test	
Species – Molybdenum	6.62
Table 6-21: Summary of Selected Toxicity Reference Values for Mammal and Bird Test	
Species – Uranium	
Table 6-22: Radiological Dose Benchmarks for Aquatic and Terrestrial Ecological Receptors .	6.63
Table 6-23: Non-radiological Risk to Ecological Receptors – Application Case and Upper	
Bound Scenario – Operations	6.68
Table 6-24: Non-radiological Risk to Ecological Receptors – Application Case and Upper	
Bound Scenario – Far-Future Projection	6.71
Table 6-25: Non-radiological Risk to Ecological Receptors – Reasonably Foreseeable	
Development Case – Operations and Far-Future Projection	6.75
Table 7-1: Average Monthly Outflows and Average Annual Outflows of the Modelled	
Waterbodies	7.3
Table 7-2: Parameters Used in Modelling Deposition of Airborne Constituents of Potential	
Concern on Soil and Plants	7.6
Table 7-3: Comparison of Arsenic Bioavailability Assumptions and the Local Intake	
Fractions of Traditional Food for Adult Subsistence Harvester	7.8
Table 7-4: Estimated Incremental Lifetime Cancer Risk from Arsenic to Subsistence	
Harvester – Sensitivity Scenario	7.8



LIST OF FIGURES

Figure 1-1: Location of the Rook I Project	1.2
Figure 1-2: Regional Area of the Rook I Project	
Figure 1-3: Layout of Infrastructure and Facilities for the Rook I Project	
Figure 1-4: Rook I Project Local Study Area and Regional Study Area Boundaries	1.8
Figure 4-1: Model Integration Diagram	4.1
Figure 4-2: Selection of Surface Water Screening Values for Constituents of Potential	
Concern for the Environmental Risk Assessment	4.5
Figure 4-3: Screening Process for Selection of Constituents of Potential Concern for the	
Environmental Risk Assessment	4.6
Figure 4-4: Comparison of Predicted Selenium Fish Tissue Concentrations against Criterion	4.16
Figure 5-1: Human Receptor Locations	5.7
Figure 5-2: Location of Rook I Site with Respect to Saskatchewan Ecozones	5.14
Figure 5-3: Proportions of Traditional Food Categories in the Boreal Shield, Boreal Plain,	
and Averaged Diets Reviewed during Engagement Activities with Local	
Communities	5.15
Figure 5-4: Proportional Distribution of Traditional Food Items in Average and High	
Traditional Diets Selected for the Human Health Risk Assessment	5.15
Figure 5-5: Human Health Conceptual Site Model	5.17
Figure 6-1: Ecological Health Conceptual Site Model	6.25
Figure 6-2: Locations of Ecological Receptors Assessed in the Ecological Risk Assessment	6.27
Figure 7-1: Concentrations of Copper and Polonium-210 in Patterson Lake North Arm –	
West Basin during Operations	7.4
Figure 7-2: Total Dose to the Subsistence Harvester (One-Year-Old) at Patterson Lake	
South Arm during Operations	7.5
LIST OF APPENDICES	
Appendix AIMPACT Model Report	9.1
Appendix B Ecological Receptor Profiles	
Appendix CModel Results in Support of the Environmental Risk Assessment	



ABBREVIATIONS AND UNITS OF MEASURE

Abbreviation	Definition
AAQC	ambient air quality criteria
ATSDR	Agency for Toxic Substances and Disease Registry
BAF	bioaccumulation factor
BC MECCS	British Columbia Ministry of Environment and Climate Change Strategy
BC MOE	British Columbia Ministry of Environment
BNDN	Birch Narrows Dene Nation
BRDN	Buffalo River Dene Nation
CAAQS	Canadian Ambient Air Quality Standards
CAC	criteria air contaminant
CCC	criterion continuous concentration
CCME	Canadian Council of Ministers of the Environment
CNSC	Canadian Nuclear Safety Commission
COPC	constituent of potential concern
COSEWIC	Committee on the Status of Endangered Wildlife in Canada
CRDN	Clearwater River Dene Nation
CSA	Canadian Standards Association
DCF	dose coefficient
EA	Environmental Assessment
EC	effect concentration
ECCC	Environment and Climate Change Canada
EcoRA	ecological risk assessment
Eco-SSL	ecological soil screening level
ECOTOX	ECOTOXicology database
EIS	Environmental Impact Statement
ENV	Saskatchewan Ministry of Environment
ERA	environmental risk assessment
ESL	effects screening level
FEQG	federal environmental quality guidelines
FNFNES	First Nations Food, Nutrition and Environment Study
HABISask	Hunting, Angling and Biodiversity Information of Saskatchewan
HHRA	human health risk assessment
HQ	hazard quotient
IC	inhibition concentration
ICRP	International Commission on Radiological Protection
IKTLU	Indigenous Knowledge and Traditional Land Use
ILCR	incremental lifetime cancer risk
ISQG	interim sediment quality guideline
JWG	Joint Working Group
LC	lethal concentration
LEL	lowest effect level



Abbreviation Definition

LOAEL lowest observed adverse effect level

LSA local study area

MN-S Métis Nation – Saskatchewan

NE2 no-effect

NexGen Energy Ltd.

NOAEL no observed adverse effect level
NPAG non-potentially acid generating
OAAQC Ontario Ambient Air Quality Criteria

OF occupancy factor

PAG potentially acid generating

PEL probable effect level PM particulate matter

PM₁₀ particulate matter with a diameter of 10 microns or less PM_{2.5} particulate matter with a diameter of 2.5 microns or less

Project Rook I Project REF reference

RFD reasonably foreseeable development

RSA regional study area

SAAQS Saskatchewan Ambient Air Quality Standards

SAR Species at Risk
SARA Species at Risk Act
SEL severe effect level

SEQG Saskatchewan environmental quality guidelines

TF transfer factor

TRV toxicity reference value
TSD Technical Support Document

TSP total suspended particulates

UGTMF underground tailings management facility

UL upper intake level

UNSCEAR United Nations Scientific Committee on the Effects of Atomic Radiation

USEPA United States Environmental Protection Agency

VC valued component

WHO World Health Organization WQO water quality objective

YNLR Ya'thi Néné Lands and Resources



UnitDefinition%percentμmmicron

μg/m³ micrograms per cubic metre Bq/m³ becquerels per cubic metre

dw dry weight fw fresh weight

g/m²/yr grams per square metre per year

h hour

h/yr hours per year kg/yr kilograms per year

km kilometre

km² square kilometre

m metre

m/s metres per second

mg CaCO₃/L milligrams of calcium carbonate per litre

mg/kg/d milligrams per kilogram per day

mg/L milligrams per litre
mGy/d milligrays per day
mGy/h milligrays per hour
mo/yr months per year
mSv/yr millisieverts per year

1.0 INTRODUCTION


NexGen Energy Ltd. (NexGen) is proposing to develop a new uranium mining and milling operation in northwestern Saskatchewan, called the Rook I Project (Project). The Project would be located approximately 40 km east of the Saskatchewan-Alberta border, 130 km north of the town of La Loche, and 640 km northwest of the city of Saskatoon (Figure 1-1). The Project would reside within Treaty 8 territory and the Métis Homeland. At a regional scale, the Project would be situated within the southern Athabasca Basin adjacent to Patterson Lake, along the upper Clearwater River system. Patterson Lake is at the interface of the Boreal Shield and Boreal Plain ecozones. Access to the Project would be from an existing road off Highway 955 (Figure 1-2), with on-site worker accommodation serviced by fly-in/fly-out access.


The Project would include the following key facilities to support the extraction and processing of uranium from the Arrow deposit for transportation off site (Figure 1-3):


- underground mine development;
- process plant buildings, including uranium concentrate packaging facilities;
- paste tailings distribution system;
- underground tailings management facility (UGTMF);
- potentially acid generating (PAG) waste rock storage area;
- non-potentially acid generating (NPAG) waste rock storage area;
- special waste rock¹ and ore storage stockpiles;
- surface and underground water management infrastructure, including water management ponds, effluent treatment plant, and sewage treatment plant;
- conventional waste management facilities and fuel storage facilities;
- ancillary infrastructure, including maintenance shop, warehouse, administration building, and camp;
- airstrip and associated infrastructure; and
- access road to Project and site roads.

_

 $^{^{1}}$ Special waste rock is mine rock that is mineralized with insufficient grade to be considered ore (i.e., greater than 0.03% of triuranium octoxide [U₃O₈] and less than 0.26% U₃O₈). All special waste would be temporarily stored in the special waste rock stockpile.







1.1 Regulatory Context

The proposed Project is subject to both federal and provincial Environmental Assessment (EA) processes, and the Environmental Impact Statement (EIS) was prepared to support the EA. This environmental risk assessment (ERA) encompasses a human health risk assessment (HHRA) and an ecological risk assessment (EcoRA), which have been prepared to be compliant with Canadian Standards Association Group (CSA) N288.6-22 Environmental Risk Assessments for Class I Nuclear Facilities and Uranium Mines and Mills (CSA 2022). It also meets the requirements for an ERA outlined in Section 4.1 of Regulatory Document-2.9.1, Environmental Principles, Assessments and Protection Measures (CNSC 2020). The ERA has been developed with current science and current regulatory attitudes in mind.

The ERA was used to inform other EA disciplines and to support the conclusions made in the EIS. The regulatory and guidance documents applicable to the EIS were discussed in the EIS.

1.2 Objectives

The objectives of this ERA were to:

- predict and assess the risk to representative human and ecological receptors resulting from exposure to radiological and non-radiological substances expected to be released to the environment throughout the phases of the Rook I Project;
- inform decision-making in the EIS; and
- inform prioritization of monitoring and mitigation measures.

1.3 Scope

The ERA used the expected sources of atmospheric and liquid releases to predict the transport of these constituents through the environment, exposure and dose to the public, and exposure and effects on representative ecological receptors.

The scope of the ERA encompassed both human and ecological health risks, and radiological and non-radiological constituents of potential concern (COPCs).

The EIS assessed valued components (VCs) which are aspects of the biophysical, cultural, and socio-economic environments that are considered to have scientific, social, cultural, economic, historical, archaeological, or aesthetic importance (Beanlands and Duinker 1983; CNSC 2021). The receptors selected for the ERA included a subset of VCs identified for the EIS, so that the results from the ERA could be used in the effects assessments for fish, vegetation, wildlife, human health, Indigenous land and resource use, and other land and resource use.

Physical stressors were considered in the assessments of other VCs in the EIS, outside of the ERA. The assessment of VCs in the EIS included a pathways analysis to determine how Project activities may interact with the environment. For primary pathways resulting in a potential effect after mitigation, a residual effects analysis was conducted. Physical stressors were considered in the

assessments of fish and fish habitat (EIS Section 11) and wildlife and wildlife habitat (EIS Section 14). With respect to the assessment of fish and fish habitat, EIS Section 11.4.1 (No Pathways) and Section 11.4.2 (Secondary Pathways) identified multiple pathways with the potential to create physical stressors on the fish and fish habitat VCs. However, these pathways were deemed to have negligible or measurable but minor effects and were not assessed further. With respect to the assessment of wildlife and wildlife habitat, EIS Section 14.4.2 (Secondary Pathways) also identified multiple pathways with the potential to create physical stressors on the wildlife and wildlife habitat VCs. However, these pathways were deemed to have measurable but minor effects and were not assessed further. In addition, sensory disturbance (e.g., lights, dust, smells, noise) to wildlife was identified as a primary pathway, which was then assessed for wildlife and wildlife habitat VCs in EIS Section 14.5 (Residual Effects Analysis) and three species at risk not selected as VCs in EIS Appendix 14A (Species at Risk Screening Assessment).

The ERA is supported by the IMPACT Model Report (Appendix A), which presents the structure and functioning of the IMPACT model as implemented for the Project. The model was used to evaluate the transport and effects of COPCs on the local environment and receptors, including humans and non-human biota.

1.3.1 Spatial Boundaries

The spatial boundaries of the ERA were consistent with the boundaries defined in the EIS for the aquatic and terrestrial environment, and included the site study area (Project footprint), the local study area (LSA) and a larger regional study area (RSA; Figure 1-4, Table 1-1). The RSA spatial boundary extends from the headwaters of the Clearwater River to the confluence of the Clearwater River and the Mirror River and includes major waterbodies along its course including Broach Lake, Patterson Lake, Forrest Lake, Beet Lake, Naomi Lake, and Lloyd Lake, as well as their contributing watersheds. The human health RSA encompasses the RSA for the aquatic and terrestrial environments, but also includes Lloyd Lake because of the Lloyd Lake Lodge, a fly-in lodge located on the western shore of Lloyd Lake that provides guided fishing and spring bear hunting trips for guests:

- **Site Study Area (Project footprint):** The Project footprint includes the camp where workers reside while at work.
- Local study area (LSA): The LSA is defined at a scale that contains the expected direct effects of the Project on selected receptors. The LSA represents the area where direct Project-related changes in the quality of air, sediment, water, and soils would be expected to occur. The LSA also represents the area where local movements in fish and wildlife could be expected to occur, and is an area where potential environmental effects from Project activities and components can be predicted or measured with a suitable degree of accuracy and confidence.
- Regional study area (RSA): The RSA is defined to be an appropriate scale for the
 assessment of cumulative effects where there is potential for spatial overlap or interactions
 with Project effects and other previous, and existing developments, and reasonably
 foreseeable developments.

Table 1-1: Spatial Boundaries for the ERA

Study Area	Area	Description/Rationale	
Site study area	228 ha (2.3 km²)	 Equivalent to the anticipated Project footprint, which includes all proposed mine infrastructure and facilities (199 ha), the access road (29 ha), and accommodations where workers reside while at work. 	
LSA	68,530 ha (685 km²)	 Encompasses the LSA for the aquatic and terrestrial environments and defines the expected extent of the direct and indirect effects from the Project on selected receptors. Includes the spatial extents of potential air quality effects. Provides local context for assessing the residual effects. 	
RSA	107,000 ha (1,070 km²)	Encompasses the RSA for the aquatic and terrestrial environments, is also includes Lloyd Lake (cumulative watershed area to Lloyd Lake outflow is 4 370 km²)	




Figure 1-4: Rook I Project Local Study Area and Regional Study Area Boundaries

1.3.2 Temporal Boundaries

The ERA assessed three phases: Construction, Operations, and Decommissioning and Reclamation (i.e., Closure), as well as a far-future projection, as described in Table 1-2. The Rook I Project is anticipated to last 43 years, after which time closure performance criteria would be fully demonstrated.

Construction is anticipated to last four years. Operations are anticipated to last 24 years. Closure is anticipated to last 15 years, which consists of the Active Closure Stage and the Transitional Monitoring Stage. The far-future projection represents the period after the closure performance criteria have been fully demonstrated.

Table 1-2: Project Phases Assessed in the Environmental Risk Assessment

Phase Description	
Project lifespan (43 years total)	Construction Phase (4 years): includes site preparation; mine, process plant, and additional infrastructure development; transportation of people and materials to and from the Project; and all activities associated with commissioning the Project up until the Operations Phase commences.
	Operations Phase (24 years) : includes all activities associated with mining and processing ore; tailings management; management of mine rock, domestic waste, and hazardous materials; water management; release of treated effluent; surface storage of clean material; site maintenance; progressive reclamation; and transportation of staff and materials to and from the Project.
	 Decommissioning and Reclamation (i.e., Closure) Phase (15 years): includes two stages: Active Closure Stage and Transitional Monitoring Stage: Active Closure Stage (5 years): includes active decommissioning and reclamation activities post-Operations such as backfilling mine workings, removal of physical infrastructure, recontouring and revegetating disturbed areas, waste disposal or removal, water management and treated effluent discharge (if required), and any other activities required to achieve decommissioning objectives and return the site to a safe and stable condition prior to the Transitional Monitoring Stage. Transitional Monitoring Stage (10 years): may take longer than 10 years; this stage will continue until monitoring and reporting verifies that the performance criteria have been met. Once performance criteria have been fully demonstrated, an application to be released from the CNSC licence would be submitted to the CNSC for approval. Once that is achieved, and upon Provincial approval, the land would be transferred back under Provincial management through the Institutional Control Program. This stage is nominally 10 years; however, NexGen acknowledges this is dependent on the achievement of performance criteria.

Phase	Description
Far-Future projection (not a Project phase)	The far-future projection encompasses the long-term timeframe during which extremely slow migration of COPCs from the UGTMF and waste rock storage area to the environment are anticipated. The far-future projection is applicable to groundwater and surface water quality intermediate components, and human health VCs and ecological receptors, which are assessed through the ERA. Far-future is a projection of the long-term return to steady-state conditions.

CNSC = Canadian Nuclear Safety Commission; VC = valued component; ERA = environmental risk assessment; COPC = constituents of potential concern; UGTMF = underground tailings management facility.

1.3.3 Assessment Cases

Assessment cases distinguish between existing, proposed, and future projects. Consistent with the EIS, the assessment cases included a Base Case, an Application Case, and a Reasonably Foreseeable Development (RFD) Case:

- The **Base Case** is represented by existing conditions. The Base Case describes the existing environment in the LSA and RSA before the application of the proposed Project to provide an understanding of the current physical, biological, economic, social, and cultural conditions that may be influenced by the Project.
- The **Application Case** represents predictions of the combined effects of the Base Case plus the effects from the proposed Project. This case also assesses incremental, Project-specific changes that are predicted to occur to human and ecological receptors:
 - A sensitivity scenario was developed as part of the Application Case. It was representative of reasonable upper bound, or 95th percentile, conditions. The purpose of this scenario was to examine a highly conservative estimation for water quality predictions to bound the uncertainty associated with the model input data. This scenario was characterized by Base Case conditions and Project effects, using reasonable upper bound source-term inputs from the site wide water balance model (SWWBM) and the groundwater solute transport model.
- The **RFD Case** includes the combined effects of developments under the Base Case and Application Case, and RFDs that have not yet been approved:
 - The RFD Case includes the Fission Patterson Lake South Property, which is planned to be located on the west shore of Patterson Lake, at the most westerly point where the North Arm and South Arm meet. The Fission Patterson Lake South Property is expected to withdraw fresh water from the Patterson Lake North Arm West Basin and discharge treated effluent and treated sewage to the Patterson Lake South Arm. Public information describes a projected three-year construction period and seven-year operating period (production and processing) (Fission 2019, 2021).

1.4 Statement of Limitations

This report has been prepared by Ecometrix Incorporated (Ecometrix) and includes environmental conclusions. In preparing the report, Ecometrix relied in good faith on data, information collected, and modelling results by others and made available to Ecometrix. Ecometrix did not independently confirm such information, unless specifically stated, and does not accept responsibility for any deficiencies, inaccuracies or misstatements in the work of others as provided, nor for conditions or issues outside of the scope of work.

Any and all conclusions made by Ecometrix in the report represent Ecometrix's professional judgement, and are based on understanding of the site conditions as described by the information made available at the time of and relied upon for the report preparation. The report is to be read in its entirety; sections or parts are not to be taken out of the context of the whole report.

Ecometrix prepared the report exercising the same standard of care, skill and diligence required by the professional practices and procedures that would normally be provided in the preparation of similar projects under similar conditions. Nothing in this report is intended to provide or constitute a legal opinion.

This report has been prepared for the sole and exclusive use of NexGen. Any use of, reliance on, or decision made by a third party on the basis of this report is the sole responsibility of such third parties. Ecometrix does not accept any responsibility for damages, if any, that are suffered by third parties because of their reliance on or use of this report to make decisions or take actions.

2.0 SITE DESCRIPTION

2.1 Project Description

The Project would include underground mining to access the uranium ore of the Arrow deposit. A process plant would be located on surface directly above the mine to process the ore on site. Tailings from the processed ore would be returned below ground to the UGTMF, which represents a key environmental design feature of the Project that avoids the permanent removal of land and associated risks to the environment.

2.2 Description of the Natural and Physical Environment

The proposed Project would be situated within the southern Athabasca Basin adjacent to Patterson Lake, along the upper Clearwater River system. The Project site would be located within the Patterson Lake Watershed, which is within the larger Clearwater River Watershed that drains to the Mackenzie River Watershed via the Athabasca and Slave rivers.

Climatic conditions at the proposed Project site are considered sub-arctic with temperatures ranging from warmer than 30°C in the summer to colder than -40°C during the winter. The proposed Project site is covered by 30 m to 100 m of glacial drift over Cretaceous mudstone, which are fine-grained clay particles that have been compressed by overlying material over a long time. The glacial drift is comprised primarily of sand with gravels, cobbles, and boulders. Drumlins, lakes, wetlands, rivers, streams, and muskegs are common in the Project vicinity. Elevations in the region range from 583 metres above sea level (masl) at the crest of major drumlins to 480 masl for some of the lowland lakes. The Project site is dominated by sandstone (i.e., sand-sized grains of rock material), with some bedrock outcroppings (i.e., rock that rises above the surface).

The broader regional area of the proposed Project intersects the Boreal Shield and Boreal Plain ecozones. At a smaller more local scale, the proposed Project is located within the Boreal Plain Ecozone of the Mid-Boreal Uplands Ecoregion. The area surrounding the proposed Project consists of recent burns with residual stands of jack pine (*Pinus banksiana*) and some black spruce (*Picea mariana*), with shrub and lichen as ground cover. Provincial wildfire data indicate that large areas in the region have been burned over the last 40 years.

The broader regional area surrounding the proposed Project is largely undisturbed by human activities and infrastructure; approximately 0.5 percent (%) of the regional area (1,000 square kilometres [km²]) encompassing the Patterson Lake Watershed has been influenced by human developments. Most human-related disturbances in this regional area include linear features such as Highway 955, cutlines, seismic lines, and trails, with some cleared areas.



2.3 Uncertainty in Site Characterization

The Project LSA and RSA are well characterized with regards to biophysical data. Baseline monitoring has been ongoing since 2018, which has provided sufficient baseline information to support the EIS and ERA. No residual uncertainties in the site characterization have been identified. Information from the baseline studies and Indigenous Knowledge and Traditional Land Use (IKTLU) Studies were used to inform the assumptions of the ERA and underlying models. Assumptions were also validated through discussions during community information sessions, Joint Working Group (JWG) meetings, and regulatory meetings, which resulted in refinements and changes to previously made assumptions.



3.0 INTEGRATION OF INDIGENOUS AND LOCAL KNOWLEDGE

The Project would be located within Treaty 8 territory, which is an area spanning 840,000 km² and covers parts of Saskatchewan, Alberta, British Columbia, and the Northwest Territories. Primary Indigenous Groups, as identified by the CNSC and the Saskatchewan Ministry of Environment (ENV), and corroborated by NexGen early engagement, who have Study Agreements with NexGen in 2019 include (in alphabetical order):

- Birch Narrows Dene Nation (BNDN);
- Buffalo River Dene Nation (BRDN);
- Clearwater River Dene Nation (CRDN); and
- Métis Nation Saskatchewan (MN-S), including as on behalf of the Locals of MN-S Northern Region 2.

In addition to the primary Indigenous Groups, the Ya'thi Néné Lands and Resources (YNLR), which represents the Fond du Lac Denesųliné First Nation and the Black Lake Denesųliné First Nation, also signed a Study Funding Agreement in 2020 and developed an Interim Report IKTLU Study that informed the ERA.

Indigenous Knowledge and Traditional Land Use Studies include all different forms of studies developed by Indigenous Groups who have completed IKTLUs for the proposed Project; these include Traditional Land Use and Occupancy studies, Traditional Knowledge and Use studies, and Indigenous Rights and Knowledge studies.

Based on information from IKTLU Studies, the Patterson Lake area is used by these Indigenous Groups for resource use activities such as hunting, fishing, trapping, and gathering (TSD II: BNDN; TSD III: BRDN; TSD IV: MN-S; TSD V.1: CRDN; TSD V.2: CRDN; TSD VI: YNLR). Feedback from engagement, JWGs, and IKTLU Studies have also identified several important themes that require consideration:

- importance of maintaining safety and health of water and fish in Patterson Lake, including consideration of potential cumulative effects from the Project and the Fission Patterson Lake South Property;
- ability to access land and resource use during and after the lifespan of the mine, and if the land would be safe to use after the mine closes;
- risks to human safety from traffic and spills due to increases in traffic and subsequent effects on the roads;
- benefits of training, employment, and contracting, but understanding the challenges (or barriers) to acquiring jobs and the negative effects associated with increased income (e.g., time away from families, loss of connection with the land); and



• importance of historical and cultural context, which includes a deep connection to the water and land for the necessities of life.

Indigenous Knowledge, as documented in the IKTLU Studies and JWG meeting minutes, continues to play a key role in developing the assumptions for the ERA, including the specific human and ecological receptors selected for the assessment, and locations and characteristics of these receptors. Where Indigenous Knowledge has been used, this is documented in the relevant section of the ERA report.

A summary of the forum for gathering Indigenous Knowledge from community information sessions, JWG meetings, and IKTLU Studies is outlined below.

3.1 Community Information Sessions

Community information sessions were hosted by NexGen in June 2019. The objectives of the sessions were to:

- inform local communities of the nature of proposed activities at the Project;
- answer questions and receive initial feedback specific to the Project for consideration during the EA, including feedback on VCs;
- provide information about the EA process; and
- introduce NexGen and the Project to the broader community.

Table 3-1 outlines the date, location and communities invited to the four community information sessions.

Table 3-1: Dates and Locations of June 2019 Community Information Sessions

Location	Date	Communities Notified of the Event
La Loche Community Hall, La Loche, SK	Monday 24 June 2019	 La Loche, SK CRDN Descharme Lake, SK Bear Creek, SK Black Point, SK Garson Lake, SK
Birch Narrows Arena, BNDN	Tuesday 25 June 2019	BNDNTurnor, SK
Buffalo River Arena, BRDN	Wednesday 26 June 2019	 BRDN Dillon, SK Michelle Village, SK St. George's Hill, SK
Lakeview Complex, Buffalo Narrows, SK	Thursday 27 June 2019	Buffalo Narrows, SK

SK = Saskatchewan; BNDN = Birch Narrows Dene Nation; BRDN = Buffalo River Dene Nation; CRDN = Clearwater River Dene Nation.



The community information sessions used a drop-in format, with a series of ten poster presentations, which were staffed by NexGen representatives. Copies of the posters were available in pamphlet form for participants to keep. The poster stations contained information about NexGen, the Project, by-product management, reclamation, water, radiation, the EA, valued components, commitment to communities land uses, and next steps.

As part of the process, community members were invited to share information about their local land and resource uses in proximity to the Project site through a map-based exercise. Information collected during the mapping exercise informed the initial selection of receptor locations in the LSA and RSA area relative to existing or previous uses.

A total of 266 participants signed in to the four sessions, with Table 3-2 outlining the participants at each session.

Table 3-2: Participation at the Community Information Sessions

Location	Date	Signed Attendance
La Loche	Monday, 24 June 2019	163
BNDN	Tuesday, 25 June 2019	32
BRDN	Wednesday, 26 June 2019	27
Buffalo Narrows	Thursday, 27 June 2019	44

BRDN = Buffalo River Dene Nation; BNDN = Birch Narrows Dene Nation.

3.2 Joint Working Groups

Joint Working Groups were assembled from representatives of the respective Indigenous Groups and NexGen and are an important engagement platform to allow for ongoing collaboration to enable the incorporation of Indigenous Group views and Indigenous and Local Knowledge into the Project.

The JWGs were formed through the Study Agreements established with each Indigenous community in 2019 and have been implemented as the agreed-upon engagement pathway for the advancement of the EA. The four JWGs are CRDN-JWG, BNDN-JWG, BRDN-JWG, and the MN-S-JWG representing Northern Region 2.

Key objectives of the JWGs are to:

- develop protocols for using and protecting Indigenous Knowledge;
- identify VCs for the Project EIS;
- identify and address community-specific information in the EA process;
- discuss potential Project effects and mitigation of those effects; and
- identify other topics that are important to the local Indigenous Groups and communities.



ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT INTEGRATION OF INDIGENOUS AND LOCAL KNOWLEDGE

Indigenous Groups selected JWG participants with consideration of diverse representation, including Elders, youth, men and women, business owners, and traditional land users in the Patterson Lake area.

The ERA was initially introduced to the JWGs on the following dates:

- BNDN on 25 October 2019 and again on 4 December 2019;
- BRDN on 1 November 2019;
- MN-S on 29 October 2019; and
- CRDN on 19 February 2020.

It was understood through the Indigenous Knowledge shared at the JWG meetings that Patterson Lake, Forrest Lake, Clearwater River, and other waterbodies and watercourses in the region are important to the local Indigenous Groups for the use of water, fish, and wildlife, cultural connections to the land, and community well-being (TSD II: BNDN; TSD III: BRDN; MN-S 2019; TSD IV: MN-S; CRDN 2019a, 2019b; TSD V.1: CRDN; TSD V.2: CRDN, YNLRO 2019; TSD VI YNLR). Key highlights from the JWG meetings strengthened the assertion that the protection of local waterbodies is paramount to the security of Indigenous well-being:

Water is the most important thing, vital to life... ensuring water remains safe is the biggest concern with the project. (BNDN-JWG 2019a)

The best lake in Saskatchewan is right here – we don't need to buy water. (BRDN-JWG 2019a)

All the lakes in the area are important, Patterson Lake especially so, as it feeds the lakes to the south and affects all the waterways from which many members fish. It is central to the river system for the entire area. Patterson Lake is an integral part of community fishing. (TSD IV: MN-S)

Additionally, discussions throughout the JWG meetings allowed Indigenous community members to share their knowledge and observations about the perceived deterioration of the environment and the local ecosystem due to increasing industrial development within the immediate and wider Athabasca Basin region. Indigenous Knowledge also indicated concerns about the cumulative effects from human development and policies on climate change in the area (TSD II: BNDN; TSD III: BRDN; TSD IV: MN-S; MN-S 2019; CRDN 2019a, 2019b; TSD V.1: CRDN; TSD V.2: CRDN; YNLRO 2019; TSD VI: YNLR). The MN-S, in particular, identified concerns that they do not want Patterson Lake to become like Lake Athabasca, which they claim has been impacted by industry to the point where the fish are inedible. They have noticed changes in the water quality and flow, aquatic life, and terrestrial life in the areas farther north of the Project and within the study areas of this assessment (i.e., RSA), due to cumulative effects of industry in the area. Some of the significant changes noted by the communities during the JWG discussions include:



Population decline:

o "In the last few years, I notice that change in the nature. Around NexGen and Fission, the animals are changing – there used to be a lot of rabbits and chickens. The last few years I've been hunting in fall, and there's not many anymore. The animals are not there like they used to be" (MN-S-JWG 2020).

Deteriorating game quality:

- "Community members have noticed a change in the quality of meat and pelts. Members have noted increased contamination in the food, especially from further north; others expressed a fear of eating fish, rabbits and moose from the north. It was further noted that rabbits around Patterson Lake are diseased, having white spots when opened; likewise, the moose are diseased, glowing when opened" (TSD IV: MN-S).
- Changes in herd migration and distribution:
 - "You see that migration routes change when infrastructure is put in. Even power lines, caribou won't cross. Communities that used to see caribou, no longer see them come through" (BRDN-JWG 2019b).

3.3 Indigenous Knowledge and Traditional Land Use Studies

Indigenous Knowledge and Traditional Land Use Studies were used to inform initial assumptions for human health receptors (i.e., people) who consume Traditional Foods or fish, hunt or gather Traditional Foods in terms of locations, residency times, and components of the Traditional Foods diet.

The following IKTLU Studies were used to inform assumptions:

- TSD II (BNDN), Birch Narrows Dene Nation Traditional Knowledge and Use Study Specific to NexGen Energy Limited's Proposed Rook I Project;
- TSD III (BRDN), Buffalo River Dene Nation Traditional Knowledge and Use Study Specific to NexGen Energy Limited's Proposed Rook I Project;
- TSD IV (MN-S), Métis Nation Saskatchewan Northern Region 2 Traditional Land Use & Diet Study for the NexGen Rook I Project;
- TSD V.1 (CRDN), Preliminary Identification of Issues and Concerns Related to the Proposed NexGen Energy Ltd. Rook I Project in the Patterson Lake Area; A Review; Clearwater River Dene Nation; Traditional Land Use and Occupancy Mapping Interviews; 2010 – 2016;
- TSD V.2 (CRDN), Clearwater River Dene Nation Indigenous Rights and Knowledge Survey related to the proposed NexGen Energy Ltd. Rook 1 Project in the Patterson Lake Area; and



ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT INTEGRATION OF INDIGENOUS AND LOCAL KNOWLEDGE

• TSD VI (YNLRO), Provision of Athabasca Denesuliné Traditional Knowledge, Land Use and Occupancy Information for the NexGen Rook I Project Environmental Assessment.

These IKTLU studies provided initial guidance for identifying locations where people may reside, areas where Traditional Foods are hunted, fished, and gathered, and mammal, bird, and plant species that are traditionally used by local Indigenous communities for food, medicine, and other traditional uses.



4.0 MODEL INTEGRATION AND EVALUATION OF SOURCES

4.1 Model Integration

The ERA integrated inputs and sources from other disciplines, as demonstrated at a high level in Figure 4-1. In general, the ERA received information directly from the site water balance, site water quality, solute transport model, and atmospheric modelling. These are described in more detail in the EIS Sections 7 to 10 and Appendices.




Figure 4-1: Model Integration Diagram

4.2 Aqueous Sources

4.2.1 Project-Related Aqueous Releases

Aqueous releases to Patterson Lake would include:

- 1. releases from the effluent treatment plant to Patterson Lake North Arm West Basin;
- 2. releases from the sewage treatment plant to Patterson Lake North Arm West Basin;
- 3. non-contact site runoff to Patterson Lake North Arm East Basin and to Patterson Lake North Arm West Basin; and
- 4. groundwater releases through the overburden and bedrock to Patterson Lake North Arm West Basin.

The effluent treatment plant would process water that would either be released to the environment through a diffuser, or further used in mill processes or underground. The effluent treatment plant would treat process plant effluents, mine water discharges, and site runoff from mineralized areas. The effluent treatment plant would involve a two-stage chemical treatment process designed to neutralize and remove metals from the effluent stream prior to pumping to the treated effluent water tank. The treated effluent water tank would overflow into one of four monitoring ponds to allow for storage of treated effluent until monitoring results confirm acceptable treated effluent quality and discharge criteria are met. Should approvals be received, and the Project be constructed and operated, treated effluent would be discharged in batches to Patterson Lake North Arm – West Basin via a diffuser.

The sewage treatment plant would be fed by the sewage collection system and the camp sewage system. The sewage collection system for the mine and process plant areas would consist of piped, underground collection systems from each building that would discharge to individual sewage holding tanks. The tank contents would be collected and hauled to the sewage treatment plant receiving station. Treated sewage from the treatment facility would be discharged to Patterson Lake North Arm – West Basin via an outfall parallel to but separate from the treated effluent diffuser.

Surface drainage refers to systems that collect or convey precipitation runoff. Surface drainage would discharge to the Patterson Lake North Arm – West Basin and Patterson Lake North Arm – East Basin. During Operations, the West Surface Runoff Discharge would include non-contact water from downstream of the mine-controlled area as well as non-mineralized contact water that has been tested and confirmed to be acceptable for release. During Operations, the East Surface Runoff would consist of non-contact water from outside the mine-controlled area. Following Closure, runoff from reclaimed, covered, and revegetated surfaces within the former mine-controlled area would be routed to the East Perimeter Diversion and discharged to the Patterson Lake North Arm – East Basin. Efforts would be made to divert natural site runoff away from site features. If water from precipitation events came in contact with contaminated areas or contact zones, the water would be captured, collected, and directed to respective site runoff points or collection areas. Contained water would be tested before release to the environment based on regulatory requirements; water that does not meet specification would report to the effluent treatment plant for treatment.

The fresh water supply intake would draw water from Patterson Lake North Arm – East Basin and would provide water for the mine area and for fire protection. Fresh water would be distributed through the fresh water pump station. Water for potable use would be pumped to the water treatment plant and then to the mine area administration building. The fresh water system would serve the mine and process plant dry areas, the wash bay, and the camp.

During Decommissioning and Reclamation, infrastructure would be removed from the site and the water management system would transition to a passive system that drains the remaining above-ground WRSAs. The active water treatment described above would be operated until passive treatment is deemed adequate to release surface water to Patterson Lake while protecting the environment and meeting applicable discharge criteria.

In the far-future projection, mine-affected groundwater would reach Patterson Lake North Arm – West Basin through groundwater flow pathways from the underground and at-grade sources. Groundwater originating at the UGTMF and stope backfill source areas is expected to migrate vertically upward, primarily through the fault and shear zones, then laterally through the sandstone, before discharging within Patterson Lake North Arm – West Basin. Based on the hydraulic gradients, hydraulic conductivities, pathway dimensions, and effective porosity values applied to the pathways (i.e., 0.015 for the fault zone and 0.098 for the sandstone), the advective groundwater travel time from the upper horizon of the mine to the discharge location at Patterson Lake is estimated to be approximately 1,000 years, as predicted by solute transport modelling completed for the EIS.

Groundwater originating beneath the waste rock area would travel through the overburden bidirectionally to the north and south, ultimately discharging in Patterson Lake for both pathways. The approximate advective groundwater travel time from the waste rock piles to Patterson Lake was estimated at 43 years to the north and 77 years to the south.

4.2.2 Aqueous Sources

In this context, sources refer to Project-related volume and mass inputs to the IMPACT model. Sources for the aquatic environment were obtained from the following models:

- Rook I site-wide water balance model for the release of runoff and treated effluent to Patterson Lake North Arm West Basin during Project phases;
- near-field water quality model for water quality in the immediate area of the Project at the boundary of the mixing zone during Project phases;
- groundwater solute transport model for the release of soluble constituents from mine sources (mine workings, UGTMF, and surface waste rock piles) to Patterson Lake North Arm – West Basin in the far-future projection; and
- regional surface water quality model for water quality after the addition of groundwater solute transport in the far-future projection.

4.2.3 Screening for Constituents of Potential Concern in Aquatic Environment

4.2.3.1 Screening Value Selection

The predicted concentrations of constituents released to Patterson Lake were compared against water quality objectives (WQOs) based on the hierarchy of water quality guidelines outlined in Figure 4-2, and the selected screening values are shown in Table 4-1. The most restrictive federal or provincial guideline for surface water quality, based on the Canadian Council of Ministers of the Environment (CCME 2021a) Canadian water quality guidelines for the protection of fresh water aquatic life, the federal environmental quality guidelines (FEQG), and the Saskatchewan environmental quality guidelines (SEQG; ENV 2021), were selected as the screening values for most surface water COPCs. Applicable guidelines were adjusted for pre-operational hardness and pH, where applicable.



For molybdenum, Saskatchewan Water Security Agency published an updated WQO for the protection of aquatic life based on current understanding of aquatic toxicity of molybdenum to fresh water aquatic organisms (WSA 2017). The WQO is based on the 5th percentile of the species sensitivity distribution and follows the CCME protocol (CCME 2007). The WQO for molybdenum of 7.6 milligrams per litre (mg/L) from BC MOE (2021) was adopted for the Project in preference over the CCME WQO of 0.073 mg/L.

Canadian drinking water quality guidelines were included for protection of human health (Health Canada 2020). These guidelines are based on current, published scientific research related to human health effects, aesthetic effects, and operational considerations. Health-based guidelines are established on the basis of a comprehensive review of the known health effects associated with each contaminant, on exposure levels, and on the availability of treatment and analytical technologies. Aesthetic effects (e.g., taste or odour) were taken into account when these play a role in determining whether consumers would consider the water drinkable, as is the case with copper. Where no Canadian drinking water quality guidelines were available, guidelines were obtained from the World Health Organization (WHO 2017) or British Columbia Ministry of Environment and Climate Change Strategy (BC MECCS 2020).

Major ions such as calcium, magnesium, and sodium were not considered COPCs. There is no evidence of adverse health effects from these major ions in drinking water (Health Canada 2020), and they were considered to be essentially non-toxic for aquatic biota. These major ions are effectively regulated on the cellular level and have not been associated with adverse effects in aquatic biota at environmental concentrations. Therefore, calcium, magnesium, and sodium were not carried forward as COPCs for further assessment in the ERA.

Phosphorus was not considered a COPC for the ERA. Phosphorus is present in the aquatic environment as phosphate, where it acts as a nutrient rather than a toxicant. The water quality guideline selected for screening is the interim Ontario Provincial Water Quality Objective, which was set to avoid nuisance concentrations of algae in lakes and is not relevant to ecological health. Therefore, phosphorus was not considered a COPC for ecological health; however, it was considered in the surface water quality section (EIS Section 10, Surface Water Quality and Sediment Quality) as a nutrient affecting primary productivity.

No formal screening was conducted for radionuclides. However, since radiation dose to human and ecological receptors is of great public and regulatory interest, the radionuclides in the uranium-238 decay series (uranium-238, uranium-234, thorium-230, radium-226, lead-210, polonium-210) were all carried forward as COPCs for further assessment in the ERA.

The screening was based on the predicted concentrations of constituents released to Patterson Lake in treated effluent or groundwater, based on the models identified in Section 4.2.2, Aqueous Sources. To verify that the screening process was sufficiently conservative, predicted upper bound concentrations were screened against WQOs based on the process outlined in Figure 4-3. As a first pass, upper bound end-of-pipe treated effluent concentrations were compared against WQOs. Those constituents with predicted upper bound treated effluent concentrations above WQOs were considered further for additional screening. Upper bound concentrations at the

boundary of the mixing zone were then compared against WQOs. Those constituents with predicted upper bound concentrations at the boundary of the mixing zone above WQOs were considered COPCs for further quantitative assessment in the ERA. Additionally, if upper bound concentrations of COPCs in runoff exceeded the WQOs, but did not exceed in the treated effluent, they were not considered COPCs in the ERA.

After the completion of Project phases (43 years total), it is assumed that groundwater solutes would start to make their way from mine-related sources (mine workings, UGTMF, and surface waste rock piles) to Patterson Lake North Arm – West Basin. A secondary screening of predicted upper bound water quality after the addition of groundwater solute loadings was performed to determine if concentrations of any additional COPCs exceeded WQOs. Maximum upper bound concentrations were used for comparison against WQOs.




Figure 4-2: Selection of Surface Water Screening Values for Constituents of Potential Concern for the Environmental Risk Assessment




Figure 4-3: Screening Process for Selection of Constituents of Potential Concern for the Environmental Risk Assessment

ERA = environmental risk assessment; COPCs = constituents of potential concern.

Table 4-1: Screening Values for the Selection of Surface Water Quality Constituents of Potential Concern for the Environmental Risk Assessment

Constituent	Units	CCME Protection Life	of Aquatic	Federal Environm Guidelines		Saskatchewan Env Quality Guideline		Othe	r	Drinkin	g Water Guidelines		Selected Screening	Source
		Long Term	Note	Long Term	Note	Long Term	Note	Long Term	Note	Health Canada ^(p)	Other Source	Note	Value	
General Parameters														
Total suspended solids	mg/L	Background + 5	n/a	n/a	n/a	Background + 5	n/a	n/a	n/a	n/a	n/a	n/a	Background + 5	SEQG/CCME
Major lons					•									•
Chloride	mg/L	1.20 × 10 ⁺⁰²	(f)	n/a	n/a	1.20 × 10 ⁺⁰²	n/a	n/a	n/a	None required	n/a	n/a	1.20 × 10 ⁺⁰²	SEQG
Sulphate	mg/L	n/a	n/a	n/a	n/a	n/a	n/a	$1.28 \times 10^{+02}$	(1)	5.00 × 10 ⁺⁰²	n/a	n/a	1.28 × 10 ⁺⁰²	вс мое
Nutrients														
Ammonia as nitrogen	mg/L	5.74 × 10 ⁺⁰⁰	(d)	n/a	n/a	5.74 × 10 ⁺⁰⁰	(d)	n/a	n/a	None required	n/a	n/a	5.74 × 10 ⁺⁰⁰	SEQG/CCME
Un-ionized ammonia as nitrogen	mg/L	1.56 × 10 ⁻⁰²	n/a	n/a	n/a	1.56 × 10 ⁻⁰²	n/a	n/a	n/a	None required	n/a	n/a	1.56 × 10 ⁻⁰²	SEQG/CCME
Nitrate as nitrogen	mg/L	3.00 × 10 ⁺⁰⁰	n/a	n/a	n/a	3.00 × 10 ⁺⁰⁰	n/a	n/a	n/a	1.00 × 10 ⁺⁰¹	n/a	n/a	3.00 × 10 ⁺⁰⁰	SEQG/CCME
Metals and Metalloids	1													•
Aluminum	mg/L	1.00 × 10 ⁻⁰¹	(e)	n/a	n/a	1.00 × 10 ⁻⁰¹	n/a	n/a	n/a	1.00 × 10 ⁻⁰¹	n/a	n/a	1.00 × 10 ⁻⁰¹	SEQG/CCME
Arsenic	mg/L	5.00 × 10 ⁻⁰³	n/a	n/a	n/a	5.00 × 10 ⁻⁰³	n/a	n/a	n/a	1.00 × 10 ⁻⁰²	n/a	n/a	5.00 × 10 ⁻⁰³	SEQG/CCME
Calcium	mg/L	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	None required	n/a	n/a	None required	n/a
Cadmium	mg/L	4.00 × 10 ⁻⁰⁵	n/a	n/a	n/a	4.00 × 10 ⁻⁰⁵	n/a	n/a	n/a	7.00 × 10 ⁻⁰³	n/a	n/a	4.00 × 10 ⁻⁰⁵	SEQG/CCME
Cobalt	mg/L	n/a	n/a	7.78 × 10 ⁻⁰⁴	(j)	n/a	n/a	n/a	n/a	n/a	1.00 × 10 ⁻⁰³	(p)	7.78 × 10 ⁻⁰⁴	FEQG
Chromium	mg/L	1.00 × 10 ⁻⁰³	n/a	n/a	n/a	1.00 × 10 ⁻⁰³	n/a	n/a	n/a	5.00 × 10 ⁻⁰²	n/a	n/a	1.00 × 10 ⁻⁰³	SEQG/CCME
Copper	mg/L	2.00 × 10 ⁻⁰³	(g)	n/a	n/a	2.00 × 10 ⁻⁰³	(b)	n/a	n/a	2.00 × 10 ⁺⁰⁰	n/a	n/a	2.00 × 10 ⁻⁰³	SEQG/CCME
Iron	mg/L	3.00 × 10 ⁻⁰¹	n/a	n/a	n/a	n/a	n/a	n/a	n/a	3.00 × 10 ⁻⁰¹	n/a	n/a	3.00 × 10 ⁻⁰¹	ССМЕ
Mercury	mg/L	2.60 × 10 ⁻⁰⁵	n/a	n/a	n/a	2.60 × 10 ⁻⁰⁵	n/a	n/a	n/a	1.00 × 10 ⁻⁰³	n/a	n/a	2.60 × 10 ⁻⁰⁵	SEQG/CCME
Magnesium	mg/L	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	None required	n/a	n/a	None required	n/a
Manganese	mg/L	2.60 × 10 ⁻⁰¹	(c)	n/a	n/a	n/a	n/a	n/a	n/a	1.20 × 10 ⁻⁰¹	n/a	n/a	1.20 × 10 ⁻⁰¹	Health Canada
Molybdenum	mg/L	7.30 × 10 ⁻⁰²	n/a	n/a	n/a	3.10 × 10 ⁺⁰¹	n/a	7.6 x 10 ⁰⁰	(r)	n/a	7.00 × 10 ⁻⁰²	(q)	7.60 × 10 ⁺⁰⁰ (Eco) 7.00 × 10 ⁻⁰² (HH)	BC MOE WHO
Sodium	mg/L	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	None required	n/a	n/a	None required	n/a
Nickel	mg/L	2.50 × 10 ⁻⁰²	(h)	n/a	n/a	2.50 × 10 ⁻⁰²	(b)	n/a	n/a	n/a	7.00 × 10 ⁻⁰²	(p)	2.50 × 10 ⁻⁰²	SEQG/CCME
Phosphorus	mg/L	n/a	n/a	n/a	n/a	n/a	n/a	2.00 × 10 ⁻⁰²	(m)	n/a	1.00 × 10 ⁻⁰²	(p)	2.00 × 10 ⁻⁰²	Ontario PWQO
Lead	mg/L	1.00 × 10 ⁻⁰³	(h)	n/a	n/a	1.00 × 10 ⁻⁰³	(b)	n/a	n/a	5.00 × 10 ⁻⁰³	n/a	n/a	1.00 × 10 ⁻⁰³	SEQG/CCME
Selenium	mg/L	1.00 × 10 ⁻⁰³	n/a	n/a	n/a	1.00 × 10 ⁻⁰³	n/a	n/a	n/a	5.00 × 10 ⁻⁰²	n/a	n/a	1.00 × 10 ⁻⁰³	SEQG/CCME
Strontium	mg/L	n/a	n/a	2.50 × 10 ⁺⁰⁰	(k)	n/a	n/a	n/a	n/a	7.00 × 10 ⁺⁰⁰	n/a	n/a	2.50 × 10 ⁺⁰⁰	FEQG
Uranium	mg/L	1.50 × 10 ⁻⁰²	n/a	n/a	n/a	1.50 × 10 ⁻⁰²	n/a	n/a	n/a	2.00 × 10 ⁻⁰²	n/a	n/a	1.50 × 10 ⁻⁰²	SEQG/CCME
Vanadium	mg/L	n/a	n/a	1.20 × 10 ⁻⁰¹	(n)	n/a	n/a	n/a	n/a	n/a	n/a	n/a	1.20 × 10 ⁻⁰¹	FEQG
Zinc	mg/L	7.00 × 10 ⁻⁰³	(i)	n/a	n/a	3.00 × 10 ⁻⁰²	n/a	n/a	n/a	5.00 × 10 ⁺⁰⁰	n/a	n/a	7.00 × 10 ⁻⁰³	ССМЕ

Constituent	Units	CCME Protection Life	CME Protection of Aquatic Fede				Saskatchewan Environmental Quality Guidelines (SEQG) ^(a)		Other		Drinking Water Guidelines			Source
		Long Term	Note	Long Term	Note	Long Term	Note	Long Term	Note	Health Canada ^(p)	Other Source	Note	Value	
Radionuclides														
Uranium-234	Bq/L	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	1.00 × 10 ⁺⁰⁰	(q)	n/a	n/a
Uranium-238	Bq/L	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	1.00 × 10 ⁺⁰¹	(q)	n/a	n/a
Thorium-230	Bq/L	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	1.00 × 10 ⁺⁰⁰	(q)	n/a	n/a
Radium-226	Bq/L	n/a	n/a	n/a	n/a	1.10 × 10 ⁻⁰¹	n/a	n/a	n/a	5.00 × 10 ⁻⁰¹	1.00 × 10 ⁺⁰⁰	(q)	n/a	n/a
Lead-210	Bq/L	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	2.00 × 10 ⁻⁰¹	1.00 × 10 ⁻⁰¹	(q)	n/a	n/a
Polonium-210	Bq/L	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	1.00 × 10 ⁻⁰¹	(q)	n/a	n/a

- (a) Saskatchewan Water Quality Objectives, SEQG on-line (https://envrbrportal.crm.saskatchewan.ca/seqg-search/), SEQG for the protection of aquatic life were selected, based on total concentrations, a temperature of 15°C and a pH of 7.0.
- (b) Hardness dependent WQOs are for very soft water (hardness <25 mg CaCO₃/L). Site-specific baseline hardness is 15 mg/L.
- (c) Scientific Criteria Document for the Development of the Canadian Water Quality Guidelines for the Protection of Aquatic Life Manganese, Appendix B Canadian Water Quality Guidelines Calculator (pH = 7.5, hardness = 15 mg/L). Guideline is based on dissolved manganese. (d) Total ammonia-N calculated from the total ammonia guideline for a temperature of 15°C and a pH of 7.0.
- (e) Based on a pH of >6.5.
- (f) Based on water hardness >0 to <17 mg/L..
- (g) Based on water hardness >0 to <82 mg/L.
- (h) Based on water hardness >0 to ≤60 mg/L.
- (i) Guideline is based on dissolved zinc.
- (j) Environment Canada 2017. Federal Environmental Quality Guidelines, Cobalt, May. Based on equation and the lowest hardness (52 mg/L) applicable to the guideline.
- (k) ECCC 2020. Federal Environmental Quality Guidelines Strontium. July.
- (I) BC MECCS 2021. British Columbia Approved Water Quality Guidelines: Aquatic Life, Wildlife & Agriculture. https://www2.gov.bc.ca/assets/gov/environment/air-land-water/water/water-quality-guidelines/approved-wqgs/wqg_summary_aquaticlife_wildlife_agri.pdf.
- (m) MOE 1994: Water management: policies, guidelines, provincial water quality objectives.
- (n) Environment Canada 2016. Federal Environmental Quality Guidelines, Vanadium. May.
- (o) Health Canada 2020. Guidelines for Canadian Drinking Water Quality Summary Table. September. https://www.canada.ca/content/dam/hc-sc/ewh-semt/alt formats/pdf/pubs/water-eau/sum guide-res_recom/summary-table-EN-2020-02-11.pdf.
- (p) BC MECCS 2020. Source Drinking Water Quality Guidelines, Guideline Summary Ministry of Environment & Climate Change Strategy Water Protection & Sustainability Branch.
- (g) WHO 2017. Guidelines for Drinking Water Quality. Fourth Edition Incorporating The First Addendum.
- (r) BC MOE (B.C. Ministry of Environment and Climate Change Strategy). 2021. Molybdenum Water Quality Guidelines for the Protection of Freshwater Aquatic Life, Livestock, Wildlife and Irrigation. Water Quality Guideline Series, WQG-07. Prov. B.C., Victoria B.C.

Bq/L = becquerels per litre; $CaCO_3 = calcium$ carbonate; CCME = Canadian Council of Ministers of the Environment; SEQG = Saskatchewan environmental quality guidelines; HH = human health; WHO = World Health Organization; PWQO = provincial water quality objective; FEQG = federal environmental quality guidelines; BC MECCS = British Columbia Ministry of Environment and Climate Change Strategy; WQO = water quality objective; > = greater than; < = less than or equal to; n/a = value not available or not applicable.



4.2.3.2 Constituents in Surface Water

The step-by-step and final results of the surface water screening are shown in Table 4-2. Based on the screening process, chloride, cobalt, and sulphate in treated effluent end-of-pipe and at the mixing zone boundary exceeded the screening values. Since the upper bound concentration of arsenic at the boundary of the mixing zone was only marginally below the screening value, it was also considered as a COPC as a precautionary measure. With the addition of groundwater solutes in the far-future projection to Patterson Lake North Arm – West Basin, copper was identified as an additional COPC.

No formal screening was conducted for radionuclides. However, since radiation dose to human and ecological receptors is of great public and regulatory interest, the radionuclides in the uranium-238 decay series (uranium-238, uranium-234, thorium-230, radium-226, lead-210, polonium-210) were all carried forward as COPCs for further assessment in the ERA.

Upper bound concentrations of uranium did not exceed the screening value. However, since uranium was evaluated as part of the radiation dose assessment and uranium-238 was estimated from the uranium concentration, uranium was also evaluated as a non-radiological COPC, because it is more chemically toxic than radiotoxic.

Radon-222 was not considered a COPC in surface water for the ERA. Radon is expected to volatilize rapidly to air. Health Canada (2020) considers that the health risk from ingesting radon-contaminated drinking water is negligible. Radon is expected to escape at the faucet or water outlet, leaving only negligible amounts in the water itself. This assumption is consistent with Clause 5.1.8 of CSA N288.1-20, *Guidelines for calculating derived release limits for radioactive material in airborne or liquid effluents for normal operation of nuclear facilities*, which indicates that noble gases, including radon-222, are not considered relevant for release to water because they do not enter environmental compartments other than air. Doses from noble gases released with treated effluent are expected to be negligible. Radon released directly to air is considered in Section 4.3, Atmospheric Sources.

Table 4-2: Surface Water Screening of Constituents of Potential Concern for the Environmental Risk Assessment

Table 4-2: Surface			Jpper-Bound Co								5.41	Is Upper	
Constituent	Units	West Runoff	Treated Effluent	Sewage	East Runoff	Selected Screening Value	Source	Is Concentration in Releases to Patterson Lake Greater than Selected Screening Value? (Y/N)	Edge of Mixing Zone - Upper Bound	Is Upper Bound Edge of Mixing Zone Concentration Greater than Selected Screening Value? (Y/N)	Patterson Lake North Arm – West Basin – Max Upper Bound (with Groundwate r Solutes)(b)	Bound Patterson Lake Concentratio n Greater than Selected Screening Value? (Y/N)	Final Selection as COPC? (Y/N)
General Parameter	rs											'	
Total suspended solids	mg/L	n/a	15	25	n/a	Background +5 mg/L	SEQG/CCME	Υ	1.4	N	n/a	n/a	N
Major lons													
Chloride	mg/L	$3.33 \times 10^{+00}$	3.31 × 10 ⁺⁰³	5.71 × 10 ⁻⁰¹	1.49 × 10 ⁺⁰⁰	1.20 × 10 ⁺⁰²	SEQG	Υ	1.66 × 10 ⁺⁰²	Υ	9.34 × 10 ⁺⁰¹	Υ	Υ
Sulphate	mg/L	$4.46 \times 10^{+00}$	3.42 × 10 ⁺⁰³	1.45 × 10 ⁺⁰⁰	1.95 × 10 ⁺⁰⁰	1.28 × 10 ⁺⁰²	BC MOE	Υ	$2.91 \times 10^{+02}$	Υ	2.00 × 10 ⁺⁰²	Υ	Υ
Nutrients													
Ammonia as nitrogen	mg/L	4.19 × 10 ⁻⁰¹	1.50 × 10 ⁺⁰¹	3.62 × 10 ⁺⁰²	1.98 × 10 ⁻⁰¹	5.74 × 10 ⁺⁰⁰	SEQG/CCME	Υ	4.98 × 10 ⁻⁰¹	N	3.59 × 10 ⁻⁰¹	N	N
Un-ionized ammonia as nitrogen ^(c)	mg/L	1.14 × 10 ⁻⁰³	4.09 × 10 ⁻⁰²	9.85 × 10 ⁻⁰¹	5.38 × 10 ⁻⁰⁴	1.56 × 10 ⁻⁰²	ССМЕ	Υ	1.36 × 10 ⁻⁰³	N	9.77 × 10 ⁻⁰⁴	N	N
Nitrate as nitrogen	mg/L	5.00 × 10 ⁻⁰¹	1.82 × 10 ⁺⁰¹	7.00 × 10 ⁺⁰²	5.00 × 10 ⁻⁰¹	3.00 × 10 ⁺⁰⁰	SEQG/CCME	Υ	5.38 × 10 ⁻⁰¹	N	3.48 × 10 ⁻⁰¹	N	N
Metals and Metall	oids												
Aluminum	mg/L	1.10×10^{-01}	1.23 × 10 ⁻⁰¹	3.83 × 10 ⁻⁰³	5.24 × 10 ⁻⁰²	1.00 × 10 ⁻⁰¹	SEQG/CCME	Υ	1.77 × 10 ⁻⁰²	N	6.09 × 10 ⁻⁰²	N	N
Arsenic	mg/L	1.25 × 10 ⁻⁰³	4.23 × 10 ⁻⁰²	9.46 × 10 ⁻⁰⁵	5.93 × 10 ⁻⁰⁴	5.00 × 10 ⁻⁰³	SEQG/CCME	Υ	3.98 × 10 ⁻⁰³	Y (marginally below)	2.73 × 10 ⁻⁰³	N	Υ
Calcium	mg/L	$4.00 \times 10^{+00}$	9.89 × 10 ⁺⁰²	3.56 × 10 ⁺⁰⁰	$4.00 \times 10^{+00}$	None required	n/a	N (non-toxic)	5.51 × 10 ⁺⁰¹	N (non-toxic)	3.43 × 10 ⁺⁰¹	N (non-toxic)	N (non-toxic)
Cadmium	mg/L	1.04×10^{-04}	1.11 × 10 ⁻⁰⁵	4.88 × 10 ⁻⁰⁶	4.94 × 10 ⁻⁰⁵	4.00× 10 ⁻⁰⁵	SEQG/CCME	Y (runoff only)	9.54 × 10 ⁻⁰⁶	N	1.22 × 10 ⁻⁰⁵	N	N
Cobalt	mg/L	2.04 × 10 ⁻⁰³	4.36 × 10 ⁻⁰³	5.00 × 10 ⁻⁰⁵	9.87 × 10 ⁻⁰⁴	4.65 × 10 ⁻⁰⁴	FEQG	Υ	4.47 × 10 ⁻⁰⁴	N	2.22 × 10 ⁻⁰³	Y	Υ
Chromium	mg/L	6.63 × 10 ⁻⁰⁴	3.84 × 10 ⁻⁰⁴	1.98 × 10 ⁻⁰⁴	5.00 × 10 ⁻⁰⁴	1.00 × 10 ⁻⁰³	SEQG/CCME	N	2.82 × 10 ⁻⁰⁴	N	2.99 × 10 ⁻⁰⁴	N	N
Copper	mg/L	5.62 × 10 ⁻⁰⁴	3.99 × 10 ⁻⁰³	1.24 × 10 ⁻⁰⁴	5.00 × 10 ⁻⁰⁴	2.00 × 10 ⁻⁰³	SEQG/CCME	Υ	4.85 × 10 ⁻⁰⁴	N	3.59 × 10 ⁻⁰³	Υ	Υ
Iron	mg/L	4.61 × 10 ⁻⁰¹	7.66 × 10 ⁻⁰²	8.59 × 10 ⁻⁰²	4.61 × 10 ⁻⁰¹	3.00 × 10 ⁻⁰¹	ССМЕ	Y (runoff only)	5.44 × 10 ⁻⁰²	N	8.84 × 10 ⁻⁰²	N	N
Mercury	mg/L	1.03 × 10 ⁻⁰⁵	3.71 × 10 ⁻⁰⁴	1.39 × 10 ⁻⁰⁶	4.49 × 10 ⁻⁰⁶	2.60 × 10 ⁻⁰⁵	SEQG/CCME	Υ	1.98 × 10 ⁻⁰⁵	N	1.30 × 10 ⁻⁰⁵	N	N
Magnesium	mg/L	$1.40 \times 10^{+00}$	1.16 × 10 ⁺⁰²	1.23 × 10 ⁺⁰⁰	1.40 × 10 ⁺⁰⁰	None required	n/a	N (non-toxic)	$7.02 \times 10^{+00}$	N (non-toxic)	$4.55 \times 10^{+00}$	N (non-toxic)	N (non-toxic)
Manganese	mg/L	1.99 × 10 ⁻⁰¹	1.50 × 10 ⁻⁰²	2.08 × 10 ⁻⁰²	8.95 × 10 ⁻⁰²	1.20 × 10 ⁻⁰¹	Health Canada	Y (runoff only)	2.55 × 10 ⁻⁰²	N	2.90 × 10 ⁻⁰²	N	N
Molybdenum	mg/L	2.33 × 10 ⁻⁰⁴	2.78 × 10 ⁻⁰¹	5.32 × 10 ⁻⁰⁵	1.00 × 10 ⁻⁰⁴	$7.60 \times 10^{+00}$ (Eco) 7.00×10^{-02} (HH)	BC MOE WHO	Y (HH – exceeds drinking water guideline only)	4.84 × 10 ⁻⁰³	N	2.96 × 10 ⁻⁰²	N	N
Sodium	mg/L	1.50 × 10 ⁺⁰⁰	1.04 × 10 ⁺⁰³	1.38 × 10 ⁺⁰⁰	1.50 × 10 ⁺⁰⁰	None required	n/a	N (non-toxic)	8.95 × 10 ⁺⁰¹	N (non-toxic)	6.19 × 10 ⁺⁰¹	N (non-toxic)	N (non-toxic)
Nickel	mg/L	2.77 × 10 ⁻⁰³	1.08 × 10 ⁻⁰²	7.26 × 10 ⁻⁰⁵	1.29 × 10 ⁻⁰³	2.50 × 10 ⁻⁰²	SEQG/CCME	N	1.08 × 10 ⁻⁰³	N	2.03 × 10 ⁻⁰³	N	N
Phosphorus	mg/L	5.00 × 10 ⁻⁰²	2.15 × 10 ⁻⁰¹	1.20 × 10 ⁺⁰⁰	5.00 × 10 ⁻⁰²	2.00 × 10 ⁻⁰²	Ontario PWQO	Υ	1.81 × 10 ⁻⁰²	N	1.40 × 10 ⁻⁰²	N	N (addressed in EIS under surface water)

		ι	Jpper-Bound Co	oncentrations ^(a)	ı					Is Upper Bound	Patterson	Is Upper	
Constituent	Units	West Runoff	Treated Effluent	Sewage	East Runoff	Selected Screening Value	Source	Is Concentration in Releases to Patterson Lake Greater than Selected Screening Value? (Y/N)	Edge of Mixing Zone - Upper Bound	Edge of Mixing Zone Concentration Greater than Selected Screening Value? (Y/N)	Upper Bound (with	Bound Patterson Lake Concentratio n Greater than Selected Screening Value? (Y/N)	Final Selection as COPC? (Y/N)
Lead	mg/L	1.33 × 10 ⁻⁰⁴	7.68 × 10 ⁻⁰⁵	4.72 × 10 ⁻⁰⁵	1.00 × 10 ⁻⁰⁴	1.00 × 10 ⁻⁰³	SEQG/CCME	N	7.68 × 10 ⁻⁰⁵	N	2.34 × 10 ⁻⁰⁴	N	N
Selenium	mg/L	1.33 × 10 ⁻⁰⁴	6.58 × 10 ⁻⁰⁴	4.72 × 10 ⁻⁰⁵	1.00 × 10 ⁻⁰⁴	1.00 × 10 ⁻⁰³	SEQG/CCME	N	1.08 × 10 ⁻⁰⁴	N	2.55 × 10 ⁻⁰⁴	N	N
Strontium	mg/L	3.24 × 10 ⁻⁰²	2.55 × 10 ⁺⁰¹	3.00 × 10 ⁻⁰²	3.21 × 10 ⁻⁰²	2.50 × 10 ⁺⁰⁰	FEQG	Υ	1.30 × 10 ⁺⁰⁰	N	7.40 × 10 ⁻⁰¹	N	N
Uranium	mg/L	1.33 × 10 ⁻⁰⁴	3.07 × 10 ⁻⁰¹	5.07 × 10 ⁻⁰⁵	1.00 × 10 ⁻⁰⁴	1.50 × 10 ⁻⁰²	SEQG/CCME	Υ	5.46 × 10 ⁻⁰³	N	3.45 × 10 ⁻⁰³	N	Y (evaluated for toxicity and radiotoxicity)
Vanadium	mg/L	5.62 × 10 ⁻⁰⁴	2.07 × 10 ⁻⁰³	1.50 × 10 ⁻⁰⁴	5.00 × 10 ⁻⁰⁴	1.20 × 10 ⁻⁰¹	FEQG	N	2.80 × 10 ⁻⁰⁴	N	2.21 × 10 ⁻⁰⁴	N	N
Zinc	mg/L	7.99 × 10 ⁻⁰³	2.05×10^{-03}	7.97 × 10 ⁻⁰⁴	3.49 × 10 ⁻⁰³	7.00 × 10 ⁻⁰³	ССМЕ	N	9.03 × 10 ⁻⁰⁴	N	2.32 × 10 ⁻⁰³	N	N

Grey shading and **bold text** indicate constituent considered a COPC for the ERA.

a) Upper-Bound concentrations for the treated effluent and runoff obtained from the site wide water balance model to the EIS (TSD XVIII Site-Wide Water Balance and Water Quality Modelling Report).

b) Maximum Upper-Bound concentrations from Patterson Lake North Arm – West Basin in the far-future projection with groundwater solutes added was obtained from the surface water quality model (EIS Appendix 10A, Surface Water Quality Modelling Report).

c) A pH of 7 and a temperature of 15°C were assumed to convert total ammonia to un-ionized ammonia in accordance with CCME (2002).

CCME = Canadian Council of Ministers of the Environment; SEQG = Saskatchewan environmental quality guidelines; HH = human health; PWQO = provincial water quality guidelines; BC MOE = British Columbia Ministry of Environment; WHO = World Health Organization; WQO = water quality objective; N = no; Y = yes; ERA = environmental risk assessment; COPC = constituent of potential concern; n/a = value not available or not applicable.

4.2.3.3 Constituents in Sediment

Predicted maximum concentrations of constituents of interest in sediment were compared against sediment quality guidelines for the protection of aquatic life and other relevant screening values. Sediment concentrations were predicted from surface water concentrations using IMPACT according to the equations outlined in the IMPACT model (Appendix A).

Sediment quality screening values were selected based on the following of sources:

- a) reference (REF) and no-effect (NE2) sediment quality values from Burnett-Seidel and Liber (Burnett-Seidel and Liber 2013);
- b) lowest effect levels (LELs) and severe effect levels (SELs) from Thompson et al. (Thompson et al. 2005); and
- c) Canadian interim sediment quality guidelines (ISQGs) and probable effect levels (PELs) from the CCME (CCME 1999a).

Burnett-Seidel and Liber (2013) was selected as the preferred source, as the reported NE2 and REF values are specifically applicable to Saskatchewan waterbodies. The REF values refer to locations upstream of mining or milling activities or located within separate but nearby drainages. Exceedances of REF values indicate that sediments downstream of predicted discharges contain elevated metal concentrations compared to natural background conditions. The NE2 values refer to exposed (lightly contaminated) areas with elevated concentrations but no significant effect on benthic invertebrate abundance, richness, and evenness. Concentrations below the NE2 values indicate that benthic invertebrate community metrics (abundance, richness, and evenness) downstream of discharges are not expected to differ significantly (less than 20% difference) from those observed at natural background conditions.

Two tiers of sediment quality guidelines are defined by Thompson et al. (2005): LELs and SELs. The CCME also provides two tiers of guidelines in sediments: ISQGs and PELs. If a predicted COPC concentration in sediment is less than the LEL or ISQG, adverse effects on benthic invertebrate communities are not anticipated for that constituent. Predicted concentrations in sediments that exceed the LEL or ISQG would not necessarily indicate that adverse effects are occurring but suggest that further investigation is warranted. These levels were, therefore, used for screening levels where there were no available REF levels.

An exceedance of a PEL or SEL is more likely to be associated with ecological effects. The SEL has been interpreted by some practitioners to be the specific COPC concentration in sediment that the majority of benthic organisms are not expected to tolerate (Persaud et al. 1993). The PEL is defined as the concentration of a COPC above which adverse effects are expected to occur frequently (more than approximately 50% of adverse effects occur above the PEL; CCME 1995).



The sediment screening (Table 4-3) focused on COPCs identified in the surface water screening as exceeding values, as well other constituents of interest from other uranium mining and milling operations. Based on comparison of maximum predicted sediment quality in Patterson Lake North Arm – West Basin in the Application Case and Upper Bound sensitivity scenario against the REF values from Burnett-Seidel and Liber (2013), arsenic and molybdenum would exceed the REF values; however, they do not exceed the NE2 values. Arsenic would exceed the REF value in Operations, while molybdenum would exceed the REF value in the far-future projection for the reasonable upper-bound sensitivity scenario.

There is no sediment screening value for cobalt; however, cobalt has already been identified as a COPC in surface water and was carried forward for further quantitative assessment in the ERA. Predicted concentrations of all other COPCs, including molybdenum, selenium, and nickel, do not exceed sediment quality guidelines and therefore were not considered sediment COPCs for further quantitative assessment in the ERA.

The maximum predicted upper bound concentrations of lead-210 and polonium-210 in sediment in Patterson Lake North Arm – West Basin exceeded the LEL values from Thompson et al. (2005); however, they did not exceed the SEL values. While exceeding the LEL does not necessarily indicate that adverse effects would occur, it indicates that further assessment is warranted. Radionuclides in the uranium-238 decay series (uranium-238, uranium-234, thorium-230, radium-226, lead-210, polonium-210) were considered sediment COPCs for further quantitative assessment in the ERA.

Table 4-3: Sediment Quality Screening for the Rook I Project

		Predicted Sedi	ment Concentrations (Patterson Lake North Arm	– West Basin) ^(a)		Sediment Quality Guidelines						ls	Final Selection
Constituent	Units	Maximum -	Maximum - Upper	Maximum - Application Case	Maximum - Upper Bound (Far-	Burnett-Se Libe		Thompson et al.(c)		ccı	CCME ^(d) Sedin Scree		eening Concentration Greater than Selected	as COPC for Aquatic Environment?
		Application Case	Bound	(Far-Future Projection)	Future)	REF	NE2	LEL	SEL	ISQG	PEL	Value	Screening Value? (Y/N)	(Y/N)
Metals and Metalloi	ds										'			
Arsenic	mg/kg dw	30.50	31.00	10.60	10.93	21	522	9.8	346	5.9	17	21	Y	Υ
Cadmium	mg/kg dw	0.27	0.27	0.29	0.30	n/d	n/d	n/d	n/d	0.6	3.5	0.6	N	N
Cobalt	mg/kg dw	1.88	1.90	3.56	4.68	n/d	n/d	n/d	n/d	n/d	n/d	n/d	n/a	Y (based on surface water)
Copper	mg/kg dw	2.93	2.94	6.51	8.52	9.1	11.3	22	269	35.7	197	9.1	N	Y (based on surface water)
Lead	mg/kg dw	10.16	10.16	10.33	11.27	16.3	19.7	37	412	35	91.3	16.3	N	N
Molybdenum	mg/kg dw	1.74	3.97	14.53	53.94	23	245	14	1,239	n/d	n/d	23	Υ	Υ
Nickel	mg/kg dw	5.67	5.74	7.39	9.79	21	326	23	484	n/d	n/d	21	N	N
Selenium	mg/kg dw	0.56	0.57	0.95	1.37	3.6	30	1.9	16	n/d	n/d	3.6	N	N
Uranium	mg/kg dw	6.33	14.20	19.30	19.32	97	2,296	104	5,874	n/d	n/d	97	N	Y (evaluated for toxicity and radiotoxicity)
Vanadium	mg/kg dw	9.48	9.51	9.17	9.18	35	32	35	160	n/d	n/d	35	N	N
Zinc	mg/kg dw	11.57	11.63	15.71	18.03	n/d	n/d	n/d	n/d	123	315	123	N	N
Radionuclides														
Uranium-234	Bq/kg dw	62	47	39	67	n/d	n/d	n/d	n/d	n/d	n/d	n/d	n/a	Y (radionuclide)
Uranium-238	Bq/kg dw	62	47	39	66	n/d	n/d	n/d	n/d	n/d	n/d	n/d	n/a	Y (radionuclide)
Thorium-230	Bq/kg dw	116	76	76	22	n/d	n/d	n/d	n/d	n/d	n/d	n/d	n/a	Y (radionuclide)
Radium-226	Bq/kg dw	104	106	85	83	n/d	n/d	600	14,400	n/d	n/d	600	N	Y (radionuclide)
Lead-210	Bq/kg dw	492	984	402	376	n/d	n/d	900	20,800	n/d	n/d	900	Υ	Y (radionuclide)
Polonium-210	Bq/kg dw	500	1002	409	382	n/d	n/d	800	12,100	n/d	n/d	800	Υ	Y (radionuclide)

Note: **Bold** and Grey shading indicates sediment concentration exceeds the REF or LEL value.

⁽a) Sediment concentrations predicted based on release of aqueous source-terms to Patterson Lake North Arm – West Basin and interaction with sediment. Modelling performed in IMPACT according to the equations outlined in Appendix A (b) Burnett-Seidel and Liber (2013) - Sediment quality values derived for application at Saskatchewan uranium operations; reference (REF) values based on reference sites unaffected by mining and milling (representing background), and no-effect level (NE2) values based on sites with no significant (difference in benthic invertebrate community effects criteria of abundance, richness and evenness between reference and exposure locations.

⁽c) Thompson et al. (2005) - Sediment quality guidelines derived for application to uranium ore bearing regions of northern Saskatchewan and Ontario; lowest effect levels (LELs) and severe effect levels (SELs) from the "weighted method".

⁽d) CCME – Canadian Sediment Quality Guidelines for the Protection of Freshwater Aquatic Life (CCME 1999a; updated September 2007; accessed July 2021: http://ceqg-rcqe.ccme.ca/).

REF = reference; NE2 = no effects; LEL = lowest effect level; CCME = Canadian Council of Ministers of the Environment; ISQG = Interim Sediment Quality Guideline; PEL = probable effect level; n/d = no guideline or data available; n/a = not applicable; Y = yes; N = no; dw = dry weight; Bq/kg = becquerels per kilogram.

4.2.4 Evaluation of Selenium in Fish Tissue

In 2016, the United States Environmental Protection Agency (USEPA) developed an aquatic criterion for selenium based on the scientific advancement that selenium toxicity in the aquatic environment is primarily from bioaccumulation of selenium in the aquatic food chain and not only from exposure to selenium in water. As such, although predicted selenium concentrations in water are below the selected water quality guideline, it is appropriate to compare predicted fish tissue concentrations in Patterson Lake North Arm – West Basin against the selenium fish tissue criterion. Fresh water fish are considered more sensitive to selenium compared to other aquatic organisms.

The recommended criterion for selenium in fish tissue is 11.3 mg/kg dry weight (dw) muscle or 8.5 mg/kg dw whole body (USEPA 2021). The whole body and muscle tissue concentrations are based on effect concentration (EC₁₀) values for selenium in egg/ovary, which caused reproductive effects in fresh water fish. The EC₁₀ is the effect concentration at which 10% effect based on the endpoint is observed. The USEPA used species-specific conversion factors to convert from egg/ovary concentrations to whole body and muscle tissue concentrations.

Environment and Climate Change Canada (ECCC; 2022) published a selenium in fish tissue criterion of 6.7 mg/kg dw whole body, which represents the 5th percentile hazard concentration of the species sensitivity distribution of whole body selenium concentrations. Environment and Climate Change Canada indicates that this value is similar to that derived by the USEPA (2021). The USEPA criterion has been adopted for the Project over the ECCC criterion as the ECCC criterion is for whole body fish tissue, whereas the USEPA criterion is for fish tissue muscle as well. The fish assessed in the ERA are large-bodied fish (i.e., northern pike and lake whitefish); therefore, a fish tissue muscle guideline is preferred over the whole body guideline.

The recommended criterion for selenium in fish tissue of 11.3 mg/kg dw muscle was used for large-bodied fish included in this assessment (i.e., northern pike and lake whitefish). To compare against predicted selenium fish tissue concentrations that are in fresh weight, the criterion needed to be adjusted from a dry weight to a fresh weight basis based on the moisture content of the fish. The dry weight (dw) to fresh weight (fw) ratio of 0.25 to 1.0 from CSA N288.1-20 was used to convert the selenium criterion to a fresh weight basis.

Fish Tissue Criterion (mg/kg fw) = Fish Tissue Criterion (mg/kg dw) * 0.25 (kg dw/kg fw)

Therefore, the recommended criterion for selenium in fish tissue on a fresh weight basis is 2.83 mg/kg fw muscle.

Predicted fish tissue concentrations of selenium in northern pike and lake whitefish over the Project phases for both the Application Case and Upper Bound sensitivity scenario are shown in Figure 4-4, along with a comparison against the selenium criterion. The selenium bioaccumulation factor (BAF) for northern pike is 9.49E+02 L/kg fw and the BAF for lake whitefish is 5.94E+03 L/kg fw based on publicly available regional data from other uranium mine sites in northern Saskatchewan. The maximum predicted upper bound selenium concentration is 0.19 mg/kg fw in lake whitefish in Patterson Lake North Arm – West Basin compared to the criterion of 2.83 mg/kg

fw. Selenium concentrations in fish at the maximum location in Patterson Lake North Arm – West Basin are all well below the recommended criterion.

Based on the above conclusions, selenium was not considered for further quantitative assessment in the ERA.




Figure 4-4: Comparison of Predicted Selenium Fish Tissue Concentrations against Criterion

Note: The predicted selenium concentrations in fish tissue for lake whitefish and northern pike are indistinguishable from each other on the figure.

USEPA = United States Environmental Protection Agency; fw = fresh weight.

4.2.5 Water Quality Constituents for Further Evaluation in the Environmental Risk Assessment

In summary, following the screening process for surface water quality (in combination with the sediment quality screening and selenium in fish tissue discussion), the following COPCs in the aquatic environment were carried forward for further quantitative assessment in the ERA for both human and ecological health:

- major ions: chloride and sulphate;
- metals and metalloids: arsenic, cobalt, copper, molybdenum, uranium; and
- radionuclides: uranium-238, uranium-234, thorium-230, radium-226, lead-210, polonium-210.



4.3 Atmospheric Sources

4.3.1 Project-Related Atmospheric Releases

The Project has the potential to change air quality through the emission of gases and particulates and deposition of particulates generated by Project activities. For emission to the atmosphere, the ERA focused on Construction and Operations when effects on air quality are expected to be the greatest due to the intensity and number of Project-related activities. Additionally, atmospheric emissions during Closure are expected to be lower than during Construction and Operations; and Closure modelling was not explicitly conducted. It was conservatively assumed that Closure air quality would be similar to Construction air quality.

The Project-related atmospheric releases considered in the ERA were consistent with the air emissions inventory detailed in the Air Dispersion Modelling Report for Project Construction and Operations (EIS Section 7.2.5, Residual Effects Analysis; EIS Appendix 7A, Air Dispersion Modelling Report). The major air emission sources considered for the ERA included:

- fossil fuel combustion emissions from mobile equipment and stationary equipment (e.g., power plant, heaters);
- fugitive dust emissions from drilling and blasting, material handling, crushing, vehicle generated road dust, and wind erosion from ore and mine rock storage piles;
- air emissions released from the milling processes (e.g., calciner, acid plant, lime silo baghouses); and
- solid waste incinerators.

Project-related atmospheric releases would include criteria air contaminants (CACs; nitrogen oxides [assessed as nitrogen dioxide, sulphur dioxide, sulphuric acid, carbon monoxide, total suspended particulates [TSP], and particulate matter [PM₁₀ and PM_{2.5}]), fugitive dust, dioxins and furans, metals and radionuclides, and radon.

Criteria air contaminants have either federal or provincial ambient air quality criteria or both. Nitrogen oxides, sulphur dioxide, carbon monoxide, and particulates (TSP, PM₁₀, and PM_{2.5}) would be CACs directly emitted by the Project from stationary and mobile sources. Sulphuric acid emissions would be associated with the acid plant.

Dust commonly refers to large particulate matter, such as fugitive dust, that tends to settle out of suspension in the air by gravity within a short distance from the source. It would be associated with such activities as aggregate generation, road dust, materials hauling, and construction activities. It would be measured in terms of TSP deposition or dustfall.

Dioxins and furans represent a family of toxic compounds formed during the combustion process that share a similar chemical structure and are persistent in the environment. They would be associated with waste incineration.



Metals and radionuclides would be emitted as a portion of dust and combustion emissions. Dust emissions from waste rock, special waste, ore, and aggregate were assumed to contain metals. Dust emissions from waste rock, special waste, and ore were assumed to contain radionuclides. Radionuclides associated with dust from aggregate materials were considered to be negligible. The radionuclides included in the ERA were uranium-238 series radionuclides and were assumed to be in secular equilibrium.

Radon emissions from a number of sources were included in the air quality assessment: mine vent; ore storage pile; special waste rock pile; ore handling, grinding circuit / process plant, and slurry vessels; paste tailings preparation; settling pond; waste rock storage piles and material handling; and low-level radioactive waste incinerator. Potential radon emissions from the calciner and uranium concentrate handling stacks were considered to be negligible. Radon emissions were estimated based on U_3O_8 grade for waste rock (0.03%), special waste (0.05%) and ore (4%).

4.3.2 Atmospheric Sources

In this context, sources refer to total concentrations of constituents in air at specific human and ecological receptor locations. Constituent predictions for ambient air quality were obtained from the Air Dispersion Modelling Report (EIS Appendix 7A). Air quality modelling followed the Saskatchewan Air Quality Modelling Guideline (ENV 2012) using the atmospheric dispersion model AERMOD (EIS Appendix 7A). The emissions inputs were calculated to represent the maximum reasonably foreseeable emissions from the Project after applying mitigation actions and policies to reduce emissions.

Air quality modelling included predictions for CACs, dust, dioxins and furans, metals and radionuclides, and radon at the fence line and at specific human health and ecological receptor locations as described in Section 5.2.1, Exposure Locations, Duration, and Frequency, for the human health risk assessment, and in Section 6.2.1, Exposure Locations, for the ecological risk assessment. For the purposes of the ERA, the fence line was assumed to be the nearest location outside the property boundary where there would be public access (ENV 2012). The predicted air concentrations were based on a single "maximum year" scenario modelled for each of Construction and Operations.

Predicted total maximum concentrations in ambient air from the Air Dispersion Modelling Report (EIS Appendix 7A) were used in Section 4.3.3, Screening for Atmospheric Constituents of Potential Concern, for the relevant averaging period for Construction and Operations. For metals and radionuclides associated with dust emissions, total concentrations include only Project-related sources. Background concentrations of metals in dust were assumed to be negligible and were set to zero in the air quality model because there are no other industrial sources within 100 km of the LSA. Regional background concentrations for CACs are summarized in Table 4-4.



Table 4-4: Background Concentrations of Air Quality Constituents Used in The Air Quality Model

Constituent (unit)	Averaging Period	Background Concentration
	1 hour	11.3
Nitrogen dioxide (µg/m³)	24 hours	9.4
	Annual	3.8
	1 hour	0
Sulphur dioxide (μg/m³)	24 hours	0
	Annual	0
Contrar = 200 = 201 = (200 / 200)	1 hour	572
Carbon monoxide (μg/m³)	8 hours	572
PM ₁₀ (μg/m ³)	24 hours	23.1
DAA ((3)	24 hours	6.5
PM _{2.5} (μg/m³)	Annual	3.1
TCD (24 hours	14.4
TSP (μg/m³)	Annual	6.2
Radon (Bq/m³)	Average	2.94

Note: Background concentrations were obtained from EIS Appendix 7A, Air Dispersion Modelling Report.

 Bq/m^3 = becquerels per cubic metre; TSP = total suspended particulates; PM_{10} = particulate matter with a diameter of 10 microns or less; $PM_{2.5}$ = particulate matter with a diameter of 2.5 microns or less.

4.3.3 Screening for Atmospheric Constituents of Potential Concern

Constituents of potential concern for air, as defined by Health Canada (Health Canada 2016a), are chemicals whose concentration(s) may become elevated in ambient air as a result of project-related activities, and which have the potential for adverse human or ecological health effects based on documented scientific evidence or suspected causal relationships. The purpose of this subsection is to identify those Project-related constituents in air that may be of concern for human and/or ecological health and require further assessment.

The screening of air quality constituents was based on maximum predicted concentrations of CACs, dioxins and furans, radon, and 28 metals and metalloids, and maximum dust deposition, at air quality model locations that correspond with receptor locations (Table 4-5), as described in Section 5.2.1, Exposure Locations, Duration, and Frequency, for the human health risk assessment, and Section 6.2.1, Exposure Locations, for the ecological risk assessment.

Table 4-5: Concordance between Air Quality Model and Receptor Locations

Air Ovelite Madel Leasting	Umman and Factorial Basedon Laudian	Air Quality Mode	el Coordinates
Air Quality Model Location	Human and Ecological Receptor Location	X (m)	Y (m)
HHRA1	Hodge Lake Reference	593,768	6,407,146
HHRA2	Broach Lake	600,359	6,398,266
HHRA3	Camp ^(a)	603,778	6,393,226
HHRA4	Patterson Lake Human Health Receptors	598,658	6,387,580
HHRA5	Patterson Lake Ecological Receptors	602,320	6,392,289
HHRA6	Forrest Lake	605,446	6,388,744
HHRA7	Forrest Lake North	605,452	6,390,021
HHRA8	Beet Lake	608,931	6,389,997
HHRA9	Naomi Lake	614,179	6,390,462
HHRA10	Clearwater River	626,340	6,380,517
HHRA11	Lloyd Lake	616,793	6,361,563

(a) HHRA3 is located at the camp; other air quality model and human and ecological receptor locations are located outside of the Project footprint.

The modelled location HHRA3 is located at the camp within the Project footprint. This location was retained for the screening as a conservative measure to ensure potential COPCs in air are captured in the ERA, since it is the closest location to the source. This location would therefore be expected to experience higher air concentrations than locations farther away.

- 1. Human and ecological receptors at receptor locations were assumed to be in contact with air emissions for prolonged periods of time, at intervals that may be long-term (i.e., annual average) or repeated and short-term (i.e., 24 hours or less) over a lifetime. For this reason, long-term and short-term screening values at the receptor locations were used for the screening of constituents in air at receptor locations. Screening of constituents in air for the receptor locations followed the following protocol using maximum predicted concentrations for all receptor locations for the relevant time period:
- 2. If the model results from the Air Dispersion Modelling Report (EIS Appendix 7A) for a constituent were below all of its relevant air quality screening values, at all receptor locations, the constituent was assumed to be below levels associated with potential human health and ecological risks and was not considered further in the ERA for direct atmospheric exposures.
- 3. If the model result for an air quality constituent was greater than any one of its relevant air quality screening values at any receptor location, the constituent was determined to be a potential COPC in air and was evaluated further in Section 4.3.3.3, Discussion of Air Quality Constituents that Exceed a Screening Value.
- 4. Potential human health and ecological risks for the air quality constituents that were determined to be COPCs in Section 4.3.4, Air Quality Constituents for Further Evaluation

in the Environmental Risk Assessment, were characterized in Section 5.4 and Section 6.4, Risk Characterization, respectively.

The ERA also considered that humans (such as the subsistence harvester) and mobile ecological receptors (e.g., birds and mammals) would be present near the fence line occasionally and for short periods (i.e., less than 24 hours) of time. Maximum predicted concentrations from the Air Dispersion Modelling Report (EIS Appendix 7A) of Project-related constituents in air at the fence line were therefore screened using short-term screening values (i.e., 1 hour, 8 hours). This screening and its results are in Section 4.3.3.2, Screening of Air Quality Constituents. If the model results were below all relevant short-term air quality screening values, the constituent was assumed to be below levels associated with potential human health and ecological risks and was not considered further in the ERA.

4.3.3.1 Screening Value Selection

Ambient air quality criteria are available for different exposure averaging periods (e.g., 1 hour, 24 hour, annual). Maximum predicted concentrations from the Air Dispersion Modelling Report (EIS Appendix 7A) of constituents in air were screened against ambient air quality criteria for the same averaging period. Ambient air quality criteria for the relevant averaging periods were selected based on the following order for selection:

- Saskatchewan Ambient Air Quality Standards (SAAQS) are maximum concentrations in ambient air from all sources as stipulated in The Clean Air Regulations (Government of Saskatchewan 2015).
- Alberta Ambient Air Quality Objectives (Alberta 2021) (AAAQO) are based on an evaluation of scientific, social, technical, and economic factors.
- Ontario Ambient Air Quality Criteria (OAAQC) are concentrations of a contaminant in air that are protective against adverse effects on health and/or the environment (MECP 2020).
- Texas effects screening levels (ESLs) are air concentrations at or below which adverse health effect in the general public, including sensitive subgroups such as children, the elderly, pregnant women, and people with pre-existing health conditions, are not likely to occur (TCEQ 2016).

Canadian Ambient Air Quality Standards (CAAQS) established under the national Air Quality Management System (CCME 2021b) were used for information purposes, but not selected as screening criteria. As indicated in the Air Quality assessment (EIS Section 7.2, Air Quality), the CAAQS achievement is determined by provinces and territories using ambient concentrations measured in the air zones for a three-year period, not by comparison of modelled predictions at or beyond a facility boundary.

Screening values for radionuclide concentrations in ambient air were not available. All relevant radionuclides were assessed in the ERA in terms of their contribution to the total radiological dose to human and ecological receptors.



A screening value for radon gas was not available from the sources identified. Health Canada's annual average corrective action trigger value of 200 becquerels per cubic metre (Bq/m³) for indoor air was used for the screening (Health Canada 2009).

The selected ambient air quality screening values for different averaging periods, their source, and their rationale in terms of potential effects are summarized in Table 4-6. Where multiple sources recommended the same criterion value, each of the relevant sources is identified. The rationale provided in Table 4-6 for each of the selected screening values describes the sensitive effect that is the basis for the value cited by the relevant source.



Table 4-6: Screening Values for the Selection of Air Quality Contaminants of Potential Concern for the Environmental Risk Assessment

Constituent	Averaging Period	Selected Screening Value	Source	Rationale
<u>CACs</u>				
	1 hour	300	SAAQS/AAAQO	Respiratory effects
Nitrogen dioxide (NO ₂)	24 hours	200	SAAQS/OAAQC	Human health
(1402)	Annual	45	SAAQS/AAAQO	Vegetation
	1 hour	450	SAAQS/AAAQO	Pulmonary effects
Sulphur dioxide (SO ₂)	24 hours	125	SAAQS/AAAQO	Human health
(302)	Annual	20	SAAQS/AAAQO	Ecosystem health
	1 hour	15,000	SAAQS/AAAQO	Oxygen carrying capacity of blood
Carbon monoxide	8 hours	6,000	SAAQS/AAAQO	Oxygen carrying capacity of blood
(CO)	Annual	n/v	n/a	n/a
	1 hour	10	AAAQO	Not stated
Sulphuric acid (H ₂ SO ₄)	24 hours	5	OAAQC	Human health
(112304)	Annual	n/v	n/a	n/a
Total suspended	24 hours	100	SAAQS/AAAQO	Human health. Pulmonary effects
particulates (TSP)	Annual	60	SAAQS/OAAQC	Visibility
Particulate matter	24 hours	50	SAAQS/OAAQC	Human health
(PM ₁₀)	Annual	n/v	n/a	n/a
Particulate matter	24 hours	27	OAAQC/CAAQS	Human health
(PM _{2.5})	Annual	8.8	OAAQC/CAAQS	Human health
<u>Dust</u>				
TSP deposition	Annual	4.6	OAAQC	Dustfall criterion. Aesthetics (g/m²/yr)
rar deposition	Monthly	5.3	AAAQO	Aesthetics (g/m²/30 days)



Constituent	Averaging Period	Selected Screening Value	Source	Rationale
Dioxins and Furans				
Dioxins and furans	24 hour	0.1	OAAQC	Human health (pg TEQ/m³)
DIOXINS and Turans	Annual	n/v	n/a	n/a
<u>Radionuclides</u>				
U-238 series	24 hour	n/v	n/a	Addressed in terms of radiation dose in the ERA
radionuclides	Annual	n/v	n/a	Addressed in terms of radiation dose in the ERA
<u>Radon</u>				
Radon	Annual	200	Government of Canada Radon Guideline (Health Canada 2009)	Addressed in terms of radiation dose in the ERA
<u>Metals</u>				
Silver (Ac)	24 hour	1	OAAQC	Human health
Silver (Ag)	Annual	n/v	n/a	n/a
Arcania (As)	24 hour	0.3	OAAQC	Human health. Applies to arsenic and arsenic compounds
Arsenic (As)	Annual	0.01	AAAQO	Human health. Carcinogenic effects
Davium (Da)	24 hour	10	OAAQC	Human health. Applies to total barium water-soluble fraction
Barium (Ba)	Annual	n/v	n/a	n/a
Damillione (Da)	24 hour	0.01	OAAQC	Human health. Applies to beryllium and beryllium compounds
Beryllium (Be)	Annual	n/v	n/a	n/a
Cadmium (Cd)	24 hour	0.025	OAAQC	Human health. Applies to cadmium and cadmium compounds. Converted from the annual AAQC to allow assessment of 24-hour air quality data
	Annual	0.005	OAAQC	Human health. Applies to cadmium and cadmium compounds



Constituent	Averaging Period	Selected Screening Value	Source	Rationale
Coholt (Co)	24 hour	0.1	OAAQC	Human health
Cobalt (Co)	Annual	n/v	n/a	n/a
Chromium (Cr)	24 hour	0.5	OAAQC	Human health. Applies to either chromium metallic, divalent, and trivalent, or to the percentage of chromium metallic, divalent, and trivalent relative to total chromium
	Annual	n/v	n/a	n/a
(C.)	24 hour	50	OAAQC	Human health
Copper (Cu)	Annual	n/v	n/a	n/a
M (11-)	24 hour	2	OAAQC	Human health
Mercury (Hg)	Annual	n/v	n/a	n/a
Molybdenum (Mo)	24 hour	120	OAAQC	Particulate - visibility; molybdenum is more likely emitted as TSP, and therefore the AAQC for TSP is applied
,	Annual	n/v	n/a	n/a
	24 hour	0.2	OAAQC	In TSP. Human health. Applies to nickel and nickel compounds. Converted from the annual criterion to allow assessment of the 24-hour data (TSP). Intended to protect from development of chronic effects
NIC L. LANCS	Annual	0.04	OAAQC	In TSP. Human health. Applies to nickel and nickel compounds
Nickel (Ni)	24 hour	0.1	OAAQC	In PM ₁₀ . Human health. Applies to nickel and nickel compounds. Converted from the annual criterion to allow assessment of the 24-hour data (TSP). Intended to protect from development of chronic effects
	Annual	0.02	OAAQC	In PM ₁₀ . Human health. Applies to nickel and nickel compounds
Lead (Pb)	24 hour	0.5	OAAQC	Human health. Applies to lead and lead compounds. Converted from the 30-day AAQC to allow assessment of 24-hour air quality data



Constituent	Averaging Period	Selected Screening Value	Source	Rationale
	Monthly	0.2	OAAQC	Human health. Applies to lead and lead compounds. As arithmetic mean of a 30-day period
	Annual	n/v	n/a	n/a
A ti (Cl-)	24 hour	25	OAAQC	Human health
Antimony (Sb)	Annual	n/v	n/a	n/a
(C-1i (C-)	24 hour	10	OAAQC	Human health
Selenium (Se)	Annual	n/v	n/a	n/a
T: (CL)	24 hour	10	OAAQC	Human health
Tin (Sb)	Annual	n/v	n/a	n/a
T (T.)	24 hour	n/v	n/a	n/a
Thorium (Th)	Annual	n/v	n/a	n/a
	24 hour	0.3	OAAQC	In TSP. Human health. Applies to uranium and uranium compounds. Converted from the annual AAQC to allow assessment of 24-hour air quality data
	Annual	0.06	OAAQC	In TSP. Human health. Applies to uranium and uranium compounds
Uranium (U)	24 hour	0.15	OAAQC	In PM ₁₀ . Human health. Applies to uranium and uranium compounds. Converted from the annual AAQC to allow assessment of 24-hour air quality data
	Annual	0.03	OAAQC	In PM ₁₀ . Human health. Applies to uranium and uranium compounds
	24 hour	2	OAAQC	Human health
Vanadium (V)	Annual	n/v	n/a	n/a
7: (7:-)	24 hour	120	OAAQC	Particulates
Zinc (Zn)	Annual	n/v	n/a	n/a



Constituent	Averaging Period	Selected Screening Value	Source	Rationale
	24 hour	n/v	n/a	n/a
Cesium (Cs)	Annual	2	Texas Interim ESL	Human health
Diagonath (Di)	24 hour	n/v	n/a	n/a
Bismuth (Bi)	Annual	5	Texas Interim ESL	Human health
(C-1-i (C-)	24 hour	n/v	Texas ESL	Particulates
Calcium (Ca)	Annual	n/v	Texas ESL	Particulates
luon (Fo)	24 hour	4	OAAQC	Human health. Applies to metallic iron
Iron (Fe)	Annual	n/v	n/a	n/a
Magnesium (Mg)	24 hour	72	OAAQC	Human health
	Annual	n/v	n/a	n/a
Manganese (Mn)	24 hour	0.1	OAAQC	Human health. OAAQC (2020): 0.4 (TSP), 0.2 (PM ₁₀) and 0.1 (PM _{2.5}). For manganese and manganese compounds in TSP
	Annual	0.2	AAAQO	Adopted from Texas (long-term ESL) and California (chronic reference exposure level for nervous system effects)
Sodium (Na)	24 hour	n/v	Texas ESL	Particulate
Sodium (Na)	Annual	n/v	Texas ESL	Particulate

Note: Units are µg/m³ unless otherwise specified.

Texas ESL = Texas Commission on Environmental Quality (TCEQ 2016); AAAQO = Alberta Ambient Air Quality Objectives (Alberta 2021); CAAQS = Canadian Ambient Air Quality Standards, air quality objectives under the *Canadian Environmental Protection Act, 1999* (CCME 2021b); OAAQC = Ontario Ambient Air Quality Criteria (MECP 2020); SAAQS = Saskatchewan Ambient Air Quality Standards (Government of Saskatchewan 2015); AAQC = ambient air quality criteria; CAC = criteria air contaminant; ESL = effects screening level; Bq/m³ = becquerels per cubic metre; TSP = total suspended particulates; ERA = environmental risk assessment; PM_{10} = particulate matter with a diameter of 10 microns or less; $PM_{2.5}$ = particulate matter with a diameter of 2.5 microns or less; $PM_{2.5}$ = porticulate matter with a diameter of 2.5 microns or less; $PM_{2.5}$ = particulate matter with a diameter of 2.5 microns or less; $PM_{2.5}$ = particulate matter with a diameter of 2.5 microns or less; $PM_{2.5}$ = particulate matter with a diameter of 2.5 microns or less; $PM_{2.5}$ = particulate matter with a diameter of 2.5 microns or less; $PM_{2.5}$ = particulate matter with a diameter of 2.5 microns or less; $PM_{2.5}$ = particulate matter with a diameter of 2.5 microns or less; $PM_{2.5}$ = particulate matter with a diameter of 2.5 microns or less; $PM_{2.5}$ = particulate matter with a diameter of 2.5 microns or less; $PM_{2.5}$ = particulate matter with a diameter of 2.5 microns or less; $PM_{2.5}$ = particulate matter with a diameter of 2.5 microns or less; $PM_{2.5}$ = particulate matter with a diameter of 2.5 microns or less; $PM_{2.5}$ = particulate matter with a diameter of 2.5 microns or less; $PM_{2.5}$ = particulate matter with a diameter of 2.5 microns or less; $PM_{2.5}$ = particulate matter with a diameter of 2.5 microns or less; $PM_{2.5}$ = particulate matter with a diameter of 2.5 microns or less; $PM_{2.5}$ = particulate matter with a diameter of 2.5 microns or less; $PM_{2.5}$ = particulate matter with a diameter of 2.5



4.3.3.2 Screening of Air Quality Constituents

The screening of air quality constituents involved the following two types of screenings:

- Comparing the predicted maximum air concentrations from the air quality model at all human and ecological receptor locations for all (short and long-term) averaging periods against the corresponding air quality criteria (Table 4-7).
- Comparing the predicted maximum air concentrations from the air quality model at the fence line for short-term averaging periods against short-term air quality criteria (Table 4-8).

The screening of air quality constituents at the human and ecological receptor locations for short- and long-term averaging periods at receptor locations is provided in Table 4-7. Both human and ecological receptors were assumed to be present for extended periods of time at these locations and therefore susceptible to both short- and long-term exposures to airborne constituents. Constituents that had only a short-term or a long-term screening value were not considered further if the maximum predicted concentration was less than the screening value, as explained in Table 4-7.

The screening of air quality constituents at the fence line for short-term averaging periods only is provided in Table 4-8. Hypothetical visitors and mobile ecological receptors were assumed to be present for short periods of time at these locations and therefore susceptible to only short-term exposures to airborne constituents. The screening was performed using the maximum predicted concentrations for all fence line locations from the air quality model (EIS Appendix 7A) for the relevant averaging period. Note that these maximum fence line concentrations are in some cases higher than those presented in EIS Section 7.2, since the air quality assessment in the EIS followed Saskatchewan air quality dispersion modelling guidance which allows for discounting the maximum modelled concentrations.

Constituents that were predicted to exceed any of their screening values based on maximum predicted values for the relevant averaging period are further discussed in Section 4.3.3.3, to determine whether or not they should be retained as COPCs and further evaluated in terms of human health and/or ecological risk. Air quality constituents with maximum concentrations that exceeded either their short- or long-term screening value at receptor locations were nitrogen dioxide, particulate matter (TSP, PM₁₀, PM_{2.5}), and uranium. Dustfall exceeded its annual criterion during Construction. Air quality parameters with maximum short-term concentrations that exceed screening values at the fence line are nitrogen dioxide, particulate matter (TSP, PM₁₀, and PM_{2.5}), and uranium.

While no identified screening level is exceeded for radionuclides (including uranium-238 series) and radon gas, potential human health and ecological risks are discussed in Section 5.4 and Section 6.4, respectively.



Table 4-7: Air Quality Screening for Short-term and Long-term Exposures to Constituents in Air at Human and Ecological Receptor Locations

		Maximum Concentration at Receptor Locations				Is Concentration Greater than		
Constituent	Construction	Operations	Screening Value	Averaging Period	Source	Selected Screening Value (Yes/No)	Is the Constituent Retained for Further Consideration as a COPC	
<u>CACs</u>								
	14.7	8.55	45	Annual	SAAQS/AAAQO			
Nitrogen dioxide (NO₂)	120	103	200	24 hours	SAAQS/OAAQC	Vac (1 hour)	Yes. No exceedance of annual value but considered further because it does exceed screening value for 1 hour during Construction at HHRA3	
	374	174	300	1 hour	SAAQS/AAAQO			
	0.0737	0.225	20	Annual	SAAQS/AAAQO			
Sulphur dioxide (SO ₂)	1.19	6.31	125	24 hours	SAAQS/AAAQO	No	No	
	8.63	25.0	450	1 hour	SAAQS/AAAQO			
Code a como de (CO)	771	994	6,000	8 hours	SAAQS/AAAQO	NI -	No. No annual value but not considered further because it does not	
Carbon monoxide (CO)	1,170	1,232	15,000	1 hour	SAAQS/AAAQO	No	exceed screening values for 8 hours or 1 hour	
	n/c	0.816	5	24 hours	OAAQC	NI.	No. No annual value but not considered further because it does not	
Sulphuric acid (H ₂ SO ₄)	n/c	3.19	10	1 hour	AAAQO	No	exceed screening values for 24 hours or 1 hour	
	15.3	10.2	60	Annual	SAAQS/OAAQC		Yes. No exceedance of annual value but considered further because in does exceed screening values for 24 hours during Construction and Operations at HHRA3	
SP	296	103	100	24 hours	SAAQS/AAAQO	Yes (24 hour)		
PM ₁₀	164	70.8	50	24 hours	SAAQS/OAAQC	Yes (24 hour)	Yes. No annual value but considered further because it does exceed screening values for 24 hours during Construction and Operations at HHRA3	
	3.10	3.84	8.8	Annual	OAAQC/CAAQS		Yes. No exceedance of annual value but considered further because it	
PM _{2.5}	65.5	28.0	27	24 hours	OAAQC/CAAQS	Yes (24 hour)	does exceed screening values for 24 hours during Construction and Operations at HHRA3	
<u>Oust</u>								
CD damanitian	87.9	5.33	7	Annual	OAAQC Dustfall Criteria (g/m²/yr)	Vac (A	Yes. Considered further because it exceeds annual value during Construction at HHRA3	
TSP deposition	1.61	1.08	5.3	Monthly	AAAQO (g/m²/30 days)	Yes (Annual)		
Dioxins and furans								
Dioxins and furans	0.000927	0.00186	0.1	24 hour	OAAQC (pg TEQ/m³)	No	No. No annual value but not considered further because it does not exceed screening value for 24 hours	
<u>Radon</u>								
Radon	n/c	44.5	200	Annual	Government of Canada Radon Guideline (Bq/m³) (Health Canada (2009)	No	Yes. Assessed in terms of radiation dose in the ERA	
<u>Metals</u>								
Silver (Ag)	n/c	3.00 × 10 ⁻⁰⁵	1	24 hour	OAAQC	No	No. No annual value but not considered further because it does not exceed screening value for 24 hours	
America (A.A.)	n/c	3.00 × 10 ⁻⁰⁵	0.01	Annual	AAAQO	No	No	
Arsenic (As)	n/c	3.90 × 10 ⁻⁰⁴	0.3	24 hour	OAAQC	No	No	
Barium (Ba)	n/c	3.70 × 10 ⁻⁰²	10	24 hour	OAAQC	No	No. No annual value but not considered further because it does not exceed screening value for 24 hours	

Constituent		Maximum Concentration at Receptor Locations		A Davied	Saura .	Is Concentration Greater than	Leather Countries and Restricted for Forether Countries and CORCO	
Constituent	Construction	Operations	Value	Averaging Period	Source	Selected Screening Value (Yes/No)	Is the Constituent Retained for Further Consideration as a COPC?	
Beryllium (Be)	n/c	6.00 × 10 ⁻⁰⁵	0.01	24 hour	OAAQC	No	No. No annual value but not considered further because it does not exceed screening value for 24 hours	
Cadmium (Cd)	n/c	5.00 × 10 ⁻⁰⁵	0.005	Annual	OAAQC	No	No	
Cadmium (Cd)	n/c	5.50 × 10 ⁻⁰⁴	0.025	24 hour	OAAQC	No	No	
Cobalt (Co)	n/c	1.05 × 10 ⁻⁰³	0.1	24 hour	OAAQC	No	No. No annual value but not considered further because it does not exceed screening value for 24 hours	
Chromium (Cr)	n/c	1.52 × 10 ⁻⁰²	0.5	24 hour	OAAQC	No	No. No annual value but not considered further because it does not exceed screening value for 24 hours	
Copper (Cu)	n/c	4.96 × 10 ⁻⁰³	50	24 hour	OAAQC	No	No. No annual value but not considered further because it does not exceed screening value for 24 hours	
Mercury (Hg)	n/c	8.00 × 10 ⁻⁰⁴	2	24 hour	OAAQC	No	No. No annual value but not considered further because it does not exceed screening value for 24 hours	
Molybdenum (Mo)	n/c	3.84 × 10 ⁻⁰³	120	24 hour	OAAQC	No	No. No annual value but not considered further because it does not exceed screening value for 24 hours	
Nickel (Ni) in TSP	n/c	1.70×10^{-04}	0.04	Annual	OAAQC	No	No	
	n/c	2.99×10^{-03}	0.2	24 hour	OAAQC	No	No	
Nickel (Ni) in PM ₁₀	n/c	1.20×10^{-04}	0.02	Annual	OAAQC	No	No	
Nickei (Ni) III Pivi ₁₀	n/c	2.75×10^{-03}	0.1	24 hour	OAAQC	No	No	
Lead (Pb)	n/c	7.40×10^{-04}	0.2	Monthly	OAAQC	No	No. No annual value but not considered further because it does not	
Lead (PD)	n/c	8.20×10^{-03}	0.5	24 hour	OAAQC	No	exceed screening values for 24 hours and monthly	
Antimony (Sb)	n/c	5.00 × 10 ⁻⁰⁵	25	24 hour	OAAQC	No	No. No annual value but not considered further because it does not exceed screening values for 24 hours and monthly	
Selenium (Se)	n/c	9.00 × 10 ⁻⁰⁵	10	24 hour	OAAQC	No	No. No annual value but not considered further because it does not exceed screening value for 24 hours	
Tin (Sb)	n/c	1.30 × 10 ⁻⁰⁴	10	24 hour	OAAQC	No	No. No annual value but not considered further because it does not exceed screening value for 24 hours	
Uranium (U) in TSP	n/c	9.38 × 10 ⁻⁰³	0.06	Annual	OAAQC	No	No	
Oranium (O) in 15P	n/c	2.68 × 10 ⁻⁰¹	0.3	24 hour	OAAQC	No	No	
Haratina (II) in DM	n/c	6.10 × 10 ⁻⁰³	0.03	Annual	OAAQC	No	Yes. Does not exceed annual value but considered further because it	
Uranium (U) in PM ₁₀	n/c	2.09 × 10 ⁻⁰¹	0.15	24 hour	OAAQC	Yes (24 hour)	exceeds the 24 hours screening value during Operations at HHRA3	
Vanadium (V)	n/c	7.82 × 10 ⁻⁰³	2	24 hour	OAAQC	No	No. No annual value but not considered further because it does not exceed screening value for 24 hours	
Zinc (Zn)	n/c	2.85 × 10 ⁻⁰³	120	24 hour	OAAQC	No	No. No annual value but not considered further because it does not exceed screening value for 24 hours	
Cesium (Cs)	n/c	<0.00001	2	Annual	Texas Interim ESL	No	No. No 24 hours value but not considered further because it does not exceed the annual screening value	
Bismuth (Bi)	n/c	<0.00001	5	Annual	Texas Interim ESL	No	No. No 24 hours value but not considered further because it does not exceed the annual screening value	

	Maximum Concent Loca	Screening				Is Concentration Greater than		
Constituent	Construction	Operations	Value	Averaging Period		Source	Selected Screening Value (Yes/No)	Is the Constituent Retained for Further Consideration as a COPC?
Iron (Fe)	n/c	1.35 × 10 ⁺⁰⁰	4	24 hour	OAAQC		No	No. No annual value but not considered further because it does not exceed screening value for 24 hours
Magnesium (Mg)	n/c	1.20 × 10 ⁺⁰⁰	72	24 hour	OAAQC		No	No. No annual value but not considered further because it does not exceed screening value for 24 hours
Manganese (Mn)	n/c	7.50 × 10 ⁻⁰⁴	0.2	Annual	AAAQO		No	No
	n/c	1.86 × 10 ⁻⁰²	0.1	24 hour	OAAQC		No	No

Notes: Air Concentrations are maximum predicted values from the Air Quality model for HHRA locations 1 through 11, inclusively, for the period indicated.

Maximum Concentration values are rounded to 3 significant figures.

For metals, where the model value was zero, the concentration was assumed to be $<1.00 \times 10^{-06} \, \mu g/m^3$. Units are $\mu g/m^3$ unless otherwise specified.

Bold represents air quality parameters predicted to exceed screening values at receptor locations or are parameters that did not exceed the screening level but are discussed further in the ERA.

n/c = not calculated; n/v = no value; n/a = not applicable; ERA = environmental risk assessment; SAAQS = Saskatchewan Ambient Air Quality Criteria (MECP 2020); CAAQS = Canadian Ambient Air Quality Standards (CCME 2021b); < = less than; Bq/m³ = becquerels per cubic metre; TSP = total suspended particulate matter with a diameter of 10 microns or less; PM_{2.5} = particulate matter with a diameter of 2.5 microns or less; ESL = effects screening level; pg TEQ/m³ = picogram toxic equivalency; CAC = criteria air contaminant.

Table 4-8: Air Quality Screening for Short-term Exposures to Constituents in Air for Hypothetical Human and Ecological Receptors at the Fence Line

Constituent		Maximum Concentration at Fence Line Locations		Averaging Period	Source	Is Concentration Greater than Selected Screening	Is the Constituent Retained for further Consideration as a COPC?	
	Construction	Operations	Value			Value (Yes/No)		
CACs								
Nitro man diavida (NO.)	95	73	200	24 hours	SAAQS/AAAQO	No	Yes. Did not exceed the 24-hour value but considered further because it exceeds its 1-hour screening	
Nitrogen dioxide (NO ₂)	458	205	300	1 hour	SAAQS/AAAQO	Yes (1 hour)	value	
	0.79	3.10	125	24 hours	SAAQS/AAAQO	No	No	
Sulphur dioxide (SO ₂)	19.74	19.7	450	1 hour	SAAQS/AAAQO	No	No	
	831	792	6,000	8 hours	SAAQS/AAAQO	No	No	
Carbon monoxide (CO)	2,693	8,759	15,000	1 hour	SAAQS/AAAQO	No	No	
C 1-1	n/c	0.391	5	24 hours	OAAQC	No	No	
Sulphuric acid (H ₂ SO ₄)	n/c	2.02	10	1 hour	AAAQO	No	No	
TSP	234	173	100	24 hours	SAAQS/AAAQO	Yes (24 hour)	Yes	
PM ₁₀	204	86.6	50	24 hours	SAAQS/OAAQC	Yes (24 hour)	Yes	
PM _{2.5}	51.5	35.1	27	24 hours	OAAQC/CAAQS	Yes (24 hour)	Yes	
Dioxins and furans	-	'						
Dioxins and furans	0.00119	0.00239	0.1	24 hour	OAAQC (pg TEQ/m³)	No	No	
<u>Metals</u>		1	1					
Silver (Ag)	n/c	1.10 × 10 ⁻⁰⁴	1	24 hour	OAAQC	No	No	
Arsenic (As)	n/c	8.70 × 10 ⁻⁰⁴	0.3	24 hour	OAAQC	No	No	
Barium (Ba)	n/c	5.47 × 10 ⁻⁰²	10	24 hour	OAAQC	No	No	
Beryllium (Be)	n/c	1.00 × 10 ⁻⁰⁴	0.01	24 hour	OAAQC	No	No	
Cadmium (Cd)	n/c	6.40 × 10 ⁻⁰⁴	0.025	24 hour	OAAQC	No	No	
Cobalt (Co)	n/c	1.66 × 10 ⁻⁰³	0.1	24 hour	OAAQC	No	Non	
Chromium (Cr)	n/c	2.84 × 10 ⁻⁰²	0.5	24 hour	OAAQC	No	No	
Copper (Cu)	n/c	1.65 × 10 ⁻⁰²	50	24 hour	OAAQC	No	No	
Mercury (Hg)	n/c	8.00 × 10 ⁻⁰⁴	2	24 hour	OAAQC	No	No	
Molybdenum (Mo)	n/c	1.34 × 10 ⁻⁰²	120	24 hour	OAAQC	No	No	
N. 1 1 0 12	n/c	3.73 × 10 ⁻⁰³	0.2	24 hour	OAAQC in TSP	No	No	
Nickel (Ni)	n/c	3.11 × 10 ⁻⁰³	0.1	24 hour	OAAQC in PM ₁₀	No	No	
Lead (Pb)	n/c	2.77 × 10 ⁻⁰²	0.5	24 hour	OAAQC	No	No	
Antimony (Sb)	n/c	1.70 × 10 ⁻⁰⁴	25	24 hour	OAAQC	No	No	
Selenium (Se)	n/c	3.10 × 10 ⁻⁰⁴	10	24 hour	OAAQC	No	No	
Tin (Sb)	n/c	2.10 × 10 ⁻⁰⁴	10	24 hour	OAAQC	No	No	
	n/c	8.55 × 10 ⁻⁰¹	0.3	24 hour	OAAQC in TSP	Yes (24 hour)	Yes	
Uranium (U)	n/c	6.74 × 10 ⁻⁰¹	0.15	24 hour	OAAQC in PM ₁₀	Yes (24 hour)	Yes	

Constituent	Maximum Concentration at Fence Line Locations		Screening Avera	Averaging Period		Is Concentration Greater than Selected Screening	Is the Constituent Retained for further Consideration as a COPC?		
	Construction	Operations				Value (Yes/No)			
Vanadium (V)	n/c	1.30 × 10 ⁻⁰²	2	24 hour	OAAQC	No No			
Zinc (Zn)	n/c	3.76 × 10 ⁻⁰³	120	24 hour	OAAQC	No No			
Iron (Fe)	n/c	1.85 × 10 ⁺⁰⁰	4	24 hour	OAAQC	No No			
Magnesium (Mg)	n/c	1.56 × 10 ⁺⁰⁰	72	24 hour	OAAQC	No No			
Manganese (Mn)	n/c	3.52 × 10 ⁻⁰²	0.1	24 hour	OAAQC	No No			

Notes: Units are µg/m³ unless otherwise specified.

Air Concentrations are maximum predicted values from the Air Quality model for HHRA fence line locations, for the period indicated.

Maximum Concentration values are rounded to 3 significant figures.

For metals, where the model value was zero, the concentration was assumed to be $<1.00 \times 10^{-06} \, \mu g/m^3$.

Bold represents air quality constituents with maximum concentrations predicted at fence line locations that exceed screening values during Construction or Operations.

< = less than; SAAQS = Saskatchewan Ambient Air Quality Standards; AAAQO = Alberta Ambient Air Quality Objectives; OAAQC = Ontario Ambient Air Quality Standards; n/c = not calculated; n/v = no value; n/a = not applicable; pg TEQ/m³ = picogram toxic equivalency; PM₁₀ = particulate matter with a diameter of 10 microns or less; PM_{2.5} = particulate matter with a diameter of 2.5 microns or less; CAC = criteria air contaminant.

4.3.3.3 Discussion of Air Quality Constituents that Exceed a Screening Value

Air quality constituents that exceeded a screening value were nitrogen dioxide, particulate matter (TSP, PM_{10} , $PM_{2.5}$, and TSP deposition), and uranium (Table 4-9). These constituents were further evaluated to determine if they require further assessment in the ERA. Additionally, radionuclides were further evaluated because a relevant screening value was not available.



Table 4-9: Summary of Air Quality Constituents that Exceed a Screening Value

	Screening Cri	teria Exceeded	Predicted Exc			
Constituent	Short-Term	Long-Term	Maximum at Human/Ecological Locations	Maximum at Fence Line	Number of Days Exceeding (Fence Line)	Frequency Exceeding (Fence Line)
Nitrogen dioxide	1 h	None	Construction: exceedance of 1 hr screening value for nitrogen dioxide at the camp (HHRA3) Operations: No exceedance	Construction: exceedance of 1 hr screening value Operations: No exceedance	Construction: n/a ^(a) Operations: n/a	Construction: n/a ^(a) Operations: n/a
Particulate matter	24 h: TSP, PM ₁₀ , PM _{2.5}	Annual: TSP deposition	Construction: exceedances of 24 hr screening values for TSP, PM ₁₀ , PM _{2.5} at the camp (HHRA3), and annual TSP deposition screening value at the camp (HHRA3) Operations: exceedance of 24 hr screening values for TSP, PM ₁₀ , PM _{2.5} at the camp (HHRA3)	Construction: exceedance of 24 hr screening values for TSP, PM ₁₀ , PM _{2.5} Operations: exceedance of 24 hr screening values for TSP, PM ₁₀ , PM _{2.5}	Construction: 10 (PM ₁₀) 4 (TSP) Operations: 2 (PM ₁₀) 2 (TSP)	Construction: 2.7% (PM ₁₀) 1.1% (TSP) Operations: 0.5% (PM ₁₀) 0.5% (TSP)
Uranium	24 h	none	Construction: not modelled for metals Operations: exceedance of 24 h screening value at the camp (HHRA3)	Construction: not modelled for metals Operations: exceedance of 24 h screening value for uranium in TSP and PM ₁₀	Construction: n/a Operations: 4 (PM ₁₀) 2 (TSP)	Construction: n/a Operations: 1.1% (PM ₁₀) 0.5% (TSP)

⁽a) The number of days and frequency of exceeding were not available for 1-hour nitrogen dioxide since no exceedance was identified in the air quality model due to guidance from Saskatchewan that allows for removal of maximum concentrations output from AERMOD (EIS Section 7.2.2.8.3, Dispersion Modelling).



ERA = environmental risk assessment; TSP = total suspended particulates; $PM_{2.5}$ = particulate matter with a diameter of 2.5 microns or less; n/a = not applicable.

4.3.3.3.1 Nitrogen Dioxide

Screening values were available for 1-hour, 24-hour, and annual averaging periods for nitrogen dioxide. The maximum predicted 1-hour concentrations of nitrogen dioxide during Construction exceeded the screening value of 300 micrograms per cubic metre (μ g/m³) only at the camp (374 μ g/m³) and at the fence line (458 μ g/m³). Maximum predicted short-term (i.e., 1-hour) levels of nitrogen dioxide at all other modelled human and ecological receptor locations farther away from the Project footprint were below Health Canada's maximum acceptable level and for this reason are not expected to be associated with adverse health effects. No exceedances were predicted for 24-hour and annual exposures; therefore, no adverse health effects are anticipated due to long-term exposure.

Potential adverse human and ecological effects from nitrogen dioxide are associated with direct atmospheric exposure through inhalation. Adverse health effects that are attributed to short-term exposures to ambient nitrogen dioxide include asthma exacerbations and possibly increased risk of cardiopulmonary effects, and to a lesser extent cardiovascular and respiratory mortality (Health Canada 2016b). Individuals with certain pre-existing diseases such as asthma appear to be sensitive to exposure to ambient nitrogen dioxide. If individuals are present during periods when ambient nitrogen dioxide concentrations exceed the screening value, it is possible that they could experience minor irritation of the respiratory system. These effects would be reversible and would subside after exposure.

Additionally, since potential adverse effects are associated with the inhalation pathway, nitrogen dioxide was not retained as a COPC for assessment of potential risk to human and ecological receptors through the food chain.

Overall, exceedance of the 1-hour short-term screening value for nitrogen dioxide at the camp site would be limited to Construction, and any health effects would be reversible and subside after exposure; therefore, nitrogen dioxide was not considered for further quantitative assessment in the ERA.

As noted in Section 4.3.3.1, Screening Value Selection, the CAAQS were not selected as screening criteria as CAAQS achievement is determined by provinces and territories using ambient concentrations measured in the air zones for a three-year period, not by comparison of modelled predictions at or beyond a facility boundary. Therefore, using the CAAQS as a screening guideline would not be appropriate. However, for information purposes, a comparison of Project predicted nitrogen dioxide emissions were compared to the annual and 1-hour nitrogen dioxide CAAQS. In summary, there are no predicted exceedances of the annual nitrogen dioxide CAAQS of 23 µg/m³ at any receptor location during any phase of the Project under the Application Case or the RFD Case. With respect to the 1-hour nitrogen dioxide CAAQS of 79 µg/m³, during Construction, exceedances are predicted to occur at the camp location approximately 7% of the time, and during Operations under both the Application Case and the RFD Case, exceedances are predicted to occur at the camp location approximately 6% of the time. Realistically, these exceedances are anticipated to occur less frequently as the predictions include multiple levels of conservatism applied in the air quality model. To facilitate comparison against the CAAQS, the predictions

ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT MODEL INTEGRATION AND EVALUATION OF SOURCES

referenced in this paragraph represent the annual average predicted concentrations and the 3-year average of the annual 98th percentile of the daily maximum 1-hour predicted concentrations.

4.3.3.3.2 Particulate Matter

Particulate matter is defined as liquid or solid particles, or a mixture of both, less than 100 μ m in diameter. Particulate matter includes TSP, particulate matter less that 10 μ m (PM₁₀), and particulate matter less than 2.5 μ m (PM_{2.5}). Particulate matter in the form of TSP, PM₁₀, PM_{2.5}, and TSP deposition were screened. Screening values were based on 24-hour and annual averaging periods for TSP, PM₁₀, PM_{2.5}, and on an annual averaging period for TSP deposition.

The discussion below demonstrates that the predicted exceedances of screening values for particulate matter in all of the forms assessed (i.e., TSP, PM₁₀, PM_{2.5}, and TSP deposition) would be short-term (i.e., 24-hour averaging period) and spatially limited to the camp and the fence line. Particulate concentrations and TSP deposition at locations outside of the Project footprint did not exceed screening values.

The predicted exceedances of screening values within the Project fence line suggest that the Project's environmental management programs would need to include measures to monitor and control particulate matter at the site during Construction and Operations.

Total Suspended Particulates and Dust Deposition

Maximum predicted concentrations of TSP exceeded its 24-hour screening value ($100 \,\mu g/m^3$) only at the camp site ($296 \,\mu g/m^3$ during Construction and $103 \,\mu g/m^3$ during Operations), and at the fence line ($234 \,\mu g/m^3$ during Construction and $173 \,\mu g/m^3$ during Operations). During Construction, the frequency of exceedance of the 24-hour TSP criterion at the fence line would be low, approximately 1.1% and during Operations, it would be 0.5%. On an annual basis, the maximum predicted concentrations for airborne TSP would not exceed screening values at any location outside of the Project footprint during any Project phase.

The 24-hour screening value for TSP is an ambient air quality standard cited by both Saskatchewan and Alberta. The 24-hour ambient air quality objective is based on potential adverse pulmonary effects (Alberta 2021). A higher 24-hour effects-based screening value of 120 $\mu g/m^3$ for TSP in ambient air is available from Ontario. The Ontario 24-hour and annual ambient air quality criteria (AAQC) are meant to be protective of chronic effects. Ontario identifies visibility as the sensitive endpoint for the TSP AAQC rather than human or ecological health. Elevated TSP concentrations are generally not considered to pose significant health risks because these particles are too large to be inhaled deep into the lungs; therefore, TSP was not considered further in the ERA.

Dust deposition (measured as dustfall) was also evaluated during this screening. Dustfall was not assessed for direct human health and ecological risks. Exceedance of the dustfall criterion is aesthetic and not health based. The maximum predicted TSP deposition rate exceeded the annual Ontario dustfall criterion of 7 g/m²/yr only during Construction and only within the Project footprint and at the camp (87.9 g/m²/yr), and did not extend to other human and ecological receptor locations at Patterson Lake, Beet Lake, and Lloyd Lake where country foods harvesting

ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT MODEL INTEGRATION AND EVALUATION OF SOURCES

may occur. The maximum predicted monthly TSP deposition did not exceed its screening value of 5.3 g/m²/30 days at any location during any Project phase. For these reasons, dustfall was not considered further in the ERA for direct human health and ecological risks. However, the potential for air quality constituents in dust to deposit to soil are considered in Section 4.3.3.4, Constituents in Soil.

Particulate Matter (PM₁₀ and PM_{2.5})

Maximum predicted concentrations of PM_{10} and $PM_{2.5}$ exceeded their 24-hour screening values (50 μ g/m³ and 27 μ g/m³, respectively) only within the Project footprint, at the camp (164 μ g/m³ and 65.5 μ g/m³ respectively during Construction, and 70.8 μ g/m³ and 28.0 μ g/m³ respectively during Operations), and at the fence line (204 μ g/m³ and 51.5 μ g/m³ respectively during Construction, and 86.6 μ g/m³ and 35.1 μ g/m³ respectively during Operations). During Construction, the frequency of exceedance of 24-hour PM_{10} at the fence line was low, approximately 2.7% and during Operations, it was 0.5%. The maximum predicted concentration for PM_{10} and $PM_{2.5}$ did not exceed their annual screening values at any modelled location during any Project phase.

Human health has been shown to be the most sensitive receptor for exposure to PM_{10} and $PM_{2.5}$ in ambient air (Health Canada 1998). Exposure to elevated concentrations of both PM_{10} and $PM_{2.5}$ are associated with various respiratory and cardiovascular effects in humans. The finer particles that can be inhaled deeply into the lungs are associated with greater risk because they are more chemically active and have more complex characteristics than larger particles (Health Canada 2016c). If individuals are present during short-term periods of elevated PM_{10} and/or $PM_{2.5}$, they may experience respiratory symptoms such as coughing or difficulty breathing, or asthma symptoms and chronic bronchitis. For most individuals, effects would be reversible and subside after exposure.

Overall, exceedances of the 24-hour short-term screening values for PM_{10} and $PM_{2.5}$ were predicted during Construction and Operation at the camp site and at the fence line. However, health effects would be infrequent, reversible and subside after exposure; therefore, PM_{10} and $PM_{2.5}$ were not considered for further quantitative assessment in the ERA.

4.3.3.3.3 Uranium

Maximum concentrations of uranium were estimated for Operations because sources would be associated with mining and waste management that would occur during Operations. Screening values for uranium were available for 24-hour and annual averaging periods that are associated with the TSP and PM₁₀ fractions of airborne particulate matter. For both uranium in TSP and in PM₁₀, the Ontario 24-hour screening values were calculated from an effects-based annual average value to allow for assessment of the 24-hour air quality data (MECP 2020). The maximum predicted concentrations of uranium did not exceed annual screening values in TSP and PM₁₀ (0.06 μ g/m³ and 0.3 μ g/m³, respectively) at any ERA location during Operations.

Ecometrix Environmental INTELLIGENCE

ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT MODEL INTEGRATION AND EVALUATION OF SOURCES

Uranium can be toxic to humans due to its chemical and radiological properties. The ambient air quality criteria for uranium (MOE 2011) is based on non-radiological effects of which kidney toxicity was the most sensitive endpoint associated with chronic exposure to uranium in air. Since the predicted maximum concentrations did not exceed the annual screening value, unacceptable levels of risk for human and ecological health are not expected from the occasional exceedances of the 24-hour value. Potential non-radiological risks from exposure to uranium in air were not assessed further.

Consideration of potential radiological effects for uranium, from uranium-238 series radionuclides in air, are discussed in Section 4.3.3.3.4, Radon and Uranium-238 Series Radionuclides.

4.3.3.3.4 Radon and Uranium-238 Series Radionuclides

Maximum concentrations for radon gas were estimated for the operational period because radon sources are associated with mining and waste management during Operations. A remedial action level of 200 Bq/m³ for radon for indoor air in dwellings (Health Canada 2014) was used as a screening value for radon. The predicted maximum annual value for radon in ambient air did not exceed the annual screening value at any human or ecological receptor location.

No concentration-based screening values for uranium-238 series radionuclides in air were available. However, because radionuclides are considered of public interest, they were assessed further in the ERA. Potential risks from radiological effects were assessed through environmental exposure pathways in terms of total radiation dose, in Section 5.0, Human Health Risk Assessment, for human health risks, and in Section 6.0, Ecological Risk Assessment, for ecological risks.

4.3.3.4 Constituents in Soil

No specific COPCs were retained from the screening of atmospheric constituents; however, as a secondary check, mine-related metals of interest were identified for further assessment in soil (Table 4-10).

Based on soil type characterization in the Terrain and Soils section (EIS Section 12), the soil type is sand. This also aligns with observations of the Patterson Lake area by BNDN and BRDN, which was characterized as having a lot of sand and supporting jackpine (TSD II: BNDN), and mostly all pine and rock (TSD III: BRDN). A member of BNDN commented that "the north" has experienced many wildfires in recent years resulting in no topsoil and only sand, which made it challenging for trees to regenerate (TSD II: BNDN).

Predicted soil concentrations were estimated from atmospheric deposition and maximum air concentrations from the AERMOD atmospheric model at the camp site (Table 4-7), along with constituent-specific deposition rates, according to the equations defined in the IMPACT Model Report (Appendix A, Section 2.3.4, Terrestrial Pathways). The camp site is represented in the IMPACT model as the terrestrial location of Patterson Lake North Arm – West Basin, and would be considered the closest human health receptor location to the Project where harvesting of traditional foods would likely occur.



ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT MODEL INTEGRATION AND EVALUATION OF SOURCES

Predicted maximum concentrations of constituents in soil from atmospheric deposition were compared against soil quality guidelines. The selected soil quality guidelines were the federal CCME (CCME 1999b) soil quality guidelines for protection of human health and environmental health. Agricultural soil quality values were used for other land types, because these guidelines account for ingestion of plants by birds and mammals. As shown in Table 4-10, all predicted soil concentrations were below the CCME soil quality guidelines. As such, no additional COPCs were identified for further quantitative assessment in the ERA based on the soil pathway.



Table 4-10: Soil Quality Screening for the Rook I Project

	Maximum Predicted Soil Concentration Concentrations(a) Atmospheric Deposition(b)			Soil Screening Guideline ^(c)					
Parameter			Agricultural	Residential/ Parkland	Commercial	Industrial	Is Concentration Greater than Selected Screening		
Non-radionuclides	Annual average µg/m³	mg/kg dw	mg/kg dw	mg/kg dw	mg/kg dw	mg/kg dw	Value? (Y/N)		
Arsenic	3.00×10^{-05}	0.59	12	12	12	12	N		
Cadmium	5.00 × 10 ⁻⁰⁵	0.11	1.4	10	22	22	N		
Cobalt	2.00 × 10 ⁻⁰⁵	0.43	40	50	300	300	N		
Copper	1.70 × 10 ⁻⁰³	0.62	63	63	91	91	N		
Lead	2.50 × 10 ⁻⁰³	1.71	70	140	260	600	N		
Molybdenum	1.20 × 10 ⁻⁰³	0.15	5	10	40	40	N		
Nickel	1.70 × 10 ⁻⁰⁴	1.05	45	45	89	89	N		
Selenium	1.00 × 10 ⁻⁰⁵	0.10	1	1	2.9	2.9	N		
Uranium	9.38 × 10 ⁻⁰³	1.30	23	23	33	300	N		
Zinc	5.00 × 10 ⁻⁰⁵	2.72	250	250	410	410	N		

Notes: **Bold** indicates soil guideline value selected for this assessment.



⁽a) Maximum annual average concentrations in the Project footprint, and the camp site from AERMOD (EIS Appendix 7A, Air Dispersion Modelling Report).

⁽b) Maximum soil concentrations estimated from maximum annual air concentrations in Table 4-7 of the HHRA and constituent-specific deposition rates (Appendix A).

⁽c) Canadian Environmental Quality Guidelines (CCME 1999b).

N = no; Y = yes; dw = dry weight; HHRA = human health risk assessment.

4.3.4 Air Quality Constituents for Further Evaluation in the Environmental Risk Assessment

Following the screening process for selecting air quality COPCs, none of the modelled air quality constituents were considered COPCs for further evaluation in the ERA for direct atmospheric exposure for human and ecological receptors. No constituents at any human or ecological receptor location exceeded their annual screening value, indicating that unacceptable chronic effects from direct exposure to air would not be expected.

Short-term exceedances, based on maximum predicted concentrations for the 24-hour averaging period, may occur at the camp and at the fence line, for nitrogen dioxide, and particulate matter, including uranium in TSP and PM_{10} . The predicted exceedances were infrequent, short-term, and limited to near the site. Unacceptable levels of risk would not be expected from infrequent direct short-term exposures to these constituents in air.

Air quality constituents would be monitored as part of an overall Environmental Protection Program, which would include ambient air monitoring and adaptive management. Additionally, on-Site health and safety requirements and mitigation measures would be developed to control dust emissions. Monitoring would be used to verify the air quality predictions and evaluate the effectiveness of mitigation measures.

No concentration-based screening values were available for uranium-238 series radionuclides in air. However, because radionuclides are considered of public interest, they were assessed further in the ERA. The ERA included the assessment of human and ecological risk from exposure to radionuclides as part of the total radiological dose from atmospheric and aquatic pathways combined in Section 5.4 and Section 6.4, respectively. While radon concentrations were predicted to be below the screening value, this constituent was considered further in the HHRA in Section 5.4, due to public interest.

Other air quality constituents were included in the evaluation of potential human health and ecological risk in Section 5.4 and Section 6.4, respectively, via indirect exposures, such as soil contact and through the food chain (including country foods). These parameters were selected based on a cumulative exposure pathways approach that included COPCs identified for exposures through aquatic pathways (Section 4.2.5, Water Quality Constituents for Further Evaluation in the Environmental Risk Assessment) and engagement with Indigenous and other communities and regulators.

4.4 Final List of Constituents of Potential Concern for the Environmental Risk Assessment

Based on evaluation of aqueous and atmospheric sources, including a conservative screening of maximum predicted concentrations in surface water, sediment, air and soil, the final list of COPCs to be evaluated further in the HHRA and EcoRA are listed in Table 4-11.



As indicated in Section 4.2.3.2, Constituents in Surface Water, radon-222 was not considered a COPC in surface water for the ERA. Radon is expected to volatilize rapidly to air. Health Canada (2020) considers that the health risk from ingesting radon-contaminated drinking water is negligible. Additionally, chloride and sulphate were identified as COPCs in the aquatic environment, but not in the terrestrial environment. As well, while deposition of COPCs from air to soil was evaluated, no COPCs in soil were identified for further evaluation; therefore, for the EcoRA, toxicity via direct contact with COPCs in soil for terrestrial plants and soil invertebrates was not quantitatively assessed.

Table 4-11: Final List of Constituents of Potential Concern for the Project Environmental Risk Assessment

Major Ions	Physical Media Where Guideline Exceeded
Chloride	Water
Sulphate	Water
Metals and Metalloids	
Arsenic	Water
Cobalt	Water
Copper	Water
Molybdenum	Water
Uranium	Water
Radionuclides	
Uranium-238	Air, Water
Uranium-234	Air, Water
Thorium-230	Air, Water
Radium-226	Air, Water
Radon-222 ^(a)	Air
Lead-210	Air, Water
Polonium-210	Air, Water

(a) Radon-222 is evaluated as a COPC for air only.

 ${\sf COPC} = {\sf constituent} \ {\sf of} \ {\sf potential} \ {\sf concern}.$

5.0 HUMAN HEALTH RISK ASSESSMENT

The components of an HHRA are Problem Formulation (Section 5.1), Exposure Assessment (Section 5.2), Toxicity Assessment (Section 5.3), and Risk Characterization (Section 5.4).

5.1 Problem Formulation

The intent of the problem formulation for an HHRA is to define the goals of the risk assessment, develop an understanding of site conditions, and develop working hypotheses as to how potential exposure of people to contaminants may result in potential risks to human health.

The assessment endpoint for the HHRA is the health of individual humans.

The problem formulation for this HHRA:

- identifies COPCs for human health risks;
- identifies and characterizes non-nuclear energy workers who may frequent the site, and the selected human health receptor groups that represent them in the ERA; and
- identifies the complete exposure pathways by which the COPCs may affect the human health receptors in a conceptual site model.

The conceptual site model for the HHRA summarizes the links between contaminant sources, exposure pathways, and receptors of concern.

5.1.1 Human Receptor Selection and Characterization

The human receptors for the HHRA were selected to be appropriate for assessment of effects on human health from both radiological and non-radiological COPCs. Off-Site members of the public would potentially be exposed to low levels of airborne or waterborne constituents being released during Project activities.

Nuclear energy workers are outside of the scope of this assessment. Consistent with CSA N288.6-22, nuclear energy workers would be classified and monitored in accordance with the requirements of the Radiation Protection Program, and therefore did not require assessment in the HHRA. Non-nuclear energy workers at the Project site that would be subject to occupational exposure and workplace monitoring outlined in the Health and Safety Program were also excluded from the HHRA. However, in the HHRA, some workers at the Project were assumed to frequent the LSA and consume Traditional Foods, and fish, hunt, and harvest from the LSA when not working. For this reason, camp workers at the Project were assessed for both radiological and non-radiological exposures. This approach is consistent with CSA N288.6-22 (CSA 2022).

The selection of human health receptor groups was based on current understanding of how people use the Project LSA and RSA, information from IKTLU Studies, which was supplemented by information obtained during community information sessions (NexGen 2019) and JWG



ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT HUMAN HEALTH RISK ASSESSMENT

meetings, as well as the potential for exposure to Project-related media (i.e., air, soil, water, sediment) concentrations during one or more Project phases. Community information sessions were held by NexGen in June 2019. As part of the process, community members were invited to share information about their local land and resource uses in proximity to the Project site through a map-based exercise. Information collected during the mapping exercise informed the initial selection of receptor locations in the study areas relative to existing or previous uses. The IKTLU Studies were also used in the ERA as a primary source of information (TSD II: BNDN; TSD III: BRDN; TSD IV: MN-S; TSD V.1: CRDN; TSD VI: YNLR). These studies provided initial guidance for identifying locations where people may reside; areas where Traditional Foods are hunted, fished, and gathered; and mammal, bird, and plant species that are traditionally used by Indigenous Groups for food, medicine, and other traditional uses. A summary of the rationale for selection is provided in Table 5-1.

Adult and one-year-old receptors were used to assess potential risk to human health. The adult represents both male and female receptors. The one-year-old is equivalent to the CSA N288.1-20 age class "infant" and falls within the Health Canada age class "toddler". The selected human health receptor groups are shown on Figure 5-1 and included:

- camp worker at the Project (adult) Project lifespan;
- subsistence harvesters at Patterson Lake South Arm, Beet Lake, and Lloyd Lake (adult and one-year-old) – Project lifespan;
- seasonal residents/lodge operators at Patterson Lake and Lloyd Lake (adult and one-year-old) Project lifespan; and
- future permanent resident at Patterson Lake North Arm area (adult and one-year-old) far-future projection only.

Table 5-1: Rationale for Selection of Human Health Receptor Groups

Receptor	Rationale for Selection
Camp worker	The camp worker would be located within the Project footprint. A camp worker such as a camp cook would reside at camp for 50% of the year.
Subsistence harvester (Patterson Lake South Arm)	The subsistence harvester represents a high consumer of Traditional Foods. Patterson Lake is important to local Indigenous Groups who use the area for water, fishing, trapping, hunting, gathering, and cultural practices (TSD II: BNDN; TSD III: BRDN; TSD IV: MN-S; TSD V.1: CRDN; TSD V.2: CRDN; TSD VI: YNLR; CRDN 2019a,b; YNLRO 2019).



Receptor	Rationale for Selection
Subsistence harvester (Beet Lake)	The subsistence harvester is used to evaluate a high consumer of Traditional Foods in the LSA. Beet Lake is important to local Indigenous Groups, who use the area for water, fishing, hunting, and gathering (TSD V.2: CRDN, TSD IV: MN-S, TSD II: BNDN). The location is consistent with cabins identified by the MN-S and CRDN (MN-S; TSD V.1: CRDN; CRDN 2019a,b).
Subsistence harvester (Lloyd Lake)	The subsistence harvester is used to evaluate a high consumer of Traditional Foods in the RSA. Lloyd Lake is important to local Indigenous Groups, who use the area for camping, hunting, trapping, fishing, and gathering (TSD V.2: CRDN, TSD IV: MN-S, TSD II: BNDN). The location is consistent with cabins identified by the MN-S and CRDN (TSD IV: MN-S; TSD V.1: CRDN; CRDN 2019a,b).
Seasonal resident (Patterson Lake South Arm)	The seasonal resident represents an average consumer of Traditional Foods. A seasonal resident at Patterson Lake South Arm would also be representative of the Forrest Lake Outfitters Camp at Beet Lake. Patterson Lake is important to local Indigenous Groups, who use the area for water, fishing, trapping, hunting, gathering, and cultural practices (TSD II: BNDN; TSD III: BRDN; TSD IV: MN-S; TSD V.1: CRDN; TSD V.2: CRDN; TSD VI: YNLR; CRDN 2019a,b;
Seasonal resident (Lloyd Lake)	YNLRO 2019). The seasonal resident represents an average consumer of Traditional Foods in the RSA. There is an existing fly-in lodge at Lloyd Lake that provides fishing and spring bear hunting excursions for guests.
Future permanent resident (Patterson Lake North Arm area)	Used to evaluate a resident living in the Patterson Lake North Arm area in the far future following Closure of the Project. This receptor is considered hypothetical as it is in the far future. The future permanent resident represents a high consumer of Traditional Foods.

MN-S = Métis Nation — Saskatchewan; CRDN = Clearwater River Dene Nation; LSA = local study area; RSA = regional study area.

The purpose of this subsection is to provide a narrative description of the key assumptions used to characterize human receptors for the Project.

Human health exposure pathways considered for the Project for each of the human receptors, including assumptions for the Traditional Foods diet, are summarized in Section 5.1.3, Selection of Exposure Pathways. The human health conceptual model is provided in Section 5.1.4, Human Health Conceptual Site Model. The residency assumptions for each of the human health receptors (locations, durations, and frequencies) are described in Section 5.1.1.1 through Section 5.1.1.4 and summarized in the exposure assessment (Section 5.2.1, Exposure Locations, Duration, and Frequency).

5.1.1.1 Camp Worker

A camp worker at the Project (i.e., a non-nuclear energy worker such as a camp cook) would represent an adult male or female. This receptor group was assumed to work and reside at the Project site (camp) for 50% of the year and away from the Project site for the other 50% of the year. A camp worker was assessed for all Project phases except for the far-future projection.

When at work, the camp worker would be exposed to Project-related COPCs through inhalation of air and ingestion of and contact with water. Fresh water for drinking and bathing would be



ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT HUMAN HEALTH RISK ASSESSMENT

obtained from Patterson Lake North Arm – East Basin while at work. While at the Project site for work (i.e., during 50% of the year), it was assumed that the camp worker would not be permitted to hunt, fish, or gather berries/plants in the area and therefore would not ingest Traditional Foods from within the LSA.

Based on discussions with NexGen, a camp worker who also does subsistence harvesting was considered representative of the northern workforce. Therefore, the camp worker was assumed to consume Traditional Foods in their overall annual diet when not at work, with an ingestion rate consistent with an Indigenous Group's high consumer of Traditional Foods (Section 5.1.3, Selection of Exposure Pathways). For the HHRA, although the camp worker would live outside of the area of influence from the Project when not at work, it was assumed that they would travel to the LSA for subsistence harvesting activities (i.e., Patterson Lake South Arm area) for a portion of the year (i.e., 3 months; 25%). This is consistent with Indigenous and Local Knowledge that suggests that individuals are likely to travel to known fishing areas to catch a large supply of fish, to be retained for later consumption (TSD IV: MN-S). During their time spent harvesting, they may have incidental ingestion of soil and sediment, they may come in direct contact with surface water through swimming and sediments through wading, and they may obtain fresh water for drinking and bathing from Patterson Lake. The remainder of the time (i.e., 3 months; 25%), when not at work or harvesting in the LSA, the camp worker would reside outside of the area of potential influence from the Project and would be exposed to constituents in the environment at background levels.

5.1.1.2 Subsistence Harvester

Subsistence harvesters would represent adults and one-year-old children who reside outside the RSA full time and come to either the LSA or Lloyd Lake occasionally to hunt, fish and gather Traditional Foods. This receptor reflects the observation that there are no documented permanent residences within the local area of the Project but based on Indigenous and Local Knowledge shared during JWG meetings and in IKTLU Studies, the area is known to be used for subsistence harvesting including fishing, hunting, trapping, and gathering. Clearwater River Dene Nation members identified Patterson Lake as a "main lake" and "good for everything" harvesting (TSD V.1: CRDN). Harvesting activities in the Patterson Lake area are also practiced by the BNDN and BRDN, and the BNDN indicated that "hunting and trapping are central to the subsistence lifeways of members of the BNDN" (TSD II: BNDN).

A subsistence harvester would be exposed to Project-related COPCs through inhalation of air and dermal (skin) contact and/or incidental ingestion of dust deposited to soil while in the LSA or Lloyd Lake. During their time in the LSA or Lloyd Lake, they would use local surface water for drinking, bathing, and swimming, and would come in contact with sediments during wading.

Subsistence harvesters would ingest locally or regionally sourced Traditional Foods that are fished, hunted, trapped, and gathered, and retain Traditional Foods collected from the same area for consumption throughout the year. The locally or regionally sourced Traditional Foods would comprise about 50% of their Traditional Foods diet. The portion of locally or regionally sourced Traditional Foods in the subsistence harvester overall annual diet would be consistent with an



ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT HUMAN HEALTH RISK ASSESSMENT

Indigenous Group's high consumer of Traditional Foods. In the HHRA, about 50% of the Traditional Foods in the overall diet of the subsistence harvesters would be sourced from either Patterson Lake South Arm or Beet Lake in the LSA or from Lloyd Lake, and the other 50% from a reference location. The Patterson Lake location is a near-field location within the LSA. The Beet Lake location is in the area of a cabin identified by the MN-S, and the Lloyd Lake location is consistent with areas identified by the MN-S.

5.1.1.3 Seasonal Resident/Lodge Operator

Seasonal residents would represent adult and one-year-old members of the public who live in the LSA or at Lloyd Lake for part of the year, such as a lodge operator or employee. The seasonal residents were assumed to reside at a lodge either on Patterson Lake or Lloyd Lake for 30% of the year (approximately 4 months) and away from the LSA and RSA for 70% of the year (approximately 8 months) during the Project phases. The Patterson Lake receptor location is a hypothetical near-field location within the LSA. The Lloyd Lake receptor location is consistent with the closest existing lodge facility to the Project with on-site operators. The Lloyd Lake Lodge, which is located on the western shore of Lloyd Lake, is a fly-in lodge that provides fishing and spring bear hunting trips for guests. A recreational visitor at Lloyd Lake would be conservatively represented by the seasonal resident who would have a longer residence time in the area.

The Forest Lake Outfitters Camp at Beet Lake and the Clearwater River Provincial Park were also evaluated as potential locations for seasonal residents. Forest Lake outfitters offer non-guided fishing camps for drive-in or fly-in customers with fishing access to six local lakes. Potable water is available for customers of the Forest Lake Outfitters Camp, but groceries must be brought in by the guests. The Clearwater River Provincial Park is a wilderness park located south of the RSA at the south end of Lloyd Lake with no services or facilities (Government of Saskatchewan, n.d.). The Patterson Lake and Lloyd Lake receptor locations would conservatively bound any seasonal resident of Forrest Lake Outfitters or Clearwater River Provincial Park, who would be expected to frequent those locations less often.

While at Patterson Lake or Lloyd Lake, seasonal residents were assumed to ingest Traditional Foods fished, hunted, and gathered local to the Patterson Lake or Lloyd Lake areas to an extent consistent with an Indigenous Group's average consumer of Traditional Foods. While at the lodge, water for drinking, bathing, and swimming was represented by water from either Patterson Lake or Lloyd Lake, and seasonal residents were assumed to come in contact with sediments of these lakes during the practice of activities such as wading. For the remainder of the time, when not at Patterson Lake or Lloyd Lake, the seasonal residents would reside at an unexposed location with constituents in the environment at background levels.

5.1.1.4 Future Permanent Resident

Indigenous Groups expressed concerns regarding the long-term potential health effects associated with mine waste materials stored at the proposed Project site following Closure (TSD II: BNDN; TSD IV: MN-S; TSD V.2: CRDN; BNDN-JWG 2019a,b; BNDN-JWG 2021a,b; BRDN-JWG 2021b; CRDN-JWG 2020b; MN-S-JWG 2019a). To address these concerns, a future permanent resident was considered for the HHRA. The permanent resident group would represent potential risks to a hypothetical family (adult and one-year-old) that would reside at the decommissioned and reclaimed Project site in the far-future projection following Closure and would be exposed to Project-related COPCs at that time. It is not anticipated that there would be a permanent resident at a former mine site; therefore, this receptor is considered hypothetical, but is assessed as a conservative assumption. The future permanent residents were assumed to reside at the former Project for 100% of the year (i.e., in the Patterson Lake North Arm area).

Similar to the subsistence harvester, the future permanent resident would consume a large proportion of locally sourced Traditional Foods in their overall annual diet, consistent with an Indigenous Group's high consumer of Traditional Foods. For the HHRA, Patterson Lake was assumed to be a preferred fishing location due to its size and proximity to the former Project location. Water for drinking, bathing, and swimming was assumed to come from Patterson Lake. Contact with sediments in Patterson Lake during activities such as wading was also assumed. Since there would be no Project-associated air emissions in the far-future projection, atmospheric exposure pathways were considered to be incomplete for the permanent resident.




Figure 5-1: Human Receptor Locations

5.1.2 Constituents of Potential Concern Retained for Further Evaluation in the Human Health Risk Assessment

The selection of chemical stressors to be evaluated in the ERA followed a tiered approach to reduce the risk of overlooking Project-related COPCs to human health that would be emitted through water and air. The selection was detailed in Section 4.0, Model Integration of Evaluation of Sources.

The final list of COPCs for assessment in the ERA is presented in Section 4.4, Final List of Constituents of Potential Concern for the Environmental Risk Assessment. While chloride and sulphate were identified as COPCs for further assessment in the ERA, they were not considered further in the HHRA. These COPCs are associated with water ingestion. Chloride does not have a drinking water standard and is not considered to present a risk to human health at concentrations found in drinking water or at concentrations predicted for Patterson Lake, even under the reasonable upper bound scenario. Sulphate in drinking water is associated with adverse physiological effects such as diarrhoea or dehydration at concentrations above 500 mg/L. The predicted upper bound concentration of sulphate at the edge of the treated effluent mixing zone would be below 500 mg/L (Table 4-2); and therefore, concentrations at exposure points farther downstream would be less than those associated with adverse physiological effects. For these reasons, chloride and sulphate were not assessed further in the HHRA.

5.1.3 Selection of Exposure Pathways

The HHRA considered potential releases of Project-related constituents to the atmosphere and to surface water. Therefore, the relevant primary media were air, water, soil, sediment, and food.

Complete exposure pathways consisted of a contaminant source, a release mechanism, a transport mechanism within the relevant environmental medium or media, a point of exposure, and an exposure route. No pathways have been removed due to controls, mitigation, or treatment.

5.1.3.1 Summary of Complete Exposure Pathways

Radiological and non-radiological exposure pathways were assessed in the HHRA. The primary exposure routes for human health included:

- ingestion of food such as fish, vegetation, game, and store-bought foods;
- incidental ingestion of soil (i.e., while harvesting) or sediment (i.e., while wading);
- ingestion of surface water as drinking water;
- dermal (i.e., skin) contact with surface water and sediment while swimming or doing other recreational activities;
- dermal contact with soil while gardening or harvesting;
- inhalation of air (i.e., vapours and/or particulates); and



external exposure to radiation from air, water, soil, and sediment.

The potential exposure pathways were expected to be the same for all human receptors assessed, with the exception of inhalation in the far-future projection when there is no ongoing source of air COPCs. Exposure pathways for selected human receptors are summarized in Table 5-2.

Table 5-2: Human Health Exposure Pathways Considered for the Rook I Project

	Environmental Exposure Pathway									
Human Receptor Group	Air	Soil	Water	Sediment	Traditional Foods ^(a)					
Non-radiological E	xposures									
Camp worker										
Subsistence harvester	Inhalation	Incidental ingestion	Ingestion	Incidental ingestion	Ingestion					
Seasonal resident		Dermal contact	Dermal contact	Dermal contact						
Permanent resident	Incomplete ^(b)									
Radiological Expos	ures									
Camp worker										
Subsistence harvester	Inhalation External	Incidental ingestion	Ingestion	Incidental ingestion	Ingestion					
Seasonal resident	EXCELLE	External	External	External						
Permanent resident	Incomplete ^(b)									

⁽a) Plants and animals that are ingested as a source of food and have been directly or indirectly exposed to Project-related COPCs. (b) Pathway is identified as incomplete as there is no ongoing source of air COPCs during the far-future projection when a future permanent resident may reside at or near the reclaimed Project.

5.1.3.2 Dietary Assumptions

The purpose of this subsection is to provide the rationale used to derive the Traditional Foods diet for the Project HHRA and how the Traditional Foods diet fits into the overall total diet for human receptors. Traditional Foods are those animals and plants that are fished, hunted, or gathered from the land and consumed as food (Health Canada 2018). The remainder of the total food diet is assumed to be from store bought foods (store foods). Store foods include grain and other agricultural products of a typical diet that are not available locally.

Each of the human health receptors described in Section 5.1.1, Human Receptor Selection and Characterization, would consume locally sourced Traditional Foods from the LSA or RSA to an extent that is consistent with either an average or a high First Nations consumer of Traditional Foods. The basis for deriving appropriate average and high Traditional Foods diets, and how the engagement process informed the derivations are presented in the following sections (Section 5.1.3.2.1, Total Food Diet General Assumptions; Section 5.1.3.2.2, Traditional Foods Diet; Section 5.1.3.2.3, Store Foods).



COPC = constituent of potential concern.

5.1.3.2.1 Total Food Diet General Assumptions

The initial assumptions for ingestion rates and components of the total foods diet for the HHRA were taken from CSA N288.1-20 Adult (Table G.9b – Central) total diet. Human receptor groups in the HHRA (Section 5.1.1) were assumed to obtain a portion of their overall diet from Traditional Foods and the rest from store bought foods.

As discussed further in Section 5.1.3.2.2, human receptors were assumed to consume lower or higher proportions of Traditional Foods in their overall diets depending on their lifestyles. These are referred to as average Traditional Foods consumers and high Traditional Foods consumers, respectively. In both cases, the proportion of Traditional Foods in the overall diet of human health receptor groups in the HHRA was based on First Nations dietary studies. The camp worker, subsistence harvester and permanent resident receptor groups represent residents of northern Saskatchewan and were assumed to have an overall diet consistent with a First Nations high consumer of Traditional Foods. The seasonal resident receptor group represents residents of Saskatchewan and was assumed to ingest Traditional Foods consistent with an average First Nations consumer of Traditional Foods. The type of traditional diet attributed to each of the human health receptor groups is listed in Table 5-3.

Table 5-3: Traditional Food Diets for Each Human Receptor Group

Human Receptor Group	Type of Traditional Diet
Camp worker	High consumer
Subsistence harvester	High consumer
Seasonal resident/lodge operator	Average consumer
Permanent resident	High consumer

The methods used to develop the Traditional Foods diet for average and high Traditional Foods consumers is described in Section 5.1.3.2.2. The store foods diet is described in Section 5.1.3.2.3.

5.1.3.2.2 Traditional Foods Diet

The initial assumptions for ingestion rates and components of the Traditional Foods diet for the HHRA used information from the First Nations Food, Nutrition and Environment Study (FNFNES) undertaken in Saskatchewan in 2015 and erratum (Chan et al. 2018, 2019). The average and 95th percentile of daily intake of Traditional Foods were used to characterize the traditional diet for an "average consumer" and "heavy consumer". The assumptions were further refined with input from Indigenous Groups and other communities primarily through JWG discussions in October 2019 and February 2020, and later in discussion with representatives from Saskatchewan Ministry of Environment, Saskatchewan Health Authority, and CNSC. As discussed in the next sections, the

ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT HUMAN HEALTH RISK ASSESSMENT

engagement activities informed changes to the initial assumptions regarding the prevalence of berries in the Traditional Foods diet, and representative fish, game, and birds to be included in the IMPACT environmental pathways model to support the Traditional Foods exposure assessment.

The starting point for the Traditional Foods diet was the FNFNES report for Saskatchewan. The FNFNES report considers First Nations diets by ecozone to reflect the types of food items available locally to consumers of Traditional Foods. The study also provides breakdowns for Traditional Food items for:

- adult male, adult female, and average adult male and female diets;
- diets for both average and high (95th percentile) consumers of Traditional Foods; and
- detailed and simplified diets for each ecozone.

Consistent with the CSA total food diet, the HHRA considers ingestion rates for adult males for both average and high Traditional Foods diets.

The Project is located near the boundary of the Boreal Shield and Boreal Plain ecozones, as shown in Figure 5-2; therefore, the food categories for Traditional Foods in the diets for each of the ecozones were initially considered, as well as a combined (averaged) diet for the two ecozones. Reliance on fish and game differed somewhat between First Nations in the Boreal Shield and Boreal Plain ecozones. The Boreal Shield diet relies on less game (38%) and more fish (50%), whereas the Boreal Plain diet relies on more game (59%) and less fish (26%). The averaged Boreal Shield / Boreal Plain diet results in a more even distribution of game (44%) and fish (43%) in the traditional diet. The three diets were presented to local First Nations communities for input.

When the communities were consulted through JWG meetings, they indicated that of the three diets presented (Figure 5-3), the averaged diet which shows approximately equal proportions for game and fish ingestion was the most representative. Therefore, further development of the Traditional Food diet considered these proportions for the four Traditional Foods categories (fish, game and game organs, birds, and berries and plants) for both average and high consumers of Traditional Foods. The community feedback indicated that locally harvested berries were more prevalent in the regional diet than what was presented, so the intake rate for berries was increased to be equivalent to 10% of the total ingestion rate of berries from all sources to accommodate that concern. During JWG meetings, the MN-S (MN-S-JWG 2019a) indicated that they are more blueberries than what was proposed since they pick and store them to eat all winter.

The increase in the Traditional Foods berry ingestion rate was balanced by a decrease in the ingestion rate for store bought foods. This resulted in an increase in the overall Traditional Foods ingestion rate when compared to FNFNES. This is reflected in the final ingestion rates developed for the ERA in Section 5.2.3.1, Annual Food Ingestion Rates for Human Receptors in the Human Health Risk Assessment.



The FNFNES report and erratum lists over 100 different Traditional Food items as being harvested across Saskatchewan. The daily intakes (average and 95th percentile) of Traditional Food items were summarized for the following food categories for each ecozone: fish; game (meat and organs); birds; and berries/plants.

Engagement through the JWG sessions confirmed general agreement with the four Traditional Foods categories (fish, game, birds, berries and plants) and their proportions in the diet (BNDN-JWG 2019a,b; BRDN-JWG. 2019b; MN-S-JWG 2019a,b). For this diet, the bulk of the Traditional Foods diet was made up of fish and game meat in almost equal proportions. Birds and plants would contribute to the Traditional Foods diet to a lesser extent, but in almost equal proportions.

Using the detailed Traditional Food data provided in the FNFNES, the food categories were expanded for the HHRA to include:

- fish:
 - o predator species; and
 - o forage species.
- game and game organs:
 - large mammals;
 - o large mammal organs; and
 - o small mammals.
- birds:
 - o aquatic birds; and
 - terrestrial birds.
- berries and plants:
 - berries; and
 - Labrador tea.

The annual Traditional Foods ingestion rates for the Boreal Shield Traditional Foods diet were selected as the basis for the HHRA Traditional Foods diet because they are higher than those estimated for Traditional Foods consumers in the Boreal Plain Ecozone. The ingestion rates for each of the Traditional Foods food categories were adjusted to maintain the distribution that the communities recognized as being more representative of their Traditional Foods diets. The ingestion rates for the various Traditional Food items within each Traditional Foods category were proportioned using the FNFNES dietary data for the Boreal Shield total diet. As a final step, the



ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT

HUMAN HEALTH RISK ASSESSMENT

amount of Traditional Foods – berries ingested was increased from approximately 5% of the fruits and berry component of the total N288.1-20 diet to 10% by reducing the store food percentage. Figure 5-4 illustrates the proportional distribution of food items in the Traditional Foods diet for average and high Traditional Foods consumers selected for the HHRA.








Figure 5-2: Location of Rook I Site with Respect to Saskatchewan Ecozones



Note: Data from Chan et al, 2018 and 2019 for average traditional diets

Figure 5-3: Proportions of Traditional Food Categories in the Boreal Shield, Boreal Plain, and Averaged Diets Reviewed during Engagement Activities with Local Communities



Note: Male average and high Traditional Foods Boreal Shield diets were initially derived using data from Chan et al. 2018, 2019. Proportions of fish species and berries in the Traditional Foods diets were adjusted based on engagement activities.

Figure 5-4: Proportional Distribution of Traditional Food Items in Average and High Traditional Diets Selected for the Human Health Risk Assessment

5.13.2.3 Store Foods

The proportion of the overall diet not accounted for by Traditional Foods was considered to be from store-bought foods. Store foods are not from the area of influence of the Project. Ingestion of COPCs in store foods would be considered part of the background dose for human health receptors.

The concentrations of constituents in the store-food diet of human receptors were based on data from Health Canada (2000, 2011a) for non-radiological constituents in foods for Canadian cities (Health Canada 2011a) and United Nations Scientific Committee on the Effects of Atomic Radiation (UNSCEAR 2000) for radiological constituents. The data for different food types were averaged to represent the HHRA food components based on the proportions of different types in the store foods of the Hatchet Lake Band (CanNorth 2000).

5.1.4 Human Health Conceptual Site Model

The human health conceptual model illustrates how receptors are exposed to COPCs. It represents the relationship between the source and receptors by identifying the source of contaminants, the receptors, and the exposure pathways to be considered in the assessment for each receptor. Exposure pathways represent the various routes by which radionuclides and/or chemicals may enter the body of the receptor, or for radionuclides, how they may exert effects from outside the body.

The complete exposure pathways for the human receptors that are considered in the HHRA are illustrated in Figure 5-5, based on the selected human receptor groups (Section 5.1.1, Human Receptor Selection and Characterization) and the potential exposure pathways (Section 5.1.3, Selection of Exposure Pathways).

As outlined in Section 4.4 and Section 5.1.2, the list of COPCs relevant to the water exposure pathway include various metals and metalloids (i.e., arsenic, cobalt, copper, molybdenum, and uranium), and select radionuclides of the uranium-238 decay chain (i.e., uranium-238, uranium-234, thorium-230, radium-226, lead-210, and polonium-210). These waterborne COPCs are also expected to partition from surface water to sediment. Human and ecological receptors may be directly exposed to these COPCs via ingestion and/or dermal contact with surface water and sediment. Human receptors would be potentially further exposed through the ingestion of aquatic and terrestrial country foods that ingest or come into contact with contaminated surface water and sediment.

The list of COPCs relevant to the air exposure pathway include the same list of radionuclides as the water exposure pathway with the addition of radon-222. Human and terrestrial ecological receptors may come into direct contact with these radionuclides via the inhalation exposure pathway. Radionuclide particles may also deposit onto soil and plants, where they may be taken up into the vascular system of plants or be incidentally ingested or come into dermal contact by humans and terrestrial biota. Human receptors would again be potentially further exposed via the consumption of terrestrial country foods that come into contact with radionuclides via inhalation or through ingestion/dermal contact with contaminated soil or plants.




Figure 5-5: Human Health Conceptual Site Model

5.1.5 Uncertainty in Problem Formulation

The assumptions used to characterize human health receptors and develop the conceptual site model followed best industry practices and CSA guidance. Where possible, region-specific information was used to develop initial assumptions for human health receptor groups, locations and frequency, and duration of exposures. Communities and regulators were engaged in the process, which resulted in adjustments to the initial assumptions to better represent affected communities and increase conservatism in areas (such as the Traditional Foods diet) where regional information was scarce.

The selection of Project-related COPCs was based on comparing maximum predicted water, sediment, air, and soil concentrations at the human health receptor locations to relevant environmental guideline concentrations (i.e., screening values). Numerous conservative measures were integrated into the models used to predict COPC concentrations used in the screening process (EIS Appendix 7A, Air Dispersion Modelling Report; EIS Appendix 9A, Hydrological Modelling Summary Report; TSD XIV, Groundwater Flow and Solute Transport Modelling Report). The predicted concentrations were compared to screening values protective of both human health

and biota. There is, therefore, a high level of confidence that the HHRA captures all Project-related COPCs that would be emitted by Project activities to water and air.

5.2 Exposure Assessment

The exposure assessment included identification of exposure locations and exposure factors for each human health receptor and presentation of exposure concentrations and doses (radiological and non-radiological). Uncertainties are discussed. This subsection presents the information used in the environmental pathways model, IMPACT, at a high level. The details of the model are included in Appendix A.

5.2.1 Exposure Locations, Duration, and Frequency

The selection of exposure locations for human health receptor groups was based on the current understanding of how people use the LSA and RSA, including information from IKTLU Studies, which was supplemented by information obtained during community information sessions and JWG meetings, as well as the potential for exposure to Project-related media concentrations during one or more Project phases. Exposure locations for each of the human health receptors are shown in Figure 5-1. The residency assumptions for each of the human health receptors are summarized in Table 5-4, which includes the exposure frequency (months per year) and fraction of time the receptor would spend at a given location. The exposure duration, which refers to the number of years of exposure, was either the duration of the Project phases (43 years) or the lifetime of the receptor.

The selected human health receptor exposure locations include:

- Patterson Lake North Arm at the Project's camp site for:
 - camp worker (adult) Project lifespan; and
 - o future permanent resident (adult and one-year-old) far-future projection only.
- Patterson Lake South Arm for:
 - o seasonal residents/lodge operators (adult and one-year-old) Project lifespan; and
 - o subsistence harvesters (adult and one-year-old) Project lifespan.
- Beet Lake for:
 - o subsistence harvesters (adult and one-year-old) Project lifespan.
- Lloyd Lake for:
 - subsistence harvesters (adult and one-year-old) Project lifespan; and
 - seasonal residents/lodge operators (adult and one-year-old) Project lifespan.



With the exception of the future permanent resident, all of the human health receptors were assumed to spend part of their time away from the LSA and RSA at a location represented in the ERA by the reference location. The reference location, represented by Broach Lake, is intended to represent a location unaffected by the proposed Project, and is characterized by the compilation of baseline data in the RSA.

Table 5-4: Summary of Residency Assumptions for Human Health Receptor Groups

Human Health Life Receptor Stage(s) Group			Resido			
		Receptor Location(s)	Fraction of Time at Location	Exposure Frequency (mo/yr)	Consumer of Traditional Foods	
All Project F	Phases: Co	nstruction, Operatior	ns, Closure			
Camp worker	Adult	Camp / Patterson Lake South Arm / reference location	0.5 / 0.25 / 0.25	6/3/3	Yes, high consumer while away from the camp	
Adult	Patterson Lake South Arm / reference location	0.5 / 0.5	6/6	Yes, high consumer		
Subsistence harvester	and one- year-old	Beet Lake / reference location	0.5 / 0.5	6/6	year-round in study area and reference location	
		Lloyd Lake / reference location	0.5 / 0.5	6/6	locution	
l resident	Adult and one-	Patterson Lake South Arm / reference location	0.3 / 0.7	4/8	Yes, average consumer year-round in study area and reference	
	year-old	Lloyd Lake / reference location	0.3 / 0.7	4/8	location	
Far-future F	Projection					
Permanent resident	Adult and one- year-old	Patterson Lake North Arm – West Basin	1	12	Yes, high consumer year-round	
	Adult	Patterson Lake South Arm / reference location	0.5 / 0.5	6/6	Yes, high consumer	
Subsistence and	and one- year-old	and one-	Beet Lake / reference location	0.5 / 0.5	6/6	year-round in study area and reference location
		Lloyd Lake / reference location	0.5 / 0.5	6/6	To Cation	
Seasonal Resident	Adult and one-	Patterson Lake South Arm / reference location	0.3 / 0.7	4/8	Yes, average consumer year-round in study area and reference	
	year-old	Lloyd Lake / reference Location	0.3 / 0.7	4/8	location	

⁽a) The exposure duration for subsistence harvesters, seasonal residents, and permanent residents is considered to be over a lifetime; the exposure duration for the camp worker is considered to be the duration of the Project (43 years).



5.2.2 Exposure and Dose Calculations

Exposure and dose calculations for human receptors were completed using IMPACT version 5.6.0. IMPACT is consistent with the COPC transport equations and radiological dose calculations outlined in CSA N288.1-20. Equations used for non-radiological dose calculations are consistent with CSA N288.6-22, which have generally been obtained from Health Canada guidance.

The inputs and assumptions used in the IMPACT model for the Rook I Project, including receptor characteristics, exposure pathways, and the derivation and identification of site-specific information used in the model, are provided in Appendix A. Relevant to the exposure and dose calculations for human receptors, the Model Report:

- describes the model structure for human health receptor assessment, specific assumptions
 made for the Project, and the generic equations used to calculate the transfer of
 constituents between environmental media and the receptors; and
- presents the input parameters for adult and one-year-old receptors, including those for dietary assumptions and intake rates, inhalation rates, and exposed skin area.

5.2.3 Exposure Factors

Exposure estimates rely on several COPC- and media-specific exposure factors for the dose calculations. These parameters include body characteristics and intake rates as well as exposure frequency (Section 5.2.1, Exposure Locations, Durations, and Frequency) and dose coefficients.

5.2.3.1 Annual Food Ingestion Rates for Human Receptors in the Human Health Risk Assessment

Adult Diet

As mentioned in Section 5.1.3.2, Dietary Assumptions, the initial assumptions for food ingestion rates and components of the total diet for the HHRA were taken from CSA N288.1-20 Adult (Table G.9b – Central) total diet. The CSA N288.1-20 dietary composition was based on the 2004 Canadian nutrition survey results (Statistics Canada 2004), processed for International Commission on Radiological Protection (ICRP) age groups and for two sexes in the adult age group. The dietary intakes for ICRP ages were adjusted to align with ICRP reference energy intakes for each age group (ICRP 2003).

The N288.1-20 human diet was selected over the Health Canada (2010a) human diet for the HHRA. Health Canada (2010a) references Richardson (Richardson 1997), which used survey results from the late 1970s. The Richardson diet was also a combined adult male/female diet. The CSA N288.1-20 total adult diet is a smaller overall diet (706 kg/yr) than the Richardson diet (808 kg/yr; by about 100 kg/yr) and is based on more recent data.



ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT HUMAN HEALTH RISK ASSESSMENT

The total food diet is the sum of Traditional Foods and store foods. Its dietary components, expressed in kilograms per year of food consumed, are shown in Table 5-5 for non-consumers, average consumers, and high consumers of Traditional Foods.

The derivation of the Traditional Foods diet is described in Section 5.1.3.2. The overall ingestion rate for the Traditional Foods diet used in the HHRA is higher that that estimated by the FNFNES due to a higher ingestion rate for Traditional Foods – berries being considered in response to input from Indigenous and local communities. The concentrations of COPCs in Traditional Food items used in the exposure assessment are estimated using the IMPACT model, as described in Appendix A, Section 3.6, Transfer of Constituents to Aquatic Receptors, and Section 3.7, Transfer of Constituents to Terrestrial Receptors and Humans. The selection of representative ecological receptors used in the IMPACT model is discussed below.

The store food intakes in each food category were estimated from the total diet intakes provided in CSA N288.1-20, minus the Traditional Food intake. If the Traditional Foods intakes were higher than the total intake for the same CSA N288.1-20 food category, then the store food intake was set to zero and the values for other store food categories were reduced as needed to obtain total dietary intakes equal to those in CSA N288.1-20. The store food diet and concentrations of COPCs in store foods used in the exposure assessment are consistent with other recent HHRAs conducted for uranium mine and mill projects in Saskatchewan. Concentrations of COPCs in store foods are provided in Appendix A (Table 2-11).

One-Year-Old Diet

Annual food consumption rates for the one-year-old diet were calculated using adult-to-one-year-old ratios from CSA N288.1-20 for each of the selected food categories. Annual ingestion rates for individual food categories were combined into relevant food categories, and the adult-to-one-year-old ratios were determined for each of these food categories.

Representative Ecological Receptors Used in the IMPACT Model

The representative ecological receptor is the specific species modelled in the ERA that represents each Traditional Food category. Based on the descriptions of the different human receptor groups and their Traditional Food diets, representative ecological receptors were selected for the IMPACT model to represent each Traditional Food category. The representative ecological receptor selected to represent the Traditional Food category was informed and modified based on feedback received from JWG meetings and engagement with regulators. Specifically, these engagement activities informed assumptions for fish species, large mammals, and small mammals.

- Northern pike and lake whitefish were included as fish species in the exposure assessment and part of the Traditional Foods diet.
- Preference for eating moose over other large mammals such as caribou.



Engagement also confirmed that the food types in the FNFNES were relevant to the Indigenous Groups, with some adjustments. Based on Indigenous Group feedback, the following adjustments were made:

- Beaver (*Castor canadensis*) was identified as a VC for the EIS and was included as part of HHRA Traditional Foods diet and as an ecological receptor, replacing muskrat (*Ondata zibethicus*).
- Spruce grouse (*Falcipennis canadensis*) replaced goose as a modelled Traditional Food item for the HHRA exposure assessment.

Once the IKTLU Studies were available, information generally confirmed selection of receptors that may be of interest to Indigenous Groups as part of the diet:

- Terrestrial vegetation (e.g., lichen, blueberry [Vaccinium myrtilloides], Labrador tea [Rhododendron groenlandicum]):
 - The CRDN identified the use of berries (i.e., blueberries, cranberries [bog (*Vaccinium vitis-idaea*), low bush (*Viburnum edule*)], gooseberries [spp.], Saskatoon (*Amelanchier alnifolia* var. *alnifolia*), cloudberry (*Rubus chamaemorus*), and strawberries [spp.]), medicines (i.e., kinnikinnick [*Arctostaphylos uva-ursi*], Labrador tea, mint [*Mentha canadensis*], spruce gum, sweet flag [rat root (*Acorus americanus*)], and mushrooms), shrubs (i.e., dogwood [*Cornus sericea*, red willow] and willows [spp.]), trees (i.e., birch [spp.], jack pine [*Pinus banksiana*], poplar [spp.], and spruce [spp.], tamarack [*Larix laricina*]), and other vegetal matter (i.e., barks, mosses, roots, and punk [rotten wood]; TSD V.1: CRDN).
 - The MN-S identified the use of berries, birch, blueberry, cranberry, mint, raspberry rat root, strawberry, and sweetgrass (*Fragaria vesca* ssp. *Americana*) (TSD IV: MN-S).
 - The BNDN identified the use of berries (spp.), birch, blueberry, bulrushes (Vaccinium myrtilloides, Juncus alpinoarticulatus), cranberry, mint, raspberry, and strawberry (TSD II: BNDN).
 - The BRDN identified the use of berries (TSD III: BRDN).
 - The Athabasca Denesųłiné identified specific plants harvested including Ts'aili Teli (Frog Tail [Sarracenia purpurea ssp. gibbosa]), Labrador tea, and blueberries, as well as berries in general (TSD VI: YNLR).
- Fish (e.g., northern pike, lake whitefish):
 - The CRDN identified use of grayling (Arctic grayling [Thymallus arcticus]), jackfish (northern pike [Esox lucius]), herring (cisco; Coregonus artedi), lake trout (Salvelinus



- namaycush), ling cod (mariah; burbot [Lota lota]), pickerel (walleye [Sander vitreus]), and suckers (spp.; TSD V.1: CRDN; TSD V.2: CRDN).
- The MN-S identified trout, whitefish, jack (jackfish, or northern pike), pickerel, suckers, and catfish² as being consumed (TSD IV: MN-S).
- The BNDN identified fishing for lake trout, whitefish (lake whitefish [Coregonus clupeaformis]), jackfish, pickerel (walleye), suckers, and mariah (burbot [Lota lota]) (TSD II: BNDN).
- The BRDN described fishing for lake whitefish, lake trout, jackfish, and pickerel (TSD III: BRDN).
- Athabasca Denesųłiné identified lake trout, northern pike, suckers, pickerel, and whitefish as important species (TSD VI: YNLR).
- Woodland caribou (Rangifer tarandus caribou):
 - The CRDN, MN-S, BNDN, BRDN, and Athabasca Denesųłiné identified woodland caribou as a culturally important species that was hunted traditionally (TSD II: BNDN; TSD III: BRDN; TSD IV: MN-S; TSD V.1: CRDN; TSD V.2: CRDN; TSD VI: YNLR). Woodland caribou was reported to still be occasionally harvested by members of the BNDN and BRDN depending on their availability (TSD II: BNDN; TSD III: BRDN).
- Beaver (Castor canadensis):
 - The CRDN, MN-S, BNDN, BRDN, and Ya'thi Néné Lands and Resources identified that they trap beaver, which is also consumed by some members (TSD II: BNDN; TSD III: BRDN; TSD IV: MN-S; TSD V.1: CRDN; TSD V.2: CRDN; TSD VI: YNLR).
- Moose (Alces alces):
 - The CRDN, MN-S, BNDN, BRDN, and Athabasca Denesyliné identified moose as a culturally important species that is harvested (TSD II: BNDN; TSD III: BRDN; TSD IV: MN-S; TSD V.1: CRDN; TSD V.2: CRDN; TSD VI: YNLR).
- Spruce grouse:

² Biological species identification is uncertain. Catfish are not known to occur in the area of the Project; however, the comment may have been referring to burbot, which have a similar appearance to catfish because of the barbel (feeler) on the chin.



ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT

HUMAN HEALTH RISK ASSESSMENT

 The CRDN, MN-S, BNDN, and Athabasca Denesyliné identified spruce grouse as hunted (TSD II: BNDN; TRSD IV: MN-S; TSD V.1: CRDN; TSD V.2: CRDN; TSD VI: YNLR).

- Mallard (Anas platyrhynchos):
 - The CRDN, MN-S, BRDN, and Athabasca Denesyliné identifed ducks as important species that are hunted (TSD III: BRDN; TSD IV: MN-S; TSD V.1: CRDN; TSD V.2: CRDN).

The final selection of representative ecological receptors for the Traditional Foods diet is shown in Table 5-6. Additional discussions on how ecological receptors were selected, their characteristics and exposure assumptions are provided in the EcoRA in Section 6.0, Ecological Risk Assessment, and Appendix B, Ecological Receptor Profiles.



Table 5-5: Annual Ingestion Rates for Traditional Food Items of the Total Food Diet for Non-consumers, and Average and High Consumers of Traditional Foods

Food Category	Non- Traditional Foods Consumer Diet ^(a)		Average Tradi Consume			High Traditional Foods Consumer ^(b, c, d, e, f)				
	Adult	Ad	lult	One-Ye	One-Year-Old		Adult		One-Year-Old	
	Total Diet	Traditional Foods	Store Foods	Traditional Foods	Store Foods	Traditional Foods	Store Foods	Traditional Foods	Store Foods	
	kg/yr	kg	kg/yr		kg/yr		kg/yr		kg/yr	
Milk	126	0	126	0	243	0	32.6	0	226	
Fish	8.25	26.8	0	5.94	0	84.8	0	18.76	0	
Northern pike	n/a	13.4	0	2.97	0	42.4	0	9.38	0	
Lake whitefish	n/a	13.4	0	2.97	0	42.4	0	9.38	0	
Meat - large mammals	67.9	6.52	24.1	0.785	2.90	20.6	0	2.48	0	
Meat - large mammal organs	2.04	3.59	0	0.897	0	9.16	0	2.29	0	
Meat - small mammals	0.230	17.4	0	1.66	0	56.9	0	5.42	0	
Meat - birds and eggs	58.1	n/a	53.1	n/a	9.48	n/a	42.3	n/a	7.55	
Aquatic bird	n/a	1.71	0	0.304	0	3.46	0	0.618	0	
Terrestrial bird	n/a	3.29	0	0.587	0	12.3	0	2.19	0	



ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT HUMAN HEALTH RISK ASSESSMENT

Food Category	Non- Traditional Foods Consumer Diet ^(a)		Average Tradi Consumer				High Tradit Consume		
	Adult	Ac	lult	One-Ye	ar-Old	Ad	ult	One-Yo	ear-Old
	Total Diet	Traditional Foods	Store Foods	Traditional Foods	Store Foods	Traditional Foods	Store Foods	Traditional Foods	Store Foods
Root vegetables	47.9	0	47.9	0	8.65	0	47.9	0	8.65
Other vegetables	297	0.844	296	0.225	82.3	2.45	294	0.681	81.9
Fruits and berries	99.4	9.94	89.5	5.46	49.2	9.94	89.5	5.46	49.2
Total diet – subtotal	n/a	70.1	636	15.9	395	200	506	37.9	373
Total diet	706	7	06	41	1	70	06	4	11

Note: Values greater than 0 are rounded to 3-significant figures.

- (a) Total Diet is derived from CSA N288.1-20 central dietary intakes Table G.9b.
- (b) Traditional Foods annual consumption rates for adults are taken from First Nations Food, Nutrition and Environment Study (Chan et al. 2018) Boreal Shield diet for Saskatchewan First Nations Communities for either average or high Traditional Food consumers (male) for the Traditional Foods diet.
- (c) Annual food consumption rates for one-year-old were calculated using 'Adult to 1-year-old' ratios (CSA N288.1-20 Adult (Central): 1-year old) derived for each food category. Annual consumption rates for individual food items were combined into relevant food category and the ratios for each food category were derived based on the total kg/yr for each category.
- (d) The store food intakes in each food category were estimated from the non-Traditional Foods intakes provided in CSA N288.1-20, minus the Traditional Foods intakes. If the Traditional Foods intakes were higher than the total intake for the food category, the store food intake was set to zero, and then the value for other store food categories were reduced as needed to obtain total dietary intakes equal to those in CSA N288.1-20.
- (e) The fractions of northern pike (predator species) and lake whitefish (forage species) were set at 50% each of the total fish diet based on input received from regulatory bodies to include both predator and forage species for a conservative assessment of exposure to constituents.
- (f) The fraction of Traditional Foods "Fruits and Berries" was increased from approximately 5% (Chan et al. 2018) to 10% of the total "fruits and berries" based on input received from local Indigenous communities. The total food diet was adjusted by reducing the store foods and increasing Traditional Foods "Fruits and Berries" fractions, respectively. n/a = value not available or not applicable.



Table 5-6: Representative Ecological Receptors used in the Traditional Foods Diet

Traditional Food Category	Representative Ecological Receptor	Traditional Food Items	Rationale ^(a)
Fish			
	Northern pike	Walleye Northern pike	 Predator species and forage species represent two ecological trophic levels Northern pike and lake whitefish are present in all lakes in the vicinity of the
Fish	Lake whitefish	Lake whitefish	 Project according to baseline studies Important Traditional Food items for Boreal Shield and Boreal Plains ecozones Both are known to be fished in the vicinity of the Project
Game Meat/Organs	3		
	Moose meat	Moose meat	
Large mammal	Moose organs	Moose kidney Moose liver Caribou kidney	 Large herbivore with linkages to the aquatic environment Generally present in the vicinity of the Project according to baseline studies Important Traditional Food items for Boreal Shield and Boreal Plains ecozones
Small mammal	Beaver	Beaver meat Rabbit meat	 Small herbivore with linkages to the aquatic environment Generally present in the vicinity of the Project according to baseline studies Important Traditional Food items for Boreal Shield ecozone
Birds			
Aquatic bird	Mallard	Mallard	 Omnivorous bird with linkages to the aquatic environment Seasonally present in the vicinity of the Project according to baseline studies Important Traditional Food items for Boreal Shield and Boreal Plains ecozones
Terrestrial bird	Grouse	Grouse Goose	 Upland herbivore bird Generally present in the vicinity of the Project according to baseline studies Important Traditional Food items for Boreal Shield and Boreal Plains ecozones
Berries/Plants			
Berries	Blueberry	Blueberry	Consults and the district of the During to a souling to be all the
Plants	Labrador tea	Mint Rat root	 Generally present in the vicinity of the Project according to baseline studies Important Traditional Food items for Boreal Shield and Boreal Plains ecozones

⁽a) The selection of representative ecological receptors to represent Traditional Food items was informed by feedback received during engagement activities.



5.2.3.2 Bioavailability

In general, the IMPACT model was used to calculate exposure doses for arsenic. However, the general assumption in IMPACT is that a COPC is 100% bioavailable from all media types, including food.

In the initial outputs from the IMPACT model, the contribution of moose organs and moose meat to total exposure assuming 100% bioavailability was relatively high and a key driver for the risk estimates, particularly the subsistence harvester. Arsenic may be present in the environment in different chemical forms such as arsenopyrite and arsenic trioxide. Some forms of arsenic can be absorbed in the gastrointestinal tract and taken up by plants, while other forms are poorly absorbed.

To account for that uncertainty, and provide a more realistic interpretation of results, the model outputs were amended to incorporate arsenic bioaccessibility into the outputs for moose meat and moose organs using data collected as part of a study based out of British Columbia (Laird and Chan 2013). Laird and Chan (2013) collected samples of various types of traditional foods considered in the FNFNES including moose organs (kidney and liver) and moose meat. The mean percent in vitro bioaccessibility (a surrogate for bioavailability) ranged from 7% to 19% for moose organs, and was 59% for moose meat. These bioaccessibilities were incorporated into the exposure assessment by adjusting the model outputs from IMPACT by 0.19 for moose organs and 0.59 for moose meat.

ATSDR (2007) indicates that in seafood (i.e., fish), approximately 10% of arsenic is present in the inorganic form, while the remainder is in an organic form (arsenobetaine) that is generally not associated with toxicity. Therefore, the estimates of exposure generated by IMPACT overestimate arsenic exposure from the fish consumption pathway. As such, the model outputs from IMPACT were adjusted such that the exposure doses from fish were multiplied by a factor of 0.10.

These are reasonable estimates of bioaccessibility and are expected to reduce the uncertainty associated with the risk estimates for those food types. However, 100% bioaccessibility was assumed for the remaining food types (i.e., terrestrial plants, beaver, mallard, and grouse), and as such, exposure and risks for those other food types may be overestimated.

5.2.3.3 Other Receptor Characteristics

Other receptor characteristics used in the exposure assessment were taken from CSA N288.1-20, and include:

- 1. air inhalation rates Table 19 in CSA N288.1-20;
- 2. water ingestion rates Table 21 in CSA N288.1-20;
- 3. soil and sediment ingestion rates Table 20 in CSA N288.1-20; and
- 4. exposed surface area (whole body) Table 22 in CSA N288.1-20.



Receptor characteristics used in the IMPACT model are provided in Appendix A, Section 2.4, Exposure of Human Receptors.

5.2.3.4 Dose Coefficients

Dose coefficients (DCFs) for all internal and external exposure routes for humans are used to estimate radiological exposure. The DCFs for ingestion and inhalation by human receptors were taken from CSA N288.1-20. The external DCFs used in this IMPACT model update were derived based on the methods described in N288.1-20. Dose coefficients used in the IMPACT model are provided in Appendix A, Section 3.7.6, Dose Coefficients for Terrestrial Animals, Birds, and Humans.

5.2.4 Exposure Point Concentration and Doses

Concentrations of COPCs in environmental media (water, sediment, air, soil, and Traditional Food items) at the receptor locations and reference location were derived using IMPACT. The concentrations of COPCs in store foods used in the exposure assessment are consistent with other recent HHRAs conducted for uranium mine and mill projects in Saskatchewan (Appendix A, Section 2.4.1, Exposure Assumptions for Human Receptors). The equations used are provided in Appendix A, Section 2.4.3, Radiological Dose and Section 2.4.4, Non-radiological Dose for radiological and non-radiological doses, respectively.

Assessment of radiation exposures to members of the public is commonly based on estimation of the incremental effects of the project or site. Assessments consider the radiation dose received from direct exposure to radiation as well as the dose received from ingestion of radionuclides. The radionuclide intake by human receptors from various pathways is converted into a dose that is presented in millisieverts per year (mSv/yr).

Assessment of non-radiological exposures to members of the public is commonly based on estimation of the total effects of the project or site. Assessments consider the dose received from ingestion of constituents of concern as well as dermal absorption due to contact with soil. This is presented as a dose in milligrams per kilogram per day (mg/kg/d) for each pathway.

Section 5.2.4, Exposure Point Concentration and Doses, presents the estimated non-radiological dose and radiological dose to human receptors due to releases from the Project during all phases of the Project, including the far-future projection once groundwater solutes have been released to Patterson Lake. The results are presented for both the Application Case as well as the reasonable upper bound sensitivity scenario (Section 5.2.4.1, Estimated Dose to Human Receptors – Application Case) and the RFD Case (Section 5.2.4.2, Estimated Dose to Human Receptors – Reasonably Foreseeable Development Case). The estimated non-radiological and radiological concentrations in environmental media and biota tissue concentrations relevant to human ingestion pathways are shown in Appendix C, Model Results in Support of the Environmental Risk Assessment.



5.2.4.1 Estimated Dose to Human Receptors – Application Case

This subsection presents the estimated doses to human receptors due to releases from the Project during all phases including the far-future projection once groundwater solutes have been released to Patterson Lake. The results are presented for both the Application Case as well as the Upper Bound sensitivity scenario. Doses are presented for non-carcinogens (cobalt, copper, molybdenum, uranium), carcinogens (arsenic), and radionuclides (uranium-238 decay series radionuclides). The doses represent the maximum dose over the Project lifespan for the COPCs of interest, which is a conservative representation as exposure varies over the different Project phases. As indicated in Section 5.1.2, Constituents of Potential Concern Retained for Further Evaluation in the Human Health Risk Assessment, chloride and sulphate were not considered COPCs for the HHRA since their predicted maximum concentrations at the edge of the mixing zone would not be considered to present a risk to human health.

5.2.4.1.1 Non-carcinogen Doses

The doses for cobalt, copper, molybdenum, and uranium to the camp worker, subsistence harvester, and seasonal resident, and future permanent resident are presented by exposure pathway for the Application Case and the reasonable upper bound sensitivity scenario during Operations and the far-future projection in Table 5-7.



Table 5-7: Estimated Non-carcinogen Doses to Human Receptors – Operations – Application Case and Upper Bound Scenario

				Dose by I	Pathway Project	Lifespan (mg/	kg/d)					Dose	by Pathway Far	Future (mg/kg	ı/d)		
	СОРС	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
						,		Ва	se Case								
	Cobalt	8.58 × 10 ⁻⁰⁶	9.04 × 10 ⁻⁰⁹	1.01 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	1.15 × 10 ⁻⁰⁵	7.21 × 10 ⁻⁰⁶	1.97×10^{-04}	2.25 x 10 ⁻⁰⁴	n/a-	n/a						
	Copper	1.24 × 10 ⁻⁰⁵	1.26 × 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	3.81 × 10 ⁻⁰⁴	1.15 × 10 ⁻⁰⁴	1.66×10^{-02}	1.71 x 10 ⁻⁰²	n/a							
	Molybdenum	3.04 × 10 ⁻⁰⁶	2.98 × 10 ⁻⁰⁹	4.54 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	3.39 × 10 ⁻⁰⁸	4.10 × 10 ⁻⁰⁶	2.30×10^{-03}	2.31 x 10 ⁻⁰³	n/a							
	Uranium	1.48 × 10 ⁻⁰⁶	5.17 × 10 ⁻⁰⁹	1.35 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	2.06 × 10 ⁻⁰⁶	1.15 × 10 ⁻⁰⁵	4.37×10^{-05}	5.88 x 10 ⁻⁰⁵	n/a							
						1			cation Case				I				
Camp Worker	Cobalt	8.63 × 10 ⁻⁰⁶	9.07 x 10 ⁻⁰⁹	1.02 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	1.25 x 10 ⁻⁰⁵	7.29 x 10 ⁻⁰⁶	2.01 x 10 ⁻⁰⁴	2.29 x 10 ⁻⁰⁴	n/a							
(Patterson Lake North Arm - West	Copper	1.24 × 10 ⁻⁰⁵	1.28 x 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	3.99 x 10 ⁻⁰⁴	1.15 x 10 ⁻⁰⁴	1.66 x 10 ⁻⁰²	1.72 x 10 ⁻⁰²	n/a							
Basin)	Molybdenum	3.04 × 10 ⁻⁰⁶	3.03 x 10 ⁻⁰⁹	4.55 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	5.00 x 10 ⁻⁰⁸	4.25 x 10 ⁻⁰⁶	2.31 x 10 ⁻⁰³	2.31 x 10 ⁻⁰³	n/a							
	Uranium	1.48 × 10 ⁻⁰⁶	1.66 x 10 ⁻⁰⁸	1.35 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.83 x 10 ⁻⁰⁶	2.44 x 10 ⁻⁰⁵	5.21 x 10 ⁻⁰⁵	8.08 x 10 ⁻⁰⁵	n/a							
						·		Upper B	ound Scenario								
	Cobalt	8.63 × 10 ⁻⁰⁶	9.07 x 10 ⁻⁰⁹	1.02 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	1.25 x 10 ⁻⁰⁵	7.29 x 10 ⁻⁰⁶	2.01 x 10 ⁻⁰⁴	2.29 x 10 ⁻⁰⁴	n/a							
	Copper	1.24 × 10 ⁻⁰⁵	1.28 x 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	3.99 x 10 ⁻⁰⁴	1.15 x 10 ⁻⁰⁴	1.66 x 10 ⁻⁰²	1.72 x 10 ⁻⁰²	n/a							
	Molybdenum	3.04 × 10 ⁻⁰⁶	3.03 x 10 ⁻⁰⁹	4.55 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	8.31 x 10 ⁻⁰⁸	4.25 x 10 ⁻⁰⁶	2.31 x 10 ⁻⁰³	2.32 x 10 ⁻⁰³	n/a							
	Uranium	1.48 × 10 ⁻⁰⁶	1.66 x 10 ⁻⁰⁸	1.35 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	4.18 x 10 ⁻⁰⁶	2.44 x 10 ⁻⁰⁵	5.24 x 10 ⁻⁰⁵	8.25 x 10 ⁻⁰⁵	n/a							
			_				ı		se Case								
	Cobalt	8.58 × 10 ⁻⁰⁶	9.04 × 10 ⁻⁰⁹	1.01 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	2.30 × 10 ⁻⁰⁵	1.44 × 10 ⁻⁰⁵	2.44 × 10 ⁻⁰⁴	2.90 x 10 ⁻⁰⁴	8.58 × 10 ⁻⁰⁶	9.04 × 10 ⁻⁰⁹	1.01 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	2.30 × 10 ⁻⁰⁵	1.44 × 10 ⁻⁰⁵	2.44 × 10 ⁻⁰⁴	2.90 x 10 ⁻⁰⁴
	Copper	1.24 × 10 ⁻⁰⁵	1.26 × 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	7.61 × 10 ⁻⁰⁴	2.29 × 10 ⁻⁰⁴	1.75 × 10 ⁻⁰²	1.86 x 10 ⁻⁰²	1.24 × 10 ⁻⁰⁵	1.26 × 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	7.61 × 10 ⁻⁰⁴	2.29 × 10 ⁻⁰⁴	1.75 × 10 ⁻⁰²	1.86 x 10 ⁻⁰²
	Molybdenum	3.04 × 10 ⁻⁰⁶	2.98 × 10 ⁻⁰⁹	4.54 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	6.77×10^{-08}	8.18 × 10 ⁻⁰⁶	1.95×10^{-03}	1.96 x 10 ⁻⁰³	3.04 × 10 ⁻⁰⁶	2.98 × 10 ⁻⁰⁹	4.54 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	6.77 × 10 ⁻⁰⁸	8.18 × 10 ⁻⁰⁶	1.95 × 10 ⁻⁰³	1.96 x 10 ⁻⁰³
	Uranium	1.48 × 10 ⁻⁰⁶	5.17 × 10 ⁻⁰⁹	1.35 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	4.11 × 10 ⁻⁰⁶	2.30×10^{-05}	4.92×10^{-05}	7.78 x 10 ⁻⁰⁵	1.48 × 10 ⁻⁰⁶	5.17 × 10 ⁻⁰⁹	1.35 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	4.11 × 10 ⁻⁰⁶	2.30×10^{-05}	4.92×10^{-05}	7.78 x 10 ⁻⁰⁵
			_					Appli	cation Case								
Subsistence	Cobalt	9.32 × 10 ⁻⁰⁶	9.05 x 10 ⁻⁰⁹	1.09 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	2.49 × 10 ⁻⁰⁵	1.46 x 10 ⁻⁰⁵	2.51 x 10 ⁻⁰⁴	3.00 x 10 ⁻⁰⁴	1.23 x 10 ⁻⁰⁵	9.05 x 10 ⁻⁰⁹	1.46 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	3.30 x 10 ⁻⁰⁵	1.44 x 10 ⁻⁰⁵	2.73 x 10 ⁻⁰⁴	3.33 x 10 ⁻⁰⁴
Harvester Adult (Patterson Lake	Copper	1.30 × 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.83 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	7.97 × 10 ⁻⁰⁴	2.30 x 10 ⁻⁰⁴	1.77 x 10 ⁻⁰²	1.87 x 10 ⁻⁰²	1.82 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	2.57 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	1.12 x 10 ⁻⁰³	2.29 x 10 ⁻⁰⁴	1.79 x 10 ⁻⁰²	1.92 x 10 ⁻⁰²
South Arm)	Molybdenum	4.51 × 10 ⁻⁰⁶	2.99 x 10 ⁻⁰⁹	6.50 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	9.99 × 10 ⁻⁰⁸	8.49 x 10 ⁻⁰⁶	1.96 x 10 ⁻⁰³	1.97 x 10 ⁻⁰³	2.18 x 10 ⁻⁰⁵	2.98 x 10 ⁻⁰⁹	3.26 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	4.86 x 10 ⁻⁰⁷	8.18 x 10 ⁻⁰⁶	1.97 x 10 ⁻⁰³	2.01 x 10 ⁻⁰³
	Uranium	2.04×10^{-06}	6.74 x 10 ⁻⁰⁹	1.75 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	5.65 × 10 ⁻⁰⁶	4.87 x 10 ⁻⁰⁵	6.59 x 10 ⁻⁰⁵	1.22 x 10 ⁻⁰⁴	3.31 x 10 ⁻⁰⁶	5.34 x 10 ⁻⁰⁹	3.02 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	9.23 x 10 ⁻⁰⁶	2.38 x 10 ⁻⁰⁵	5.17 x 10 ⁻⁰⁵	8.81 x 10 ⁻⁰⁵
						1			ound Scenario				I				
	Cobalt	9.35 × 10 ⁻⁰⁶	9.05 x 10 ⁻⁰⁹	1.10 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.50 x 10 ⁻⁰⁵	1.46 x 10 ⁻⁰⁵	2.51 x 10 ⁻⁰⁴	3.00 x 10 ⁻⁰⁴	1.44 x 10 ⁻⁰⁵	9.05 x 10 ⁻⁰⁹	1.70 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	3.84 x 10 ⁻⁰⁵	1.44 x 10 ⁻⁰⁵	2.89 x 10 ⁻⁰⁴	3.56 x 10 ⁻⁰⁴
	Copper	1.30 × 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.83 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	7.98 x 10 ⁻⁰⁴	2.30 x 10 ⁻⁰⁴	1.77 x 10 ⁻⁰²	1.87 x 10 ⁻⁰²	2.11 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	2.99 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	1.29 x 10 ⁻⁰³	2.29 x 10 ⁻⁰⁴	1.80 x 10 ⁻⁰²	1.96 x 10 ⁻⁰²
	Molybdenum	7.51 × 10 ⁻⁰⁶	2.99 x 10 ⁻⁰⁹	1.06 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	1.66 x 10 ⁻⁰⁷	8.49 x 10 ⁻⁰⁶	1.96 x 10 ⁻⁰³	1.98 x 10 ⁻⁰³	7.51 x 10 ⁻⁰⁵	2.98 x 10 ⁻⁰⁹	1.12 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.68 x 10 ⁻⁰⁶	8.18 x 10 ⁻⁰⁶	2.04 x 10 ⁻⁰³	2.12 x 10 ⁻⁰³
	Uranium	3.01×10^{-06}	6.74 x 10 ⁻⁰⁹	2.48 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	8.34 x 10 ⁻⁰⁶	4.87 x 10 ⁻⁰⁵	6.66 x 10 ⁻⁰⁵	1.27 x 10 ⁻⁰⁴	3.32 x 10 ⁻⁰⁶	5.34 x 10 ⁻⁰⁹	3.02 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	9.24 x 10 ⁻⁰⁶	2.38 x 10 ⁻⁰⁵	5.17 x 10 ⁻⁰⁵	8.81 x 10 ⁻⁰⁵
									se Case								
	Cobalt	9.58 × 10 ⁻⁰⁶	5.91 × 10 ⁻⁰⁷	6.62 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	2.18 × 10 ⁻⁰⁵	2.84 × 10 ⁻⁰⁵	4.78 × 10 ⁻⁰⁴	5.39 x 10 ⁻⁰⁴	9.58 × 10 ⁻⁰⁶	5.91 × 10 ⁻⁰⁷	6.62 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	2.18 × 10 ⁻⁰⁵	2.84 × 10 ⁻⁰⁵	4.78 × 10 ⁻⁰⁴	5.39 x 10 ⁻⁰⁴
Subsistence	Copper	1.38 × 10 ⁻⁰⁵	8.21 × 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	7.21 × 10 ⁻⁰⁴	4.52 × 10 ⁻⁰⁴	4.13 × 10 ⁻⁰²	4.25 x 10 ⁻⁰²	1.38 × 10 ⁻⁰⁵	8.21 × 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	7.21 × 10 ⁻⁰⁴	4.52 × 10 ⁻⁰⁴	4.13 × 10 ⁻⁰²	4.25 x 10 ⁻⁰²
Harvester One- Year-Old	Molybdenum	3.39 × 10 ⁻⁰⁶	1.95 × 10 ⁻⁰⁷	2.97 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	6.42 × 10 ⁻⁰⁸	1.61 × 10 ⁻⁰⁵	6.06×10^{-03}	6.08 x 10 ⁻⁰³	3.39 × 10 ⁻⁰⁶	1.95 × 10 ⁻⁰⁷	2.97 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	6.42 × 10 ⁻⁰⁸	1.61 × 10 ⁻⁰⁵	6.06 × 10 ⁻⁰³	6.08 x 10 ⁻⁰³
(Patterson Lake	Uranium	1.65 × 10 ⁻⁰⁶	3.38 × 10 ⁻⁰⁷	8.79 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	3.90 × 10 ⁻⁰⁶	4.54 × 10 ⁻⁰⁵	1.02 × 10 ⁻⁰⁴	1.54 x 10 ⁻⁰⁴	1.65 × 10 ⁻⁰⁶	3.38 × 10 ⁻⁰⁷	8.79 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	3.90 × 10 ⁻⁰⁶	4.54×10^{-05}	1.02 × 10 ⁻⁰⁴	1.54 x 10 ⁻⁰⁴
South Arm)									cation Case								
	Cobalt	1.04 × 10 ⁻⁰⁵	5.91 x 10 ⁻⁰⁷	7.14 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	2.36 × 10 ⁻⁰⁵	2.86 x 10 ⁻⁰⁵	4.85 x 10 ⁻⁰⁴	5.49 x 10 ⁻⁰⁴	1.38 x 10 ⁻⁰⁵	5.91 x 10 ⁻⁰⁷	9.54 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	3.13 × 10 ⁻⁰⁵	2.84 x 10 ⁻⁰⁵	5.08 x 10 ⁻⁰⁴	5.83 x 10 ⁻⁰⁴
	Copper	1.45 × 10 ⁻⁰⁵	8.23 x 10 ⁻⁰⁷	1.20 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	7.55 × 10 ⁻⁰⁴	4.54 x 10 ⁻⁰⁴	4.15 x 10 ⁻⁰²	4.27 x 10 ⁻⁰²	2.03 x 10 ⁻⁰⁵	8.21 x 10 ⁻⁰⁷	1.68 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	1.06×10^{-03}	4.52 x 10 ⁻⁰⁴	4.16 x 10 ⁻⁰²	4.32 x 10 ⁻⁰²

				Dose by F	Pathway Project	Lifespan (mg/	kg/d)					Dose	by Pathway Fa	r Future (mg/k	g/d)		
	СОРС	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
	Molybdenum	5.04 × 10 ⁻⁰⁶	1.95 x 10 ⁻⁰⁷	4.25 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	9.47 × 10 ⁻⁰⁸	1.66 x 10 ⁻⁰⁵	6.07 x 10 ⁻⁰³	6.09 x 10 ⁻⁰³	2.43 x 10 ⁻⁰⁵	1.95 x 10 ⁻⁰⁷	2.13 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	4.61 × 10 ⁻⁰⁷	1.61 x 10 ⁻⁰⁵	6.08 x 10 ⁻⁰³	6.13 x 10 ⁻⁰³
	Uranium	2.27 × 10 ⁻⁰⁶	4.41 x 10 ⁻⁰⁷	1.14 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	5.36 × 10 ⁻⁰⁶	8.40 x 10 ⁻⁰⁵	1.14 x 10 ⁻⁰⁴	2.08 x 10 ⁻⁰⁴	3.70 x 10 ⁻⁰⁶	3.49 x 10 ⁻⁰⁷	1.98 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	8.75 × 10 ⁻⁰⁶	4.68 x 10 ⁻⁰⁵	1.04 x 10 ⁻⁰⁴	1.66 x 10 ⁻⁰⁴
								Upper Bo	und Scenario								
	Cobalt	1.04 × 10 ⁻⁰⁵	5.91 x 10 ⁻⁰⁷	7.16 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.37 × 10 ⁻⁰⁵	2.86 x 10 ⁻⁰⁵	4.85 x 10 ⁻⁰⁴	5.49 x 10 ⁻⁰⁴	1.60 × 10 ⁻⁰⁵	5.91 x 10 ⁻⁰⁷	1.11 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	3.64 × 10 ⁻⁰⁵	2.84 x 10 ⁻⁰⁵	5.25 x 10 ⁻⁰⁴	6.07 x 10 ⁻⁰⁴
	Copper	1.45 × 10 ⁻⁰⁵	8.23 x 10 ⁻⁰⁷	1.20 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	7.56 × 10 ⁻⁰⁴	4.54 x 10 ⁻⁰⁴	4.15 x 10 ⁻⁰²	4.27 x 10 ⁻⁰²	2.35 × 10 ⁻⁰⁵	8.21 x 10 ⁻⁰⁷	1.95 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	1.23 × 10 ⁻⁰³	4.52 x 10 ⁻⁰⁴	4.18 x 10 ⁻⁰²	4.35 x 10 ⁻⁰²
	Molybdenum	8.39 × 10 ⁻⁰⁶	1.95 x 10 ⁻⁰⁷	6.94 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	1.57 × 10 ⁻⁰⁷	1.66 x 10 ⁻⁰⁵	6.07 x 10 ⁻⁰³	6.10 x 10 ⁻⁰³	8.39 × 10 ⁻⁰⁵	1.95 x 10 ⁻⁰⁷	7.35 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	1.59 × 10 ⁻⁰⁶	1.61 x 10 ⁻⁰⁵	6.14 x 10 ⁻⁰³	6.25 x 10 ⁻⁰³
	Uranium	3.37×10^{-06}	4.41 x 10 ⁻⁰⁷	1.62 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	7.91 × 10 ⁻⁰⁶	8.40 x 10 ⁻⁰⁵	1.15 x 10 ⁻⁰⁴	2.12 x 10 ⁻⁰⁴	3.70 × 10 ⁻⁰⁶	3.49 x 10 ⁻⁰⁷	1.98 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	8.76 × 10 ⁻⁰⁶	4.68 x 10 ⁻⁰⁵	1.04 x 10 ⁻⁰⁴	1.66 x 10 ⁻⁰⁴
						1			se Case					1	1		
	Cobalt	8.58 × 10 ⁻⁰⁶	9.04 × 10 ⁻⁰⁹	1.01 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	2.30 × 10 ⁻⁰⁵	1.44 × 10 ⁻⁰⁵	2.44 × 10 ⁻⁰⁴	2.90 x 10 ⁻⁰⁴	8.58 × 10 ⁻⁰⁶	9.04 × 10 ⁻⁰⁹	1.01 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	2.30 × 10 ⁻⁰⁵	1.44 × 10 ⁻⁰⁵	2.44 × 10 ⁻⁰⁴	2.90 x 10 ⁻⁰⁴
	Copper	1.24 × 10 ⁻⁰⁵	1.26 × 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	7.61 × 10 ⁻⁰⁴	2.29 × 10 ⁻⁰⁴	1.75 × 10 ⁻⁰²	1.86 x 10 ⁻⁰²	1.24 × 10 ⁻⁰⁵	1.26 × 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	7.61 × 10 ⁻⁰⁴	2.29 × 10 ⁻⁰⁴	1.75 × 10 ⁻⁰²	1.86 x 10 ⁻⁰²
	Molybdenum	3.04 × 10 ⁻⁰⁶	2.98 × 10 ⁻⁰⁹	4.54 x 10 ⁻⁰⁹	0.00 × 10 ⁺⁰⁰	6.77 × 10 ⁻⁰⁸	8.18 × 10 ⁻⁰⁶	1.95 × 10 ⁻⁰³	1.96 x 10 ⁻⁰³	3.04 × 10 ⁻⁰⁶	2.98 × 10 ⁻⁰⁹	4.54 x 10 ⁻⁰⁹	0.00 × 10 ⁺⁰⁰	6.77 × 10 ⁻⁰⁸	8.18 × 10 ⁻⁰⁶	1.95 × 10 ⁻⁰³	1.96 x 10 ⁻⁰³
	Uranium	1.48 × 10 ⁻⁰⁶	5.17 × 10 ⁻⁰⁹	1.35 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	4.11 × 10 ⁻⁰⁶	2.30 × 10 ⁻⁰⁵	4.92 × 10 ⁻⁰⁵	7.78 x 10 ⁻⁰⁵	1.48 × 10 ⁻⁰⁶	5.17 × 10 ⁻⁰⁹	1.35 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	4.11 × 10 ⁻⁰⁶	2.30 × 10 ⁻⁰⁵	4.92 × 10 ⁻⁰⁵	7.78 x 10 ⁻⁰⁵
	6.1.11	0.07 40-06	0.05 - 40-09	4.00 4.0+08	0.00 40+00	2.40 40-05	4.40 - 40.05		ation Case	4.00 40*05	0.0540.09	4.05 - 4.0-08	0.00 40+00	2.02 40:05	4.4440.05	0.54 4.0-04	0.00 4.0:04
Subsistence	Cobalt	8.97 × 10 ⁻⁰⁶	9.05 x 10 ⁻⁰⁹	1.06 x 10 ⁻⁰⁸ 1.79 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	2.40 × 10 ⁻⁰⁵	1.46 x 10 ⁻⁰⁵ 2.30 x 10 ⁻⁰⁴	2.47 x 10 ⁻⁰⁴ 1.76 x 10 ⁻⁰²	2.94 x 10 ⁻⁰⁴ 1.86 x 10 ⁻⁰²	1.06 x 10 ⁻⁰⁵ 1.54 x 10 ⁻⁰⁵	9.05 x 10 ⁻⁰⁹ 1.26 x 10 ⁻⁰⁸	1.25 x 10 ⁻⁰⁸ 2.18 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	2.83 × 10 ⁻⁰⁵	1.44 x 10 ⁻⁰⁵ 2.29 x 10 ⁻⁰⁴	2.54 x 10 ⁻⁰⁴ 1.77 x 10 ⁻⁰²	3.08 x 10 ⁻⁰⁴
Harvester Adult (Beet Lake)	Copper	1.27 × 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸ 2.98 x 10 ⁻⁰⁹	5.55 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	7.79×10^{-04} 8.42×10^{-08}	8.34 x 10 ⁻⁰⁶	1.76 x 10 ⁻⁰²	1.86 X 10 ⁻⁰²	1.54 x 10 °5 1.28 x 10 °5	2.98 x 10 ⁻⁰⁹	1.92 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	9.46 × 10 ⁻⁰⁴	8.18 x 10 ⁻⁰⁶	1.77 x 10 ⁻⁰²	1.89 x 10 ⁻⁰² 1.98 x 10 ⁻⁰³
	Molybdenum Uranium	3.79×10^{-06} 1.68×10^{-06}	6.10 x 10 ⁻⁰⁹	1.49 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	4.68 × 10 ⁻⁰⁶	3.82 x 10 ⁻⁰⁵	5.55 x 10 ⁻⁰⁵	1.97 x 10 ⁻⁰⁴	2.16 x 10 ⁻⁰⁶	5.27 x 10 ⁻⁰⁹	1.92 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	2.86×10^{-07} 6.01×10^{-06}	2.35 x 10 ⁻⁰⁵	5.01 x 10 ⁻⁰⁵	8.17 x 10 ⁻⁰⁵
	Oranium	1.68 × 10 ··	6.10 X 10 **	1.49 X 10 **	0.00 × 10 ···	4.00 × 10 °°	3.62 X 10 **		ound Scenario	2.16 % 10 **	5.27 X 10 ···	1.97 X 10 **	0.00 x 10 ···	6.01 × 10 ···	2.35 X 10 **	5.01 X 10 **	6.17 X 10 ··
	Cobalt	8.98 × 10 ⁻⁰⁶	9.05 x 10 ⁻⁰⁹	1.06 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	2.40 × 10 ⁻⁰⁵	1.46 x 10 ⁻⁰⁵	2.47 x 10 ⁻⁰⁴	2.94 x 10 ⁻⁰⁴	1.16 x 10 ⁻⁰⁵	9.05 x 10 ⁻⁰⁹	1.38 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	3.11 × 10 ⁻⁰⁵	1.44 x 10 ⁻⁰⁵	2.60 x 10 ⁻⁰⁴	3.17 x 10 ⁻⁰⁴
	Copper	1.27 × 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.79 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	7.80 × 10 ⁻⁰⁴	2.30 x 10 ⁻⁰⁴	1.76 x 10 ⁻⁰²	1.86 x 10 ⁻⁰²	1.69 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	2.40 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	1.04 × 10 ⁻⁰³	2.29 x 10 ⁻⁰⁴	1.77 x 10 ⁻⁰²	1.90 x 10 ⁻⁰²
	Molybdenum	5.35 × 10 ⁻⁰⁶	2.98 x 10 ⁻⁰⁹	7.66 x 10 ⁻⁰⁹	0.00 × 10 ⁺⁰⁰	1.18 × 10 ⁻⁰⁷	8.34 x 10 ⁻⁰⁶	1.96 x 10 ⁻⁰³	1.97 x 10 ⁻⁰³	4.06 x 10 ⁻⁰⁵	2.98 x 10 ⁻⁰⁹	6.07 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	9.05 × 10 ⁻⁰⁷	8.18 x 10 ⁻⁰⁶	1.98 x 10 ⁻⁰³	2.03 x 10 ⁻⁰³
	Uranium	2.04 × 10 ⁻⁰⁶	6.10 x 10 ⁻⁰⁹	1.76 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	5.67 × 10 ⁻⁰⁶	3.82 x 10 ⁻⁰⁵	5.57 x 10 ⁻⁰⁵	1.02 x 10 ⁻⁰⁴	2.16 x 10 ⁻⁰⁶	5.27 x 10 ⁻⁰⁹	1.97 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	6.02 × 10 ⁻⁰⁶	2.35 x 10 ⁻⁰⁵	5.01 x 10 ⁻⁰⁵	8.17 x 10 ⁻⁰⁵
									se Case								
	Cobalt	9.58 × 10 ⁻⁰⁶	5.91 × 10 ⁻⁰⁷	6.62 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	2.18 × 10 ⁻⁰⁵	2.84 × 10 ⁻⁰⁵	4.78 × 10 ⁻⁰⁴	5.39 x 10 ⁻⁰⁴	9.58 × 10 ⁻⁰⁶	5.91 × 10 ⁻⁰⁷	6.62 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	2.18 × 10 ⁻⁰⁵	2.84 × 10 ⁻⁰⁵	4.78 × 10 ⁻⁰⁴	5.39 x 10 ⁻⁰⁴
	Copper	1.38 × 10 ⁻⁰⁵	8.21 × 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	7.21 × 10 ⁻⁰⁴	4.52 × 10 ⁻⁰⁴	4.13 × 10 ⁻⁰²	4.25 x 10 ⁻⁰²	1.38 × 10 ⁻⁰⁵	8.21 × 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	7.21 × 10 ⁻⁰⁴	4.52 × 10 ⁻⁰⁴	4.13 × 10 ⁻⁰²	4.25 x 10 ⁻⁰²
	Molybdenum	3.39 × 10 ⁻⁰⁶	1.95 × 10 ⁻⁰⁷	2.97 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	6.42 × 10 ⁻⁰⁸	1.61 × 10 ⁻⁰⁵	6.06 × 10 ⁻⁰³	6.08 x 10 ⁻⁰³	3.39 × 10 ⁻⁰⁶	1.95 × 10 ⁻⁰⁷	2.97 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	6.42 × 10 ⁻⁰⁸	1.61 × 10 ⁻⁰⁵	6.06 × 10 ⁻⁰³	6.08 x 10 ⁻⁰³
	Uranium	1.65 × 10 ⁻⁰⁶	3.38 × 10 ⁻⁰⁷	8.79 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	3.90 × 10 ⁻⁰⁶	4.54 × 10 ⁻⁰⁵	1.02 × 10 ⁻⁰⁴	1.54 x 10 ⁻⁰⁴	1.65 × 10 ⁻⁰⁶	3.38 × 10 ⁻⁰⁷	8.79 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	3.90 × 10 ⁻⁰⁶	4.54 × 10 ⁻⁰⁵	1.02 × 10 ⁻⁰⁴	1.54 x 10 ⁻⁰⁴
								Applic	ation Case								
Subsistence	Cobalt	1.00×10^{-05}	5.91 x 10 ⁻⁰⁷	6.89 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.27 × 10 ⁻⁰⁵	2.86 x 10 ⁻⁰⁵	4.80 x 10 ⁻⁰⁴	5.43 x 10 ⁻⁰⁴	1.18 x 10 ⁻⁰⁵	5.91 x 10 ⁻⁰⁷	8.17 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.68 × 10 ⁻⁰⁵	2.84 x 10 ⁻⁰⁵	4.89 x 10 ⁻⁰⁴	5.57 x 10 ⁻⁰⁴
Harvester One- Year-Old (Beet	Copper	1.42 × 10 ⁻⁰⁵	8.23 x 10 ⁻⁰⁷	1.17 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	7.39 × 10 ⁻⁰⁴	4.53 x 10 ⁻⁰⁴	4.14 x 10 ⁻⁰²	4.26 x 10 ⁻⁰²	1.72 x 10 ⁻⁰⁵	8.21 x 10 ⁻⁰⁷	1.43 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	8.97 × 10 ⁻⁰⁴	4.52 x 10 ⁻⁰⁴	4.14 x 10 ⁻⁰²	4.28 x 10 ⁻⁰²
Lake)	Molybdenum	4.23 × 10 ⁻⁰⁶	1.95 x 10 ⁻⁰⁷	3.62 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	7.98 × 10 ⁻⁰⁸	1.63 x 10 ⁻⁰⁵	6.06 x 10 ⁻⁰³	6.08 x 10 ⁻⁰³	1.43 x 10 ⁻⁰⁵	1.95 x 10 ⁻⁰⁷	1.25 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.71 × 10 ⁻⁰⁷	1.61 x 10 ⁻⁰⁵	6.07 x 10 ⁻⁰³	6.10 x 10 ⁻⁰³
	Uranium	1.88 × 10 ⁻⁰⁶	3.98 x 10 ⁻⁰⁷	9.77 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	4.43×10^{-06}	6.82 x 10 ⁻⁰⁵	1.07 x 10 ⁻⁰⁴	1.83 x 10 ⁻⁰⁴	2.41 x 10 ⁻⁰⁶	3.44 x 10 ⁻⁰⁷	1.29 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	5.70 × 10 ⁻⁰⁶	4.63 x 10 ⁻⁰⁵	1.03 x 10 ⁻⁰⁴	1.59 x 10 ⁻⁰⁴
									und Scenario								
	Cobalt	1.00 × 10 ⁻⁰⁵	5.91 x 10 ⁻⁰⁷	6.91 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	2.28 × 10 ⁻⁰⁵	2.86 x 10 ⁻⁰⁵	4.81 x 10 ⁻⁰⁴	5.43 x 10 ⁻⁰⁴	1.30 x 10 ⁻⁰⁵	5.91 x 10 ⁻⁰⁷	8.99 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	2.95 × 10 ⁻⁰⁵	2.84 × 10 ⁻⁰⁵	4.94 × 10 ⁻⁰⁴	5.66 × 10 ⁻⁰⁴
	Copper	1.42 × 10 ⁻⁰⁵	8.23 x 10 ⁻⁰⁷	1.17 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	7.39 × 10 ⁻⁰⁴	4.53 x 10 ⁻⁰⁴	4.14 x 10 ⁻⁰²	4.26 x 10 ⁻⁰²	1.89 x 10 ⁻⁰⁵	8.21 x 10 ⁻⁰⁷	1.57 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	9.85 × 10 ⁻⁰⁴	4.52 × 10 ⁻⁰⁴	4.15 × 10 ⁻⁰²	4.30 × 10 ⁻⁰²
	Molybdenum	5.97 × 10 ⁻⁰⁶	1.95 x 10 ⁻⁰⁷	5.01 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	1.12 × 10 ⁻⁰⁷	1.63 x 10 ⁻⁰⁵	6.06 x 10 ⁻⁰³	6.09 x 10 ⁻⁰³	4.53 x 10 ⁻⁰⁵	1.95 x 10 ⁻⁰⁷	3.97 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	8.58 × 10 ⁻⁰⁷	1.61 × 10 ⁻⁰⁵	6.09 × 10 ⁻⁰³	6.16 × 10 ⁻⁰³
	Uranium	2.28 × 10 ⁻⁰⁶	3.98 x 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	5.37 × 10 ⁻⁰⁶	6.82 x 10 ⁻⁰⁵	1.07 x 10 ⁻⁰⁴	1.84 x 10 ⁻⁰⁴	2.41 x 10 ⁻⁰⁶	3.44 x 10 ⁻⁰⁷	1.29 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	5.70 × 10 ⁻⁰⁶	4.63 × 10 ⁻⁰⁵	1.03 × 10 ⁻⁰⁴	1.59 × 10 ⁻⁰⁴
		2.20		1.04 := 00					se Case		- 00	4.04 := 00					0.00 15.01
	Cobalt	8.58 × 10 ⁻⁰⁶	9.04 × 10 ⁻⁰⁹	1.01 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	2.30 × 10 ⁻⁰⁵	1.44 × 10 ⁻⁰⁵	2.44 × 10 ⁻⁰⁴	2.90 x 10 ⁻⁰⁴	8.58 × 10 ⁻⁰⁶	9.04 × 10 ⁻⁰⁹	1.01 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	2.30 × 10 ⁻⁰⁵	1.44 × 10 ⁻⁰⁵	2.44 × 10 ⁻⁰⁴	2.90 x 10 ⁻⁰⁴

				Dose by F	athway Project	Lifespan (mg/	kg/d)					Dose	by Pathway Fa	r Future (mg/k	g/d)		
	СОРС	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
	Copper	1.24 × 10 ⁻⁰⁵	1.26 × 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	7.61 × 10 ⁻⁰⁴	2.29 × 10 ⁻⁰⁴	1.75 × 10 ⁻⁰²	1.86 x 10 ⁻⁰²	1.24 × 10 ⁻⁰⁵	1.26 × 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	7.61 × 10 ⁻⁰⁴	2.29 × 10 ⁻⁰⁴	1.75 × 10 ⁻⁰²	1.86 x 10 ⁻⁰²
	Molybdenum	3.04 × 10 ⁻⁰⁶	2.98 × 10 ⁻⁰⁹	4.54 x 10 ⁻⁰⁹	0.00 × 10 ⁺⁰⁰	6.77 × 10 ⁻⁰⁸	8.18 × 10 ⁻⁰⁶	1.95 × 10 ⁻⁰³	1.96 x 10 ⁻⁰³	3.04 × 10 ⁻⁰⁶	2.98 × 10 ⁻⁰⁹	4.54 x 10 ⁻⁰⁹	0.00 × 10 ⁺⁰⁰	6.77 × 10 ⁻⁰⁸	8.18 × 10 ⁻⁰⁶	1.95 × 10 ⁻⁰³	1.96 x 10 ⁻⁰³
	Uranium	1.48 × 10 ⁻⁰⁶	5.17 × 10 ⁻⁰⁹	1.35 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	4.11 × 10 ⁻⁰⁶	2.30 × 10 ⁻⁰⁵	4.92 × 10 ⁻⁰⁵	7.78 x 10 ⁻⁰⁵	1.48 × 10 ⁻⁰⁶	5.17 × 10 ⁻⁰⁹	1.35 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	4.11 × 10 ⁻⁰⁶	2.30 × 10 ⁻⁰⁵	4.92 × 10 ⁻⁰⁵	7.78 x 10 ⁻⁰⁵
								Applio	cation Case								
	Cobalt	8.62 × 10 ⁻⁰⁶	9.05 x 10 ⁻⁰⁹	1.02 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	2.31 × 10 ⁻⁰⁵	1.46 x 10 ⁻⁰⁵	2.45 x 10 ⁻⁰⁴	2.91 x 10 ⁻⁰⁴	8.78 x 10 ⁻⁰⁶	9.05 x 10 ⁻⁰⁹	1.04 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	2.35 × 10 ⁻⁰⁵	1.44 x 10 ⁻⁰⁵	2.45 x 10 ⁻⁰⁴	2.92 x 10 ⁻⁰⁴
Subsistence	Copper	1.24 × 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	7.63 × 10 ⁻⁰⁴	2.30 x 10 ⁻⁰⁴	1.76 x 10 ⁻⁰²	1.86 x 10 ⁻⁰²	1.27 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.80 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	7.80 × 10 ⁻⁰⁴	2.29 x 10 ⁻⁰⁴	1.76 x 10 ⁻⁰²	1.86 x 10 ⁻⁰²
Harvester Adult	Molybdenum	3.11 × 10 ⁻⁰⁶	2.98 x 10 ⁻⁰⁹	4.65 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	6.94 × 10 ⁻⁰⁸	8.34 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰³	1.97 x 10 ⁻⁰³	4.03 x 10 ⁻⁰⁶	2.98 x 10 ⁻⁰⁹	6.03 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	8.99 × 10 ⁻⁰⁸	8.18 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰³	1.97 x 10 ⁻⁰³
(Lloyd Lake)	Uranium	1.49 × 10 ⁻⁰⁶	5.21 x 10 ⁻⁰⁹	1.36 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	4.16 × 10 ⁻⁰⁶	2.36 x 10 ⁻⁰⁵	4.94 x 10 ⁻⁰⁵	7.87 x 10 ⁻⁰⁵	1.54 x 10 ⁻⁰⁶	5.18 x 10 ⁻⁰⁹	1.40 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	4.28 × 10 ⁻⁰⁶	2.30 x 10 ⁻⁰⁵	4.92 x 10 ⁻⁰⁵	7.81 x 10 ⁻⁰⁵
			T	1			I	Upper Bo	ound Scenario				T		ı	ı	ı
	Cobalt	8.62 × 10 ⁻⁰⁶	9.05 x 10 ⁻⁰⁹	1.02 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	2.31 × 10 ⁻⁰⁵	1.46 x 10 ⁻⁰⁵	2.45 x 10 ⁻⁰⁴	2.91 x 10 ⁻⁰⁴	8.89 x 10 ⁻⁰⁶	9.05 x 10 ⁻⁰⁹	1.05 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	2.38 × 10 ⁻⁰⁵	1.44 x 10 ⁻⁰⁵	2.46 x 10 ⁻⁰⁴	2.93 x 10 ⁻⁰⁴
	Copper	1.24 × 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	7.63 × 10 ⁻⁰⁴	2.30 x 10 ⁻⁰⁴	1.76 x 10 ⁻⁰²	1.86 x 10 ⁻⁰²	1.29 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.82 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	7.89 × 10 ⁻⁰⁴	2.29 x 10 ⁻⁰⁴	1.76 x 10 ⁻⁰²	1.86 x 10 ⁻⁰²
	Molybdenum	3.27 × 10 ⁻⁰⁶	2.98 x 10 ⁻⁰⁹	4.86 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	7.29 × 10 ⁻⁰⁸	8.34 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰³	1.97 x 10 ⁻⁰³	6.86 x 10 ⁻⁰⁶	2.98 x 10 ⁻⁰⁹	1.03 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	1.53 × 10 ⁻⁰⁷	8.18 x 10 ⁻⁰⁶	1.96 x 10 ⁻⁰³	1.97 x 10 ⁻⁰³
	Uranium	1.53 × 10 ⁻⁰⁶	5.21 x 10 ⁻⁰⁹	1.38 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	4.25×10^{-06}	2.36 x 10 ⁻⁰⁵	4.95 x 10 ⁻⁰⁵	7.89 x 10 ⁻⁰⁵	1.54 x 10 ⁻⁰⁶	5.18 x 10 ⁻⁰⁹	1.40 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	4.28 × 10 ⁻⁰⁶	2.30 x 10 ⁻⁰⁵	4.92 x 10 ⁻⁰⁵	7.81 x 10 ⁻⁰⁵
			Ī	1			I		se Case		1		I	ı	T	T	T
	Cobalt	9.58 × 10 ⁻⁰⁶	5.91 × 10 ⁻⁰⁷	6.62 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.18 × 10 ⁻⁰⁵	2.84 × 10 ⁻⁰⁵	4.78 × 10 ⁻⁰⁴	5.39 x 10 ⁻⁰⁴	9.58 × 10 ⁻⁰⁶	5.91 × 10 ⁻⁰⁷	6.62 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	2.18 × 10 ⁻⁰⁵	2.84 × 10 ⁻⁰⁵	4.78 × 10 ⁻⁰⁴	5.39 x 10 ⁻⁰⁴
	Copper	1.38 × 10 ⁻⁰⁵	8.21 × 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	7.21 × 10 ⁻⁰⁴	4.52 × 10 ⁻⁰⁴	4.13 × 10 ⁻⁰²	4.25 x 10 ⁻⁰²	1.38 × 10 ⁻⁰⁵	8.21 × 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	7.21 × 10 ⁻⁰⁴	4.52 × 10 ⁻⁰⁴	4.13 × 10 ⁻⁰²	4.25 x 10 ⁻⁰²
	Molybdenum	3.39 × 10 ⁻⁰⁶	1.95 × 10 ⁻⁰⁷	2.97 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	6.42 × 10 ⁻⁰⁸	1.61 × 10 ⁻⁰⁵	6.06 × 10 ⁻⁰³	6.08 x 10 ⁻⁰³	3.39 × 10 ⁻⁰⁶	1.95 × 10 ⁻⁰⁷	2.97 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	6.42 × 10 ⁻⁰⁸	1.61 × 10 ⁻⁰⁵	6.06 × 10 ⁻⁰³	6.08 x 10 ⁻⁰³
	Uranium	1.65 × 10 ⁻⁰⁶	3.38 × 10 ⁻⁰⁷	8.79 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	3.90 × 10 ⁻⁰⁶	4.54 × 10 ⁻⁰⁵	1.02 × 10 ⁻⁰⁴	1.54 x 10 ⁻⁰⁴	1.65 × 10 ⁻⁰⁶	3.38 × 10 ⁻⁰⁷	8.79 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	3.90 × 10 ⁻⁰⁶	4.54 × 10 ⁻⁰⁵	1.02 × 10 ⁻⁰⁴	1.54 x 10 ⁻⁰⁴
			I						cation Case				I	-			
Subsistence Harvester One-	Cobalt	9.62 × 10 ⁻⁰⁶	5.91 x 10 ⁻⁰⁷	6.65 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	2.19 × 10 ⁻⁰⁵	2.86 x 10 ⁻⁰⁵	4.79 x 10 ⁻⁰⁴	5.40 x 10 ⁻⁰⁴	9.80 x 10 ⁻⁰⁶	5.91 x 10 ⁻⁰⁷	6.78 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	2.23 × 10 ⁻⁰⁵	2.84 x 10 ⁻⁰⁵	4.79 x 10 ⁻⁰⁴	5.41 x 10 ⁻⁰⁴
Year-Old (Lloyd	Copper	1.39 × 10 ⁻⁰⁵	8.22 x 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	7.23 × 10 ⁻⁰⁴	4.53 x 10 ⁻⁰⁴	4.13 x 10 ⁻⁰²	4.25 x 10 ⁻⁰²	1.42 x 10 ⁻⁰⁵	8.21 x 10 ⁻⁰⁷	1.18 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	7.39 × 10 ⁻⁰⁴	4.52 x 10 ⁻⁰⁴	4.13 x 10 ⁻⁰²	4.25 x 10 ⁻⁰²
Lake)	Molybdenum	3.48 × 10 ⁻⁰⁶	1.95 x 10 ⁻⁰⁷	3.04 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	6.58 × 10 ⁻⁰⁸	1.63 x 10 ⁻⁰⁵	6.06 x 10 ⁻⁰³	6.08 x 10 ⁻⁰³	4.50 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰⁷	3.94 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	8.52 × 10 ⁻⁰⁸	1.61 x 10 ⁻⁰⁵	6.06 x 10 ⁻⁰³	6.08 x 10 ⁻⁰³
	Uranium	1.67 × 10 ⁻⁰⁶	3.40 x 10 ⁻⁰⁷	8.88 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	3.95 × 10 ⁻⁰⁶	4.63 x 10 ⁻⁰⁵	1.02 x 10 ⁻⁰⁴	1.55 x 10 ⁻⁰⁴	1.72 x 10 ⁻⁰⁶	3.38 x 10 ⁻⁰⁷	9.16 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	4.06 × 10 ⁻⁰⁶	4.54 x 10 ⁻⁰⁵	1.02 x 10 ⁻⁰⁴	1.55 x 10 ⁻⁰⁴
		10.00	- 04 40 07	0.07	12:00	- 12 12 05	2.22 12.05		ound Scenario	2.00 40.06	- 04 40 07				0.04 40.05	1.00 10.01	- 40 40 04
	Cobalt	9.62 × 10 ⁻⁰⁶	5.91 x 10 ⁻⁰⁷	6.65 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	2.19 × 10 ⁻⁰⁵	2.86 x 10 ⁻⁰⁵	4.79 x 10 ⁻⁰⁴	5.40 x 10 ⁻⁰⁴	9.93 x 10 ⁻⁰⁶	5.91 x 10 ⁻⁰⁷	6.87 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	2.26 × 10 ⁻⁰⁵	2.84 x 10 ⁻⁰⁵	4.80 x 10 ⁻⁰⁴	5.42 x 10 ⁻⁰⁴
	Copper	1.39 × 10 ⁻⁰⁵	8.22 x 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	7.23 × 10 ⁻⁰⁴	4.53 x 10 ⁻⁰⁴	4.13 x 10 ⁻⁰²	4.25 x 10 ⁻⁰²	1.44 x 10 ⁻⁰⁵	8.21 x 10 ⁻⁰⁷	1.19 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	7.48 × 10 ⁻⁰⁴	4.52 x 10 ⁻⁰⁴	4.13 x 10 ⁻⁰²	4.26 x 10 ⁻⁰²
	Molybdenum	3.66 × 10 ⁻⁰⁶	1.95 x 10 ⁻⁰⁷	3.18 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	6.91 × 10 ⁻⁰⁸	1.63 x 10 ⁻⁰⁵	6.06 x 10 ⁻⁰³	6.08 x 10 ⁻⁰³	7.66 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰⁷	6.71 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	1.45 × 10 ⁻⁰⁷	1.61 x 10 ⁻⁰⁵	6.06 x 10 ⁻⁰³	6.09 x 10 ⁻⁰³
	Uranium	1.71 × 10 ⁻⁰⁶	3.40 x 10 ⁻⁰⁷	9.04 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	4.03 × 10 ⁻⁰⁶	4.63 x 10 ⁻⁰⁵	1.02 x 10 ⁻⁰⁴	1.56 x 10 ⁻⁰⁴	1.72 x 10 ⁻⁰⁶	3.38 x 10 ⁻⁰⁷	9.16 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	4.06 × 10 ⁻⁰⁶	4.54 x 10 ⁻⁰⁵	1.02 x 10 ⁻⁰⁴	1.55 x 10 ⁻⁰⁴
	Cobalt	8.58 × 10 ⁻⁰⁶	9.04 × 10 ⁻⁰⁹	1.01 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	7.27 × 10 ⁻⁰⁶	1.13 × 10 ⁻⁰⁵	1.88 × 10 ⁻⁰⁴	2.15 x 10 ⁻⁰⁴	8.58 × 10 ⁻⁰⁶	9.04 × 10 ⁻⁰⁹	1.01 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	7.27 × 10 ⁻⁰⁶	1.13 × 10 ⁻⁰⁵	1.88 × 10 ⁻⁰⁴	2.15 x 10 ⁻⁰⁴
	Copper	1.24 × 10 ⁻⁰⁵	1.26 × 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	2.41 × 10 ⁻⁰⁴	1.80 × 10 ⁻⁰⁴	1.65 × 10 ⁻⁰²	1.69 x 10 ⁻⁰²	1.24 × 10 ⁻⁰⁵	1.26 × 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	2.41 × 10 ⁻⁰⁴	1.80 × 10 ⁻⁰⁴	1.65 × 10 ⁻⁰²	1.69 x 10 ⁻⁰²
	Molybdenum	3.04 × 10 ⁻⁰⁶	2.98 × 10 ⁻⁰⁹	4.54 x 10 ⁻⁰⁹	0.00 × 10 ⁺⁰⁰	2.41×10^{-08}	6.42 × 10 ⁻⁰⁶	2.41 × 10 ⁻⁰³	2.42 x 10 ⁻⁰³	3.04 × 10 ⁻⁰⁶	2.98 × 10 ⁻⁰⁹	4.54 x 10 ⁻⁰⁹	0.00 × 10 · · · · · · · · · · · · · · · · · ·	2.41 × 10 2.14 × 10 ⁻⁰⁸	6.42×10^{-06}	2.41×10^{-03}	2.42 x 10 ⁻⁰³
	Uranium	1.48 × 10 ⁻⁰⁶	5.17 × 10 ⁻⁰⁹	1.35 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	1.30 × 10 ⁻⁰⁶	1.81 × 10 ⁻⁰⁵	4.08 × 10 ⁻⁰⁵	6.16 x 10 ⁻⁰⁵	1.48 × 10 ⁻⁰⁶	5.17 × 10 ⁻⁰⁹	1.35 x 10 ⁻⁰⁸	0.00 × 10 · · · · · · · · · · · · · · · · · ·	1.30 × 10 ⁻⁰⁶	1.81 × 10 ⁻⁰⁵	4.08×10^{-05}	6.16 x 10 ⁻⁰⁵
Seasonal Resident Adult (Patterson	- Oraniam	1.70 ^ 10	3.17 ^ 10	1.00 X 10	0.00 × 10	1.50 ^ 10	1.01 ^ 10		cation Case	1.70 ^ 10	3.17 ^ 10	1.00 X 10	0.00 × 10	1.50 ^ 10	1.01 ^ 10	7.00 ^ 10	0.10 × 10
Lake South Arm)	Cobalt	9.02 × 10 ⁻⁰⁶	9.05 x 10 ⁻⁰⁹	1.06 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	7.65 × 10 ⁻⁰⁶	1.13 x 10 ⁻⁰⁵	1.89 x 10 ⁻⁰⁴	2.17 x 10 ⁻⁰⁴	1.08 x 10 ⁻⁰⁵	9.04 x 10 ⁻⁰⁹	1.28 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	9.19 × 10 ⁻⁰⁶	1.13 x 10 ⁻⁰⁵	1.95 x 10 ⁻⁰⁴	2.26 x 10 ⁻⁰⁴
	Copper	1.27 × 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.80 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	2.48 × 10 ⁻⁰⁴	1.80 x 10 ⁻⁰⁴	1.65 x 10 ⁻⁰²	1.69 x 10 ⁻⁰²	1.59 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	2.25 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	3.08 × 10 ⁻⁰⁴	1.80 x 10 ⁻⁰⁴	1.65 x 10 ⁻⁰²	1.70 x 10 ⁻⁰²
	Molybdenum	3.92 × 10 ⁻⁰⁶	2.98 x 10 ⁻⁰⁹	5.72 x 10 ⁻⁰⁹	0.00 × 10 0.00 × 10+00	2.76 × 10 ⁻⁰⁸	6.50 x 10 ⁻⁰⁶	2.41 x 10 ⁻⁰³	2.42 x 10 ⁻⁰³	1.43 x 10 ⁻⁰⁵	2.98 x 10 ⁻⁰⁹	2.14 x 10 ⁻⁰⁸	0.00 × 10 · · · · · · · · · · · · · · · · · ·	1.01 × 10 ⁻⁰⁷	6.42×10^{-06}	2.41×10^{-03}	2.43 x 10 ⁻⁰³
	Uranium	1.81 × 10 ⁻⁰⁶	6.11 x 10 ⁻⁰⁹	1.59 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	1.59 × 10 ⁻⁰⁶	2.61 x 10 ⁻⁰⁵	4.39 x 10 ⁻⁰⁵	7.34 x 10 ⁻⁰⁵	2.58 x 10 ⁻⁰⁶	5.27 x 10 ⁻⁰⁹	2.35 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	2.28 × 10 ⁻⁰⁶	1.84 x 10 ⁻⁰⁵	4.14 x 10 ⁻⁰⁵	6.47 x 10 ⁻⁰⁵
	J'amani	1.01 ^ 10	0.11 X 10	1.00 X 10	5.00 A 10	1.33 × 10	x 10		ound Scenario	2.00 X 10	5.21 X 10	2.00 X 10	0.00 × 10	2.20 ^ 10	1.0 1 % 10		0. 11 X 10

				Dose by F	Pathway Project	Lifespan (mg/	kg/d)					Dose	by Pathway Fa	r Future (mg/k	g/d)		
	СОРС	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
	Cobalt	9.04 × 10 ⁻⁰⁶	9.05 x 10 ⁻⁰⁹	1.06 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	7.66 × 10 ⁻⁰⁶	1.13 x 10 ⁻⁰⁵	1.89 x 10 ⁻⁰⁴	2.18 x 10 ⁻⁰⁴	1.20 x 10 ⁻⁰⁵	9.04 x 10 ⁻⁰⁹	1.42 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	1.02 × 10 ⁻⁰⁵	1.13 x 10 ⁻⁰⁵	1.99 x 10 ⁻⁰⁴	2.32 x 10 ⁻⁰⁴
	Copper	1.28 × 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.80 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	2.48 × 10 ⁻⁰⁴	1.80 x 10 ⁻⁰⁴	1.65 x 10 ⁻⁰²	1.69 x 10 ⁻⁰²	1.76 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	2.49 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	3.42 × 10 ⁻⁰⁴	1.80 x 10 ⁻⁰⁴	1.66 x 10 ⁻⁰²	1.71 x 10 ⁻⁰²
	Molybdenum	5.72 × 10 ⁻⁰⁶	2.98 x 10 ⁻⁰⁹	8.19 x 10 ⁻⁰⁹	0.00 × 10 ⁺⁰⁰	4.01 × 10 ⁻⁰⁸	6.50 x 10 ⁻⁰⁶	2.41 x 10 ⁻⁰³	2.42 x 10 ⁻⁰³	4.63 x 10 ⁻⁰⁵	2.98 x 10 ⁻⁰⁹	6.93 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	3.27 × 10 ⁻⁰⁷	6.42 x 10 ⁻⁰⁶	2.43 x 10 ⁻⁰³	2.48 x 10 ⁻⁰³
	Uranium	2.40×10^{-06}	6.11 x 10 ⁻⁰⁹	2.03 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	2.11 × 10 ⁻⁰⁶	2.61 x 10 ⁻⁰⁵	4.41 x 10 ⁻⁰⁵	7.47 x 10 ⁻⁰⁵	2.58 x 10 ⁻⁰⁶	5.27 x 10 ⁻⁰⁹	2.35 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	2.28 × 10 ⁻⁰⁶	1.84 x 10 ⁻⁰⁵	4.14 x 10 ⁻⁰⁵	6.47 x 10 ⁻⁰⁵
								Ва	se Case						_		
	Cobalt	9.58 × 10 ⁻⁰⁶	5.91 × 10 ⁻⁰⁷	6.62 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	6.89 × 10 ⁻⁰⁶	2.47 × 10 ⁻⁰⁵	4.13 × 10 ⁻⁰⁴	4.55 × 10 ⁻⁰⁴	9.58 × 10 ⁻⁰⁶	5.91 × 10 ⁻⁰⁷	6.62 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	6.89 × 10 ⁻⁰⁶	2.47 × 10 ⁻⁰⁵	4.13 × 10 ⁻⁰⁴	4.56 x 10 ⁻⁰⁴
	Copper	1.38 × 10 ⁻⁰⁵	8.21 × 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.29 × 10 ⁻⁰⁴	3.93 × 10 ⁻⁰⁴	3.97 × 10 ⁻⁰²	4.03 × 10 ⁻⁰²	1.38 × 10 ⁻⁰⁵	8.21 × 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.29 × 10 ⁻⁰⁴	3.93 × 10 ⁻⁰⁴	3.97 × 10 ⁻⁰²	4.03 x 10 ⁻⁰²
	Molybdenum	3.39 × 10 ⁻⁰⁶	1.95 × 10 ⁻⁰⁷	2.97 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.03 × 10 ⁻⁰⁸	1.40 × 10 ⁻⁰⁵	6.38 × 10 ⁻⁰³	6.40 × 10 ⁻⁰³	3.39×10^{-06}	1.95 × 10 ⁻⁰⁷	2.97 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.03 × 10 ⁻⁰⁸	1.40 × 10 ⁻⁰⁵	6.38 × 10 ⁻⁰³	6.40 x 10 ⁻⁰³
	Uranium	1.65 × 10 ⁻⁰⁶	3.38 × 10 ⁻⁰⁷	8.79 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	1.23 × 10 ⁻⁰⁶	3.95 × 10 ⁻⁰⁵	9.62 × 10 ⁻⁰⁵	1.40 × 10 ⁻⁰⁴	1.65 × 10 ⁻⁰⁶	3.38 × 10 ⁻⁰⁷	8.79 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	1.23 × 10 ⁻⁰⁶	3.95 × 10 ⁻⁰⁵	9.62 × 10 ⁻⁰⁵	1.40 x 10 ⁻⁰⁴
								Applic	ation Case								
Seasonal Resident	Cobalt	1.01 × 10 ⁻⁰⁵	5.91 x 10 ⁻⁰⁷	6.94 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	7.25 × 10 ⁻⁰⁶	2.48 x 10 ⁻⁰⁵	4.15 x 10 ⁻⁰⁴	4.59 x 10 ⁻⁰⁴	1.21 × 10 ⁻⁰⁵	5.91 x 10 ⁻⁰⁷	8.37 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	8.71 × 10 ⁻⁰⁶	2.47 x 10 ⁻⁰⁵	4.21 x 10 ⁻⁰⁴	4.68 x 10 ⁻⁰⁴
One-Year-Old (Patterson Lake	Copper	1.42×10^{-05}	8.23 x 10 ⁻⁰⁷	1.18 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.35 × 10 ⁻⁰⁴	3.94 x 10 ⁻⁰⁴	3.97 x 10 ⁻⁰²	4.04 x 10 ⁻⁰²	1.77×10^{-05}	8.21 x 10 ⁻⁰⁷	1.47 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.92 × 10 ⁻⁰⁴	3.93 x 10 ⁻⁰⁴	3.98 x 10 ⁻⁰²	4.05 x 10 ⁻⁰²
South Arm)	Molybdenum	4.38×10^{-06}	1.95 x 10 ⁻⁰⁷	3.74 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.61 × 10 ⁻⁰⁸	1.42 x 10 ⁻⁰⁵	6.38 x 10 ⁻⁰³	6.40 x 10 ⁻⁰³	1.60×10^{-05}	1.95 x 10 ⁻⁰⁷	1.40 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	9.58 × 10 ⁻⁰⁸	1.40 x 10 ⁻⁰⁵	6.38 x 10 ⁻⁰³	6.41 x 10 ⁻⁰³
	Uranium	2.02×10^{-06}	4.00 x 10 ⁻⁰⁷	1.04 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	1.51 × 10 ⁻⁰⁶	5.38 x 10 ⁻⁰⁵	9.85 x 10 ⁻⁰⁵	1.57 x 10 ⁻⁰⁴	2.88×10^{-06}	3.45 x 10 ⁻⁰⁷	1.54 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.16 × 10 ⁻⁰⁶	4.02 x 10 ⁻⁰⁵	9.67 x 10 ⁻⁰⁵	1.44 x 10 ⁻⁰⁴
								Upper Bo	ound Scenario						_		
	Cobalt	1.01 × 10 ⁻⁰⁵	5.91 x 10 ⁻⁰⁷	6.95 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	7.26 × 10 ⁻⁰⁶	2.48 x 10 ⁻⁰⁵	4.15 x 10 ⁻⁰⁴	4.59 x 10 ⁻⁰⁴	1.34 x 10 ⁻⁰⁵	5.91 x 10 ⁻⁰⁷	9.30 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	9.68 × 10 ⁻⁰⁶	2.47 x 10 ⁻⁰⁵	4.25 x 10 ⁻⁰⁴	4.74 x 10 ⁻⁰⁴
	Copper	1.42 × 10 ⁻⁰⁵	8.23 x 10 ⁻⁰⁷	1.18 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.35 × 10 ⁻⁰⁴	3.94 x 10 ⁻⁰⁴	3.97 x 10 ⁻⁰²	4.04 x 10 ⁻⁰²	1.97 x 10 ⁻⁰⁵	8.21 x 10 ⁻⁰⁷	1.63 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	3.25 × 10 ⁻⁰⁴	3.93 x 10 ⁻⁰⁴	3.98 x 10 ⁻⁰²	4.05 x 10 ⁻⁰²
	Molybdenum	6.39 × 10 ⁻⁰⁶	1.95 x 10 ⁻⁰⁷	5.35 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	3.80 × 10 ⁻⁰⁸	1.42 x 10 ⁻⁰⁵	6.38 x 10 ⁻⁰³	6.40 x 10 ⁻⁰³	5.17 x 10 ⁻⁰⁵	1.95 x 10 ⁻⁰⁷	4.53 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	3.10 × 10 ⁻⁰⁷	1.40 x 10 ⁻⁰⁵	6.40 x 10 ⁻⁰³	6.47 x 10 ⁻⁰³
	Uranium	2.68×10^{-06}	4.00 x 10 ⁻⁰⁷	1.33 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.00 × 10 ⁻⁰⁶	5.38 x 10 ⁻⁰⁵	9.87 x 10 ⁻⁰⁵	1.59 x 10 ⁻⁰⁴	2.88 x 10 ⁻⁰⁶	3.45 x 10 ⁻⁰⁷	1.54 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.16 × 10 ⁻⁰⁶	4.02 x 10 ⁻⁰⁵	9.67 x 10 ⁻⁰⁵	1.44 x 10 ⁻⁰⁴
								Ва	se Case								
	Cobalt	8.58 × 10 ⁻⁰⁶	9.04 × 10 ⁻⁰⁹	1.01 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	7.27 × 10 ⁻⁰⁶	1.13 × 10 ⁻⁰⁵	1.88 × 10 ⁻⁰⁴	2.15 x 10 ⁻⁰⁴	8.58 x 10 ⁻⁰⁶	9.04 x 10 ⁻⁰⁹	1.01 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	7.27 × 10 ⁻⁰⁶	1.13 × 10 ⁻⁰⁵	1.88 × 10 ⁻⁰⁴	2.15 x 10 ⁻⁰⁴
	Copper	1.24 × 10 ⁻⁰⁵	1.26 × 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	2.41 × 10 ⁻⁰⁴	1.80 × 10 ⁻⁰⁴	1.65 × 10 ⁻⁰²	1.69 x 10 ⁻⁰²	1.24 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	2.41 × 10 ⁻⁰⁴	1.80 × 10 ⁻⁰⁴	1.65 × 10 ⁻⁰²	1.69 x 10 ⁻⁰²
	Molybdenum	3.04 × 10 ⁻⁰⁶	2.98 × 10 ⁻⁰⁹	4.54 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	2.14 × 10 ⁻⁰⁸	6.42 × 10 ⁻⁰⁶	2.41 × 10 ⁻⁰³	2.42 x 10 ⁻⁰³	3.04 x 10 ⁻⁰⁶	2.98 x 10 ⁻⁰⁹	4.54 x 10 ⁻⁰⁹	0.00 × 10 ⁺⁰⁰	2.14 × 10 ⁻⁰⁸	6.42 × 10 ⁻⁰⁶	2.41 × 10 ⁻⁰³	2.42 x 10 ⁻⁰³
	Uranium	1.48×10^{-06}	5.17 × 10 ⁻⁰⁹	1.35 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	1.30 × 10 ⁻⁰⁶	1.81 × 10 ⁻⁰⁵	4.08 × 10 ⁻⁰⁵	6.16 x 10 ⁻⁰⁵	1.48 x 10 ⁻⁰⁶	5.17 x 10 ⁻⁰⁹	1.35 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	1.30 × 10 ⁻⁰⁶	1.81 × 10 ⁻⁰⁵	4.08 × 10 ⁻⁰⁵	6.16 x 10 ⁻⁰⁵
									ation Case								
	Cobalt	8.60 × 10 ⁻⁰⁶	9.05 x 10 ⁻⁰⁹	1.02 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	7.31 × 10 ⁻⁰⁶	1.13 x 10 ⁻⁰⁵	1.88 x 10 ⁻⁰⁴	2.15 x 10 ⁻⁰⁴	8.70 x 10 ⁻⁰⁶	9.04 x 10 ⁻⁰⁹	1.03 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	7.45 × 10 ⁻⁰⁶	1.13 x 10 ⁻⁰⁵	1.88 x 10 ⁻⁰⁴	2.16 x 10 ⁻⁰⁴
Seasonal Resident Adult (Lloyd Lake)	Copper	1.24 × 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	3.21 × 10 ⁻⁰⁴	1.80 x 10 ⁻⁰⁴	1.65 x 10 ⁻⁰²	1.70 x 10 ⁻⁰²	1.26 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.78 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	3.28 × 10 ⁻⁰⁴	1.80 x 10 ⁻⁰⁴	1.65 x 10 ⁻⁰²	1.70 x 10 ⁻⁰²
(, ,	Molybdenum	3.08×10^{-06}	2.98 x 10 ⁻⁰⁹	4.60 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	2.20 × 10 ⁻⁰⁸	6.46 x 10 ⁻⁰⁶	2.41 x 10 ⁻⁰³	2.42 x 10 ⁻⁰³	3.63 x 10 ⁻⁰⁶	2.98 x 10 ⁻⁰⁹	5.44 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	2.85 × 10 ⁻⁰⁸	6.42 x 10 ⁻⁰⁶	2.41 x 10 ⁻⁰³	2.42 x 10 ⁻⁰³
	Uranium	1.49 × 10 ⁻⁰⁶	5.20 x 10 ⁻⁰⁹	1.35 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	1.64 × 10 ⁻⁰⁶	1.82 x 10 ⁻⁰⁵	4.08 x 10 ⁻⁰⁵	6.22 x 10 ⁻⁰⁵	1.51 x 10 ⁻⁰⁶	5.18 x 10 ⁻⁰⁹	1.38 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	1.68 × 10 ⁻⁰⁶	1.81 x 10 ⁻⁰⁵	4.08 x 10 ⁻⁰⁵	6.21 x 10 ⁻⁰⁵
								Upper Bo	ound Scenario								
	Cobalt	8.60 × 10 ⁻⁰⁶	9.05 x 10 ⁻⁰⁹	1.02 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	7.31 × 10 ⁻⁰⁶	1.13 x 10 ⁻⁰⁵	1.88 x 10 ⁻⁰⁴	2.15 x 10 ⁻⁰⁴	8.76 x 10 ⁻⁰⁶	9.04 x 10 ⁻⁰⁹	1.04 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	7.54 × 10 ⁻⁰⁶	1.13 x 10 ⁻⁰⁵	1.88 x 10 ⁻⁰⁴	2.16 x 10 ⁻⁰⁴
	Copper	1.24×10^{-05}	1.26 x 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	3.21 × 10 ⁻⁰⁴	1.80 x 10 ⁻⁰⁴	1.65 x 10 ⁻⁰²	1.70 x 10 ⁻⁰²	1.27 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.80 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	3.32 × 10 ⁻⁰⁴	1.80 x 10 ⁻⁰⁴	1.65 x 10 ⁻⁰²	1.70 x 10 ⁻⁰²
	Molybdenum	3.18×10^{-06}	2.98 x 10 ⁻⁰⁹	4.73 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	2.31 × 10 ⁻⁰⁸	6.46 x 10 ⁻⁰⁶	2.41 x 10 ⁻⁰³	2.42 x 10 ⁻⁰³	5.33 x 10 ⁻⁰⁶	2.98 x 10 ⁻⁰⁹	7.98 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	4.85 × 10 ⁻⁰⁸	6.42 x 10 ⁻⁰⁶	2.41 x 10 ⁻⁰³	2.42 x 10 ⁻⁰³
	Uranium	1.51 × 10 ⁻⁰⁶	5.20 x 10 ⁻⁰⁹	1.37 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	1.67 × 10 ⁻⁰⁶	1.82 x 10 ⁻⁰⁵	4.08 x 10 ⁻⁰⁵	6.23 x 10 ⁻⁰⁵	1.51 x 10 ⁻⁰⁶	5.18 x 10 ⁻⁰⁹	1.38 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	1.68 × 10 ⁻⁰⁶	1.81 x 10 ⁻⁰⁵	4.08 x 10 ⁻⁰⁵	6.21 x 10 ⁻⁰⁵

				Dose by F	Pathway Project	Lifespan (mg/	kg/d)					Dose	by Pathway Fa	r Future (mg/k	g/d)		
	СОРС	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
								Ва	se Case								
	Cobalt	9.58 × 10 ⁻⁰⁶	5.91 × 10 ⁻⁰⁷	6.62 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	6.89 × 10 ⁻⁰⁶	2.47 × 10 ⁻⁰⁵	4.13 × 10 ⁻⁰⁴	4.56 x 10 ⁻⁰⁴	9.58 × 10 ⁻⁰⁶	5.91 × 10 ⁻⁰⁷	6.62 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	6.89 × 10 ⁻⁰⁶	2.47 × 10 ⁻⁰⁵	4.13 × 10 ⁻⁰⁴	4.56 x 10 ⁻⁰⁴
	Copper	1.38 × 10 ⁻⁰⁵	8.21 × 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.29 × 10 ⁻⁰⁴	3.93 × 10 ⁻⁰⁴	3.97 × 10 ⁻⁰²	4.03 x 10 ⁻⁰²	1.38 × 10 ⁻⁰⁵	8.21 × 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.29 × 10 ⁻⁰⁴	3.93 × 10 ⁻⁰⁴	3.97 × 10 ⁻⁰²	4.03 x 10 ⁻⁰²
	Molybdenum	3.39×10^{-06}	1.95 × 10 ⁻⁰⁷	2.97 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.03 × 10 ⁻⁰⁸	1.40 × 10 ⁻⁰⁵	6.38 × 10 ⁻⁰³	6.40 x 10 ⁻⁰³	3.39 × 10 ⁻⁰⁶	1.95 × 10 ⁻⁰⁷	2.97 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.03 × 10 ⁻⁰⁸	1.40 × 10 ⁻⁰⁵	6.38 × 10 ⁻⁰³	6.40 x 10 ⁻⁰³
	Uranium	1.65 × 10 ⁻⁰⁶	3.38 × 10 ⁻⁰⁷	8.79 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	1.23 × 10 ⁻⁰⁶	3.95 × 10 ⁻⁰⁵	9.62 × 10 ⁻⁰⁵	1.40 x 10 ⁻⁰⁴	1.65 × 10 ⁻⁰⁶	3.38 × 10 ⁻⁰⁷	8.79 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	1.23 × 10 ⁻⁰⁶	3.95 × 10 ⁻⁰⁵	9.62 × 10 ⁻⁰⁵	1.40 x 10 ⁻⁰⁴
			ı	T				Applio	cation Case							Ī	
Seasonal Resident	Cobalt	9.60 × 10 ⁻⁰⁶	5.91 x 10 ⁻⁰⁷	6.64 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	6.91 × 10 ⁻⁰⁶	2.48 x 10 ⁻⁰⁵	4.14 x 10 ⁻⁰⁴	4.56 x 10 ⁻⁰⁴	9.71 x 10 ⁻⁰⁶	5.91 x 10 ⁻⁰⁷	6.72 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	6.99 × 10 ⁻⁰⁶	2.47 x 10 ⁻⁰⁵	4.14 x 10 ⁻⁰⁴	4.56 x 10 ⁻⁰⁴
One-Year-Old	Copper	1.39 × 10 ⁻⁰⁵	8.22 x 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.29 × 10 ⁻⁰⁴	3.93 x 10 ⁻⁰⁴	3.97 x 10 ⁻⁰²	4.03 x 10 ⁻⁰²	1.40 x 10 ⁻⁰⁵	8.21 x 10 ⁻⁰⁷	1.16 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.32 × 10 ⁻⁰⁴	3.93 x 10 ⁻⁰⁴	3.97 x 10 ⁻⁰²	4.03 x 10 ⁻⁰²
(Lloyd Lake)	Molybdenum	3.44 × 10 ⁻⁰⁶	1.95 x 10 ⁻⁰⁷	3.01 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.06 × 10 ⁻⁰⁸	1.41 x 10 ⁻⁰⁵	6.38 x 10 ⁻⁰³	6.40 x 10 ⁻⁰³	4.06 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰⁷	3.55 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.43 × 10 ⁻⁰⁸	1.40 x 10 ⁻⁰⁵	6.38 x 10 ⁻⁰³	6.40 x 10 ⁻⁰³
	Uranium	1.66 × 10 ⁻⁰⁶	3.39 x 10 ⁻⁰⁷	8.85 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	1.24 × 10 ⁻⁰⁶	3.98 x 10 ⁻⁰⁵	9.63 x 10 ⁻⁰⁵	1.40 x 10 ⁻⁰⁴	1.69 x 10 ⁻⁰⁶	3.38 x 10 ⁻⁰⁷	9.02 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	1.27×10^{-06}	3.95 x 10 ⁻⁰⁵	9.63 x 10 ⁻⁰⁵	1.40 x 10 ⁻⁰⁴
			1	1		1	1	Upper Bo	ound Scenario		ı		ı	1		ı	
	Cobalt	9.60 × 10 ⁻⁰⁶	5.91 x 10 ⁻⁰⁷	6.64 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	6.91 × 10 ⁻⁰⁶	2.48 x 10 ⁻⁰⁵	4.14 x 10 ⁻⁰⁴	4.56 x 10 ⁻⁰⁴	9.79 x 10 ⁻⁰⁶	5.91 x 10 ⁻⁰⁷	6.77 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	7.05 × 10 ⁻⁰⁶	2.47 x 10 ⁻⁰⁵	4.14 x 10 ⁻⁰⁴	4.57 x 10 ⁻⁰⁴
	Copper	1.39 × 10 ⁻⁰⁵	8.22 x 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.29 × 10 ⁻⁰⁴	3.93 x 10 ⁻⁰⁴	3.97 x 10 ⁻⁰²	4.03 x 10 ⁻⁰²	1.42 x 10 ⁻⁰⁵	8.21 x 10 ⁻⁰⁷	1.17 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.34 × 10 ⁻⁰⁴	3.93 x 10 ⁻⁰⁴	3.97 x 10 ⁻⁰²	4.03 x 10 ⁻⁰²
	Molybdenum	3.55 × 10 ⁻⁰⁶	1.95 x 10 ⁻⁰⁷	3.09 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.13 × 10 ⁻⁰⁸	1.41 x 10 ⁻⁰⁵	6.38 x 10 ⁻⁰³	6.40 x 10 ⁻⁰³	5.95 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰⁷	5.21 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	3.57 × 10 ⁻⁰⁸	1.40 x 10 ⁻⁰⁵	6.38 x 10 ⁻⁰³	6.40 x 10 ⁻⁰³
	Uranium	1.68 × 10 ⁻⁰⁶	3.39 x 10 ⁻⁰⁷	8.94 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	1.26 × 10 ⁻⁰⁶	3.98 x 10 ⁻⁰⁵	9.63 x 10 ⁻⁰⁵	1.40 x 10 ⁻⁰⁴	1.69 x 10 ⁻⁰⁶	3.38 x 10 ⁻⁰⁷	9.02 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	1.27×10^{-06}	3.95 x 10 ⁻⁰⁵	9.63 x 10 ⁻⁰⁵	1.40 x 10 ⁻⁰⁴
			ı	1		ı	ı	Ва	se Case		ı		ı	1	1	T	
	Cobalt	n/a	8.58 × 10 ⁻⁰⁶	9.04 × 10 ⁻⁰⁹	1.01 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	2.30 × 10 ⁻⁰⁵	1.44 × 10 ⁻⁰⁵	2.44 × 10 ⁻⁰⁴	2.90 x 10 ⁻⁰⁴							
	Copper	n/a	1.24 × 10 ⁻⁰⁵	1.26 × 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	7.61 × 10 ⁻⁰⁴	2.29 × 10 ⁻⁰⁴	1.75 × 10 ⁻⁰²	1.86 x 10 ⁻⁰²							
	Molybdenum	n/a	3.04 × 10 ⁻⁰⁶	2.98 × 10 ⁻⁰⁹	4.54 x 10 ⁻⁰⁹	0.00 × 10 ⁺⁰⁰	6.77 × 10 ⁻⁰⁸	8.18 × 10 ⁻⁰⁶	1.95 × 10 ⁻⁰³	1.96 x 10 ⁻⁰³							
	Uranium	n/a	1.48 × 10 ⁻⁰⁶	5.17 × 10 ⁻⁰⁹	1.35 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	4.11 × 10 ⁻⁰⁶	2.30 × 10 ⁻⁰⁵	4.92 × 10 ⁻⁰⁵	7.78 x 10 ⁻⁰⁵							
Permanent			l			1	1	Applio	cation Case		ı		ı	1	1		
Resident Adult	Cobalt	n/a	2.10 × 10 ⁻⁰⁵	9.07 x 10 ⁻⁰⁹	2.48 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	5.62 x 10 ⁻⁰⁵	1.44 x 10 ⁻⁰⁵	3.08 x 10 ⁻⁰⁴	4.00 x 10 ⁻⁰⁴							
(Patterson Lake North Arm - West	Copper	n/a	3.21 × 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	4.54 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	1.97 x 10 ⁻⁰³	2.30 x 10 ⁻⁰⁴	1.83 x 10 ⁻⁰²	2.05 x 10 ⁻⁰²							
Basin)	Molybdenum	n/a	6.77 × 10 ⁻⁰⁵	2.98 x 10 ⁻⁰⁹	1.01 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.51 x 10 ⁻⁰⁶	8.18 x 10 ⁻⁰⁶	2.00 x 10 ⁻⁰³	2.08 x 10 ⁻⁰³							
	Uranium	n/a	1.48 × 10 ⁻⁰⁵	7.48 x 10 ⁻⁰⁹	1.35 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	4.11 x 10 ⁻⁰⁵	3.33 x 10 ⁻⁰⁵	6.76 x 10 ⁻⁰⁵	1.57 x 10 ⁻⁰⁴							
			1				1	• • •	ound Scenario								
	Cobalt	n/a	2.76 × 10 ⁻⁰⁵	9.07 x 10 ⁻⁰⁹	3.26 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	7.39 × 10 ⁻⁰⁵	1.44 x 10 ⁻⁰⁵	3.42 x 10 ⁻⁰⁴	4.58 x 10 ⁻⁰⁴							
	Copper	n/a	4.20 × 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	5.94 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	2.58 × 10 ⁻⁰³	2.30 x 10 ⁻⁰⁴	1.87 x 10 ⁻⁰²	2.15 x 10 ⁻⁰²							
	Molybdenum	n/a	2.51 × 10 ⁻⁰⁴	2.98 x 10 ⁻⁰⁹	3.76 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	5.61 × 10 ⁻⁰⁶	8.18 x 10 ⁻⁰⁶	2.15 x 10 ⁻⁰³	2.41 x 10 ⁻⁰³							
	Uranium	n/a	1.48 × 10 ⁻⁰⁵	7.48 x 10 ⁻⁰⁹	1.35 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	4.12 × 10 ⁻⁰⁵	3.33 x 10 ⁻⁰⁵	6.76 x 10 ⁻⁰⁵	1.57 x 10 ⁻⁰⁴							

ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT HUMAN HEALTH RISK ASSESSMENT

				Dose by F	Pathway Project	: Lifespan (mg,	/kg/d)					Dose	by Pathway Fa	r Future (mg/k	g/d)		
	СОРС	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
									Base C	ase							
	Cobalt	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	9.58 × 10 ⁻⁰⁶	5.91 × 10 ⁻⁰⁷	6.62 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	2.18 × 10 ⁻⁰⁵	2.84 × 10 ⁻⁰⁵	4.78 × 10 ⁻⁰⁴	5.39 x 10 ⁻⁰⁴
	Copper	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	1.38 × 10 ⁻⁰⁵	8.21 × 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	7.21 × 10 ⁻⁰⁴	4.52 × 10 ⁻⁰⁴	4.13 × 10 ⁻⁰²	4.25 x 10 ⁻⁰²
	Molybdenum	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	3.39 × 10 ⁻⁰⁶	1.95 × 10 ⁻⁰⁷	2.97 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	6.42 × 10 ⁻⁰⁸	1.61 × 10 ⁻⁰⁵	6.06 × 10 ⁻⁰³	6.08 x 10 ⁻⁰³
	Uranium	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	1.65 × 10 ⁻⁰⁶	3.38 × 10 ⁻⁰⁷	8.79 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	3.90 × 10 ⁻⁰⁶	4.54 × 10 ⁻⁰⁵	1.02 × 10 ⁻⁰⁴	1.54 x 10 ⁻⁰⁴
Permanent								Applic	ation Case								
Resident One-	Cobalt	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	2.34 × 10 ⁻⁰⁵	5.92 x 10 ⁻⁰⁷	1.62 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	5.33 × 10 ⁻⁰⁵	2.85 x 10 ⁻⁰⁵	5.43 x 10 ⁻⁰⁴	6.50 x 10 ⁻⁰⁴
Year-Old (Patterson Lake	Copper	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	3.58 × 10 ⁻⁰⁵	8.23 x 10 ⁻⁰⁷	2.97 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	1.87 × 10 ⁻⁰³	4.53 x 10 ⁻⁰⁴	4.20 x 10 ⁻⁰²	4.44 x 10 ⁻⁰²
North Arm - West	Molybdenum	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	7.56 × 10 ⁻⁰⁵	1.95 x 10 ⁻⁰⁷	6.62 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	1.43 × 10 ⁻⁰⁶	1.61 x 10 ⁻⁰⁵	6.11 x 10 ⁻⁰³	6.21 x 10 ⁻⁰³
Basin)	Uranium	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	1.65 × 10 ⁻⁰⁵	4.89 x 10 ⁻⁰⁷	8.80 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	3.90 × 10 ⁻⁰⁵	6.56 x 10 ⁻⁰⁵	1.16 x 10 ⁻⁰⁴	2.46 x 10 ⁻⁰⁴
								Upper Bo	und Scenario								
	Cobalt	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	3.08 × 10 ⁻⁰⁵	5.92 x 10 ⁻⁰⁷	2.13 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	7.01 × 10 ⁻⁰⁵	2.85 x 10 ⁻⁰⁵	5.77 x 10 ⁻⁰⁴	7.09 x 10 ⁻⁰⁴
	Copper	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	4.69 × 10 ⁻⁰⁵	8.23 x 10 ⁻⁰⁷	3.88 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.44 × 10 ⁻⁰³	4.53 x 10 ⁻⁰⁴	4.24 x 10 ⁻⁰²	4.53 x 10 ⁻⁰²
	Molybdenum	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	2.81 × 10 ⁻⁰⁴	1.95 x 10 ⁻⁰⁷	2.46 x 10 ⁻⁰⁵	0.00 × 10 ⁺⁰⁰	5.32 × 10 ⁻⁰⁶	1.61 x 10 ⁻⁰⁵	6.26 x 10 ⁻⁰³	6.58 x 10 ⁻⁰³
	Uranium	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	1.65 × 10 ⁻⁰⁵	4.89 x 10 ⁻⁰⁷	8.81 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	3.90 × 10 ⁻⁰⁵	6.56 x 10 ⁻⁰⁵	1.16 x 10 ⁻⁰⁴	2.46 x 10 ⁻⁰⁴

COPC = constituent of potential concern; n/a = receptor not assessed in that phase.

5.2.4.1.2 Carcinogen Doses

Arsenic was evaluated in the HHRA as a non-threshold carcinogen (i.e., a linear dose-response relationship); therefore, predicted exposure was averaged over the receptor's lifetime to estimate a lifetime average daily dose representing a combination of all life stages (Health Canada 2021a). For this assessment, the lifetime average daily dose was estimated for various age groups (toddler, child, teen, adult) to permit estimation of the lifetime risk to a composite receptor for each of the subsistence harvester, seasonal resident, and permanent resident (Table 5-8). Therefore, a composite receptor was calculated assuming 4.5 years as a toddler, 7 years as a child, 8 years as a teen and 60 years as an adult. For the camp worker, an adult receptor was considered appropriate. The composite receptor represents a person exposed to the constituent throughout all stages of a lifetime.



Ref. 19-2574 APRIL 2024

Table 5-8: Estimated Carcinogen Doses for Arsenic to Human Receptors – Operations – Application Case and Upper Bound Scenario

				Lifetime A	Average Daily Dose (m	g/kg/d)						
		А	pplication Case – Opei	rations				U	pper Bound Scenario	- Operations		
Age Group	Camp Worker	Subsistence Harvester - Patterson Lake South Arm	Subsistence Harvester - Beet Lake	Subsistence Harvester - Lloyd Lake	Seasonal Resident - Patterson Lake South Arm	Seasonal Resident - Lloyd Lake	Camp Worker	Subsistence Harvester - Patterson Lake South Arm	Subsistence Harvester - Beet Lake	Subsistence Harvester - Lloyd Lake	Seasonal Resident - Patterson Lake South Arm	Seasonal Resident - Lloyd Lake
1 year old	n/a	9.61 × 10 ⁻⁰⁷	5.55 × 10 ⁻⁰⁸	2.59 × 10 ⁻⁰⁸	2.13 × 10 ⁻⁰⁷	6.14 × 10 ⁻⁰⁹	n/a	9.84 × 10 ⁻⁰⁷	5.63 × 10 ⁻⁰⁸	2.60 × 10 ⁻⁰⁸	2.18 × 10 ⁻⁰⁷	6.15 × 10 ⁻⁰⁹
Child	n/a	2.05 × 10 ⁻⁰⁶	1.26 × 10 ⁻⁰⁷	5.76 × 10 ⁻⁰⁸	4.55 × 10 ⁻⁰⁷	1.27 × 10 ⁻⁰⁸	n/a	2.10 × 10 ⁻⁰⁶	1.28 × 10 ⁻⁰⁷	5.77 × 10 ⁻⁰⁸	4.66 × 10 ⁻⁰⁷	1.28 × 10 ⁻⁰⁸
Teen	n/a	1.60 × 10 ⁻⁰⁶	1.01 × 10 ⁻⁰⁷	4.78 × 10 ⁻⁰⁸	3.47 × 10 ⁻⁰⁷	9.93 × 10 ⁻⁰⁹	n/a	1.64 × 10 ⁻⁰⁶	1.02 × 10 ⁻⁰⁷	4.79 × 10 ⁻⁰⁸	3.55 × 10 ⁻⁰⁷	9.95 × 10 ⁻⁰⁹
Adult	4.27 × 10 ⁻⁰⁵	1.61 × 10 ⁻⁰⁵	1.02 × 10 ⁻⁰⁶	4.67 × 10 ⁻⁰⁷	3.43 × 10 ⁻⁰⁶	9.62 × 10 ⁻⁰⁸	4.36 × 10 ⁻⁰⁵	1.65 × 10 ⁻⁰⁵	1.03 × 10 ⁻⁰⁶	4.68 × 10 ⁻⁰⁷	3.52 × 10 ⁻⁰⁶	9.64 × 10 ⁻⁰⁸

				Lifetime .	Average Daily Dose (m	ng/kg/d)						
		Appl	ication Case - Far-Future	Projection				Upper	Bound Scenario - Fa	r-Future Project	ion	
Age Group	Permanent Resident	Subsistence Harvester - Patterson Lake South Arm	Subsistence Harvester - Beet Lake	Subsistence Harvester - Lloyd Lake	Seasonal Resident - Patterson Lake South Arm	Seasonal Resident - Lloyd Lake	Permanent Resident	Subsistence Harvester - Patterson Lake South Arm	Subsistence Harvester - Beet Lake	Subsistence Harvester - Lloyd Lake	Seasonal Resident - Patterson Lake South Arm	Seasonal Resident - Lloyd Lake
1 year old	1.25 × 10 ⁻⁰⁷	2.37 × 10 ⁻⁰⁸	8.31 × 10 ⁻¹⁰	4.79 × 10 ⁻¹¹	5.24 × 10 ⁻⁰⁹	1.04 × 10 ⁻¹¹	1.93 × 10 ⁻⁰⁷	3.74 × 10 ⁻⁰⁸	1.28 × 10 ⁻⁰⁹	7.36 × 10 ⁻¹¹	8.28 × 10 ⁻⁰⁹	1.60 × 10 ⁻¹¹
Child	2.90 × 10 ⁻⁰⁷	5.05 × 10 ⁻⁰⁸	1.92 × 10 ⁻⁰⁹	1.11 × 10 ⁻¹⁰	1.12 × 10 ⁻⁰⁸	2.40 × 10 ⁻¹¹	4.46 × 10 ⁻⁰⁷	7.97 × 10 ⁻⁰⁸	2.96 × 10 ⁻⁰⁹	1.70 × 10 ⁻¹⁰	1.77 × 10 ⁻⁰⁸	3.70 × 10 ⁻¹¹
Teen	2.26 × 10 ⁻⁰⁷	3.94 × 10 ⁻⁰⁸	1.49 × 10 ⁻⁰⁹	8.62 × 10 ⁻¹¹	8.56 × 10 ⁻⁰⁹	1.85 × 10 ⁻¹¹	3.47 × 10 ⁻⁰⁷	6.22 × 10 ⁻⁰⁸	2.30 × 10 ⁻⁰⁹	1.32 × 10 ⁻¹⁰	1.35 × 10 ⁻⁰⁸	2.84 × 10 ⁻¹¹
Adult	2.35 × 10 ⁻⁰⁶	3.97 × 10 ⁻⁰⁷	1.56 × 10 ⁻⁰⁸	8.98 × 10 ⁻¹⁰	8.50 × 10 ⁻⁰⁸	1.89 × 10 ⁻¹⁰	3.61 × 10 ⁻⁰⁶	6.28 × 10 ⁻⁰⁷	2.40 × 10 ⁻⁰⁸	1.38 × 10 ⁻⁰⁹	1.34 × 10 ⁻⁰⁷	2.91 × 10 ⁻¹⁰

n/a = not applicable or not assessed.

5.2.4.1.3 Radiological Dose

The estimated radiation doses to human receptors during the Project phases and far-future projection resulting from the Application Case and reasonable upper bound sensitivity scenario are shown in Table 5-9 and Table 5-10. The doses shown represent the maximum annual dose over the assessment period. The tables present the dose breakdown by radionuclide and exposure pathway, as well as the total dose. The radiation dose is presented as an incremental dose (i.e., only considering Project effects).

The maximum estimated radiation dose in the Application Case was predicted for the subsistence harvester (one-year-old) who eats Traditional Foods gathered at Patterson Lake South Arm during Operations. The maximum estimated radiation dose to the subsistence harvester (one-year-old) was predicted to be 0.1 mSv/yr for the Application Case and 0.1 mSv/yr for the reasonable upper bound sensitivity scenario. The main contribution to total dose would be from polonium-210 from eating terrestrial animals in the Traditional Food diet, including moose (meat and organs), beaver, grouse, and mallard.

In the far-future projection, the maximum estimated dose was predicted to be 0.1 mSv/yr for the future permanent resident residing at the decommissioned and reclaimed Project site well beyond Closure, both for the Application Case and reasonable upper bound sensitivity scenario.



Ref. 19-2574 APRIL 2024

Table 5-9: Estimated Radiation Doses to Human Receptors – Operations – Application Case, Upper Bound Scenario

							Incremental Dose b	y Pathway - Proje	ct Lifespan (mSv/	yr)				
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by Radionuclide
		0.44 40.04	2.24 40-12	2.20 40-08	4.67. 40-12	4.22 40-07	4.47. 40.03	Application Case		0.00 40+00	4.44 40.05	4.60 40-05	6.25 40.05	2.42.40.03
	Uranium-238	8.41 x 10 ⁻⁰⁴	2.31 x 10 ⁻¹²	2.20 x 10 ⁻⁰⁸	1.67 x 10 ⁻¹³	1.32 x 10 ⁻⁰⁷	1.47 x 10 ⁻⁰³	7.65 x 10 ⁻¹¹	8.15 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	1.11 x 10 ⁻⁰⁵	4.68 x 10 ⁻⁰⁵	6.25 x 10 ⁻⁰⁵	2.43 x 10 ⁻⁰³
	Uranium-234	1.01 x 10 ⁻⁰³	5.64 x 10 ⁻¹²	2.40 x 10 ⁻⁰⁸	2.92 x 10 ⁻¹⁴	1.44 x 10 ⁻⁰⁷	7.40 x 10 ⁻⁰⁶	8.36 x 10 ⁻¹¹	2.97 x 10 ⁻¹²	0.00 x 10 ⁺⁰⁰	1.22 x 10 ⁻⁰⁵	5.10 x 10 ⁻⁰⁵	6.81 x 10 ⁻⁰⁵	1.15 x 10 ⁻⁰³
	Thorium-230 Radium-226	4.06 x 10 ⁻⁰³ 1.01 x 10 ⁻⁰³	1.37 x 10 ⁻¹¹ 2.62 x 10 ⁻¹⁰	1.44 x 10 ⁻⁰⁶ 6.77 x 10 ⁻⁰⁷	9.77 x 10 ⁻¹³	6.91 x 10 ⁻⁰⁷ 9.34 x 10 ⁻⁰⁷	1.06 x 10 ⁻⁰⁵ 2.42 x 10 ⁻⁰²	1.03 x 10 ⁻⁰⁹ 1.61 x 10 ⁻⁰⁹	2.54 x 10 ⁻¹¹ 1.93 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	6.62 x 10 ⁻⁰⁴ 9.97 x 10 ⁻⁰⁵	5.09 x 10 ⁻⁰⁴ 8.83 x 10 ⁻⁰⁴	6.56 x 10 ⁻⁰³ 7.43 x 10 ⁻⁰⁴	1.18 x 10 ⁻⁰² 2.69 x 10 ⁻⁰²
	Lead-210	3.19 x 10 ⁻⁰⁴	4.12 x 10 ⁻¹¹	2.22 x 10 ⁻⁰⁶	3.02 x 10 ⁻¹⁰ 1.95 x 10 ⁻¹²	1.48 x 10 ⁻⁰⁶	3.44 x 10 ⁻⁰⁴	1.96 x 10 ⁻⁰⁸	1.93 x 10 °° 1.11 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	8.90 x 10 ⁻⁰⁵	1.90 x 10 ⁻⁰³	6.37 x 10 ⁻⁰³	9.03 x 10 ⁻⁰³
	Polonium-210	9.56 x 10 ⁻⁰⁴	3.59 x 10 ⁻¹³	1.58 x 10 ⁻⁰⁶	4.74 x 10 ⁻¹⁵	2.62 x 10 ⁻⁰⁶	7.77 x 10 ⁻⁰⁸	3.26 x 10 ⁻⁰⁸	4.48 x 10 ⁻¹²	0.00 x 10 ⁺⁰⁰	5.49 x 10 ⁻⁰⁴	3.04 x 10 ⁻⁰³	9.16 x 10 ⁻⁰³	1.37 x 10 ⁻⁰²
Camp Worker	Total by Pathway	8.20 x 10 ⁻⁰³	3.25 x 10 ⁻¹⁰	5.96 x 10 ⁻⁰⁶	3.05 x 10 ⁻¹⁰	6.01 x 10 ⁻⁰⁶	2.60 x 10 ⁻⁰²	5.50 x 10 ⁻⁰⁸	1.95 x 10 ⁻⁰⁷	0.00 x 10 0.00 x 10+00	1.42 x 10 ⁻⁰³	6.43 x 10 ⁻⁰³	2.30 x 10 ⁻⁰²	6.50 x 10 ⁻⁰²
(Patterson Lake North Arm - West		8.20 X 10	3.23 X 10	3.90 X 10	3.03 X 10	0.01 X 10		pper Bound Scena		0.00 X 10	1.42 X 10	0.43 X 10	2.30 X 10	0.30 X 10
Basin)	Uranium-238	8.41 x 10 ⁻⁰⁴	2.31 x 10 ⁻¹²	2.27 x 10 ⁻⁰⁸	1.72 x 10 ⁻¹³	1.32 x 10 ⁻⁰⁷	1.47 x 10 ⁻⁰³	7.91 x 10 ⁻¹¹	8.43 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	3.05 x 10 ⁻⁰⁵	4.68 x 10 ⁻⁰⁵	6.75 x 10 ⁻⁰⁵	2.45 x 10 ⁻⁰³
	Uranium-234	1.01 x 10 ⁻⁰³	5.64 x 10 ⁻¹²	2.48 x 10 ⁻⁰⁸	3.02 x 10 ⁻¹⁴	1.44 x 10 ⁻⁰⁷	7.40 x 10 ⁻⁰⁶	8.64 x 10 ⁻¹¹	3.07 x 10 ⁻¹²	0.00 x 10 ⁺⁰⁰	3.33 x 10 ⁻⁰⁵	5.10 x 10 ⁻⁰⁵	7.35 x 10 ⁻⁰⁵	1.18 x 10 ⁻⁰³
	Thorium-230	4.06 x 10 ⁻⁰³	1.37 x 10 ⁻¹¹	1.48 x 10 ⁻⁰⁶	1.01 x 10 ⁻¹²	6.91 x 10 ⁻⁰⁷	1.06 x 10 ⁻⁰⁵	1.06 x 10 ⁻⁰⁹	2.61 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	6.70 x 10 ⁻⁰⁴	5.09 x 10 ⁻⁰⁴	6.60 x 10 ⁻⁰³	1.19 x 10 ⁻⁰²
	Radium-226	1.01 x 10 ⁻⁰³	2.62 x 10 ⁻¹⁰	7.12 x 10 ⁻⁰⁷	3.18 x 10 ⁻¹⁰	9.34 x 10 ⁻⁰⁷	2.42 x 10 ⁻⁰²	1.72 x 10 ⁻⁰⁹	2.06 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.04 x 10 ⁻⁰⁴	8.83 x 10 ⁻⁰⁴	7.45 x 10 ⁻⁰⁴	2.69 x 10 ⁻⁰²
	Lead-210	3.19 x 10 ⁻⁰⁴	4.12 x 10 ⁻¹¹	2.29 x 10 ⁻⁰⁶	2.02 x 10 ⁻¹²	1.48 x 10 ⁻⁰⁶	3.44 x 10 ⁻⁰⁴	2.09 x 10 ⁻⁰⁸	1.18 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	4.54 x 10 ⁻⁰⁴	1.90 x 10 ⁻⁰³	1.58 x 10 ⁻⁰²	1.89 x 10 ⁻⁰²
	Polonium-210	9.56 x 10 ⁻⁰⁴	3.59 x 10 ⁻¹³	1.73 x 10 ⁻⁰⁶	5.20 x 10 ⁻¹⁵	2.62 x 10 ⁻⁰⁶	7.77 x 10 ⁻⁰⁸	3.49 x 10 ⁻⁰⁸	4.79 x 10 ⁻¹²	0.00 x 10 ⁺⁰⁰	2.84 x 10 ⁻⁰³	3.04 x 10 ⁻⁰³	1.43 x 10 ⁻⁰²	2.12 x 10 ⁻⁰²
	Total by Pathway	8.20 x 10 ⁻⁰³	3.25 x 10 ⁻¹⁰	6.27 x 10 ⁻⁰⁶	3.21 x 10 ⁻¹⁰	6.01 x 10 ⁻⁰⁶	2.60 x 10 ⁻⁰²	5.88 x 10 ⁻⁰⁸	2.08 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	4.13 x 10 ⁻⁰³	6.43 x 10 ⁻⁰³	3.77 x 10 ⁻⁰²	8.24 x 10 ⁻⁰²
								Application Case						
	Uranium-238	1.30 x 10 ⁻⁰⁴	3.57 x 10 ⁻¹³	8.09 x 10 ⁻⁰⁶	6.14 x 10 ⁻¹¹	2.05 x 10 ⁻⁰⁸	2.27 x 10 ⁻⁰⁴	5.87 x 10 ⁻⁰⁸	6.26 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	2.22 x 10 ⁻⁰⁵	9.36 x 10 ⁻⁰⁵	1.25 x 10 ⁻⁰⁴	6.07 x 10 ⁻⁰⁴
	Uranium-234	1.57 x 10 ⁻⁰⁴	8.74 x 10 ⁻¹³	8.84 x 10 ⁻⁰⁶	1.07 x 10 ⁻¹¹	2.23 x 10 ⁻⁰⁸	1.15 x 10 ⁻⁰⁶	6.42 x 10 ⁻⁰⁸	2.28 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	2.43 x 10 ⁻⁰⁵	1.02 x 10 ⁻⁰⁴	1.36 x 10 ⁻⁰⁴	4.30 x 10 ⁻⁰⁴
	Thorium-230	6.29 x 10 ⁻⁰⁴	2.12 x 10 ⁻¹²	9.97 x 10 ⁻⁰⁴	6.76 x 10 ⁻¹⁰	1.07 x 10 ⁻⁰⁷	1.64 x 10 ⁻⁰⁶	1.02 x 10 ⁻⁰⁶	2.53 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	1.32 x 10 ⁻⁰³	1.02 x 10 ⁻⁰³	1.31 x 10 ⁻⁰²	1.71 x 10 ⁻⁰²
	Radium-226	1.57 x 10 ⁻⁰⁴	4.06 x 10 ⁻¹¹	7.70 x 10 ⁻⁰⁵	3.44 x 10 ⁻⁰⁸	1.45 x 10 ⁻⁰⁷	3.75 x 10 ⁻⁰³	3.40 x 10 ⁻⁰⁷	4.08 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	1.99 x 10 ⁻⁰⁴	1.76 x 10 ⁻⁰³	1.49 x 10 ⁻⁰³	7.47 x 10 ⁻⁰³
	Lead-210	4.94 x 10 ⁻⁰⁵	6.39 x 10 ⁻¹²	2.03 x 10 ⁻⁰⁵	1.78 x 10 ⁻¹¹	2.30 x 10 ⁻⁰⁷	5.34 x 10 ⁻⁰⁵	5.51 x 10 ⁻⁰⁷	3.12 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	1.78 x 10 ⁻⁰⁴	3.80 x 10 ⁻⁰³	1.27 x 10 ⁻⁰²	1.68 x 10 ⁻⁰²
Subsistence	Polonium-210	1.48 x 10 ⁻⁰⁴	5.57 x 10 ⁻¹⁴	3.54 x 10 ⁻⁰⁵	1.06 x 10 ⁻¹³	4.07 x 10 ⁻⁰⁷	1.20 x 10 ⁻⁰⁸	9.74 x 10 ⁻⁰⁷	1.34 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	1.10 x 10 ⁻⁰³	6.07 x 10 ⁻⁰³	1.83 x 10 ⁻⁰²	2.57 x 10 ⁻⁰²
Harvester	Total by Pathway	1.27 x 10 ⁻⁰³	5.04 x 10 ⁻¹¹	1.15 x 10 ⁻⁰³	3.52 x 10 ⁻⁰⁸	9.31 x 10 ⁻⁰⁷	4.03 x 10 ⁻⁰³	3.01 x 10 ⁻⁰⁶	4.15 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	2.84 x 10 ⁻⁰³	1.28 x 10 ⁻⁰²	4.59 x 10 ⁻⁰²	6.80 x 10 ⁻⁰²
(Patterson Lake							U	pper Bound Scena	rio					
South Arm)	Uranium-238	1.30 x 10 ⁻⁰⁴	3.57 x 10 ⁻¹³	2.21 x 10 ⁻⁰⁵	1.68 x 10 ⁻¹⁰	2.05 x 10 ⁻⁰⁸	2.27 x 10 ⁻⁰⁴	1.64 x 10 ⁻⁰⁷	1.74 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	6.09 x 10 ⁻⁰⁵	9.36 x 10 ⁻⁰⁵	1.35 x 10 ⁻⁰⁴	6.71 x 10 ⁻⁰⁴
	Uranium-234	1.57 x 10 ⁻⁰⁴	8.74 x 10 ⁻¹³	2.42 x 10 ⁻⁰⁵	2.94 x 10 ⁻¹¹	2.23 x 10 ⁻⁰⁸	1.15 x 10 ⁻⁰⁶	1.79 x 10 ⁻⁰⁷	6.35 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	6.66 x 10 ⁻⁰⁵	1.02 x 10 ⁻⁰⁴	1.47 x 10 ⁻⁰⁴	4.98 x 10 ⁻⁰⁴
	Thorium-230	6.29 x 10 ⁻⁰⁴	2.12 x 10 ⁻¹²	1.01 x 10 ⁻⁰³	6.85 x 10 ⁻¹⁰	1.07 x 10 ⁻⁰⁷	1.64 x 10 ⁻⁰⁶	1.04 x 10 ⁻⁰⁶	2.56 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	1.34 x 10 ⁻⁰³	1.02 x 10 ⁻⁰³	1.32 x 10 ⁻⁰²	1.72 x 10 ⁻⁰²
	Radium-226	1.57 x 10 ⁻⁰⁴	4.06 x 10 ⁻¹¹	8.05 x 10 ⁻⁰⁵	3.59 x 10 ⁻⁰⁸	1.45 x 10 ⁻⁰⁷	3.75 x 10 ⁻⁰³	3.56 x 10 ⁻⁰⁷	4.27 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	2.08 x 10 ⁻⁰⁴	1.76 x 10 ⁻⁰³	1.49 x 10 ⁻⁰³	7.49 x 10 ⁻⁰³
	Lead-210	4.94 x 10 ⁻⁰⁵	6.39 x 10 ⁻¹²	1.03 x 10 ⁻⁰⁴	9.12 x 10 ⁻¹¹	2.30 x 10 ⁻⁰⁷	5.34 x 10 ⁻⁰⁵	2.60 x 10 ⁻⁰⁶	1.47 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	9.08 x 10 ⁻⁰⁴	3.80 x 10 ⁻⁰³	3.16 x 10 ⁻⁰²	3.65 x 10 ⁻⁰²
	Polonium-210	1.48 x 10 ⁻⁰⁴	5.57 x 10 ⁻¹⁴	1.84 x 10 ⁻⁰⁴	5.53 x 10 ⁻¹³	4.07 x 10 ⁻⁰⁷	1.20 x 10 ⁻⁰⁸	4.59 x 10 ⁻⁰⁶	6.30 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	5.68 x 10 ⁻⁰³	6.07 x 10 ⁻⁰³	2.87 x 10 ⁻⁰²	4.08 x 10 ⁻⁰²
	Total by Pathway	1.27 x 10 ⁻⁰³	5.04 x 10 ⁻¹¹	1.42 x 10 ⁻⁰³	3.69 x 10 ⁻⁰⁸	9.31 x 10 ⁻⁰⁷	4.03 x 10 ⁻⁰³	8.92 x 10 ⁻⁰⁶	4.46 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	8.26 x 10 ⁻⁰³	1.28 x 10 ⁻⁰²	7.52 x 10 ⁻⁰²	1.03 x 10 ⁻⁰¹
		1.20 10-04	4.65 40-12	F 62 40-06	4.52 40-11	0.24 40-07	2.04 40:04	Application Case		0.00 40+00	4.24 40.05	4.4.5 40-04	F. C.A. 40:05	6.20 40.04
	Uranium-238	1.30 x 10 ⁻⁰⁴	4.65 x 10 ⁻¹³	5.62 x 10 ⁻⁰⁶	1.52 x 10 ⁻¹¹	8.34 x 10 ⁻⁰⁷	3.04 x 10 ⁻⁰⁴	2.39 x 10 ⁻⁰⁶	8.14 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.31 x 10 ⁻⁰⁵	1.16 x 10 ⁻⁰⁴	5.64 x 10 ⁻⁰⁵	6.29 x 10 ⁻⁰⁴
	Uranium-234	1.52 x 10 ⁻⁰⁴	1.14 x 10 ⁻¹²	6.11 x 10 ⁻⁰⁶	2.66 x 10 ⁻¹²	9.03 x 10 ⁻⁰⁷	1.53 x 10 ⁻⁰⁶	2.60 x 10 ⁻⁰⁶	2.96 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	1.43 x 10 ⁻⁰⁵	1.25 x 10 ⁻⁰⁴	6.12 x 10 ⁻⁰⁵	3.64 x 10 ⁻⁰⁴
Subsistence	Thorium-230	4.84 x 10 ⁻⁰⁴	2.75 x 10 ⁻¹²	5.07 x 10 ⁻⁰⁴	1.67 x 10 ⁻¹⁰	3.19 x 10 ⁻⁰⁶	1.87 x 10 ⁻⁰⁶	3.04 x 10 ⁻⁰⁵	3.28 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	5.71 x 10 ⁻⁰⁴	5.57 x 10 ⁻⁰⁴	6.32 x 10 ⁻⁰³	8.48 x 10 ⁻⁰³
Harvester One-	Radium-226	1.52 x 10 ⁻⁰⁴	5.27 x 10 ⁻¹¹	6.88 x 10 ⁻⁰⁵	8.51 x 10 ⁻⁰⁹	7.57 x 10 ⁻⁰⁶	5.02 x 10 ⁻⁰³	1.78 x 10 ⁻⁰⁵	5.32 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	1.51 x 10 ⁻⁰⁴	2.00 x 10 ⁻⁰³	6.71 x 10 ⁻⁰⁴	8.14 x 10 ⁻⁰³
Year-Old	Lead-210	5.11 x 10 ⁻⁰⁵	8.29 x 10 ⁻¹²	2.75 x 10 ⁻⁰⁵	3.62 x 10 ⁻¹²	1.83 x 10 ⁻⁰⁵	7.14 x 10 ⁻⁰⁵	4.38 x 10 ⁻⁰⁵	4.05 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.05 x 10 ⁻⁰⁴	6.21 x 10 ⁻⁰³	1.57 x 10 ⁻⁰²	2.24 x 10 ⁻⁰²
(Patterson Lake	Polonium-210	1.52 x 10 ⁻⁰⁴ 1.12 x 10 ⁻⁰³	7.25 x 10 ⁻¹⁴ 6.55 x 10 ⁻¹¹	6.76 x 10 ⁻⁰⁵ 6.83 x 10 ⁻⁰⁴	2.63 x 10 ⁻¹⁴ 8.70 x 10 ⁻⁰⁹	4.55 x 10 ⁻⁰⁵ 7.63 x 10 ⁻⁰⁵	1.57 x 10 ⁻⁰⁸ 5.39 x 10 ⁻⁰³	1.09 x 10 ⁻⁰⁴ 2.06 x 10 ⁻⁰⁴	1.74 x 10 ⁻¹⁰ 5.41 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰ 0.00 x 10 ⁺⁰⁰	1.78 x 10 ⁻⁰³ 2.74 x 10 ⁻⁰³	1.44 x 10 ⁻⁰² 2.34 x 10 ⁻⁰²	1.80 x 10 ⁻⁰² 4.09 x 10 ⁻⁰²	3.46 x 10 ⁻⁰² 7.45 x 10 ⁻⁰²
South Arm)	Total by Pathway	1, 12 X 10 **	0.35 X 10 ··	0.03 X 10	0.70 X 10	7.05 X 10		pper Bound Scena		0.00 x 10 · · ·	2.14 X 10 **	2.34 X IU	4.03 X 10	7.45 X IU
	Uranium-238	1.30 x 10 ⁻⁰⁴	4.65 x 10 ⁻¹³	1.54 x 10 ⁻⁰⁵	4.15 x 10 ⁻¹¹	8.34 x 10 ⁻⁰⁷	3.04 x 10 ⁻⁰⁴	6.65 x 10 ⁻⁰⁶	2.27 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	3.59 x 10 ⁻⁰⁵	1.16 x 10 ⁻⁰⁴	6.11 x 10 ⁻⁰⁵	6.72 x 10 ⁻⁰⁴
	Uranium-234	1.52 x 10 ⁻⁰⁴	1.14 x 10 ⁻¹²	1.54 x 10 1.67 x 10 ⁻⁰⁵	7.27 x 10 ⁻¹²	9.03 x 10 ⁻⁰⁷	1.53 x 10 ⁻⁰⁶	7.23 x 10 ⁻⁰⁶	8.25 x 10 ⁻⁰⁹	0.00 x 10 0.00 x 10+00	3.91 x 10 ⁻⁰⁵	1.16 x 10 1.25 x 10 ⁻⁰⁴	6.62 x 10 ⁻⁰⁵	4.09 x 10 ⁻⁰⁴
	Thorium-230	4.84 x 10 ⁻⁰⁴	2.75 x 10 ⁻¹²	5.14 x 10 ⁻⁰⁴	1.70 x 10 ⁻¹⁰	3.19 x 10 ⁻⁰⁶	1.87 x 10 ⁻⁰⁶	3.08 x 10 ⁻⁰⁵	3.33 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	5.79 x 10 ⁻⁰⁴	5.57 x 10 ⁻⁰⁴	6.37 x 10 ⁻⁰³	8.54 x 10 ⁻⁰³
	Filorium-230	7.04 X 10	2.13 × 10	J. 17 A 10	1.70 x 10	J. 1J A 10	1.07 × 10	J.00 X 10	J.JJ X 10	0.00 X 10	J.13 X 10	J.J1 A 10	0.57 X 10	0.54 × 10

							Incremental Dose b	y Pathway - Proje	ct Lifespan (mSv/y	/r)				
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by Radionuclide
	Radium-226	1.52 x 10 ⁻⁰⁴	5.27 x 10 ⁻¹¹	7.19 x 10 ⁻⁰⁵	8.90 x 10 ⁻⁰⁹	7.57 x 10 ⁻⁰⁶	5.02 x 10 ⁻⁰³	1.86 x 10 ⁻⁰⁵	5.57 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	1.58 x 10 ⁻⁰⁴	2.00 x 10 ⁻⁰³	6.72 x 10 ⁻⁰⁴	8.15 x 10 ⁻⁰³
	Lead-210	5.11 x 10 ⁻⁰⁵	8.29 x 10 ⁻¹²	1.41 x 10 ⁻⁰⁴	1.85 x 10 ⁻¹¹	1.83 x 10 ⁻⁰⁵	7.14 x 10 ⁻⁰⁵	2.07 x 10 ⁻⁰⁴	1.91 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.05 x 10 ⁻⁰³	6.21 x 10 ⁻⁰³	3.94 x 10 ⁻⁰²	4.71 x 10 ⁻⁰²
	Polonium-210	1.52 x 10 ⁻⁰⁴	7.25 x 10 ⁻¹⁴	3.52 x 10 ⁻⁰⁴	1.37 x 10 ⁻¹³	4.55 x 10 ⁻⁰⁵	1.57 x 10 ⁻⁰⁸	5.14 x 10 ⁻⁰⁴	8.19 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	9.22 x 10 ⁻⁰³	1.44 x 10 ⁻⁰²	2.99 x 10 ⁻⁰²	5.45 x 10 ⁻⁰²
	Total by Pathway	1.12 x 10 ⁻⁰³	6.55 x 10 ⁻¹¹	1.11 x 10 ⁻⁰³	9.14 x 10 ⁻⁰⁹	7.63 x 10 ⁻⁰⁵	5.39 x 10 ⁻⁰³	7.84 x 10 ⁻⁰⁴	5.82 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	1.11 x 10 ⁻⁰²	2.34 x 10 ⁻⁰²	7.64 x 10 ⁻⁰²	1.19 x 10 ⁻⁰¹
								Application Case						
	Uranium-238	7.59 x 10 ⁻⁰⁵	2.08 x 10 ⁻¹³	2.98 x 10 ⁻⁰⁶	2.26 x 10 ⁻¹¹	1.19 x 10 ⁻⁰⁸	1.33 x 10 ⁻⁰⁴	2.15 x 10 ⁻⁰⁸	2.30 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	8.18 x 10 ⁻⁰⁶	5.45 x 10 ⁻⁰⁵	4.94 x 10 ⁻⁰⁵	3.24 x 10 ⁻⁰⁴
	Uranium-234	9.16 x 10 ⁻⁰⁵	5.10 x 10 ⁻¹³	3.25 x 10 ⁻⁰⁶	3.95 x 10 ⁻¹²	1.30 x 10 ⁻⁰⁸	6.68 x 10 ⁻⁰⁷	2.35 x 10 ⁻⁰⁸	8.36 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	8.94 x 10 ⁻⁰⁶	5.94 x 10 ⁻⁰⁵	5.38 x 10 ⁻⁰⁵	2.18 x 10 ⁻⁰⁴
	Thorium-230	3.67 x 10 ⁻⁰⁴	1.24 x 10 ⁻¹²	5.24 x 10 ⁻⁰⁴	3.55 x 10 ⁻¹⁰	6.24 x 10 ⁻⁰⁸	9.56 x 10 ⁻⁰⁷	5.37 x 10 ⁻⁰⁷	1.33 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	6.94 x 10 ⁻⁰⁴	5.93 x 10 ⁻⁰⁴	2.56 x 10 ⁻⁰³	4.74 x 10 ⁻⁰³
	Radium-226	9.16 x 10 ⁻⁰⁵	2.36 x 10 ⁻¹¹	3.28 x 10 ⁻⁰⁵	1.46 x 10 ⁻⁰⁸	8.44 x 10 ⁻⁰⁸	2.18 x 10 ⁻⁰³	1.45 x 10 ⁻⁰⁷	1.73 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	8.47 x 10 ⁻⁰⁵	1.03 x 10 ⁻⁰³	3.58 x 10 ⁻⁰⁴	3.80 x 10 ⁻⁰³
	Lead-210	2.88 x 10 ⁻⁰⁵	3.72 x 10 ⁻¹²	2.67 x 10 ⁻⁰⁶	2.35 x 10 ⁻¹²	1.34 x 10 ⁻⁰⁷	3.11 x 10 ⁻⁰⁵	7.27 x 10 ⁻⁰⁸	4.11 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	2.35 x 10 ⁻⁰⁵	2.22 x 10 ⁻⁰³	1.00 x 10 ⁻⁰³	3.31 x 10 ⁻⁰³
	Polonium-210	8.64 x 10 ⁻⁰⁵	3.25 x 10 ⁻¹⁴	4.67 x 10 ⁻⁰⁶	1.40 x 10 ⁻¹⁴	2.37 x 10 ⁻⁰⁷	7.02 x 10 ⁻⁰⁹	1.28 x 10 ⁻⁰⁷	1.76 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	1.45 x 10 ⁻⁰⁴	3.54 x 10 ⁻⁰³	5.68 x 10 ⁻⁰³	9.45 x 10 ⁻⁰³
Subsistence	Total by Pathway	7.41 x 10 ⁻⁰⁴	2.94 x 10 ⁻¹¹	5.71 x 10 ⁻⁰⁴	1.50 x 10 ⁻⁰⁸	5.43 x 10 ⁻⁰⁷	2.35 x 10 ⁻⁰³	9.28 x 10 ⁻⁰⁷	1.76 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	9.64 x 10 ⁻⁰⁴	7.49 x 10 ⁻⁰³	9.70 x 10 ⁻⁰³	2.18 x 10 ⁻⁰²
Harvester (Beet Lake)							U	pper Bound Scena	rio					
,	Uranium-238	7.59 x 10 ⁻⁰⁵	2.08 x 10 ⁻¹³	8.14 x 10 ⁻⁰⁶	6.18 x 10 ⁻¹¹	1.19 x 10 ⁻⁰⁸	1.33 x 10 ⁻⁰⁴	6.00 x 10 ⁻⁰⁸	6.40 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	2.24 x 10 ⁻⁰⁵	5.45 x 10 ⁻⁰⁵	5.23 x 10 ⁻⁰⁵	3.47 x 10 ⁻⁰⁴
	Uranium-234	9.16 x 10 ⁻⁰⁵	5.10 x 10 ⁻¹³	8.90 x 10 ⁻⁰⁶	1.08 x 10 ⁻¹¹	1.30 x 10 ⁻⁰⁸	6.68 x 10 ⁻⁰⁷	6.56 x 10 ⁻⁰⁸	2.33 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	2.45 x 10 ⁻⁰⁵	5.94 x 10 ⁻⁰⁵	5.70 x 10 ⁻⁰⁵	2.42 x 10 ⁻⁰⁴
	Thorium-230	3.67 x 10 ⁻⁰⁴	1.24 x 10 ⁻¹²	5.31 x 10 ⁻⁰⁴	3.60 x 10 ⁻¹⁰	6.24 x 10 ⁻⁰⁸	9.56 x 10 ⁻⁰⁷	5.45 x 10 ⁻⁰⁷	1.35 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	7.04 x 10 ⁻⁰⁴	5.93 x 10 ⁻⁰⁴	2.59 x 10 ⁻⁰³	4.78 x 10 ⁻⁰³
	Radium-226	9.16 x 10 ⁻⁰⁵	2.36 x 10 ⁻¹¹	3.43 x 10 ⁻⁰⁵	1.53 x 10 ⁻⁰⁸	8.44 x 10 ⁻⁰⁸	2.18 x 10 ⁻⁰³	1.51 x 10 ⁻⁰⁷	1.81 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	8.85 x 10 ⁻⁰⁵	1.03 x 10 ⁻⁰³	3.59 x 10 ⁻⁰⁴	3.80 x 10 ⁻⁰³
	Lead-210	2.88 x 10 ⁻⁰⁵	3.72 x 10 ⁻¹²	1.34 x 10 ⁻⁰⁵	1.18 x 10 ⁻¹¹	1.34 x 10 ⁻⁰⁷	3.11 x 10 ⁻⁰⁵	3.54 x 10 ⁻⁰⁷	2.00 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	1.18 x 10 ⁻⁰⁴	2.22 x 10 ⁻⁰³	1.15 x 10 ⁻⁰³	3.55 x 10 ⁻⁰³
	Polonium-210	8.64 x 10 ⁻⁰⁵	3.25 x 10 ⁻¹⁴	2.38 x 10 ⁻⁰⁵	7.16 x 10 ⁻¹⁴	2.37 x 10 ⁻⁰⁷	7.02 x 10 ⁻⁰⁹	6.27 x 10 ⁻⁰⁷	8.60 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	7.37 x 10 ⁻⁰⁴	3.54 x 10 ⁻⁰³	6.66 x 10 ⁻⁰³	1.10 x 10 ⁻⁰²
	Total by Pathway	7.41 x 10 ⁻⁰⁴	2.94 x 10 ⁻¹¹	6.20 x 10 ⁻⁰⁴	1.57 x 10 ⁻⁰⁸	5.43 x 10 ⁻⁰⁷	2.35 x 10 ⁻⁰³	1.80 x 10 ⁻⁰⁶	1.88 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	1.69 x 10 ⁻⁰³	7.49 x 10 ⁻⁰³	1.09 x 10 ⁻⁰²	2.38 x 10 ⁻⁰²
								Application Case						
	Uranium-238	7.57 x 10 ⁻⁰⁵	2.71 x 10 ⁻¹³	2.07 x 10 ⁻⁰⁶	5.58 x 10 ⁻¹²	4.86 x 10 ⁻⁰⁷	1.77 x 10 ⁻⁰⁴	8.76 x 10 ⁻⁰⁷	2.99 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	4.83 x 10 ⁻⁰⁶	6.73 x 10 ⁻⁰⁵	2.19 x 10 ⁻⁰⁵	3.51 x 10 ⁻⁰⁴
	Uranium-234	8.86 x 10 ⁻⁰⁵	6.62 x 10 ⁻¹³	2.25 x 10 ⁻⁰⁶	9.78 x 10 ⁻¹³	5.26 x 10 ⁻⁰⁷	8.94 x 10 ⁻⁰⁷	9.53 x 10 ⁻⁰⁷	1.09 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	5.25 x 10 ⁻⁰⁶	7.29 x 10 ⁻⁰⁵	2.37 x 10 ⁻⁰⁵	1.95 x 10 ⁻⁰⁴
	Thorium-230	2.82 x 10 ⁻⁰⁴	1.61 x 10 ⁻¹²	2.67 x 10 ⁻⁰⁴	8.80 x 10 ⁻¹¹	1.86 x 10 ⁻⁰⁶	1.09 x 10 ⁻⁰⁶	1.60 x 10 ⁻⁰⁵	1.73 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	3.00 x 10 ⁻⁰⁴	3.24 x 10 ⁻⁰⁴	1.23 x 10 ⁻⁰³	2.42 x 10 ⁻⁰³
	Radium-226	8.86 x 10 ⁻⁰⁵	3.07 x 10 ⁻¹¹	2.93 x 10 ⁻⁰⁵	3.63 x 10 ⁻⁰⁹	4.41 x 10 ⁻⁰⁶	2.92 x 10 ⁻⁰³	7.57 x 10 ⁻⁰⁶	2.26 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	6.42 x 10 ⁻⁰⁵	1.17 x 10 ⁻⁰³	1.38 x 10 ⁻⁰⁴	4.44 x 10 ⁻⁰³
	Lead-210	2.98 x 10 ⁻⁰⁵	4.83 x 10 ⁻¹²	3.63 x 10 ⁻⁰⁶	4.78 x 10 ⁻¹³	1.07 x 10 ⁻⁰⁵	4.16 x 10 ⁻⁰⁵	5.78 x 10 ⁻⁰⁶	5.34 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	2.71 x 10 ⁻⁰⁵	3.62 x 10 ⁻⁰³	1.13 x 10 ⁻⁰³	4.87 x 10 ⁻⁰³
Subsistence	Polonium-210	8.86 x 10 ⁻⁰⁵	4.22 x 10 ⁻¹⁴	8.93 x 10 ⁻⁰⁶	3.48 x 10 ⁻¹⁵	2.65 x 10 ⁻⁰⁵	9.18 x 10 ⁻⁰⁹	1.44 x 10 ⁻⁰⁵	2.29 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	2.35 x 10 ⁻⁰⁴	8.37 x 10 ⁻⁰³	5.76 x 10 ⁻⁰³	1.45 x 10 ⁻⁰²
Harvester One-	Total by Pathway	6.53 x 10 ⁻⁰⁴	3.81 x 10 ⁻¹¹	3.13 x 10 ⁻⁰⁴	3.72 x 10 ⁻⁰⁹	4.44 x 10 ⁻⁰⁵	3.14 x 10 ⁻⁰³	4.55 x 10 ⁻⁰⁵	2.29 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	6.36 x 10 ⁻⁰⁴	1.36 x 10 ⁻⁰²	8.31 x 10 ⁻⁰³	2.68 x 10 ⁻⁰²
Year-Old (Beet								pper Bound Scena					ı	
Lake)	Uranium-238	7.57 x 10 ⁻⁰⁵	2.71 x 10 ⁻¹³	5.66 x 10 ⁻⁰⁶	1.53 x 10 ⁻¹¹	4.86 x 10 ⁻⁰⁷	1.77 x 10 ⁻⁰⁴	2.44 x 10 ⁻⁰⁶	8.32 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.32 x 10 ⁻⁰⁵	6.73 x 10 ⁻⁰⁵	2.32 x 10 ⁻⁰⁵	3.66 x 10 ⁻⁰⁴
	Uranium-234	8.86 x 10 ⁻⁰⁵	6.62 x 10 ⁻¹³	6.15 x 10 ⁻⁰⁶	2.68 x 10 ⁻¹²	5.26 x 10 ⁻⁰⁷	8.94 x 10 ⁻⁰⁷	2.65 x 10 ⁻⁰⁶	3.03 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	1.44 x 10 ⁻⁰⁵	7.29 x 10 ⁻⁰⁵	2.51 x 10 ⁻⁰⁵	2.11 x 10 ⁻⁰⁴
	Thorium-230	2.82 x 10 ⁻⁰⁴	1.61 x 10 ⁻¹²	2.70 x 10 ⁻⁰⁴	8.92 x 10 ⁻¹¹	1.86 x 10 ⁻⁰⁶	1.09 x 10 ⁻⁰⁶	1.62 x 10 ⁻⁰⁵	1.75 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	3.04 x 10 ⁻⁰⁴	3.24 x 10 ⁻⁰⁴	1.24 x 10 ⁻⁰³	2.44 x 10 ⁻⁰³
	Radium-226	8.86 x 10 ⁻⁰⁵	3.07 x 10 ⁻¹¹	3.06 x 10 ⁻⁰⁵	3.79 x 10 ⁻⁰⁹	4.41 x 10 ⁻⁰⁶	2.92 x 10 ⁻⁰³	7.92 x 10 ⁻⁰⁶	2.37 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	6.71 x 10 ⁻⁰⁵	1.17 x 10 ⁻⁰³	1.38 x 10 ⁻⁰⁴	4.45 x 10 ⁻⁰³
	Lead-210	2.98 x 10 ⁻⁰⁵	4.83 x 10 ⁻¹²	1.83 x 10 ⁻⁰⁵	2.40 x 10 ⁻¹²	1.07 x 10 ⁻⁰⁵	4.16 x 10 ⁻⁰⁵	2.82 x 10 ⁻⁰⁵	2.60 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	1.36 x 10 ⁻⁰⁴	3.62 x 10 ⁻⁰³	1.30 x 10 ⁻⁰³	5.18 x 10 ⁻⁰³
	Polonium-210	8.86 x 10 ⁻⁰⁵	4.22 x 10 ⁻¹⁴	4.55 x 10 ⁻⁰⁵	1.77 x 10 ⁻¹⁴	2.65 x 10 ⁻⁰⁵	9.18 x 10 ⁻⁰⁹	7.01 x 10 ⁻⁰⁵	1.12 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	1.20 x 10 ⁻⁰³	8.37 x 10 ⁻⁰³	7.01 x 10 ⁻⁰³	1.68 x 10 ⁻⁰²
	Total by Pathway	6.53 x 10 ⁻⁰⁴	3.81 x 10 ⁻¹¹	3.77 x 10 ⁻⁰⁴	3.90 x 10 ⁻⁰⁹	4.44 x 10 ⁻⁰⁵	3.14 x 10 ⁻⁰³	1.27 x 10 ⁻⁰⁴	2.46 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	1.73 x 10 ⁻⁰³	1.36 x 10 ⁻⁰²	9.73 x 10 ⁻⁰³	2.94 x 10 ⁻⁰²
	11 - 1 - 220	2.54 40-06	6.07 10-15	2.72 40-07	2.06 10:12	4.00 4.0-10	4.42 4.0-06	Application Case		0.00 10+00	7.47 40-07	1.02 10:06	1.75 10-06	1.16 10:05
	Uranium-238	2.54 x 10 ⁻⁰⁶	6.97 x 10 ⁻¹⁵	2.72 x 10 ⁻⁰⁷	2.06 x 10 ⁻¹²	4.00 x 10 ⁻¹⁰	4.43 x 10 ⁻⁰⁶	1.96 x 10 ⁻⁰⁹	2.09 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	7.47 x 10 ⁻⁰⁷	1.82 x 10 ⁻⁰⁶	1.75 x 10 ⁻⁰⁶	1.16 x 10 ⁻⁰⁵
	Uranium-234	3.07 x 10 ⁻⁰⁶	1.70 x 10 ⁻¹⁴	2.97 x 10 ⁻⁰⁷	3.61 x 10 ⁻¹³	4.35 x 10 ⁻¹⁰	2.23 x 10 ⁻⁰⁸	2.14 x 10 ⁻⁰⁹	7.59 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	8.16 x 10 ⁻⁰⁷	1.99 x 10 ⁻⁰⁶	1.90 x 10 ⁻⁰⁶	8.09 x 10 ⁻⁰⁶
Subsistence	Thorium-230	1.23 x 10 ⁻⁰⁵	4.13 x 10 ⁻¹⁴	5.40 x 10 ⁻⁰⁵	3.66 x 10 ⁻¹¹	2.09 x 10 ⁻⁰⁹	3.20 x 10 ⁻⁰⁸	5.53 x 10 ⁻⁰⁸	1.37 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	7.14 x 10 ⁻⁰⁵	1.98 x 10 ⁻⁰⁵	2.17 x 10 ⁻⁰⁴	3.75 x 10 ⁻⁰⁴
Harvester (Lloyd	Radium-226	3.07 x 10 ⁻⁰⁶	7.91 x 10 ⁻¹³	3.17 x 10 ⁻⁰⁶	1.41 x 10 ⁻⁰⁹	2.82 x 10 ⁻⁰⁹	7.31 x 10 ⁻⁰⁵	1.39 x 10 ⁻⁰⁸	1.67 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	8.19 x 10 ⁻⁰⁶	3.44 x 10 ⁻⁰⁵	1.31 x 10 ⁻⁰⁵	1.37 x 10 ⁻⁰⁴
Lake)	Lead-210	9.63 x 10 ⁻⁰⁷	1.25 x 10 ⁻¹³	1.40 x 10 ⁻⁰⁷	1.24 x 10 ⁻¹³	4.48 x 10 ⁻⁰⁹	1.04 x 10 ⁻⁰⁶	3.64 x 10 ⁻¹⁰	2.07 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	1.20 x 10 ⁻⁰⁶	7.41 x 10 ⁻⁰⁵	3.41 x 10 ⁻⁰⁵	1.12 x 10 ⁻⁰⁴
	Polonium-210	2.89 x 10 ⁻⁰⁶	1.09 x 10 ⁻¹⁵	2.08 x 10 ⁻⁰⁷	6.26 x 10 ⁻¹⁶	7.92 x 10 ⁻⁰⁹	2.35 x 10 ⁻¹⁰	5.68 x 10 ⁻¹⁰	7.82 x 10 ⁻¹⁴	0.00 x 10 ⁺⁰⁰	6.12 x 10 ⁻⁰⁶	1.18 x 10 ⁻⁰⁴	1.83 x 10 ⁻⁰⁴	3.11 x 10 ⁻⁰⁴
	Total by Pathway	2.48 x 10 ⁻⁰⁵	9.82 x 10 ⁻¹³	5.81 x 10 ⁻⁰⁵	1.45 x 10 ⁻⁰⁹	1.82 x 10 ⁻⁰⁸	7.86 x 10 ⁻⁰⁵	7.42 x 10 ⁻⁰⁸	1.69 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	8.85 x 10 ⁻⁰⁵	2.50 x 10 ⁻⁰⁴	4.51 x 10 ⁻⁰⁴	9.53 x 10 ⁻⁰⁴
	Hranium 220	2.54 x 10 ⁻⁰⁶	6.97 x 10 ⁻¹⁵	7.48 x 10 ⁻⁰⁷	5.68 x 10 ⁻¹²	4.00 x 10 ⁻¹⁰	4.43 x 10 ⁻⁰⁶	pper Bound Scena 5.45 x 10 ⁻⁰⁹		0.00 x 10 ⁺⁰⁰	2.06 x 10 ⁻⁰⁶	1.82 x 10 ⁻⁰⁶	2.01 x 10 ⁻⁰⁶	1.37 x 10 ⁻⁰⁵
	Uranium-238	2.54 X 10 °°	0.91 X 1U 'S	7.40 X IU "	5.00 X IU	4.00 X 10 ···	4.45 X 10 °°	5.45 X IU **	5.81 x 10 ⁻⁰⁸	0.00 X 10 · ° °	2.00 X 10 °°	1.02 X 1U **	2.01 X 10 °°	1.57 X 10 °

							Incremental Dose b	y Pathway - Proje	ct Lifespan (mSv/y	r)				
	Radionuclide	Air (Internal)	Air (External)	Water	Water	Soil (Internal)	Soil (External)	Sediment	Sediment	Aquatic	Aquatic	Terrestrial	Terrestrial	Total by
				(Internal)	(External)			(Internal)	(External)	Plants	Animals	Plants	Animals	Radionuclide
	Uranium-234	3.07 x 10 ⁻⁰⁶	1.70 x 10 ⁻¹⁴	8.18 x 10 ⁻⁰⁷	9.93 x 10 ⁻¹³	4.35 x 10 ⁻¹⁰	2.23 x 10 ⁻⁰⁸	5.96 x 10 ⁻⁰⁹	2.11 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	2.25 x 10 ⁻⁰⁶	1.99 x 10 ⁻⁰⁶	2.19 x 10 ⁻⁰⁶	1.03 x 10 ⁻⁰⁵
	Thorium-230	1.23 x 10 ⁻⁰⁵	4.13 x 10 ⁻¹⁴	5.48 x 10 ⁻⁰⁵	3.72 x 10 ⁻¹¹	2.09 x 10 ⁻⁰⁹	3.20 x 10 ⁻⁰⁸	5.60 x 10 ⁻⁰⁸	1.39 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	7.25 x 10 ⁻⁰⁵	1.98 x 10 ⁻⁰⁵	2.20 x 10 ⁻⁰⁴	3.80 x 10 ⁻⁰⁴
	Radium-226	3.07 x 10 ⁻⁰⁶	7.91 x 10 ⁻¹³	3.31 x 10 ⁻⁰⁶	1.48 x 10 ⁻⁰⁹	2.82 x 10 ⁻⁰⁹	7.31 x 10 ⁻⁰⁵	1.46 x 10 ⁻⁰⁸	1.74 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	8.56 x 10 ⁻⁰⁶	3.44 x 10 ⁻⁰⁵	1.32 x 10 ⁻⁰⁵	1.37 x 10 ⁻⁰⁴
	Lead-210	9.63 x 10 ⁻⁰⁷	1.25 x 10 ⁻¹³	6.89 x 10 ⁻⁰⁷	6.07 x 10 ⁻¹³	4.48 x 10 ⁻⁰⁹	1.04 x 10 ⁻⁰⁶	1.48 x 10 ⁻⁰⁸	8.39 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	6.02 x 10 ⁻⁰⁶	7.41 x 10 ⁻⁰⁵	4.14 x 10 ⁻⁰⁵	1.24 x 10 ⁻⁰⁴
	Polonium-210	2.89 x 10 ⁻⁰⁶	1.09 x 10 ⁻¹⁵	1.18 x 10 ⁻⁰⁶	3.54 x 10 ⁻¹⁵	7.92 x 10 ⁻⁰⁹	2.35 x 10 ⁻¹⁰	2.62 x 10 ⁻⁰⁸	3.59 x 10 ⁻¹²	0.00 x 10 ⁺⁰⁰	3.62 x 10 ⁻⁰⁵	1.18 x 10 ⁻⁰⁴	2.33 x 10 ⁻⁰⁴	3.92 x 10 ⁻⁰⁴
	Total by Pathway	2.48 x 10 ⁻⁰⁵	9.82 x 10 ⁻¹³	6.16 x 10 ⁻⁰⁵	1.52 x 10 ⁻⁰⁹	1.82 x 10 ⁻⁰⁸	7.86 x 10 ⁻⁰⁵	1.23 x 10 ⁻⁰⁷	1.80 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	1.28 x 10 ⁻⁰⁴	2.50 x 10 ⁻⁰⁴	5.12 x 10 ⁻⁰⁴	1.06 x 10 ⁻⁰³
	Uranium-238	2.53 x 10 ⁻⁰⁶	9.06 x 10 ⁻¹⁵	1.89 x 10 ⁻⁰⁷	5.10 x 10 ⁻¹³	1.63 x 10 ⁻⁰⁸	5.93 x 10 ⁻⁰⁶	Application Case 7.95 x 10 ⁻⁰⁸	2.71 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	4.41 x 10 ⁻⁰⁷	2.25 x 10 ⁻⁰⁶	7.75 x 10 ⁻⁰⁷	1.22 x 10 ⁻⁰⁵
	Uranium-234	2.96 x 10 ⁻⁰⁶	2.22 x 10 ⁻¹⁴	2.05 x 10 ⁻⁰⁷	8.93 x 10 ⁻¹⁴	1.76 x 10 ⁻⁰⁸	2.99 x 10 ⁻⁰⁸	8.65 x 10 ⁻⁰⁸	9.87 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	4.79 x 10 ⁻⁰⁷	2.44 x 10 ⁻⁰⁶	8.40 x 10 ⁻⁰⁷	7.06 x 10 ⁻⁰⁶
	Thorium-230	9.43 x 10 ⁻⁰⁶	5.37 x 10 ⁻¹⁴	2.75 x 10 ⁻⁰⁵	9.07 x 10 ⁻¹²	6.22 x 10 ⁻⁰⁸	3.64 x 10 ⁻⁰⁸	1.65 x 10 ⁻⁰⁶	1.78 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	3.09 x 10 ⁻⁰⁵	1.09 x 10 ⁻⁰⁵	1.05 x 10 ⁻⁰⁴	1.85 x 10 ⁻⁰⁴
	Radium-226	2.96 x 10 ⁻⁰⁶	1.03 x 10 ⁻¹²	2.83 x 10 ⁻⁰⁶	3.50 x 10 ⁻¹⁰	1.48 x 10 ⁻⁰⁷	9.78 x 10 ⁻⁰⁵	7.27 x 10 ⁻⁰⁷	2.17 x 10 ⁻⁰⁶	0.00 x 10 0.00 x 10+00	6.21 x 10 ⁻⁰⁶	3.90 x 10 ⁻⁰⁵	5.07 x 10 ⁻⁰⁶	1.57 x 10 ⁻⁰⁴
	Lead-210	9.97 x 10 ⁻⁰⁷	1.62 x 10 ⁻¹³	1.91 x 10 ⁻⁰⁷	2.51 x 10 ⁻¹⁴	3.57 x 10 ⁻⁰⁷	1.39 x 10 ⁻⁰⁶	2.89 x 10 ⁻⁰⁸	2.68 x 10 ⁻¹¹	0.00 x 10 0.00 x 10+00	1.38 x 10 ⁻⁰⁶	1.21 x 10 ⁻⁰⁴	3.83 x 10 ⁻⁰⁵	1.64 x 10 ⁻⁰⁴
	Polonium-210	2.96 x 10 ⁻⁰⁶	1.41 x 10 ⁻¹⁵	3.98 x 10 ⁻⁰⁷	1.55 x 10 ⁻¹⁶	8.86 x 10 ⁻⁰⁷	3.07 x 10 ⁻¹⁰	6.24 x 10 ⁻⁰⁸	9.95 x 10 ⁻¹⁴	0.00 x 10 0.00 x 10+00	9.94 x 10 ⁻⁰⁶	2.80 x 10 ⁻⁰⁴	1.84 x 10 ⁻⁰⁴	4.78 x 10 ⁻⁰⁴
Subsistence	Total by Pathway	2.18 x 10 ⁻⁰⁵	1.28 x 10 ⁻¹²	3.13 x 10 ⁻⁰⁵	3.60 x 10 ⁻¹⁰	1.49 x 10 ⁻⁰⁶	1.05 x 10 ⁻⁰⁴	2.63 x 10 ⁻⁰⁶	2.20 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	4.93 x 10 ⁻⁰⁵	4.55 x 10 ⁻⁰⁴	3.33 x 10 ⁻⁰⁴	1.00 x 10 ⁻⁰³
Harvester One- Year-Old (Lloyd	Total by Fathway	2.10 X 10 **	1.26 X 10	5.15 X 10 **	5.00 X 10	1.49 X 10		pper Bound Scena		0.00 x 10 **	4.95 X 10 **	4.55 X 10 °	5.55 X 10 °	1.00 x 10 **
Lake)	Uranium-238	2.53 x 10 ⁻⁰⁶	9.06 x 10 ⁻¹⁵	5.20 x 10 ⁻⁰⁷	1.40 x 10 ⁻¹²	1.63 x 10 ⁻⁰⁸	5.93 x 10 ⁻⁰⁶	2.22 x 10 ⁻⁰⁷	7.55 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	1.21 x 10 ⁻⁰⁶	2.25 x 10 ⁻⁰⁶	8.95 x 10 ⁻⁰⁷	1.37 x 10 ⁻⁰⁵
	Uranium-234	2.96 x 10 ⁻⁰⁶	2.22 x 10 ⁻¹⁴	5.65 x 10 ⁻⁰⁷	2.46 x 10 ⁻¹³	1.76 x 10 ⁻⁰⁸	2.99 x 10 ⁻⁰⁸	2.41 x 10 ⁻⁰⁷	2.75 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	1.32 x 10 ⁻⁰⁶	2.44 x 10 ⁻⁰⁶	9.70 x 10 ⁻⁰⁷	8.55 x 10 ⁻⁰⁶
	Thorium-230	9.43 x 10 ⁻⁰⁶	5.37 x 10 ⁻¹⁴	2.79 x 10 ⁻⁰⁵	9.20 x 10 ⁻¹²	6.22 x 10 ⁻⁰⁸	3.64 x 10 ⁻⁰⁸	1.67 x 10 ⁻⁰⁶	1.80 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	3.13 x 10 ⁻⁰⁵	1.09 x 10 ⁻⁰⁵	1.06 x 10 ⁻⁰⁴	1.87 x 10 ⁻⁰⁴
	Radium-226	2.96 x 10 ⁻⁰⁶	1.03 x 10 ⁻¹²	2.79 x 10 ⁻⁰⁶	3.66 x 10 ⁻¹⁰	1.48 x 10 ⁻⁰⁷	9.78 x 10 ⁻⁰⁵	7.61 x 10 ⁻⁰⁷	2.28 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	6.50 x 10 ⁻⁰⁶	3.90 x 10 ⁻⁰⁵	5.11 x 10 ⁻⁰⁶	1.57 x 10 ⁻⁰⁴
	Lead-210	9.97 x 10 ⁻⁰⁷	1.62 x 10 ⁻¹³	9.37 x 10 ⁻⁰⁷	1.23 x 10 ⁻¹³	3.57 x 10 ⁻⁰⁷	1.39 x 10 ⁻⁰⁶	1.18 x 10 ⁻⁰⁶	1.09 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	6.95 x 10 ⁻⁰⁶	1.21 x 10 ⁻⁰⁴	4.70 x 10 ⁻⁰⁵	1.80 x 10 ⁻⁰⁴
	Polonium-210	2.96 x 10 ⁻⁰⁶	1.41 x 10 ⁻¹⁵	2.25 x 10 ⁻⁰⁶	8.77 x 10 ⁻¹⁶	8.86 x 10 ⁻⁰⁷	3.07 x 10 ⁻¹⁰	2.93 x 10 ⁻⁰⁶	4.66 x 10 ⁻¹²	0.00 x 10 ⁺⁰⁰	5.87 x 10 ⁻⁰⁵	2.80 x 10 ⁻⁰⁴	2.48 x 10 ⁻⁰⁴	5.95 x 10 ⁻⁰⁴
		2.18 x 10 ⁻⁰⁵	1.28 x 10 ⁻¹²	3.51 x 10 ⁻⁰⁵	3.77 x 10 ⁻¹⁰	1.49 x 10 ⁻⁰⁶	1.05 x 10 ⁻⁰⁴	7.00 x 10 ⁻⁰⁶	2.35 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	1.06 x 10 ⁻⁰⁴	4.55 x 10 ⁻⁰⁴	4.08 x 10 ⁻⁰⁴	1.14 x 10 ⁻⁰³
	Total by Pathway	2.10 X 10	1.20 X 10	3.31 X 10	3.77 X 10	1.49 X 10	1.03 X 10	Application Case		0.00 x 10	1.00 x 10	4.55 X 10	4.00 X 10	1.14 x 10
	Uranium-238	7.82 x 10 ⁻⁰⁵	2.14 x 10 ⁻¹³	4.85 x 10 ⁻⁰⁶	3.68 x 10 ⁻¹¹	1.23 x 10 ⁻⁰⁸	1.36 x 10 ⁻⁰⁴	3.52 x 10 ⁻⁰⁸	3.76 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	4.23 x 10 ⁻⁰⁶	4.44 x 10 ⁻⁰⁵	2.33 x 10 ⁻⁰⁵	2.92 x 10 ⁻⁰⁴
	Uranium-234	9.43 x 10 ⁻⁰⁵	5.25 x 10 ⁻¹³	5.31 x 10 ⁻⁰⁶	6.44 x 10 ⁻¹²	1.34 x 10 ⁻⁰⁸	6.88 x 10 ⁻⁰⁷	3.85 x 10 ⁻⁰⁸	1.37 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	4.62 x 10 ⁻⁰⁶	4.84 x 10 ⁻⁰⁵	2.54 x 10 ⁻⁰⁵	1.79 x 10 ⁻⁰⁴
	Thorium-230	3.77 x 10 ⁻⁰⁴	1.27 x 10 ⁻¹²	5.98 x 10 ⁻⁰⁴	4.06 x 10 ⁻¹⁰	6.43 x 10 ⁻⁰⁸	9.84 x 10 ⁻⁰⁷	6.13 x 10 ⁻⁰⁷	1.52 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.51 x 10 ⁻⁰⁴	2.14 x 10 ⁻⁰⁴	3.05 x 10 ⁻⁰³	4.50 x 10 ⁻⁰³
	Radium-226	9.43 x 10 ⁻⁰⁵	2.43 x 10 ⁻¹¹	4.62 x 10 ⁻⁰⁵	2.06 x 10 ⁻⁰⁸	8.68 x 10 ⁻⁰⁸	2.25 x 10 ⁻⁰³	2.04 x 10 ⁻⁰⁷	2.45 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	3.79 x 10 ⁻⁰⁵	5.00 x 10 ⁻⁰⁴	2.87 x 10 ⁻⁰⁴	3.24 x 10 ⁻⁰³
	Lead-210	2.97 x 10 ⁻⁰⁵	3.83 x 10 ⁻¹²	1.21 x 10 ⁻⁰⁵	1.07 x 10 ⁻¹¹	1.38 x 10 ⁻⁰⁷	3.20 x 10 ⁻⁰⁵	3.29 x 10 ⁻⁰⁷	1.86 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	3.38 x 10 ⁻⁰⁵	9.79 x 10 ⁻⁰⁴	2.92 × 10 ⁻⁰³	4.01 x 10 ⁻⁰³
	Polonium-210	8.90 x 10 ⁻⁰⁵	3.34 x 10 ⁻¹⁴	2.12 x 10 ⁻⁰⁵	6.38 x 10 ⁻¹⁴	2.44 x 10 ⁻⁰⁷	7.23 x 10 ⁻⁰⁹	5.82 x 10 ⁻⁰⁷	7.98 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	2.09 x 10 ⁻⁰⁴	1.65 x 10 ⁻⁰³	3.53 x 10 ⁻⁰³	5.50 x 10 ⁻⁰³
Seasonal Resident	Total by Pathway	7.63 x 10 ⁻⁰⁴	3.02 x 10 ⁻¹¹	6.88 x 10 ⁻⁰⁴	2.11 x 10 ⁻⁰⁸	5.59 x 10 ⁻⁰⁷	2.42 x 10 ⁻⁰³	1.80 x 10 ⁻⁰⁶	2.49 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	5.40 x 10 ⁻⁰⁴	3.43 x 10 ⁻⁰³	9.84 x 10 ⁻⁰³	1.77 x 10 ⁻⁰²
(Patterson Lake	, ,							1.00 X 10	L.73 X 10					
South Arm								pper Bound Scena						
South Arm)	Uranium-238	7.82 x 10 ⁻⁰⁵	2.14 x 10 ⁻¹³	1.33 x 10 ⁻⁰⁵	1.01 x 10 ⁻¹⁰	1.23 x 10 ⁻⁰⁸			rio	0.00 x 10 ⁺⁰⁰	1.16 x 10 ⁻⁰⁵	4.44 x 10 ⁻⁰⁵	2.60 x 10 ⁻⁰⁵	3.11 x 10 ⁻⁰⁴
South Arm)	Uranium-238 Uranium-234	7.82 x 10 ⁻⁰⁵ 9.43 x 10 ⁻⁰⁵	2.14 x 10 ⁻¹³ 5.25 x 10 ⁻¹³	1.33 x 10 ⁻⁰⁵ 1.45 x 10 ⁻⁰⁵	1.01 x 10 ⁻¹⁰ 1.76 x 10 ⁻¹¹	1.23 x 10 ⁻⁰⁸ 1.34 x 10 ⁻⁰⁸	UĮ	per Bound Scena			1.16 x 10 ⁻⁰⁵ 1.27 x 10 ⁻⁰⁵	4.44 x 10 ⁻⁰⁵ 4.84 x 10 ⁻⁰⁵	2.60 x 10 ⁻⁰⁵ 2.83 x 10 ⁻⁰⁵	3.11 x 10 ⁻⁰⁴ 1.99 x 10 ⁻⁰⁴
South Arm)						+	Մլ 1.36 x 10 ⁻⁰⁴	9.81 x 10 ⁻⁰⁸	rio 1.05 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰				
South Arm)	Uranium-234	9.43 x 10 ⁻⁰⁵	5.25 x 10 ⁻¹³	1.45 x 10 ⁻⁰⁵	1.76 x 10 ⁻¹¹	1.34 x 10 ⁻⁰⁸	1.36 x 10 ⁻⁰⁴ 6.88 x 10 ⁻⁰⁷	9.81 x 10 ⁻⁰⁸ 1.07 x 10 ⁻⁰⁷	1.05 x 10 ⁻⁰⁶ 3.81 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰ 0.00 x 10 ⁺⁰⁰	1.27 x 10 ⁻⁰⁵	4.84 x 10 ⁻⁰⁵	2.83 x 10 ⁻⁰⁵	1.99 x 10 ⁻⁰⁴
South Arm)	Uranium-234 Thorium-230	9.43 x 10 ⁻⁰⁵ 3.77 x 10 ⁻⁰⁴	5.25 x 10 ⁻¹³ 1.27 x 10 ⁻¹²	1.45 x 10 ⁻⁰⁵ 6.06 x 10 ⁻⁰⁴	1.76 x 10 ⁻¹¹ 4.11 x 10 ⁻¹⁰	1.34 x 10 ⁻⁰⁸ 6.43 x 10 ⁻⁰⁸	1.36 x 10 ⁻⁰⁴ 6.88 x 10 ⁻⁰⁷ 9.84 x 10 ⁻⁰⁷	9.81 x 10 ⁻⁰⁸ 1.07 x 10 ⁻⁰⁷ 6.22 x 10 ⁻⁰⁷	1.05 x 10 ⁻⁰⁶ 3.81 x 10 ⁻⁰⁹ 1.54 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰ 0.00 x 10 ⁺⁰⁰ 0.00 x 10 ⁺⁰⁰	1.27 x 10 ⁻⁰⁵ 2.55 x 10 ⁻⁰⁴	4.84 x 10 ⁻⁰⁵ 2.14 x 10 ⁻⁰⁴	2.83 x 10 ⁻⁰⁵ 3.08 x 10 ⁻⁰³	1.99 x 10 ⁻⁰⁴ 4.53 x 10 ⁻⁰³
South Arm)	Uranium-234 Thorium-230 Radium-226	9.43 x 10 ⁻⁰⁵ 3.77 x 10 ⁻⁰⁴ 9.43 x 10 ⁻⁰⁵	5.25 x 10 ⁻¹³ 1.27 x 10 ⁻¹² 2.43 x 10 ⁻¹¹	1.45 x 10 ⁻⁰⁵ 6.06 x 10 ⁻⁰⁴ 4.83 x 10 ⁻⁰⁵	1.76 x 10 ⁻¹¹ 4.11 x 10 ⁻¹⁰ 2.16 x 10 ⁻⁰⁸	1.34 x 10 ⁻⁰⁸ 6.43 x 10 ⁻⁰⁸ 8.68 x 10 ⁻⁰⁸	1.36 x 10 ⁻⁰⁴ 6.88 x 10 ⁻⁰⁷ 9.84 x 10 ⁻⁰⁷ 2.25 x 10 ⁻⁰³	9.81 x 10 ⁻⁰⁸ 1.07 x 10 ⁻⁰⁷ 6.22 x 10 ⁻⁰⁷ 2.14 x 10 ⁻⁰⁷	1.05 x 10 ⁻⁰⁶ 3.81 x 10 ⁻⁰⁹ 1.54 x 10 ⁻⁰⁸ 2.56 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰ 0.00 x 10 ⁺⁰⁰ 0.00 x 10 ⁺⁰⁰ 0.00 x 10 ⁺⁰⁰	1.27 x 10 ⁻⁰⁵ 2.55 x 10 ⁻⁰⁴ 3.96 x 10 ⁻⁰⁵	4.84 x 10 ⁻⁰⁵ 2.14 x 10 ⁻⁰⁴ 5.00 x 10 ⁻⁰⁴	2.83 x 10 ⁻⁰⁵ 3.08 x 10 ⁻⁰³ 2.87 x 10 ⁻⁰⁴	1.99 x 10 ⁻⁰⁴ 4.53 x 10 ⁻⁰³ 3.24 x 10 ⁻⁰³
South Arm)	Uranium-234 Thorium-230 Radium-226 Lead-210	9.43 x 10 ⁻⁰⁵ 3.77 x 10 ⁻⁰⁴ 9.43 x 10 ⁻⁰⁵ 2.97 x 10 ⁻⁰⁵	5.25 x 10 ⁻¹³ 1.27 x 10 ⁻¹² 2.43 x 10 ⁻¹¹ 3.83 x 10 ⁻¹²	1.45 x 10 ⁻⁰⁵ 6.06 x 10 ⁻⁰⁴ 4.83 x 10 ⁻⁰⁵ 6.21 x 10 ⁻⁰⁵	1.76 x 10 ⁻¹¹ 4.11 x 10 ⁻¹⁰ 2.16 x 10 ⁻⁰⁸ 5.47 x 10 ⁻¹¹	1.34 x 10 ⁻⁰⁸ 6.43 x 10 ⁻⁰⁸ 8.68 x 10 ⁻⁰⁸ 1.38 x 10 ⁻⁰⁷	1.36 x 10 ⁻⁰⁴ 6.88 x 10 ⁻⁰⁷ 9.84 x 10 ⁻⁰⁷ 2.25 x 10 ⁻⁰³ 3.20 x 10 ⁻⁰⁵	9.81 x 10 ⁻⁰⁸ 1.07 x 10 ⁻⁰⁷ 6.22 x 10 ⁻⁰⁷ 2.14 x 10 ⁻⁰⁷ 1.56 x 10 ⁻⁰⁶	1.05 x 10 ⁻⁰⁶ 3.81 x 10 ⁻⁰⁹ 1.54 x 10 ⁻⁰⁸ 2.56 x 10 ⁻⁰⁵ 8.80 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰ 0.00 x 10 ⁺⁰⁰ 0.00 x 10 ⁺⁰⁰ 0.00 x 10 ⁺⁰⁰ 0.00 x 10 ⁺⁰⁰	1.27 x 10 ⁻⁰⁵ 2.55 x 10 ⁻⁰⁴ 3.96 x 10 ⁻⁰⁵ 1.73 x 10 ⁻⁰⁴	4.84 x 10 ⁻⁰⁵ 2.14 x 10 ⁻⁰⁴ 5.00 x 10 ⁻⁰⁴ 9.79 x 10 ⁻⁰⁴	2.83 x 10 ⁻⁰⁵ 3.08 x 10 ⁻⁰³ 2.87 x 10 ⁻⁰⁴ 7.28 x 10 ⁻⁰³	1.99 x 10 ⁻⁰⁴ 4.53 x 10 ⁻⁰³ 3.24 x 10 ⁻⁰³ 8.55 x 10 ⁻⁰³
South Arm)	Uranium-234 Thorium-230 Radium-226 Lead-210 Polonium-210	9.43 x 10 ⁻⁰⁵ 3.77 x 10 ⁻⁰⁴ 9.43 x 10 ⁻⁰⁵ 2.97 x 10 ⁻⁰⁵ 8.90 x 10 ⁻⁰⁵	5.25 x 10 ⁻¹³ 1.27 x 10 ⁻¹² 2.43 x 10 ⁻¹¹ 3.83 x 10 ⁻¹² 3.34 x 10 ⁻¹⁴	1.45 x 10 ⁻⁰⁵ 6.06 x 10 ⁻⁰⁴ 4.83 x 10 ⁻⁰⁵ 6.21 x 10 ⁻⁰⁵ 1.10 x 10 ⁻⁰⁴	1.76 x 10 ⁻¹¹ 4.11 x 10 ⁻¹⁰ 2.16 x 10 ⁻⁰⁸ 5.47 x 10 ⁻¹¹ 3.32 x 10 ⁻¹³	1.34 x 10 ⁻⁰⁸ 6.43 x 10 ⁻⁰⁸ 8.68 x 10 ⁻⁰⁸ 1.38 x 10 ⁻⁰⁷ 2.44 x 10 ⁻⁰⁷	1.36 x 10 ⁻⁰⁴ 6.88 x 10 ⁻⁰⁷ 9.84 x 10 ⁻⁰⁷ 2.25 x 10 ⁻⁰³ 3.20 x 10 ⁻⁰⁵ 7.23 x 10 ⁻⁰⁹ 2.42 x 10 ⁻⁰³	9.81 x 10 ⁻⁰⁸ 1.07 x 10 ⁻⁰⁷ 6.22 x 10 ⁻⁰⁷ 2.14 x 10 ⁻⁰⁷ 1.56 x 10 ⁻⁰⁶ 2.75 x 10 ⁻⁰⁶	1.05 x 10 ⁻⁰⁶ 3.81 x 10 ⁻⁰⁹ 1.54 x 10 ⁻⁰⁸ 2.56 x 10 ⁻⁰⁵ 8.80 x 10 ⁻⁰⁸ 3.78 x 10 ⁻¹⁰ 2.68 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰ 0.00 x 10 ⁺⁰⁰	1.27 x 10 ⁻⁰⁵ 2.55 x 10 ⁻⁰⁴ 3.96 x 10 ⁻⁰⁵ 1.73 x 10 ⁻⁰⁴ 1.08 x 10 ⁻⁰³	4.84 x 10 ⁻⁰⁵ 2.14 x 10 ⁻⁰⁴ 5.00 x 10 ⁻⁰⁴ 9.79 x 10 ⁻⁰⁴ 1.65 x 10 ⁻⁰³	2.83 x 10 ⁻⁰⁵ 3.08 x 10 ⁻⁰³ 2.87 x 10 ⁻⁰⁴ 7.28 x 10 ⁻⁰³ 6.26 x 10 ⁻⁰³	1.99 x 10 ⁻⁰⁴ 4.53 x 10 ⁻⁰³ 3.24 x 10 ⁻⁰³ 8.55 x 10 ⁻⁰³ 9.19 x 10 ⁻⁰³ 2.60 x 10 ⁻⁰²
South Arm)	Uranium-234 Thorium-230 Radium-226 Lead-210 Polonium-210 Total by Pathway Uranium-238	9.43 x 10 ⁻⁰⁵ 3.77 x 10 ⁻⁰⁴ 9.43 x 10 ⁻⁰⁵ 2.97 x 10 ⁻⁰⁵ 8.90 x 10 ⁻⁰⁵ 7.63 x 10 ⁻⁰⁴	5.25 x 10 ⁻¹³ 1.27 x 10 ⁻¹² 2.43 x 10 ⁻¹¹ 3.83 x 10 ⁻¹² 3.34 x 10 ⁻¹⁴	1.45 x 10 ⁻⁰⁵ 6.06 x 10 ⁻⁰⁴ 4.83 x 10 ⁻⁰⁵ 6.21 x 10 ⁻⁰⁵ 1.10 x 10 ⁻⁰⁴ 8.55 x 10 ⁻⁰⁴	1.76 x 10 ⁻¹¹ 4.11 x 10 ⁻¹⁰ 2.16 x 10 ⁻⁰⁸ 5.47 x 10 ⁻¹¹ 3.32 x 10 ⁻¹³	1.34 x 10 ⁻⁰⁸ 6.43 x 10 ⁻⁰⁸ 8.68 x 10 ⁻⁰⁸ 1.38 x 10 ⁻⁰⁷ 2.44 x 10 ⁻⁰⁷ 5.59 x 10 ⁻⁰⁷	1.36 x 10 ⁻⁰⁴ 6.88 x 10 ⁻⁰⁷ 9.84 x 10 ⁻⁰⁷ 2.25 x 10 ⁻⁰³ 3.20 x 10 ⁻⁰⁵ 7.23 x 10 ⁻⁰⁹ 2.42 x 10 ⁻⁰³	9.81 x 10 ⁻⁰⁸ 1.07 x 10 ⁻⁰⁷ 6.22 x 10 ⁻⁰⁷ 2.14 x 10 ⁻⁰⁷ 1.56 x 10 ⁻⁰⁶ 2.75 x 10 ⁻⁰⁶ 5.35 x 10 ⁻⁰⁶	1.05 x 10 ⁻⁰⁶ 3.81 x 10 ⁻⁰⁹ 1.54 x 10 ⁻⁰⁸ 2.56 x 10 ⁻⁰⁵ 8.80 x 10 ⁻⁰⁸ 3.78 x 10 ⁻¹⁰ 2.68 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰ 0.00 x 10 ⁺⁰⁰	1.27 x 10 ⁻⁰⁵ 2.55 x 10 ⁻⁰⁴ 3.96 x 10 ⁻⁰⁵ 1.73 x 10 ⁻⁰⁴ 1.08 x 10 ⁻⁰³	4.84 x 10 ⁻⁰⁵ 2.14 x 10 ⁻⁰⁴ 5.00 x 10 ⁻⁰⁴ 9.79 x 10 ⁻⁰⁴ 1.65 x 10 ⁻⁰³	2.83 x 10 ⁻⁰⁵ 3.08 x 10 ⁻⁰³ 2.87 x 10 ⁻⁰⁴ 7.28 x 10 ⁻⁰³ 6.26 x 10 ⁻⁰³	1.99 x 10 ⁻⁰⁴ 4.53 x 10 ⁻⁰³ 3.24 x 10 ⁻⁰³ 8.55 x 10 ⁻⁰³ 9.19 x 10 ⁻⁰³
	Uranium-234 Thorium-230 Radium-226 Lead-210 Polonium-210 Total by Pathway	9.43 x 10 ⁻⁰⁵ 3.77 x 10 ⁻⁰⁴ 9.43 x 10 ⁻⁰⁵ 2.97 x 10 ⁻⁰⁵ 8.90 x 10 ⁻⁰⁵ 7.63 x 10 ⁻⁰⁴	5.25 x 10 ⁻¹³ 1.27 x 10 ⁻¹² 2.43 x 10 ⁻¹¹ 3.83 x 10 ⁻¹² 3.34 x 10 ⁻¹⁴ 3.02 x 10 ⁻¹¹	1.45 x 10 ⁻⁰⁵ 6.06 x 10 ⁻⁰⁴ 4.83 x 10 ⁻⁰⁵ 6.21 x 10 ⁻⁰⁵ 1.10 x 10 ⁻⁰⁴ 8.55 x 10 ⁻⁰⁴	1.76 x 10 ⁻¹¹ 4.11 x 10 ⁻¹⁰ 2.16 x 10 ⁻⁰⁸ 5.47 x 10 ⁻¹¹ 3.32 x 10 ⁻¹³ 2.21 x 10 ⁻⁰⁸	1.34 x 10 ⁻⁰⁸ 6.43 x 10 ⁻⁰⁸ 8.68 x 10 ⁻⁰⁸ 1.38 x 10 ⁻⁰⁷ 2.44 x 10 ⁻⁰⁷ 5.59 x 10 ⁻⁰⁷	1.36 x 10 ⁻⁰⁴ 6.88 x 10 ⁻⁰⁷ 9.84 x 10 ⁻⁰⁷ 2.25 x 10 ⁻⁰³ 3.20 x 10 ⁻⁰⁵ 7.23 x 10 ⁻⁰⁹ 2.42 x 10 ⁻⁰³	9.81 x 10 ⁻⁰⁸ 1.07 x 10 ⁻⁰⁷ 6.22 x 10 ⁻⁰⁷ 2.14 x 10 ⁻⁰⁶ 1.56 x 10 ⁻⁰⁶ 2.75 x 10 ⁻⁰⁶ 5.35 x 10 ⁻⁰⁶ Application Case 1.43 x 10 ⁻⁰⁶ 1.56 x 10 ⁻⁰⁶	1.05 x 10 ⁻⁰⁶ 3.81 x 10 ⁻⁰⁹ 1.54 x 10 ⁻⁰⁸ 2.56 x 10 ⁻⁰⁵ 8.80 x 10 ⁻⁰⁸ 3.78 x 10 ⁻¹⁰ 2.68 x 10 ⁻⁰⁵ 4.88 x 10 ⁻⁰⁷ 1.78 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰ 0.00 x 10 ⁺⁰⁰	1.27 x 10 ⁻⁰⁵ 2.55 x 10 ⁻⁰⁴ 3.96 x 10 ⁻⁰⁵ 1.73 x 10 ⁻⁰⁴ 1.08 x 10 ⁻⁰³ 1.57 x 10 ⁻⁰³ 2.50 x 10 ⁻⁰⁶ 2.71 x 10 ⁻⁰⁶	4.84 x 10 ⁻⁰⁵ 2.14 x 10 ⁻⁰⁴ 5.00 x 10 ⁻⁰⁴ 9.79 x 10 ⁻⁰⁴ 1.65 x 10 ⁻⁰³ 3.43 x 10 ⁻⁰³	2.83 x 10 ⁻⁰⁵ 3.08 x 10 ⁻⁰³ 2.87 x 10 ⁻⁰⁴ 7.28 x 10 ⁻⁰³ 6.26 x 10 ⁻⁰³ 1.70 x 10 ⁻⁰²	1.99 x 10 ⁻⁰⁴ 4.53 x 10 ⁻⁰³ 3.24 x 10 ⁻⁰³ 8.55 x 10 ⁻⁰³ 9.19 x 10 ⁻⁰³ 2.60 x 10 ⁻⁰² 3.40 x 10 ⁻⁰⁴ 1.78 x 10 ⁻⁰⁴
	Uranium-234 Thorium-230 Radium-226 Lead-210 Polonium-210 Total by Pathway Uranium-238	9.43 x 10 ⁻⁰⁵ 3.77 x 10 ⁻⁰⁴ 9.43 x 10 ⁻⁰⁵ 2.97 x 10 ⁻⁰⁵ 8.90 x 10 ⁻⁰⁵ 7.63 x 10 ⁻⁰⁴	5.25 x 10 ⁻¹³ 1.27 x 10 ⁻¹² 2.43 x 10 ⁻¹¹ 3.83 x 10 ⁻¹² 3.34 x 10 ⁻¹⁴ 3.02 x 10 ⁻¹¹	1.45 x 10 ⁻⁰⁵ 6.06 x 10 ⁻⁰⁴ 4.83 x 10 ⁻⁰⁵ 6.21 x 10 ⁻⁰⁵ 1.10 x 10 ⁻⁰⁴ 8.55 x 10 ⁻⁰⁴	1.76 x 10 ⁻¹¹ 4.11 x 10 ⁻¹⁰ 2.16 x 10 ⁻⁰⁸ 5.47 x 10 ⁻¹¹ 3.32 x 10 ⁻¹³ 2.21 x 10 ⁻⁰⁸ 9.11 x 10 ⁻¹² 1.60 x 10 ⁻¹² 1.00 x 10 ⁻¹⁰	1.34 x 10 ⁻⁰⁸ 6.43 x 10 ⁻⁰⁸ 8.68 x 10 ⁻⁰⁸ 1.38 x 10 ⁻⁰⁷ 2.44 x 10 ⁻⁰⁷ 5.59 x 10 ⁻⁰⁷	1.36 x 10 ⁻⁰⁴ 6.88 x 10 ⁻⁰⁷ 9.84 x 10 ⁻⁰⁷ 2.25 x 10 ⁻⁰³ 3.20 x 10 ⁻⁰⁵ 7.23 x 10 ⁻⁰⁹ 2.42 x 10 ⁻⁰³	9.81 x 10 ⁻⁰⁸ 1.07 x 10 ⁻⁰⁷ 6.22 x 10 ⁻⁰⁷ 2.14 x 10 ⁻⁰⁶ 1.56 x 10 ⁻⁰⁶ 2.75 x 10 ⁻⁰⁶ 5.35 x 10 ⁻⁰⁶ Application Case 1.43 x 10 ⁻⁰⁶ 1.56 x 10 ⁻⁰⁶ 1.83 x 10 ⁻⁰⁵	1.05 x 10 ⁻⁰⁶ 3.81 x 10 ⁻⁰⁹ 1.54 x 10 ⁻⁰⁸ 2.56 x 10 ⁻⁰⁵ 8.80 x 10 ⁻⁰⁸ 3.78 x 10 ⁻¹⁰ 2.68 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	1.27 x 10 ⁻⁰⁵ 2.55 x 10 ⁻⁰⁴ 3.96 x 10 ⁻⁰⁵ 1.73 x 10 ⁻⁰⁴ 1.08 x 10 ⁻⁰³ 1.57 x 10 ⁻⁰³	4.84 x 10 ⁻⁰⁵ 2.14 x 10 ⁻⁰⁴ 5.00 x 10 ⁻⁰⁴ 9.79 x 10 ⁻⁰⁴ 1.65 x 10 ⁻⁰³ 3.43 x 10 ⁻⁰³ 6.06 x 10 ⁻⁰⁵ 6.56 x 10 ⁻⁰⁵ 1.18 x 10 ⁻⁰⁴	2.83 x 10 ⁻⁰⁵ 3.08 x 10 ⁻⁰³ 2.87 x 10 ⁻⁰⁴ 7.28 x 10 ⁻⁰³ 6.26 x 10 ⁻⁰³ 1.70 x 10 ⁻⁰²	1.99 x 10 ⁻⁰⁴ 4.53 x 10 ⁻⁰³ 3.24 x 10 ⁻⁰³ 8.55 x 10 ⁻⁰³ 9.19 x 10 ⁻⁰³ 2.60 x 10 ⁻⁰² 3.40 x 10 ⁻⁰⁴ 1.78 x 10 ⁻⁰⁴ 2.33 x 10 ⁻⁰³
Seasonal Resident One-Year-Old (Patterson Lake	Uranium-234 Thorium-230 Radium-226 Lead-210 Polonium-210 Total by Pathway Uranium-238 Uranium-234	9.43 x 10 ⁻⁰⁵ 3.77 x 10 ⁻⁰⁴ 9.43 x 10 ⁻⁰⁵ 2.97 x 10 ⁻⁰⁵ 8.90 x 10 ⁻⁰⁵ 7.63 x 10 ⁻⁰⁴ 7.79 x 10 ⁻⁰⁵ 9.12 x 10 ⁻⁰⁵	5.25 x 10 ⁻¹³ 1.27 x 10 ⁻¹² 2.43 x 10 ⁻¹¹ 3.83 x 10 ⁻¹² 3.34 x 10 ⁻¹⁴ 3.02 x 10 ⁻¹¹ 2.79 x 10 ⁻¹³ 6.82 x 10 ⁻¹³	1.45 x 10 ⁻⁰⁵ 6.06 x 10 ⁻⁰⁴ 4.83 x 10 ⁻⁰⁵ 6.21 x 10 ⁻⁰⁵ 1.10 x 10 ⁻⁰⁴ 8.55 x 10 ⁻⁰⁴ 3.37 x 10 ⁻⁰⁶ 3.67 x 10 ⁻⁰⁶	1.76 x 10 ⁻¹¹ 4.11 x 10 ⁻¹⁰ 2.16 x 10 ⁻⁰⁸ 5.47 x 10 ⁻¹¹ 3.32 x 10 ⁻¹³ 2.21 x 10 ⁻⁰⁸ 9.11 x 10 ⁻¹²	1.34 x 10 ⁻⁰⁸ 6.43 x 10 ⁻⁰⁸ 8.68 x 10 ⁻⁰⁸ 1.38 x 10 ⁻⁰⁷ 2.44 x 10 ⁻⁰⁷ 5.59 x 10 ⁻⁰⁷ 5.00 x 10 ⁻⁰⁷ 5.42 x 10 ⁻⁰⁷	1.36 x 10 ⁻⁰⁴ 6.88 x 10 ⁻⁰⁷ 9.84 x 10 ⁻⁰⁷ 2.25 x 10 ⁻⁰³ 3.20 x 10 ⁻⁰⁵ 7.23 x 10 ⁻⁰⁹ 2.42 x 10 ⁻⁰³ 1.83 x 10 ⁻⁰⁴ 9.21 x 10 ⁻⁰⁷	9.81 x 10 ⁻⁰⁸ 1.07 x 10 ⁻⁰⁷ 6.22 x 10 ⁻⁰⁷ 2.14 x 10 ⁻⁰⁶ 1.56 x 10 ⁻⁰⁶ 2.75 x 10 ⁻⁰⁶ 5.35 x 10 ⁻⁰⁶ Application Case 1.43 x 10 ⁻⁰⁶ 1.56 x 10 ⁻⁰⁶	1.05 x 10 ⁻⁰⁶ 3.81 x 10 ⁻⁰⁹ 1.54 x 10 ⁻⁰⁸ 2.56 x 10 ⁻⁰⁵ 8.80 x 10 ⁻⁰⁸ 3.78 x 10 ⁻¹⁰ 2.68 x 10 ⁻⁰⁵ 4.88 x 10 ⁻⁰⁷ 1.78 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰ 0.00 x 10 ⁺⁰⁰	1.27 x 10 ⁻⁰⁵ 2.55 x 10 ⁻⁰⁴ 3.96 x 10 ⁻⁰⁵ 1.73 x 10 ⁻⁰⁴ 1.08 x 10 ⁻⁰³ 1.57 x 10 ⁻⁰³ 2.50 x 10 ⁻⁰⁶ 2.71 x 10 ⁻⁰⁶	4.84 x 10 ⁻⁰⁵ 2.14 x 10 ⁻⁰⁴ 5.00 x 10 ⁻⁰⁴ 9.79 x 10 ⁻⁰⁴ 1.65 x 10 ⁻⁰³ 3.43 x 10 ⁻⁰³ 6.06 x 10 ⁻⁰⁵ 6.56 x 10 ⁻⁰⁵	2.83 x 10 ⁻⁰⁵ 3.08 x 10 ⁻⁰³ 2.87 x 10 ⁻⁰⁴ 7.28 x 10 ⁻⁰³ 6.26 x 10 ⁻⁰³ 1.70 x 10 ⁻⁰² 1.06 x 10 ⁻⁰⁵ 1.15 x 10 ⁻⁰⁵	1.99 x 10 ⁻⁰⁴ 4.53 x 10 ⁻⁰³ 3.24 x 10 ⁻⁰³ 8.55 x 10 ⁻⁰³ 9.19 x 10 ⁻⁰³ 2.60 x 10 ⁻⁰² 3.40 x 10 ⁻⁰⁴ 1.78 x 10 ⁻⁰⁴
Seasonal Resident One-Year-Old	Uranium-234 Thorium-230 Radium-226 Lead-210 Polonium-210 Total by Pathway Uranium-238 Uranium-234 Thorium-230	9.43 x 10 ⁻⁰⁵ 3.77 x 10 ⁻⁰⁴ 9.43 x 10 ⁻⁰⁵ 2.97 x 10 ⁻⁰⁵ 8.90 x 10 ⁻⁰⁵ 7.63 x 10 ⁻⁰⁴ 7.79 x 10 ⁻⁰⁵ 9.12 x 10 ⁻⁰⁵ 2.90 x 10 ⁻⁰⁴	5.25 x 10 ⁻¹³ 1.27 x 10 ⁻¹² 2.43 x 10 ⁻¹¹ 3.83 x 10 ⁻¹² 3.34 x 10 ⁻¹⁴ 3.02 x 10 ⁻¹¹ 2.79 x 10 ⁻¹³ 6.82 x 10 ⁻¹³ 1.65 x 10 ⁻¹²	1.45 x 10 ⁻⁰⁵ 6.06 x 10 ⁻⁰⁴ 4.83 x 10 ⁻⁰⁵ 6.21 x 10 ⁻⁰⁵ 1.10 x 10 ⁻⁰⁴ 8.55 x 10 ⁻⁰⁴ 3.37 x 10 ⁻⁰⁶ 3.67 x 10 ⁻⁰⁶ 3.04 x 10 ⁻⁰⁴	1.76 x 10 ⁻¹¹ 4.11 x 10 ⁻¹⁰ 2.16 x 10 ⁻⁰⁸ 5.47 x 10 ⁻¹¹ 3.32 x 10 ⁻¹³ 2.21 x 10 ⁻⁰⁸ 9.11 x 10 ⁻¹² 1.60 x 10 ⁻¹² 1.00 x 10 ⁻¹⁰	1.34 x 10 ⁻⁰⁸ 6.43 x 10 ⁻⁰⁸ 8.68 x 10 ⁻⁰⁸ 1.38 x 10 ⁻⁰⁷ 2.44 x 10 ⁻⁰⁷ 5.59 x 10 ⁻⁰⁷ 5.00 x 10 ⁻⁰⁷ 5.42 x 10 ⁻⁰⁷ 1.91 x 10 ⁻⁰⁶	1.36 x 10 ⁻⁰⁴ 6.88 x 10 ⁻⁰⁷ 9.84 x 10 ⁻⁰⁷ 2.25 x 10 ⁻⁰³ 3.20 x 10 ⁻⁰⁵ 7.23 x 10 ⁻⁰⁹ 2.42 x 10 ⁻⁰³ 1.83 x 10 ⁻⁰⁴ 9.21 x 10 ⁻⁰⁷ 1.12 x 10 ⁻⁰⁶	9.81 x 10 ⁻⁰⁸ 1.07 x 10 ⁻⁰⁷ 6.22 x 10 ⁻⁰⁷ 2.14 x 10 ⁻⁰⁶ 1.56 x 10 ⁻⁰⁶ 2.75 x 10 ⁻⁰⁶ 5.35 x 10 ⁻⁰⁶ Application Case 1.43 x 10 ⁻⁰⁶ 1.56 x 10 ⁻⁰⁶ 1.83 x 10 ⁻⁰⁵	1.05 x 10 ⁻⁰⁶ 3.81 x 10 ⁻⁰⁹ 1.54 x 10 ⁻⁰⁸ 2.56 x 10 ⁻⁰⁵ 8.80 x 10 ⁻⁰⁸ 3.78 x 10 ⁻¹⁰ 2.68 x 10 ⁻⁰⁵ 4.88 x 10 ⁻⁰⁷ 1.78 x 10 ⁻⁰⁹ 1.97 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	1.27 x 10 ⁻⁰⁵ 2.55 x 10 ⁻⁰⁴ 3.96 x 10 ⁻⁰⁵ 1.73 x 10 ⁻⁰⁴ 1.08 x 10 ⁻⁰³ 1.57 x 10 ⁻⁰³ 2.50 x 10 ⁻⁰⁶ 2.71 x 10 ⁻⁰⁶ 1.09 x 10 ⁻⁰⁴	4.84 x 10 ⁻⁰⁵ 2.14 x 10 ⁻⁰⁴ 5.00 x 10 ⁻⁰⁴ 9.79 x 10 ⁻⁰⁴ 1.65 x 10 ⁻⁰³ 3.43 x 10 ⁻⁰³ 6.06 x 10 ⁻⁰⁵ 6.56 x 10 ⁻⁰⁵ 1.18 x 10 ⁻⁰⁴	2.83 x 10 ⁻⁰⁵ 3.08 x 10 ⁻⁰³ 2.87 x 10 ⁻⁰⁴ 7.28 x 10 ⁻⁰³ 6.26 x 10 ⁻⁰³ 1.70 x 10 ⁻⁰² 1.06 x 10 ⁻⁰⁵ 1.15 x 10 ⁻⁰⁵ 1.49 x 10 ⁻⁰³	1.99 x 10 ⁻⁰⁴ 4.53 x 10 ⁻⁰³ 3.24 x 10 ⁻⁰³ 8.55 x 10 ⁻⁰³ 9.19 x 10 ⁻⁰³ 2.60 x 10 ⁻⁰² 3.40 x 10 ⁻⁰⁴ 1.78 x 10 ⁻⁰⁴ 2.33 x 10 ⁻⁰³
Seasonal Resident One-Year-Old (Patterson Lake	Uranium-234 Thorium-230 Radium-226 Lead-210 Polonium-210 Total by Pathway Uranium-238 Uranium-234 Thorium-230 Radium-226	9.43 x 10 ⁻⁰⁵ 3.77 x 10 ⁻⁰⁴ 9.43 x 10 ⁻⁰⁵ 2.97 x 10 ⁻⁰⁵ 8.90 x 10 ⁻⁰⁵ 7.63 x 10 ⁻⁰⁴ 7.79 x 10 ⁻⁰⁵ 9.12 x 10 ⁻⁰⁵ 2.90 x 10 ⁻⁰⁴ 9.12 x 10 ⁻⁰⁵	5.25 x 10 ⁻¹³ 1.27 x 10 ⁻¹² 2.43 x 10 ⁻¹¹ 3.83 x 10 ⁻¹² 3.34 x 10 ⁻¹⁴ 3.02 x 10 ⁻¹¹ 2.79 x 10 ⁻¹³ 6.82 x 10 ⁻¹³ 1.65 x 10 ⁻¹² 3.16 x 10 ⁻¹¹	1.45 x 10 ⁻⁰⁵ 6.06 x 10 ⁻⁰⁴ 4.83 x 10 ⁻⁰⁵ 6.21 x 10 ⁻⁰⁵ 1.10 x 10 ⁻⁰⁴ 8.55 x 10 ⁻⁰⁴ 3.37 x 10 ⁻⁰⁶ 3.67 x 10 ⁻⁰⁶ 3.04 x 10 ⁻⁰⁴ 4.13 x 10 ⁻⁰⁵	1.76 x 10 ⁻¹¹ 4.11 x 10 ⁻¹⁰ 2.16 x 10 ⁻⁰⁸ 5.47 x 10 ⁻¹¹ 3.32 x 10 ⁻¹³ 2.21 x 10 ⁻⁰⁸ 9.11 x 10 ⁻¹² 1.60 x 10 ⁻¹² 1.00 x 10 ⁻¹⁰ 5.11 x 10 ⁻⁰⁹	1.34 x 10 ⁻⁰⁸ 6.43 x 10 ⁻⁰⁸ 8.68 x 10 ⁻⁰⁸ 1.38 x 10 ⁻⁰⁷ 2.44 x 10 ⁻⁰⁷ 5.59 x 10 ⁻⁰⁷ 5.00 x 10 ⁻⁰⁷ 5.42 x 10 ⁻⁰⁶ 4.54 x 10 ⁻⁰⁶	1.36 x 10 ⁻⁰⁴ 6.88 x 10 ⁻⁰⁷ 9.84 x 10 ⁻⁰⁷ 2.25 x 10 ⁻⁰³ 3.20 x 10 ⁻⁰⁵ 7.23 x 10 ⁻⁰⁹ 2.42 x 10 ⁻⁰³ 1.83 x 10 ⁻⁰⁴ 9.21 x 10 ⁻⁰⁷ 1.12 x 10 ⁻⁰⁶ 3.01 x 10 ⁻⁰³	9.81 x 10 ⁻⁰⁸ 1.07 x 10 ⁻⁰⁷ 6.22 x 10 ⁻⁰⁷ 2.14 x 10 ⁻⁰⁷ 1.56 x 10 ⁻⁰⁶ 2.75 x 10 ⁻⁰⁶ 5.35 x 10 ⁻⁰⁶ Application Case 1.43 x 10 ⁻⁰⁶ 1.56 x 10 ⁻⁰⁶ 1.83 x 10 ⁻⁰⁵	1.05 x 10 ⁻⁰⁶ 3.81 x 10 ⁻⁰⁹ 1.54 x 10 ⁻⁰⁸ 2.56 x 10 ⁻⁰⁵ 8.80 x 10 ⁻⁰⁸ 3.78 x 10 ⁻¹⁰ 2.68 x 10 ⁻⁰⁵ 4.88 x 10 ⁻⁰⁷ 1.78 x 10 ⁻⁰⁹ 1.97 x 10 ⁻⁰⁸ 3.19 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	1.27 x 10 ⁻⁰⁵ 2.55 x 10 ⁻⁰⁴ 3.96 x 10 ⁻⁰⁵ 1.73 x 10 ⁻⁰⁴ 1.08 x 10 ⁻⁰³ 1.57 x 10 ⁻⁰³ 2.50 x 10 ⁻⁰⁶ 2.71 x 10 ⁻⁰⁶ 1.09 x 10 ⁻⁰⁴ 2.87 x 10 ⁻⁰⁵	4.84 x 10 ⁻⁰⁵ 2.14 x 10 ⁻⁰⁴ 5.00 x 10 ⁻⁰⁴ 9.79 x 10 ⁻⁰⁴ 1.65 x 10 ⁻⁰³ 3.43 x 10 ⁻⁰³ 6.06 x 10 ⁻⁰⁵ 6.56 x 10 ⁻⁰⁵ 1.18 x 10 ⁻⁰⁴ 6.68 x 10 ⁻⁰⁴	2.83 x 10 ⁻⁰⁵ 3.08 x 10 ⁻⁰³ 2.87 x 10 ⁻⁰⁴ 7.28 x 10 ⁻⁰³ 6.26 x 10 ⁻⁰³ 1.70 x 10 ⁻⁰² 1.06 x 10 ⁻⁰⁵ 1.15 x 10 ⁻⁰⁵ 1.49 x 10 ⁻⁰³ 1.35 x 10 ⁻⁰⁴	1.99 x 10 ⁻⁰⁴ 4.53 x 10 ⁻⁰³ 3.24 x 10 ⁻⁰³ 8.55 x 10 ⁻⁰³ 9.19 x 10 ⁻⁰³ 2.60 x 10 ⁻⁰² 3.40 x 10 ⁻⁰⁴ 1.78 x 10 ⁻⁰⁴ 2.33 x 10 ⁻⁰³ 4.02 x 10 ⁻⁰³

							Incremental Dose b	ov Pathwav - Projec	ct Lifespan (mSv/v	vr)				
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by Radionuclide
		7.70 40:05	2.70 40.13	0.22 40.06	2.4040-11	5.00 40-07		pper Bound Scena		0.00 40+00	6.02 40-06	6.06 40.05	1.10 10.05	2.55 40:04
	Uranium-238	7.79 x 10 ⁻⁰⁵	2.79 x 10 ⁻¹³	9.23 x 10 ⁻⁰⁶	2.49 x 10 ⁻¹¹	5.00 x 10 ⁻⁰⁷	1.83 x 10 ⁻⁰⁴	3.99 x 10 ⁻⁰⁶	1.36 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	6.83 x 10 ⁻⁰⁶	6.06 x 10 ⁻⁰⁵	1.19 x 10 ⁻⁰⁵	3.55 x 10 ⁻⁰⁴
	Uranium-234	9.12 x 10 ⁻⁰⁵	6.82 x 10 ⁻¹³	1.00 x 10 ⁻⁰⁵	4.36 x 10 ⁻¹²	5.42 x 10 ⁻⁰⁷	9.21 x 10 ⁻⁰⁷	4.34 x 10 ⁻⁰⁶	4.95 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	7.43 x 10 ⁻⁰⁶	6.56 x 10 ⁻⁰⁵	1.29 x 10 ⁻⁰⁵	1.93 x 10 ⁻⁰⁴
	Thorium-230	2.90 x 10 ⁻⁰⁴	1.65 x 10 ⁻¹²	3.08 x 10 ⁻⁰⁴	1.02 x 10 ⁻¹⁰	1.91 x 10 ⁻⁰⁶	1.12 x 10 ⁻⁰⁶	1.85 x 10 ⁻⁰⁵	2.00 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	1.10 x 10 ⁻⁰⁴	1.18 x 10 ⁻⁰⁴	1.50 x 10 ⁻⁰³	2.35 x 10 ⁻⁰³
	Radium-226	9.12 x 10 ⁻⁰⁵	3.16 x 10 ⁻¹¹	4.32 x 10 ⁻⁰⁵	5.34 x 10 ⁻⁰⁹	4.54 x 10 ⁻⁰⁶	3.01 x 10 ⁻⁰³	1.12 x 10 ⁻⁰⁵	3.34 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	3.00 x 10 ⁻⁰⁵	6.68 x 10 ⁻⁰⁴	1.35 x 10 ⁻⁰⁴	4.03 x 10 ⁻⁰³
	Lead-210	3.07 x 10 ⁻⁰⁵	4.97 x 10 ⁻¹²	8.44 x 10 ⁻⁰⁵	1.11 x 10 ⁻¹¹	1.10 x 10 ⁻⁰⁵	4.28 x 10 ⁻⁰⁵	1.24 x 10 ⁻⁰⁴	1.14 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.99 x 10 ⁻⁰⁴	1.83 x 10 ⁻⁰³	9.22 x 10 ⁻⁰³	1.15 x 10 ⁻⁰²
	Polonium-210	9.12 x 10 ⁻⁰⁵	4.35 x 10 ⁻¹⁴	2.11 x 10 ⁻⁰⁴	8.22 x 10 ⁻¹⁴	2.73 x 10 ⁻⁰⁵	9.45 x 10 ⁻⁰⁹	3.08 x 10 ⁻⁰⁴	4.91 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	1.75 x 10 ⁻⁰³	4.54 x 10 ⁻⁰³	6.64 x 10 ⁻⁰³	1.36 x 10 ⁻⁰²
	Total by Pathway	6.72 x 10 ⁻⁰⁴	3.93 x 10 ⁻¹¹	6.66 x 10 ⁻⁰⁴	5.48 x 10 ⁻⁰⁹	4.58 x 10 ⁻⁰⁵	3.24 x 10 ⁻⁰³	4.70 x 10 ⁻⁰⁴	3.49 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	2.11 x 10 ⁻⁰³	7.29 x 10 ⁻⁰³	1.75 x 10 ⁻⁰²	3.20 x 10 ⁻⁰²
		0.5	45	07		10	0.5	Application Case						
	Uranium-238	1.52 x 10 ⁻⁰⁶	4.18 x 10 ⁻¹⁵	1.63 x 10 ⁻⁰⁷	1.24 x 10 ⁻¹²	2.40 x 10 ⁻¹⁰	2.66 x 10 ⁻⁰⁶	1.17 x 10 ⁻⁰⁹	1.25 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	1.59 x 10 ⁻⁰⁶	8.66 x 10 ⁻⁰⁷	3.15 x 10 ⁻⁰⁷	7.13 x 10 ⁻⁰⁶
	Uranium-234	1.84 x 10 ⁻⁰⁶	1.02 x 10 ⁻¹⁴	1.78 x 10 ⁻⁰⁷	2.16 x 10 ⁻¹³	2.61 x 10 ⁻¹⁰	1.34 x 10 ⁻⁰⁸	1.28 x 10 ⁻⁰⁹	4.55 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	1.73 x 10 ⁻⁰⁶	9.43 x 10 ⁻⁰⁷	3.43 x 10 ⁻⁰⁷	5.05 x 10 ⁻⁰⁶
	Thorium-230	7.36 x 10 ⁻⁰⁶	2.48 x 10 ⁻¹⁴	3.24 x 10 ⁻⁰⁵	2.20 x 10 ⁻¹¹	1.25 x 10 ⁻⁰⁹	1.92 x 10 ⁻⁰⁸	3.32 x 10 ⁻⁰⁸	8.20 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	2.28 x 10 ⁻⁰⁵	4.17 x 10 ⁻⁰⁶	5.07 x 10 ⁻⁰⁵	1.18 x 10 ⁻⁰⁴
	Radium-226	1.84 x 10 ⁻⁰⁶	4.75 x 10 ⁻¹³	1.90 x 10 ⁻⁰⁶	8.49 x 10 ⁻¹⁰	1.69 x 10 ⁻⁰⁹	4.38 x 10 ⁻⁰⁵	8.34 x 10 ⁻⁰⁹	1.00 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	3.31 x 10 ⁻⁰⁶	9.75 x 10 ⁻⁰⁶	2.46 x 10 ⁻⁰⁶	6.41 x 10 ⁻⁰⁵
	Lead-210	5.78 x 10 ⁻⁰⁷	7.47 x 10 ⁻¹⁴	8.31 x 10 ⁻⁰⁸	7.33 x 10 ⁻¹⁴	2.69 x 10 ⁻⁰⁹	6.24 x 10 ⁻⁰⁷	1.49 x 10 ⁻¹⁰	8.64 x 10 ⁻¹²	0.00 x 10 ⁺⁰⁰	7.56 x 10 ⁻⁰⁴	1.91 x 10 ⁻⁰⁵	7.49 x 10 ⁻⁰⁶	7.84 x 10 ⁻⁰⁴
	Polonium-210	1.73 x 10 ⁻⁰⁶	6.52 x 10 ⁻¹⁶	1.24 x 10 ⁻⁰⁷	3.71 x 10 ⁻¹⁶	4.76 x 10 ⁻⁰⁹	1.41 x 10 ⁻¹⁰	2.26 x 10 ⁻¹⁰	2.93 x 10 ⁻¹⁴	0.00 x 10 ⁺⁰⁰	5.12 x 10 ⁻⁰⁴	3.21 x 10 ⁻⁰⁵	2.97 x 10 ⁻⁰⁵	5.75 x 10 ⁻⁰⁴
Seasonal Resident	Total by Pathway	1.49 x 10 ⁻⁰⁵	5.89 x 10 ⁻¹³	3.49 x 10 ⁻⁰⁵	8.72 x 10 ⁻¹⁰	1.09 x 10 ⁻⁰⁸	4.71 x 10 ⁻⁰⁵	4.43 x 10 ⁻⁰⁸	1.01 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	1.30 x 10 ⁻⁰³	6.69 x 10 ⁻⁰⁵	9.11 x 10 ⁻⁰⁵	1.55 x 10 ⁻⁰³
(Lloyd Lake)								pper Bound Scena		•				
	Uranium-238	1.52 x 10 ⁻⁰⁶	4.18 x 10 ⁻¹⁵	4.49 x 10 ⁻⁰⁷	3.41 x 10 ⁻¹²	2.40 x 10 ⁻¹⁰	2.66 x 10 ⁻⁰⁶	3.27 x 10 ⁻⁰⁹	3.49 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.11 x 10 ⁻⁰⁶	8.66 x 10 ⁻⁰⁷	3.91 x 10 ⁻⁰⁷	8.04 x 10 ⁻⁰⁶
	Uranium-234	1.84 x 10 ⁻⁰⁶	1.02 x 10 ⁻¹⁴	4.91 x 10 ⁻⁰⁷	5.96 x 10 ⁻¹³	2.61 x 10 ⁻¹⁰	1.34 x 10 ⁻⁰⁸	3.57 x 10 ⁻⁰⁹	1.27 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	2.30 x 10 ⁻⁰⁶	9.43 x 10 ⁻⁰⁷	4.26 x 10 ⁻⁰⁷	6.02 x 10 ⁻⁰⁶
	Thorium-230	7.36 x 10 ⁻⁰⁶	2.48 x 10 ⁻¹⁴	3.29 x 10 ⁻⁰⁵	2.23 x 10 ⁻¹¹	1.25 x 10 ⁻⁰⁹	1.92 x 10 ⁻⁰⁸	3.36 x 10 ⁻⁰⁸	8.31 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	2.32 x 10 ⁻⁰⁵	4.17 x 10 ⁻⁰⁶	5.14 x 10 ⁻⁰⁵	1.19 x 10 ⁻⁰⁴
	Radium-226	1.84 x 10 ⁻⁰⁶	4.75 x 10 ⁻¹³	1.99 x 10 ⁻⁰⁶	8.87 x 10 ⁻¹⁰	1.69 x 10 ⁻⁰⁹	4.38 x 10 ⁻⁰⁵	8.73 x 10 ⁻⁰⁹	1.05 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	3.43 x 10 ⁻⁰⁶	9.75 x 10 ⁻⁰⁶	2.47 x 10 ⁻⁰⁶	6.44 x 10 ⁻⁰⁵
	Lead-210	5.78 x 10 ⁻⁰⁷	7.47 x 10 ⁻¹⁴	4.12 x 10 ⁻⁰⁷	3.63 x 10 ⁻¹³	2.69 x 10 ⁻⁰⁹	6.24 x 10 ⁻⁰⁷	7.40 x 10 ⁻⁰⁹	4.19 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	7.58 x 10 ⁻⁰⁴	1.91 x 10 ⁻⁰⁵	9.17 x 10 ⁻⁰⁶	7.88 x 10 ⁻⁰⁴
	Polonium-210	1.73 x 10 ⁻⁰⁶	6.52 x 10 ⁻¹⁶	7.06 x 10 ⁻⁰⁷	2.12 x 10 ⁻¹⁵	4.76 x 10 ⁻⁰⁹	1.41 x 10 ⁻¹⁰	1.30 x 10 ⁻⁰⁸	1.79 x 10 ⁻¹²	0.00 x 10 ⁺⁰⁰	5.21 x 10 ⁻⁰⁴	3.21 x 10 ⁻⁰⁵	4.48 x 10 ⁻⁰⁵	6.01 x 10 ⁻⁰⁴
	Total by Pathway	1.49 x 10 ⁻⁰⁵	5.89 x 10 ⁻¹³	3.69 x 10 ⁻⁰⁵	9.14 x 10 ⁻¹⁰	1.09 x 10 ⁻⁰⁸	4.71 x 10 ⁻⁰⁵	6.96 x 10 ⁻⁰⁸	1.08 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	1.31 x 10 ⁻⁰³	6.69 x 10 ⁻⁰⁵	1.09 x 10 ⁻⁰⁴	1.59 x 10 ⁻⁰³
						_		Application Case						
	Uranium-238	1.52 x 10 ⁻⁰⁶	5.44 x 10 ⁻¹⁵	1.13 x 10 ⁻⁰⁷	3.06 x 10 ⁻¹³	9.75 x 10 ⁻⁰⁹	3.56 x 10 ⁻⁰⁶	4.77 x 10 ⁻⁰⁸	1.63 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	8.37 x 10 ⁻⁰⁸	1.18 x 10 ⁻⁰⁶	1.38 x 10 ⁻⁰⁷	6.67 x 10 ⁻⁰⁶
	Uranium-234	1.78 x 10 ⁻⁰⁶	1.33 x 10 ⁻¹⁴	1.23 x 10 ⁻⁰⁷	5.36 x 10 ⁻¹⁴	1.06 x 10 ⁻⁰⁸	1.79 x 10 ⁻⁰⁸	5.19 x 10 ⁻⁰⁸	5.92 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	9.11 x 10 ⁻⁰⁸	1.28 x 10 ⁻⁰⁶	1.49 x 10 ⁻⁰⁷	3.50 x 10 ⁻⁰⁶
	Thorium-230	5.66 x 10 ⁻⁰⁶	3.22 x 10 ⁻¹⁴	1.65 x 10 ⁻⁰⁵	5.44 x 10 ⁻¹²	3.73 x 10 ⁻⁰⁸	2.18 x 10 ⁻⁰⁸	9.87 x 10 ⁻⁰⁷	1.07 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	5.87 x 10 ⁻⁰⁶	2.31 x 10 ⁻⁰⁶	2.46 x 10 ⁻⁰⁵	5.60 x 10 ⁻⁰⁵
	Radium-226	1.78 x 10 ⁻⁰⁶	6.17 x 10 ⁻¹³	1.70 x 10 ⁻⁰⁶	2.10 x 10 ⁻¹⁰	8.85 x 10 ⁻⁰⁸	5.87 x 10 ⁻⁰⁵	4.36 x 10 ⁻⁰⁷	1.30 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	1.18 x 10 ⁻⁰⁶	1.30 x 10 ⁻⁰⁵	9.76 x 10 ⁻⁰⁷	7.91 x 10 ⁻⁰⁵
	Lead-210	5.98 x 10 ⁻⁰⁷	9.70 x 10 ⁻¹⁴	1.13 x 10 ⁻⁰⁷	1.49 x 10 ⁻¹⁴	2.14 x 10 ⁻⁰⁷	8.35 x 10 ⁻⁰⁷	1.19 x 10 ⁻⁰⁸	1.09 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	2.55 x 10 ⁻⁰⁷	3.57 x 10 ⁻⁰⁵	8.72 x 10 ⁻⁰⁶	4.65 x 10 ⁻⁰⁵
	Polonium-210	1.78 x 10 ⁻⁰⁶	8.48 x 10 ⁻¹⁶	2.36 x 10 ⁻⁰⁷	9.19 x 10 ⁻¹⁷	5.32 x 10 ⁻⁰⁷	1.84 x 10 ⁻¹⁰	2.33 x 10 ⁻⁰⁸	3.73 x 10 ⁻¹⁴	0.00 x 10 ⁺⁰⁰	1.77 x 10 ⁻⁰⁶	8.86 x 10 ⁻⁰⁵	2.83 x 10 ⁻⁰⁵	1.21 x 10 ⁻⁰⁴
Seasonal Resident One-Year-Old	Total by Pathway	1.31 x 10 ⁻⁰⁵	7.66 x 10 ⁻¹³	1.88 x 10 ⁻⁰⁵	2.16 x 10 ⁻¹⁰	8.92 x 10 ⁻⁰⁷	6.31 x 10 ⁻⁰⁵	1.56 x 10 ⁻⁰⁶	1.32 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	9.25 x 10 ⁻⁰⁶	1.42 x 10 ⁻⁰⁴	6.30 x 10 ⁻⁰⁵	3.13 x 10 ⁻⁰⁴
(Lloyd Lake)							U	pper Bound Scena	rio					
	Uranium-238	1.52 x 10 ⁻⁰⁶	5.44 x 10 ⁻¹⁵	3.12 x 10 ⁻⁰⁷	8.42 x 10 ⁻¹³	9.75 x 10 ⁻⁰⁹	3.56 x 10 ⁻⁰⁶	1.33 x 10 ⁻⁰⁷	4.53 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.31 x 10 ⁻⁰⁷	1.18 x 10 ⁻⁰⁶	1.72 x 10 ⁻⁰⁷	7.16 x 10 ⁻⁰⁶
	Uranium-234	1.78 x 10 ⁻⁰⁶	1.33 x 10 ⁻¹⁴	3.39 x 10 ⁻⁰⁷	1.48 x 10 ⁻¹³	1.06 x 10 ⁻⁰⁸	1.79 x 10 ⁻⁰⁸	1.45 x 10 ⁻⁰⁷	1.65 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	2.51 x 10 ⁻⁰⁷	1.28 x 10 ⁻⁰⁶	1.86 x 10 ⁻⁰⁷	4.01 x 10 ⁻⁰⁶
	Thorium-230	5.66 x 10 ⁻⁰⁶	3.22 x 10 ⁻¹⁴	1.67 x 10 ⁻⁰⁵	5.52 x 10 ⁻¹²	3.73 x 10 ⁻⁰⁸	2.18 x 10 ⁻⁰⁸	1.00 x 10 ⁻⁰⁶	1.08 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	5.95 x 10 ⁻⁰⁶	2.31 x 10 ⁻⁰⁶	2.50 x 10 ⁻⁰⁵	5.67 x 10 ⁻⁰⁵
	Radium-226	1.78 x 10 ⁻⁰⁶	6.17 x 10 ⁻¹³	1.78 x 10 ⁻⁰⁶	2.20 x 10 ⁻¹⁰	8.85 x 10 ⁻⁰⁸	5.87 x 10 ⁻⁰⁵	4.56 x 10 ⁻⁰⁷	1.37 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	1.23 x 10 ⁻⁰⁶	1.30 x 10 ⁻⁰⁵	9.83 x 10 ⁻⁰⁷	7.94 x 10 ⁻⁰⁵
	Lead-210	5.98 x 10 ⁻⁰⁷	9.70 x 10 ⁻¹⁴	5.61 x 10 ⁻⁰⁷	7.37 x 10 ⁻¹⁴	2.14 x 10 ⁻⁰⁷	8.35 x 10 ⁻⁰⁷	5.89 x 10 ⁻⁰⁷	5.44 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	1.31 x 10 ⁻⁰⁶	3.57 x 10 ⁻⁰⁵	1.08 x 10 ⁻⁰⁵	5.06 x 10 ⁻⁰⁵
	Polonium-210	1.78 x 10 ⁻⁰⁶	8.48 x 10 ⁻¹⁶	1.35 x 10 ⁻⁰⁶	5.25 x 10 ⁻¹⁶	5.32 x 10 ⁻⁰⁷	1.84 x 10 ⁻¹⁰	1.46 x 10 ⁻⁰⁶	2.32 x 10 ⁻¹²	0.00 x 10 ⁺⁰⁰	1.10 x 10 ⁻⁰⁵	8.86 x 10 ⁻⁰⁵	4.68 x 10 ⁻⁰⁵	1.52 x 10 ⁻⁰⁴
	Total by Pathway	1.31 x 10 ⁻⁰⁵	7.66 x 10 ⁻¹³	2.11 x 10 ⁻⁰⁵	2.26 x 10 ⁻¹⁰	8.92 x 10 ⁻⁰⁷	6.31 x 10 ⁻⁰⁵	3.78 x 10 ⁻⁰⁶	1.41 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	2.00 x 10 ⁻⁰⁵	1.42 x 10 ⁻⁰⁴	8.39 x 10 ⁻⁰⁵	3.49 x 10 ⁻⁰⁴
								Application Case						
Permanent	Uranium-238	n/a												
Resident	Uranium-234	n/a												
(Patterson Lake North Arm - West	Thorium-230	n/a												
Basin)	Radium-226	n/a												
Dasiii)														

							Incremental Dose b	y Pathway - Proje	ct Lifespan (mSv/y	r)				
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by Radionuclide
	Polonium-210	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Total by Pathway	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
							U _l	per Bound Scena	rio					
	Uranium-238	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Uranium-234	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Thorium-230	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Radium-226	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Lead-210	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Polonium-210	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Total by Pathway	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
								Application Case						
	Uranium-238	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Uranium-234	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Thorium-230	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Radium-226	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
Permanent	Lead-210	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
Resident One-	Polonium-210	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
Year-Old	Total by Pathway	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
(Patterson Lake						_	Ul	pper Bound Scena	rio					
North Arm - West Basin)	Uranium-238	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
Dasilly	Uranium-234	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Thorium-230	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Radium-226	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Lead-210	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Polonium-210	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Total by Pathway	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a

mSv/yr = millisieverts per year; n/a = not applicable or not assessed.

Table 5-10: Estimated Radiation Doses to Human Receptors – Far-Future – Application Case, Upper Bound Scenario

						lr	cremental Dose by	Pathway - Far-Fu	ıture (mSv/yr)					
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by Radionuclide
			1	. ,	,	,	•	plication Case	,					
	Uranium-238	n/a												
	Uranium-234	n/a												
	Thorium-230	n/a												
	Radium-226	n/a												
	Lead-210	n/a												
Camp Worker	Polonium-210	n/a												
(Patterson Lake	Total by Pathway	n/a												
North Arm - West Basin)		:- /-	/-	- /-	/-	/-		r Bound Scenario	1	/	- /-	/-	/-	
west basili)	Uranium-238	n/a												
	Uranium-234	n/a												
	Thorium-230	n/a												
	Radium-226	n/a												
	Lead-210	n/a												
	Polonium-210	n/a												
	Total by Pathway	n/a												
		0.00 40+00	0.00 4.0+00	2.64 40:05	2.00 4.0-10	2.46 40-09		plication Case	2.57 40.06	0.00 40+00	7.25 40-05	0.50 40-06	2.47. 40-05	4.74 40.04
	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.64 x 10 ⁻⁰⁵	2.00 x 10 ⁻¹⁰	2.16 x 10 ⁻⁰⁹	2.39 x 10 ⁻⁰⁵	2.41 x 10 ⁻⁰⁷	2.57 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	7.35 x 10 ⁻⁰⁵	9.59 x 10 ⁻⁰⁶	3.47 x 10 ⁻⁰⁵	1.71 x 10 ⁻⁰⁴
	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.88 x 10 ⁻⁰⁵	3.50 x 10 ⁻¹¹	2.35 x 10 ⁻⁰⁹	1.21 x 10 ⁻⁰⁷	2.63 x 10 ⁻⁰⁷	9.34 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	8.03 x 10 ⁻⁰⁵	1.04 x 10 ⁻⁰⁵	3.79 x 10 ⁻⁰⁵	1.58 x 10 ⁻⁰⁴
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.54 x 10 ⁻⁰⁵	1.04 x 10 ⁻¹¹	5.98 x 10 ⁻⁰⁸	9.16 x 10 ⁻⁰⁷	1.68 x 10 ⁻⁰⁸	4.16 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	2.06 x 10 ⁻⁰⁵	5.13 x 10 ⁻⁰⁶	1.90 x 10 ⁻⁰⁴	2.32 x 10 ⁻⁰⁴
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	3.14 x 10 ⁻⁰⁵	1.40 x 10 ⁻⁰⁸	9.75 x 10 ⁻⁰⁸	2.53 x 10 ⁻⁰³	1.74 x 10 ⁻⁰⁷	2.08 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	8.37 x 10 ⁻⁰⁵	1.61 x 10 ⁻⁰⁴	3.76 x 10 ⁻⁰⁴	3.20 x 10 ⁻⁰³
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.02 x 10 ⁻⁰⁶	1.78 x 10 ⁻¹²	1.56 x 10 ⁻¹⁰	3.63 x 10 ⁻⁰⁸	6.81 x 10 ⁻⁰⁸	3.85 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	1.80 x 10 ⁻⁰⁵	3.45 x 10 ⁻⁰⁸	4.02 x 10 ⁻⁰⁴	4.22 x 10 ⁻⁰⁴
Subsistence	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	3.55 x 10 ⁻⁰⁶	1.07 x 10 ⁻¹⁴	2.78 x 10 ⁻¹⁰	8.24 x 10 ⁻¹²	1.20 x 10 ⁻⁰⁷	1.65 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	1.13 x 10 ⁻⁰⁴	2.95 x 10 ⁻⁰⁷	3.22 x 10 ⁻⁰⁴	4.39 x 10 ⁻⁰⁴
Harvester	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.08 x 10 ⁻⁰⁴	1.42 x 10 ⁻⁰⁸	1.62 x 10 ⁻⁰⁷	2.55 x 10 ⁻⁰³	8.82 x 10 ⁻⁰⁷	2.34 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	3.89 x 10 ⁻⁰⁴	1.86 x 10 ⁻⁰⁴	1.36 x 10 ⁻⁰³	4.62 x 10 ⁻⁰³
(Patterson Lake			1		10	1 00		r Bound Scenario		1 00	0.5		0.5	-
South Arm)	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.64 x 10 ⁻⁰⁵	2.00 x 10 ⁻¹⁰	2.16 x 10 ⁻⁰⁹	2.39 x 10 ⁻⁰⁵	2.41 x 10 ⁻⁰⁷	2.57 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	7.35 x 10 ⁻⁰⁵	9.59 x 10 ⁻⁰⁶	3.47 x 10 ⁻⁰⁵	1.71 x 10 ⁻⁰⁴
	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.89 x 10 ⁻⁰⁵	3.51 x 10 ⁻¹¹	2.35 x 10 ⁻⁰⁹	1.21 x 10 ⁻⁰⁷	2.63 x 10 ⁻⁰⁷	9.35 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	8.04 x 10 ⁻⁰⁵	1.04 x 10 ⁻⁰⁵	3.79 x 10 ⁻⁰⁵	1.58 x 10 ⁻⁰⁴
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.54 x 10 ⁻⁰⁵	1.04 x 10 ⁻¹¹	5.98 x 10 ⁻⁰⁸	9.16 x 10 ⁻⁰⁷	1.68 x 10 ⁻⁰⁸	4.16 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	2.06 x 10 ⁻⁰⁵	5.13 x 10 ⁻⁰⁶	1.90 x 10 ⁻⁰⁴	2.32 x 10 ⁻⁰⁴
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	3.14 x 10 ⁻⁰⁵	1.40 x 10 ⁻⁰⁸	9.75 x 10 ⁻⁰⁸	2.53 x 10 ⁻⁰³	1.74 x 10 ⁻⁰⁷	2.09 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	8.39 x 10 ⁻⁰⁵	1.61 x 10 ⁻⁰⁴	3.76 x 10 ⁻⁰⁴	3.20 x 10 ⁻⁰³
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.03 x 10 ⁻⁰⁶	1.79 x 10 ⁻¹²	1.56 x 10 ⁻¹⁰	3.63 x 10 ⁻⁰⁸	6.84 x 10 ⁻⁰⁸	3.87 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	1.81 x 10 ⁻⁰⁵	3.45 x 10 ⁻⁰⁸	4.03 x 10 ⁻⁰⁴	4.24 x 10 ⁻⁰⁴
	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	3.57 x 10 ⁻⁰⁶	1.07 x 10 ⁻¹⁴	2.78 x 10 ⁻¹⁰	8.24 x 10 ⁻¹²	1.21 x 10 ⁻⁰⁷	1.66 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	1.13 x 10 ⁻⁰⁴	2.95 x 10 ⁻⁰⁷	3.24 x 10 ⁻⁰⁴	4.41 x 10 ⁻⁰⁴
	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.08 x 10 ⁻⁰⁴	1.43 x 10 ⁻⁰⁸	1.62 x 10 ⁻⁰⁷	2.55 x 10 ⁻⁰³	8.84 x 10 ⁻⁰⁷	2.34 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	3.90 x 10 ⁻⁰⁴	1.86 x 10 ⁻⁰⁴	1.37 x 10 ⁻⁰³	4.62 x 10 ⁻⁰³
					İ			plication Case		1		•		
	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.83 x 10 ⁻⁰⁵	4.95 x 10 ⁻¹¹	8.76 x 10 ⁻⁰⁸	3.20 x 10 ⁻⁰⁵	9.78 x 10 ⁻⁰⁶	3.34 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	4.34 x 10 ⁻⁰⁵	1.18 x 10 ⁻⁰⁵	1.59 x 10 ⁻⁰⁵	1.35 x 10 ⁻⁰⁴
	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.99 x 10 ⁻⁰⁵	8.67 x 10 ⁻¹²	9.49 x 10 ⁻⁰⁸	1.61 x 10 ⁻⁰⁷	1.06 x 10 ⁻⁰⁵	1.21 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	4.72 x 10 ⁻⁰⁵	1.27 x 10 ⁻⁰⁵	1.73 x 10 ⁻⁰⁵	1.08 x 10 ⁻⁰⁴
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	7.83 x 10 ⁻⁰⁶	2.58 x 10 ⁻¹²	1.78 x 10 ⁻⁰⁶	1.04 x 10 ⁻⁰⁶	5.00 x 10 ⁻⁰⁷	5.40 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	8.91 x 10 ⁻⁰⁶	4.61 x 10 ⁻⁰⁶	9.15 x 10 ⁻⁰⁵	1.16 x 10 ⁻⁰⁴
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.80 x 10 ⁻⁰⁵	3.47 x 10 ⁻⁰⁹	5.10 x 10 ⁻⁰⁶	3.38 x 10 ⁻⁰³	9.08 x 10 ⁻⁰⁶	2.71 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	6.35 x 10 ⁻⁰⁵	2.54 x 10 ⁻⁰⁴	1.70 x 10 ⁻⁰⁴	3.94 x 10 ⁻⁰³
Subsistence	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.75 x 10 ⁻⁰⁶	3.62 x 10 ⁻¹³	1.26 x 10 ⁻⁰⁸	4.89 x 10 ⁻⁰⁸	5.42 x 10 ⁻⁰⁶	5.01 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	2.08 x 10 ⁻⁰⁵	8.29 x 10 ⁻⁰⁸	5.02 x 10 ⁻⁰⁴	5.32 x 10 ⁻⁰⁴
Harvester One-	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	6.79 x 10 ⁻⁰⁶	2.64 x 10 ⁻¹⁵	3.10 x 10 ⁻⁰⁸	1.08 x 10 ⁻¹¹	1.35 x 10 ⁻⁰⁵	2.15 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	1.83 x 10 ⁻⁰⁴	9.95 x 10 ⁻⁰⁷	3.77 x 10 ⁻⁰⁴	5.81 x 10 ⁻⁰⁴
Year-Old	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	8.37 x 10 ⁻⁰⁵	3.53 x 10 ⁻⁰⁹	7.11 x 10 ⁻⁰⁶	3.41 x 10 ⁻⁰³	4.89 x 10 ⁻⁰⁵	3.05 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	3.67 x 10 ⁻⁰⁴	2.84 x 10 ⁻⁰⁴	1.17 x 10 ⁻⁰³	5.41 x 10 ⁻⁰³
(Patterson Lake					•			r Bound Scenario				•		
South Arm)	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.84 x 10 ⁻⁰⁵	4.95 x 10 ⁻¹¹	8.76 x 10 ⁻⁰⁸	3.20 x 10 ⁻⁰⁵	9.79 x 10 ⁻⁰⁶	3.34 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	4.34 x 10 ⁻⁰⁵	1.18 x 10 ⁻⁰⁵	1.59 x 10 ⁻⁰⁵	1.35 x 10 ⁻⁰⁴
	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.00 x 10 ⁻⁰⁵	8.68 x 10 ⁻¹²	9.49 x 10 ⁻⁰⁸	1.61 x 10 ⁻⁰⁷	1.07 x 10 ⁻⁰⁵	1.22 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	4.72 x 10 ⁻⁰⁵	1.27 x 10 ⁻⁰⁵	1.73 x 10 ⁻⁰⁵	1.08 x 10 ⁻⁰⁴
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	7.83 x 10 ⁻⁰⁶	2.58 x 10 ⁻¹²	1.78 x 10 ⁻⁰⁶	1.04 x 10 ⁻⁰⁶	5.00 x 10 ⁻⁰⁷	5.40 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	8.91 x 10 ⁻⁰⁶	4.61 x 10 ⁻⁰⁶	9.15 x 10 ⁻⁰⁵	1.16 x 10 ⁻⁰⁴
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.81 x 10 ⁻⁰⁵	3.47 x 10 ⁻⁰⁹	5.10 x 10 ⁻⁰⁶	3.38 x 10 ⁻⁰³	9.10 x 10 ⁻⁰⁶	2.72 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	6.37 x 10 ⁻⁰⁵	2.54 x 10 ⁻⁰⁴	1.70 x 10 ⁻⁰⁴	3.94 x 10 ⁻⁰³
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.76 x 10 ⁻⁰⁶	3.63 x 10 ⁻¹³	1.26 x 10 ⁻⁰⁸	4.89 x 10 ⁻⁰⁸	5.44 x 10 ⁻⁰⁶	5.03 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	2.09 x 10 ⁻⁰⁵	8.29 x 10 ⁻⁰⁸	5.05 x 10 ⁻⁰⁴	5.34 x 10 ⁻⁰⁴
	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	6.82 x 10 ⁻⁰⁶	2.66 x 10 ⁻¹⁵	3.10 x 10 ⁻⁰⁸	1.08 x 10 ⁻¹¹	1.35 x 10 ⁻⁰⁵	2.16 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	1.84 x 10 ⁻⁰⁴	9.95 x 10 ⁻⁰⁷	3.78 x 10 ⁻⁰⁴	5.84 x 10 ⁻⁰⁴
	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	8.38 x 10 ⁻⁰⁵	3.54 x 10 ⁻⁰⁹	7.11 x 10 ⁻⁰⁶	3.41 x 10 ⁻⁰³	4.90 x 10 ⁻⁰⁵	3.06 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	3.68 x 10 ⁻⁰⁴	2.84 x 10 ⁻⁰⁴	1.18 x 10 ⁻⁰³	5.41 x 10 ⁻⁰³
							Ар	plication Case						
Subsistence	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	9.81 x 10 ⁻⁰⁶	7.44 x 10 ⁻¹¹	1.26 x 10 ⁻⁰⁹	1.39 x 10 ⁻⁰⁵	8.94 x 10 ⁻⁰⁸	9.54 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	2.73 x 10 ⁻⁰⁵	5.59 x 10 ⁻⁰⁶	1.18 x 10 ⁻⁰⁵	6.95 x 10 ⁻⁰⁵
Harvester (Beet	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.07 x 10 ⁻⁰⁵	1.30 x 10 ⁻¹¹	1.37 x 10 ⁻⁰⁹	7.02 x 10 ⁻⁰⁸	9.78 x 10 ⁻⁰⁸	3.47 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	2.99 x 10 ⁻⁰⁵	6.09 x 10 ⁻⁰⁶	1.29 x 10 ⁻⁰⁵	5.97 x 10 ⁻⁰⁵
Lake)	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	8.19 x 10 ⁻⁰⁶	5.55 x 10 ⁻¹²	3.49 x 10 ⁻⁰⁸	5.34 x 10 ⁻⁰⁷	8.93 x 10 ⁻⁰⁹	2.21 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	1.10 x 10 ⁻⁰⁵	2.99 x 10 ⁻⁰⁶	4.72 x 10 ⁻⁰⁵	7.00 x 10 ⁻⁰⁵
Lake)	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.35 x 10 ⁻⁰⁵	6.02 x 10 ⁻⁰⁹	5.68 x 10 ⁻⁰⁸	1.47 x 10 ⁻⁰³	7.47 x 10 ⁻⁰⁸	8.95 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	3.60 x 10 ⁻⁰⁵	9.37 x 10 ⁻⁰⁵	9.19 x 10 ⁻⁰⁵	1.72 x 10 ⁻⁰³
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.79 x 10 ⁻⁰⁷	2.46 x 10 ⁻¹³	9.19 x 10 ⁻¹¹	2.12 x 10 ⁻⁰⁸	9.37 x 10 ⁻⁰⁹	5.30 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	2.48 x 10 ⁻⁰⁶	2.00 x 10 ⁻⁰⁸	4.16 x 10 ⁻⁰⁶	6.97 x 10 ⁻⁰⁶

							ncremental Dose by	Pathway - Far-Fu	ture (mSv/vr)					
	Radionuclide	Ata (lasta and al)	Air (Fortower)	Water	Water			Sediment	Sediment	Ati - Dianta	Aquatic	Terrestrial	Terrestrial	Total by
		Air (Internal)	Air (External)	(Internal)	(External)	Soil (Internal)	Soil (External)	(Internal)	(External)	Aquatic Plants	Animals	Plants	Animals	Radionuclide
	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	4.75 x 10 ⁻⁰⁷	1.42 x 10 ⁻¹⁵	1.62 x 10 ⁻¹⁰	4.80 x 10 ⁻¹²	1.65 x 10 ⁻⁰⁸	2.27 x 10 ⁻¹²	0.00 x 10 ⁺⁰⁰	1.51 x 10 ⁻⁰⁵	1.72 x 10 ⁻⁰⁷	3.27 x 10 ⁻⁰⁵	4.85 x 10 ⁻⁰⁵
	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	4.30 x 10 ⁻⁰⁵	6.11 x 10 ⁻⁰⁹	9.46 x 10 ⁻⁰⁸	1.49 x 10 ⁻⁰³	2.97 x 10 ⁻⁰⁷	9.91 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	1.22 x 10 ⁻⁰⁴	1.09 x 10 ⁻⁰⁴	2.01 x 10 ⁻⁰⁴	1.97 x 10 ⁻⁰³
								Bound Scenario						
	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	9.82 x 10 ⁻⁰⁶	7.45 x 10 ⁻¹¹	1.26 x 10 ⁻⁰⁹	1.39 x 10 ⁻⁰⁵	8.95 x 10 ⁻⁰⁸	9.55 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	2.73 x 10 ⁻⁰⁵	5.59 x 10 ⁻⁰⁶	1.18 x 10 ⁻⁰⁵	6.95 x 10 ⁻⁰⁵
	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.07 x 10 ⁻⁰⁵	1.30 x 10 ⁻¹¹	1.37 x 10 ⁻⁰⁹	7.02 x 10 ⁻⁰⁸	9.79 x 10 ⁻⁰⁸	3.48 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	2.99 x 10 ⁻⁰⁵	6.09 x 10 ⁻⁰⁶	1.29 x 10 ⁻⁰⁵	5.98 x 10 ⁻⁰⁵
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	8.19 x 10 ⁻⁰⁶	5.55 x 10 ⁻¹²	3.49 x 10 ⁻⁰⁸	5.34 x 10 ⁻⁰⁷	8.93 x 10 ⁻⁰⁹	2.21 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	1.10 x 10 ⁻⁰⁵	2.99 x 10 ⁻⁰⁶	4.72 x 10 ⁻⁰⁵	7.00 x 10 ⁻⁰⁵
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.35 x 10 ⁻⁰⁵	6.03 x 10 ⁻⁰⁹	5.68 x 10 ⁻⁰⁸	1.47 x 10 ⁻⁰³	7.49 x 10 ⁻⁰⁸	8.97 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	3.61 x 10 ⁻⁰⁵	9.37 x 10 ⁻⁰⁵	9.19 x 10 ⁻⁰⁵	1.72 x 10 ⁻⁰³
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.80 x 10 ⁻⁰⁷	2.47 x 10 ⁻¹³	9.19 x 10 ⁻¹¹	2.12 x 10 ⁻⁰⁸	9.41 x 10 ⁻⁰⁹	5.32 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	2.49 x 10 ⁻⁰⁶	2.00 x 10 ⁻⁰⁸	4.18 x 10 ⁻⁰⁶	7.00 x 10 ⁻⁰⁶
	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	4.76 x 10 ⁻⁰⁷	1.43 x 10 ⁻¹⁵	1.62 x 10 ⁻¹⁰	4.80 x 10 ⁻¹² 1.49 x 10 ⁻⁰³	1.66 x 10 ⁻⁰⁸	2.28 x 10 ⁻¹²	0.00 x 10 ⁺⁰⁰	1.52 x 10 ⁻⁰⁵	1.72 x 10 ⁻⁰⁷	3.28 x 10 ⁻⁰⁵	4.87 x 10 ⁻⁰⁵
	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	4.30 x 10 ⁻⁰⁵	6.13 x 10 ⁻⁰⁹	9.46 x 10 ⁻⁰⁸		2.97 x 10 ⁻⁰⁷ plication Case	9.93 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	1.22 x 10 ⁻⁰⁴	1.09 x 10 ⁻⁰⁴	2.01 x 10 ⁻⁰⁴	1.97 x 10 ⁻⁰³
	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	6.81 x 10 ⁻⁰⁶	1.84 x 10 ⁻¹¹	5.11 x 10 ⁻⁰⁸	1.86 x 10 ⁻⁰⁵	3.64 x 10 ⁻⁰⁶	1.24 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	1.61 x 10 ⁻⁰⁵	6.86 x 10 ⁻⁰⁶	5.35 x 10 ⁻⁰⁶	5.87 x 10 ⁻⁰⁵
	Uranium-234	0.00 x 10 40 0.00 x 10+00	0.00 x 10 40 0.00 x 10 400	7.41 x 10 ⁻⁰⁶	3.22 x 10 ⁻¹²	5.53 x 10 ⁻⁰⁸	9.40 x 10 ⁻⁰⁸	3.96 x 10 ⁻⁰⁶	4.51 x 10 ⁻⁰⁹	0.00 x 10 40 0.00 x 10+00	1.75 x 10 ⁻⁰⁵	7.43 x 10 ⁻⁰⁶	5.81 x 10 ⁻⁰⁶	4.23 x 10 ⁻⁰⁵
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	4.17 x 10 ⁻⁰⁶	1.37 x 10 ⁻¹²	1.04 x 10 ⁻⁰⁶	6.07 x 10 ⁻⁰⁷	2.66 x 10 ⁻⁰⁷	2.87 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	4.74 x 10 ⁻⁰⁶	2.68 x 10 ⁻⁰⁶	2.27 x 10 ⁻⁰⁵	3.62 x 10 ⁻⁰⁵
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.20 x 10 ⁻⁰⁵	1.49 x 10 ⁻⁰⁹	2.97 x 10 ⁻⁰⁶	1.97 x 10 ⁻⁰³	3.90 x 10 ⁻⁰⁶	1.17 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	2.73 x 10 ⁻⁰⁵	1.48 x 10 ⁻⁰⁴	3.56 x 10 ⁻⁰⁵	2.21 x 10 ⁻⁰³
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	3.79 x 10 ⁻⁰⁷	4.98 x 10 ⁻¹⁴	7.33 x 10 ⁻⁰⁹	2.89 x 10 ⁻⁰⁸	7.46 x 10 ⁻⁰⁷	6.89 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	2.86 x 10 ⁻⁰⁶	4.94 x 10 ⁻⁰⁸	4.94 x 10 ⁻⁰⁶	9.02 x 10 ⁻⁰⁶
Subsistence	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	9.06 x 10 ⁻⁰⁷	3.53 x 10 ⁻¹⁶	1.81 x 10 ⁻⁰⁸	6.28 x 10 ⁻¹²	1.85 x 10 ⁻⁰⁶	2.95 x 10 ⁻¹²	0.00 x 10 ⁺⁰⁰	2.45 x 10 ⁻⁰⁵	5.79 x 10 ⁻⁰⁷	4.15 x 10 ⁻⁰⁵	6.94 x 10 ⁻⁰⁵
Harvester One-	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	3.17 x 10 ⁻⁰⁵	1.51 x 10 ⁻⁰⁹	4.14 x 10 ⁻⁰⁶	1.99 x 10 ⁻⁰³	1.44 x 10 ⁻⁰⁵	1.29 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	9.31 x 10 ⁻⁰⁵	1.65 x 10 ⁻⁰⁴	1.16 x 10 ⁻⁰⁴	2.43 x 10 ⁻⁰³
Year-Old (Beet	, , , , , , , , , , , , , , , , , , , ,							Bound Scenario						
Lake)	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	6.82 x 10 ⁻⁰⁶	1.84 x 10 ⁻¹¹	5.11 x 10 ⁻⁰⁸	1.86 x 10 ⁻⁰⁵	3.64 x 10 ⁻⁰⁶	1.24 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	1.61 x 10 ⁻⁰⁵	6.86 x 10 ⁻⁰⁶	5.35 x 10 ⁻⁰⁶	5.87 x 10 ⁻⁰⁵
	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	7.42 x 10 ⁻⁰⁶	3.23 x 10 ⁻¹²	5.53 x 10 ⁻⁰⁸	9.40 x 10 ⁻⁰⁸	3.96 x 10 ⁻⁰⁶	4.52 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	1.75 x 10 ⁻⁰⁵	7.43 x 10 ⁻⁰⁶	5.81 x 10 ⁻⁰⁶	4.23 x 10 ⁻⁰⁵
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	4.17 x 10 ⁻⁰⁶	1.37 x 10 ⁻¹²	1.04 x 10 ⁻⁰⁶	6.07 x 10 ⁻⁰⁷	2.66 x 10 ⁻⁰⁷	2.87 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	4.74 x 10 ⁻⁰⁶	2.68 x 10 ⁻⁰⁶	2.27 x 10 ⁻⁰⁵	3.62 x 10 ⁻⁰⁵
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.21 x 10 ⁻⁰⁵	1.49 x 10 ⁻⁰⁹	2.97 x 10 ⁻⁰⁶	1.97 x 10 ⁻⁰³	3.91 x 10 ⁻⁰⁶	1.17 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	2.74 x 10 ⁻⁰⁵	1.48 x 10 ⁻⁰⁴	3.56 x 10 ⁻⁰⁵	2.21 x 10 ⁻⁰³
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	3.81 x 10 ⁻⁰⁷	5.00 x 10 ⁻¹⁴	7.33 x 10 ⁻⁰⁹	2.89 x 10 ⁻⁰⁸	7.49 x 10 ⁻⁰⁷	6.92 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	2.88 x 10 ⁻⁰⁶	4.94 x 10 ⁻⁰⁸	4.95 x 10 ⁻⁰⁶	9.05 x 10 ⁻⁰⁶
	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	9.10 x 10 ⁻⁰⁷	3.55 x 10 ⁻¹⁶	1.81 x 10 ⁻⁰⁸	6.28 x 10 ⁻¹²	1.86 x 10 ⁻⁰⁶	2.96 x 10 ⁻¹²	0.00 x 10 ⁺⁰⁰	2.46 x 10 ⁻⁰⁵	5.79 x 10 ⁻⁰⁷	4.17 x 10 ⁻⁰⁵	6.97 x 10 ⁻⁰⁵
	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	3.18 x 10 ⁻⁰⁵	1.52 x 10 ⁻⁰⁹	4.14 x 10 ⁻⁰⁶	1.99 x 10 ⁻⁰³	1.44 x 10 ⁻⁰⁵	1.29 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	9.33 x 10 ⁻⁰⁵	1.65 x 10 ⁻⁰⁴	1.16 x 10 ⁻⁰⁴	2.43 x 10 ⁻⁰³
								plication Case						
	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	8.88 x 10 ⁻⁰⁷	6.74 x 10 ⁻¹²	4.20 x 10 ⁻¹¹	4.66 x 10 ⁻⁰⁷	8.10 x 10 ⁻⁰⁹	8.64 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.47 x 10 ⁻⁰⁶	1.87 x 10 ⁻⁰⁷	7.85 x 10 ⁻⁰⁷	4.89 x 10 ⁻⁰⁶
	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	9.71 x 10 ⁻⁰⁷	1.18 x 10 ⁻¹²	4.58 x 10 ⁻¹¹	2.35 x 10 ⁻⁰⁹	8.85 x 10 ⁻⁰⁹	3.14 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	2.70 x 10 ⁻⁰⁶	2.04 x 10 ⁻⁰⁷	8.57 x 10 ⁻⁰⁷	4.75 x 10 ⁻⁰⁶
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	8.40 x 10 ⁻⁰⁷	5.69 x 10 ⁻¹³	1.17 x 10 ⁻⁰⁹	1.79 x 10 ⁻⁰⁸	9.16 x 10 ⁻¹⁰	2.26 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	1.12 x 10 ⁻⁰⁶	1.00 x 10 ⁻⁰⁷	3.79 x 10 ⁻⁰⁶	5.87 x 10 ⁻⁰⁶
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.29 x 10 ⁻⁰⁶	5.76 x 10 ⁻¹⁰	1.90 x 10 ⁻⁰⁹	4.92 x 10 ⁻⁰⁵	7.15 x 10 ⁻⁰⁹	8.56 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	3.45 x 10 ⁻⁰⁶	3.13 x 10 ⁻⁰⁶	3.55 x 10 ⁻⁰⁶	6.15 x 10 ⁻⁰⁵
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.04 x 10 ⁻⁰⁸	1.81 x 10 ⁻¹⁴	2.73 x 10 ⁻¹²	6.98 x 10 ⁻¹⁰	6.88 x 10 ⁻¹⁰	3.89 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	1.83 x 10 ⁻⁰⁷	4.66 x 10 ⁻¹⁰	3.09 x 10 ⁻⁰⁷	5.14 x 10 ⁻⁰⁷
Subsistence	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	5.91 x 10 ⁻¹²	1.71 x 10 ⁻¹³	1.14 x 10 ⁻⁰⁹	1.56 x 10 ⁻¹³	0.00 x 10 ⁺⁰⁰	5.96 x 10 ⁻⁰⁸	6.05 x 10 ⁻⁰⁹	2.18 x 10 ⁻⁰⁶	2.24 x 10 ⁻⁰⁶
Harvester (Lloyd	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	4.01 x 10 ⁻⁰⁶	5.84 x 10 ⁻¹⁰	3.16 x 10 ⁻⁰⁹	4.97 x 10 ⁻⁰⁵	2.68 x 10 ⁻⁰⁸	9.43 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	9.99 x 10 ⁻⁰⁶	3.63 x 10 ⁻⁰⁶	1.15 x 10 ⁻⁰⁵	7.98 x 10 ⁻⁰⁵
Lake)	H	0.00 10+00	0.00 x 10 ⁺⁰⁰	0.00 10-07	C 75 10-12	4 20 10-11		Bound Scenario	0.65 10-08	0.00 10+00	2.40 10-06	1.07 10-07	7.05 10-07	4.00 10-06
	Uranium-238 Uranium-234	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	0.00 x 10 ⁺⁰⁰	8.89 x 10 ⁻⁰⁷ 9.72 x 10 ⁻⁰⁷	6.75 x 10 ⁻¹² 1.18 x 10 ⁻¹²	4.20 x 10 ⁻¹¹ 4.58 x 10 ⁻¹¹	4.66 x 10 ⁻⁰⁷ 2.35 x 10 ⁻⁰⁹	8.11 x 10 ⁻⁰⁹ 8.86 x 10 ⁻⁰⁹	8.65 x 10 ⁻⁰⁸ 3.15 x 10 ⁻¹⁰	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	2.48 x 10 ⁻⁰⁶ 2.71 x 10 ⁻⁰⁶	1.87 x 10 ⁻⁰⁷ 2.04 x 10 ⁻⁰⁷	7.85 x 10 ⁻⁰⁷ 8.57 x 10 ⁻⁰⁷	4.90 x 10 ⁻⁰⁶ 4.75 x 10 ⁻⁰⁶
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	9.72 x 10 ⁻⁰⁷	5.69 x 10 ⁻¹³	1.17 x 10 ⁻⁰⁹	1.79 x 10 ⁻⁰⁸	9.16 x 10 ⁻¹⁰	2.26 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	1.12 x 10 ⁻⁰⁶	1.00 x 10 ⁻⁰⁷	3.79 x 10 ⁻⁰⁶	5.87 x 10 ⁻⁰⁶
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 0.00 x 10 ⁺⁰⁰	1.29 x 10 ⁻⁰⁶	5.77 x 10 ⁻¹⁰	1.90 x 10 ⁻⁰⁹	4.92 x 10 ⁻⁰⁵	7.16 x 10 ⁻⁰⁹	8.58 x 10 ⁻⁰⁷	0.00 x 10 0.00 x 10 ⁺⁰⁰	3.45 x 10 ⁻⁰⁶	3.13 x 10 ⁻⁰⁶	3.55 x 10 ⁻⁰⁶	6.15 x 10 ⁻⁰⁵
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.05 x 10 ⁻⁰⁸	1.82 x 10 ⁻¹⁴	2.73 x 10 ⁻¹²	6.98 x 10 ⁻¹⁰	6.91 x 10 ⁻¹⁰	3.91 x 10 ⁻¹¹	0.00 x 10 0.00 x 10+00	1.83 x 10 ⁻⁰⁷	4.66 x 10 ⁻¹⁰	3.09 x 10 ⁻⁰⁷	5.15 x 10 ⁻⁰⁷
	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	5.91 x 10 ⁻¹²	1.71 x 10 ⁻¹³	1.14 x 10 ⁻⁰⁹	1.57 x 10 ⁻¹³	0.00 x 10 ⁺⁰⁰	5.96 x 10 ⁻⁰⁸	6.05 x 10 ⁻⁰⁹	2.18 x 10 ⁻⁰⁶	2.24 x 10 ⁻⁰⁶
	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	4.01 x 10 ⁻⁰⁶	5.86 x 10 ⁻¹⁰	3.16 x 10 ⁻⁰⁹	4.97 x 10 ⁻⁰⁵	2.69 x 10 ⁻⁰⁸	9.45 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.00 x 10 ⁻⁰⁵	3.63 x 10 ⁻⁰⁶	1.15 x 10 ⁻⁰⁵	7.98 x 10 ⁻⁰⁵
		0.00 X 10	0.00 % 10	no i x i o	Side X 10	5110 X 10		plication Case	31.10 X 10	616 G X 10	1,00 X 10	5.05 X 10	1113 % 10	7.50 X 10
	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	6.17 x 10 ⁻⁰⁷	1.67 x 10 ⁻¹²	1.71 x 10 ⁻⁰⁹	6.24 x 10 ⁻⁰⁷	3.29 x 10 ⁻⁰⁷	1.12 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.46 x 10 ⁻⁰⁶	2.29 x 10 ⁻⁰⁷	3.58 x 10 ⁻⁰⁷	3.73 x 10 ⁻⁰⁶
	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	6.71 x 10 ⁻⁰⁷	2.92 x 10 ⁻¹³	1.85 x 10 ⁻⁰⁹	3.14 x 10 ⁻⁰⁹	3.58 x 10 ⁻⁰⁷	4.09 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	1.59 x 10 ⁻⁰⁶	2.49 x 10 ⁻⁰⁷	3.89 x 10 ⁻⁰⁷	3.26 x 10 ⁻⁰⁶
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	4.28 x 10 ⁻⁰⁷	1.41 x 10 ⁻¹³	3.47 x 10 ⁻⁰⁸	2.03 x 10 ⁻⁰⁸	2.73 x 10 ⁻⁰⁸	2.94 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	4.86 x 10 ⁻⁰⁷	8.98 x 10 ⁻⁰⁸	1.82 x 10 ⁻⁰⁶	2.91 x 10 ⁻⁰⁶
6 1	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.15 x 10 ⁻⁰⁶	1.43 x 10 ⁻¹⁰	9.94 x 10 ⁻⁰⁸	6.59 x 10 ⁻⁰⁵	3.74 x 10 ⁻⁰⁷	1.12 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	2.61 x 10 ⁻⁰⁶	4.94 x 10 ⁻⁰⁶	1.39 x 10 ⁻⁰⁶	7.76 x 10 ⁻⁰⁵
Subsistence	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.79 x 10 ⁻⁰⁸	3.66 x 10 ⁻¹⁵	2.33 x 10 ⁻¹⁰	9.31 x 10 ⁻¹⁰	5.49 x 10 ⁻⁰⁸	5.07 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	2.11 x 10 ⁻⁰⁷	1.86 x 10 ⁻⁰⁹	3.65 x 10 ⁻⁰⁷	6.62 x 10 ⁻⁰⁷
Harvester One- Year-Old (Lloyd	Dalanium 210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	5.82 x 10 ⁻¹⁰	2.27 x 10 ⁻¹³	1.29 x 10 ⁻⁰⁷	2.03 x 10 ⁻¹³	0.00 x 10 ⁺⁰⁰	9.69 x 10 ⁻⁰⁸	2.05 x 10 ⁻⁰⁸	2.80 x 10 ⁻⁰⁶	3.05 x 10 ⁻⁰⁶
7/42/16 B1/6 B1/6 V/6	Polonium-210			00	4.5 40-10	1.39 x 10 ⁻⁰⁷	6.65 x 10 ⁻⁰⁵	1.27 x 10 ⁻⁰⁶	1.23 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	6.45 x 10 ⁻⁰⁶	5.53 x 10 ⁻⁰⁶	7.12 x 10 ⁻⁰⁶	9.12 x 10 ⁻⁰⁵
	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.89 x 10 ⁻⁰⁶	1.45 x 10 ⁻¹⁰	1.59 X 10	0.03 X 1	1.27 X 10	1120 % 10		01 15 X 10	5155 X 10	7.12 X 10	
Lake)					1.45 x 10 ⁻¹⁰	1.59 X 10	Upper	Bound Scenario			0.15 % .0	5.55 X . C		
		$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	6.18 x 10 ⁻⁰⁷	1.45 x 10 ⁻¹⁰	1.71 x 10 ⁻⁰⁹		Bound Scenario 3.30 x 10 ⁻⁰⁷	1.12 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.46 x 10 ⁻⁰⁶	2.29 x 10 ⁻⁰⁷	3.58 x 10 ⁻⁰⁷	3.73 x 10 ⁻⁰⁶
	Total by Pathway						Upper	Bound Scenario			1.46 x 10 ⁻⁰⁶ 1.59 x 10 ⁻⁰⁶			
	Total by Pathway Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	6.18 x 10 ⁻⁰⁷	1.67 x 10 ⁻¹²	1.71 x 10 ⁻⁰⁹	Upper 6.24 x 10 ⁻⁰⁷	Bound Scenario 3.30 x 10 ⁻⁰⁷	1.12 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.46 x 10 ⁻⁰⁶	2.29 x 10 ⁻⁰⁷	3.58 x 10 ⁻⁰⁷	3.73 x 10 ⁻⁰⁶

							ncremental Dose by	Pathway - Far-Fu	iture (mSv/vr)					
	Radionuclide			Water	Water			Sediment	Sediment		Aquatic	Terrestrial	Terrestrial	Total by
		Air (Internal)	Air (External)	(Internal)	(External)	Soil (Internal)	Soil (External)	(Internal)	(External)	Aquatic Plants	Animals	Plants	Animals	Radionuclide
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.79 x 10 ⁻⁰⁸	3.69 x 10 ⁻¹⁵	2.33 x 10 ⁻¹⁰	9.31 x 10 ⁻¹⁰	5.52 x 10 ⁻⁰⁸	5.07 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	2.11 x 10 ⁻⁰⁷	1.86 x 10 ⁻⁰⁹	3.65 x 10 ⁻⁰⁷	6.63 x 10 ⁻⁰⁷
	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	5.82 x 10 ⁻¹⁰	2.27 x 10 ⁻¹³	1.29 x 10 ⁻⁰⁷	2.03 x 10 ⁻¹³	0.00 x 10 ⁺⁰⁰	1.04 x 10 ⁻⁰⁷	2.05 x 10 ⁻⁰⁸	2.80 x 10 ⁻⁰⁶	3.06 x 10 ⁻⁰⁶
	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.90 x 10 ⁻⁰⁶	1.45 x 10 ⁻¹⁰	1.39 x 10 ⁻⁰⁷	6.65 x 10 ⁻⁰⁵	1.27 x 10 ⁻⁰⁶	1.23 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	6.47 x 10 ⁻⁰⁶	5.53 x 10 ⁻⁰⁶	7.12 x 10 ⁻⁰⁶	9.12 x 10 ⁻⁰⁵
		12:00	10.00	1.70	1 2 2 10	1 2 2 1 2 00		plication Case		12:00	1.2.05	. = 1	12.00	
	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.58 x 10 ⁻⁰⁵	1.20 x 10 ⁻¹⁰	1.29 x 10 ⁻⁰⁹	1.43 x 10 ⁻⁰⁵	1.44 x 10 ⁻⁰⁷	1.54 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	1.40 x 10 ⁻⁰⁵	4.51 x 10 ⁻⁰⁶	8.43 x 10 ⁻⁰⁶	5.88 x 10 ⁻⁰⁵
	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.73 x 10 ⁻⁰⁵	2.10 x 10 ⁻¹¹	1.41 x 10 ⁻⁰⁹ 3.59 x 10 ⁻⁰⁸	7.23 x 10 ⁻⁰⁸	1.58 x 10 ⁻⁰⁷	5.60 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	1.53 x 10 ⁻⁰⁵	4.92 x 10 ⁻⁰⁶	9.21 x 10 ⁻⁰⁶	4.69 x 10 ⁻⁰⁵
	Thorium-230 Radium-226	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	9.24 x 10 ⁻⁰⁶ 1.88 x 10 ⁻⁰⁵	6.26 x 10 ⁻¹² 8.40 x 10 ⁻⁰⁹	5.85 x 10 ⁻⁰⁸	5.50 x 10 ⁻⁰⁷ 1.52 x 10 ⁻⁰³	1.01 x 10 ⁻⁰⁸ 1.04 x 10 ⁻⁰⁷	2.49 x 10 ⁻¹⁰ 1.25 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	3.92 x 10 ⁻⁰⁶ 1.59 x 10 ⁻⁰⁵	2.41 x 10 ⁻⁰⁶ 7.57 x 10 ⁻⁰⁵	4.42 x 10 ⁻⁰⁵ 7.26 x 10 ⁻⁰⁵	6.04 x 10 ⁻⁰⁵ 1.71 x 10 ⁻⁰³
	Lead-210	0.00 x 10 40 0.00 x 10+00	0.00 x 10 40 0.00 x 10 400	1.21 x 10 ⁻⁰⁶	1.07 x 10 ⁻¹²	9.46 x 10 ⁻¹¹	2.19 x 10 ⁻⁰⁸	4.08 x 10 ⁻⁰⁸	2.31 x 10 ⁻⁰⁹	0.00 x 10 40 0.00 x 10+00	3.43 x 10 ⁻⁰⁶	1.61 x 10 ⁻⁰⁸	9.27 x 10 ⁻⁰⁵	9.74 x 10 ⁻⁰⁵
Seasonal	Polonium-210	0.00 x 10	0.00 x 10 ⁺⁰⁰	2.13 x 10 ⁻⁰⁶	6.40 x 10 ⁻¹⁵	1.67 x 10 ⁻¹⁰	4.95 x 10 ⁻¹²	7.22 x 10 ⁻⁰⁸	9.91 x 10 ⁻¹²	0.00 x 10	2.15 x 10 ⁻⁰⁵	1.39 x 10 ⁻⁰⁷	8.72 x 10 ⁻⁰⁵	1.11 x 10 ⁻⁰⁴
Resident	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	6.45 x 10 ⁻⁰⁵	8.55 x 10 ⁻⁰⁹	9.74 x 10 ⁻⁰⁸	1.53 x 10 ⁻⁰³	5.29 x 10 ⁻⁰⁷	1.40 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	7.40 x 10 ⁻⁰⁵	8.77 x 10 ⁻⁰⁵	3.14 x 10 ⁻⁰⁴	2.09 x 10 ⁻⁰³
(Patterson Lake		0.00 % 10	0.00 % 10	0.15 x 10	0.00 X 10	311 1 1 1 1	•	Bound Scenario	11.0 % 10	0.00 % 10	7110 % 10	GILL X 10	5111710	Zies X ie
South Arm)	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.58 x 10 ⁻⁰⁵	1.20 x 10 ⁻¹⁰	1.29 x 10 ⁻⁰⁹	1.43 x 10 ⁻⁰⁵	1.44 x 10 ⁻⁰⁷	1.54 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	1.40 x 10 ⁻⁰⁵	4.51 x 10 ⁻⁰⁶	8.43 x 10 ⁻⁰⁶	5.88 x 10 ⁻⁰⁵
	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.73 x 10 ⁻⁰⁵	2.10 x 10 ⁻¹¹	1.41 x 10 ⁻⁰⁹	7.23 x 10 ⁻⁰⁸	1.58 x 10 ⁻⁰⁷	5.61 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	1.53 x 10 ⁻⁰⁵	4.92 x 10 ⁻⁰⁶	9.21 x 10 ⁻⁰⁶	4.70 x 10 ⁻⁰⁵
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	9.24 x 10 ⁻⁰⁶	6.26 x 10 ⁻¹²	3.59 x 10 ⁻⁰⁸	5.50 x 10 ⁻⁰⁷	1.01 x 10 ⁻⁰⁸	2.49 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	3.92 x 10 ⁻⁰⁶	2.41 x 10 ⁻⁰⁶	4.42 x 10 ⁻⁰⁵	6.04 x 10 ⁻⁰⁵
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.89 x 10 ⁻⁰⁵	8.42 x 10 ⁻⁰⁹	5.85 x 10 ⁻⁰⁸	1.52 x 10 ⁻⁰³	1.04 x 10 ⁻⁰⁷	1.25 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	1.60 x 10 ⁻⁰⁵	7.57 x 10 ⁻⁰⁵	7.26 x 10 ⁻⁰⁵	1.71 x 10 ⁻⁰³
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.22 x 10 ⁻⁰⁶	1.07 x 10 ⁻¹²	9.46 x 10 ⁻¹¹	2.19 x 10 ⁻⁰⁸	4.10 x 10 ⁻⁰⁸	2.32 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	3.44 x 10 ⁻⁰⁶	1.61 x 10 ⁻⁰⁸	9.31 x 10 ⁻⁰⁵	9.78 x 10 ⁻⁰⁵
	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.14 x 10 ⁻⁰⁶	6.43 x 10 ⁻¹⁵	1.67 x 10 ⁻¹⁰	4.95 x 10 ⁻¹²	7.25 x 10 ⁻⁰⁸	9.95 x 10 ⁻¹²	0.00 x 10 ⁺⁰⁰	2.16 x 10 ⁻⁰⁵	1.39 x 10 ⁻⁰⁷	8.76 x 10 ⁻⁰⁵	1.11 x 10 ⁻⁰⁴
	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	6.46 x 10 ⁻⁰⁵	8.57 x 10 ⁻⁰⁹	9.74 x 10 ⁻⁰⁸	1.53 x 10 ⁻⁰³	5.31 x 10 ⁻⁰⁷	1.41 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	7.42 x 10 ⁻⁰⁵	8.77 x 10 ⁻⁰⁵	3.15 x 10 ⁻⁰⁴	2.09 x 10 ⁻⁰³
		12:00	10.00	10.05				plication Case		12:00	1.00	12.00	1000	
	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.10 x 10 ⁻⁰⁵	2.97 x 10 ⁻¹¹	5.26 x 10 ⁻⁰⁸	1.92 x 10 ⁻⁰⁵	5.87 x 10 ⁻⁰⁶	2.00 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	8.24 x 10 ⁻⁰⁶	6.14 x 10 ⁻⁰⁶	3.84 x 10 ⁻⁰⁶	5.63 x 10 ⁻⁰⁵
	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.20 x 10 ⁻⁰⁵	5.20 x 10 ⁻¹²	5.70 x 10 ⁻⁰⁸	9.68 x 10 ⁻⁰⁸	6.39 x 10 ⁻⁰⁶	7.29 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	8.96 x 10 ⁻⁰⁶	6.65 x 10 ⁻⁰⁶	4.17 x 10 ⁻⁰⁶	3.83 x 10 ⁻⁰⁵
	Thorium-230	0.00 x 10 ⁺⁰⁰ 0.00 x 10 ⁺⁰⁰	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	4.70 x 10 ⁻⁰⁶ 1.68 x 10 ⁻⁰⁵	1.55 x 10 ⁻¹²	1.07 x 10 ⁻⁰⁶	6.25 x 10 ⁻⁰⁷ 2.03 x 10 ⁻⁰³	3.00 x 10 ⁻⁰⁷ 5.45 x 10 ⁻⁰⁶	3.24 x 10 ⁻¹⁰	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	1.69 x 10 ⁻⁰⁶ 1.21 x 10 ⁻⁰⁵	2.40 x 10 ⁻⁰⁶ 1.32 x 10 ⁻⁰⁴	2.16 x 10 ⁻⁰⁵ 3.41 x 10 ⁻⁰⁵	3.24 x 10 ⁻⁰⁵ 2.25 x 10 ⁻⁰³
	Radium-226 Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.65 x 10 ⁻⁰⁶	2.08 x 10 ⁻⁰⁹ 2.17 x 10 ⁻¹³	3.06 x 10 ⁻⁰⁶ 7.45 x 10 ⁻⁰⁹	2.93 x 10 ⁻⁰⁸	3.25 x 10 ⁻⁰⁶	1.63 x 10 ⁻⁰⁵ 3.00 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	3.95 x 10 ⁻⁰⁶	4.33 x 10 ⁻⁰⁸	1.18 x 10 ⁻⁰⁴	1.27 x 10 ⁻⁰⁴
Seasonal	Polonium-210	0.00 x 10	0.00 x 10 ⁺⁰⁰	4.07 x 10 ⁻⁰⁶	1.59 x 10 ⁻¹⁵	1.86 x 10 ⁻⁰⁸	6.48 x 10 ⁻¹²	8.08 x 10 ⁻⁰⁶	1.29 x 10 ⁻¹¹	0.00 x 10	3.48 x 10 ⁻⁰⁵	5.20 x 10 ⁻⁰⁷	1.02 x 10 ⁻⁰⁴	1.49 x 10 ⁻⁰⁴
Resident One-	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	5.02 x 10 ⁻⁰⁵	2.12 x 10 ⁻⁰⁹	4.26 x 10 ⁻⁰⁶	2.05 x 10 ⁻⁰³	2.93 x 10 ⁻⁰⁵	1.83 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	6.98 x 10 ⁻⁰⁵	1.48 x 10 ⁻⁰⁴	2.83 x 10 ⁻⁰⁴	2.65 x 10 ⁻⁰³
Year-Old	Total by Fathway	0.00 X 10	0.00 X 10	3.02 x 10	2.12 X 10	1.20 X 10		Bound Scenario	1.03 X 10	0.00 X 10	0.50 X 10	1.10 X 10	2.03 X 10	2.03 X 10
(Patterson Lake	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.10 x 10 ⁻⁰⁵	2.97 x 10 ⁻¹¹	5.26 x 10 ⁻⁰⁸	1.92 x 10 ⁻⁰⁵	5.88 x 10 ⁻⁰⁶	2.00 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	8.25 x 10 ⁻⁰⁶	6.14 x 10 ⁻⁰⁶	3.84 x 10 ⁻⁰⁶	5.64 x 10 ⁻⁰⁵
South Arm)	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.20 x 10 ⁻⁰⁵	5.21 x 10 ⁻¹²	5.70 x 10 ⁻⁰⁸	9.68 x 10 ⁻⁰⁸	6.39 x 10 ⁻⁰⁶	7.29 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	8.97 x 10 ⁻⁰⁶	6.65 x 10 ⁻⁰⁶	4.17 x 10 ⁻⁰⁶	3.83 x 10 ⁻⁰⁵
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	4.70 x 10 ⁻⁰⁶	1.55 x 10 ⁻¹²	1.07 x 10 ⁻⁰⁶	6.25 x 10 ⁻⁰⁷	3.00 x 10 ⁻⁰⁷	3.24 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	1.69 x 10 ⁻⁰⁶	2.40 x 10 ⁻⁰⁶	2.16 x 10 ⁻⁰⁵	3.24 x 10 ⁻⁰⁵
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.69 x 10 ⁻⁰⁵	2.08 x 10 ⁻⁰⁹	3.06 x 10 ⁻⁰⁶	2.03 x 10 ⁻⁰³	5.46 x 10 ⁻⁰⁶	1.63 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	1.21 x 10 ⁻⁰⁵	1.32 x 10 ⁻⁰⁴	3.41 x 10 ⁻⁰⁵	2.25 x 10 ⁻⁰³
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.66 x 10 ⁻⁰⁶	2.18 x 10 ⁻¹³	7.45 x 10 ⁻⁰⁹	2.93 x 10 ⁻⁰⁸	3.26 x 10 ⁻⁰⁶	3.02 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	3.97 x 10 ⁻⁰⁶	4.33 x 10 ⁻⁰⁸	1.18 x 10 ⁻⁰⁴	1.27 x 10 ⁻⁰⁴
	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	4.09 x 10 ⁻⁰⁶	1.59 x 10 ⁻¹⁵	1.86 x 10 ⁻⁰⁸	6.48 x 10 ⁻¹²	8.11 x 10 ⁻⁰⁶	1.29 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	3.50 x 10 ⁻⁰⁵	5.20 x 10 ⁻⁰⁷	1.02 x 10 ⁻⁰⁴	1.50 x 10 ⁻⁰⁴
	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	5.03 x 10 ⁻⁰⁵	2.12 x 10 ⁻⁰⁹	4.26 x 10 ⁻⁰⁶	2.05 x 10 ⁻⁰³	2.94 x 10 ⁻⁰⁵	1.83 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	7.00 x 10 ⁻⁰⁵	1.48 x 10 ⁻⁰⁴	2.84 x 10 ⁻⁰⁴	2.65 x 10 ⁻⁰³
		0.00 10+00	0.00 10+00	F 22 10-07	4.04 4.0-12	2.52 40-11		plication Case	F 10 10 10 108	0.00 10+00	2.27 10-06	0.00 10-08	2.00 10-07	2.42. 40-06
	Uranium-238 Uranium-234	0.00 x 10 ⁺⁰⁰ 0.00 x 10 ⁺⁰⁰	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	5.33 x 10 ⁻⁰⁷ 5.82 x 10 ⁻⁰⁷	4.04 x 10 ⁻¹² 7.07 x 10 ⁻¹³	2.52 x 10 ⁻¹¹ 2.74 x 10 ⁻¹¹	2.80 x 10 ⁻⁰⁷ 1.41 x 10 ⁻⁰⁹	4.86 x 10 ⁻⁰⁹ 5.31 x 10 ⁻⁰⁹	5.18 x 10 ⁻⁰⁸ 1.89 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰ 0.00 x 10 ⁺⁰⁰	2.27 x 10 ⁻⁰⁶ 2.47 x 10 ⁻⁰⁶	8.80 x 10 ⁻⁰⁸ 9.58 x 10 ⁻⁰⁸	2.08 x 10 ⁻⁰⁷ 2.28 x 10 ⁻⁰⁷	3.43 x 10 ⁻⁰⁶ 3.39 x 10 ⁻⁰⁶
	Thorium-230	0.00 x 10 40 0.00 x 10+00	0.00 x 10 ⁺⁰⁰	5.04 x 10 ⁻⁰⁷	3.42 x 10 ⁻¹³	6.99 x 10 ⁻¹⁰	1.41 x 10 ⁻⁰⁸	5.50 x 10 ⁻¹⁰	1.36 x 10 ⁻¹¹	0.00 x 10 40 0.00 x 10+00	4.47 x 10 ⁻⁰⁷	4.70 x 10 ⁻⁰⁸	8.83 x 10 ⁻⁰⁷	1.89 x 10 ⁻⁰⁶
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	7.74 x 10 ⁻⁰⁷	3.46 x 10 ⁻¹⁰	1.14 x 10 ⁻⁰⁹	2.95 x 10 ⁻⁰⁵	4.29 x 10 ⁻⁰⁹	5.14 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.77 x 10 ⁻⁰⁶	1.47 x 10 ⁻⁰⁶	6.70 x 10 ⁻⁰⁷	3.47 x 10 ⁻⁰⁵
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.22 x 10 ⁻⁰⁸	1.09 x 10 ⁻¹⁴	9.09 x 10 ⁻¹³	2.33 x 10 ⁻¹⁰	4.15 x 10 ⁻¹⁰	2.32 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	7.56 x 10 ⁻⁰⁴	0.00 x 10 ⁺⁰⁰	7.08 x 10 ⁻⁰⁸	7.56 x 10 ⁻⁰⁴
	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	3.18 x 10 ⁻¹²	1.14 x 10 ⁻¹³	6.84 x 10 ⁻¹⁰	9.33 x 10 ⁻¹⁴	0.00 x 10 ⁺⁰⁰	5.10 x 10 ⁻⁰⁴	3.26 x 10 ⁻⁰⁹	6.56 x 10 ⁻⁰⁷	5.10 x 10 ⁻⁰⁴
Seasonal	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.41 x 10 ⁻⁰⁶	3.51 x 10 ⁻¹⁰	1.90 x 10 ⁻⁰⁹	2.98 x 10 ⁻⁰⁵	1.61 x 10 ⁻⁰⁸	5.66 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.27 x 10 ⁻⁰³	1.71 x 10 ⁻⁰⁶	2.72 x 10 ⁻⁰⁶	1.31 x 10 ⁻⁰³
Resident (Lloyd							Uppe	Bound Scenario						
Lake)	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	5.33 x 10 ⁻⁰⁷	4.05 x 10 ⁻¹²	2.52 x 10 ⁻¹¹	2.80 x 10 ⁻⁰⁷	4.86 x 10 ⁻⁰⁹	5.19 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.27 x 10 ⁻⁰⁶	8.80 x 10 ⁻⁰⁸	2.08 x 10 ⁻⁰⁷	3.43 x 10 ⁻⁰⁶
	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	5.83 x 10 ⁻⁰⁷	7.08 x 10 ⁻¹³	2.74 x 10 ⁻¹¹	1.41 x 10 ⁻⁰⁹	5.32 x 10 ⁻⁰⁹	1.89 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	2.47 x 10 ⁻⁰⁶	9.58 x 10 ⁻⁰⁸	2.28 x 10 ⁻⁰⁷	3.39 x 10 ⁻⁰⁶
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	5.04 x 10 ⁻⁰⁷	3.42 x 10 ⁻¹³	6.99 x 10 ⁻¹⁰	1.07 x 10 ⁻⁰⁸	5.50 x 10 ⁻¹⁰	1.36 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	4.47 x 10 ⁻⁰⁷	4.70 x 10 ⁻⁰⁸	8.83 x 10 ⁻⁰⁷	1.89 x 10 ⁻⁰⁶
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	7.76 x 10 ⁻⁰⁷	3.46 x 10 ⁻¹⁰	1.14 x 10 ⁻⁰⁹	2.95 x 10 ⁻⁰⁵	4.30 x 10 ⁻⁰⁹	5.15 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.77 x 10 ⁻⁰⁶	1.47 x 10 ⁻⁰⁶	6.70 x 10 ⁻⁰⁷	3.48 x 10 ⁻⁰⁵
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.22 x 10 ⁻⁰⁸	1.09 x 10 ⁻¹⁴	9.09 x 10 ⁻¹³	2.33 x 10 ⁻¹⁰	4.15 x 10 ⁻¹⁰	2.34 x 10 ⁻¹¹	$0.00 \times 10^{+00}$	7.56 x 10 ⁻⁰⁴	0.00 x 10 ⁺⁰⁰	7.08 x 10 ⁻⁰⁸	7.56 x 10 ⁻⁰⁴
	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	3.18 x 10 ⁻¹²	1.14 x 10 ⁻¹³	6.84 x 10 ⁻¹⁰	9.41 x 10 ⁻¹⁴	0.00 x 10 ⁺⁰⁰	5.10 x 10 ⁻⁰⁴	3.26 x 10 ⁻⁰⁹	6.56 x 10 ⁻⁰⁷	5.10 x 10 ⁻⁰⁴
Carran	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.41 x 10 ⁻⁰⁶	3.52 x 10 ⁻¹⁰	1.90 x 10 ⁻⁰⁹	2.98 x 10 ⁻⁰⁵	1.61 x 10 ⁻⁰⁸	5.67 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.27 x 10 ⁻⁰³	1.71 x 10 ⁻⁰⁶	2.72 x 10 ⁻⁰⁶	1.31 x 10 ⁻⁰³
Seasonal Resident One-	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	3.70 x 10 ⁻⁰⁷	9.99 x 10 ⁻¹³	1.03 x 10 ⁻⁰⁹	3.74 x 10 ⁻⁰⁷	plication Case 1.98 x 10 ⁻⁰⁷	6.74 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.77 x 10 ⁻⁰⁷	1.20 x 10 ⁻⁰⁷	9.31 x 10 ⁻⁰⁸	1.50 x 10 ⁻⁰⁶
Year-Old (Lloyd	Uranium-234	0.00 x 10 40 0.00 x 10+00	0.00 x 10 ⁺⁰⁰	4.03 x 10 ⁻⁰⁷	1.75 x 10 ⁻¹³	1.11 x 10 ⁻⁰⁹	1.89 x 10 ⁻⁰⁹	2.15 x 10 ⁻⁰⁷	2.45 x 10 ⁻¹⁰	0.00 x 10 40 0.00 x 10+00	3.02 x 10 ⁻⁰⁷	1.30 x 10 ⁻⁰⁷	1.01 x 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁶
Lake)	Thorium-230	0.00 x 10	0.00 x 10 ⁺⁰⁰	2.57 x 10 ⁻⁰⁷	8.46 x 10 ⁻¹⁴	2.08 x 10 ⁻⁰⁸	1.22 x 10 ⁻⁰⁸	1.64 x 10 ⁻⁰⁸	1.77 x 10 ⁻¹¹	0.00 x 10 0.00 x 10 ⁺⁰⁰	9.24 x 10 ⁻⁰⁸	4.68 x 10 ⁻⁰⁸	4.29 x 10 ⁻⁰⁷	8.74 x 10 ⁻⁰⁷
	Thoriam 250	0.00 X 10	0.00 X 10	L.37 X 10	0. 10 X 10	L.00 X 10	1.22 / 10	1.017.10	1.77 / 10	0.00 X 10	J.L 1 X 10	1.00 A 10	1.LJ X 10	0.7 T X TO

						İr	ncremental Dose by	Pathway - Far-Fu	ıture (mSv/yr)					
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by Radionuclide
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	6.92 x 10 ⁻⁰⁷	8.56 x 10 ⁻¹¹	5.96 x 10 ⁻⁰⁸	3.95 x 10 ⁻⁰⁵	2.24 x 10 ⁻⁰⁷	6.70 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	4.97 x 10 ⁻⁰⁷	2.58 x 10 ⁻⁰⁶	2.70 x 10 ⁻⁰⁷	4.45 x 10 ⁻⁰⁵
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.66 x 10 ⁻⁰⁸	2.22 x 10 ⁻¹⁵	1.16 x 10 ⁻¹⁰	4.66 x 10 ⁻¹⁰	3.28 x 10 ⁻⁰⁸	3.05 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	4.00 x 10 ⁻⁰⁸	9.31 x 10 ⁻¹⁰	8.94 x 10 ⁻⁰⁸	1.80 x 10 ⁻⁰⁷
	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	3.49 x 10 ⁻¹⁰	1.42 x 10 ⁻¹³	7.73 x 10 ⁻⁰⁸	1.22 x 10 ⁻¹³	0.00 x 10 ⁺⁰⁰	2.24 x 10 ⁻⁰⁸	1.12 x 10 ⁻⁰⁸	8.05 x 10 ⁻⁰⁷	9.15 x 10 ⁻⁰⁷
	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.74 x 10 ⁻⁰⁶	8.68 x 10 ⁻¹¹	8.31 x 10 ⁻⁰⁸	3.99 x 10 ⁻⁰⁵	7.63 x 10 ⁻⁰⁷	7.38 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.23 x 10 ⁻⁰⁶	2.89 x 10 ⁻⁰⁶	1.79 x 10 ⁻⁰⁶	4.91 x 10 ⁻⁰⁵
		2.22 12.100	0.00 10.00	2.71 10.07	1.00 10.13	1.00 10.00		Bound Scenario	674 40.00	0.00 10.00	0.70 10.07	1 0 0 10 07	0.24 10.09	1.50 10.06
	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	3.71 x 10 ⁻⁰⁷	1.00 x 10 ⁻¹²	1.03 x 10 ⁻⁰⁹	3.74 x 10 ⁻⁰⁷	1.98 x 10 ⁻⁰⁷	6.74 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.78 x 10 ⁻⁰⁷	1.20 x 10 ⁻⁰⁷	9.31 x 10 ⁻⁰⁸	1.50 x 10 ⁻⁰⁶
	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	4.03 x 10 ⁻⁰⁷	1.75 x 10 ⁻¹³	1.11 x 10 ⁻⁰⁹	1.89 x 10 ⁻⁰⁹	2.15 x 10 ⁻⁰⁷	2.45 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	3.02 x 10 ⁻⁰⁷	1.30 x 10 ⁻⁰⁷	1.01 x 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁶
	Thorium-230	0.00 x 10 ⁺⁰⁰ 0.00 x 10 ⁺⁰⁰	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	2.57 x 10 ⁻⁰⁷ 6.94 x 10 ⁻⁰⁷	8.46 x 10 ⁻¹⁴ 8.58 x 10 ⁻¹¹	2.08 x 10 ⁻⁰⁸ 5.96 x 10 ⁻⁰⁸	1.22 x 10 ⁻⁰⁸ 3.95 x 10 ⁻⁰⁵	1.64 x 10 ⁻⁰⁸ 2.25 x 10 ⁻⁰⁷	1.77 x 10 ⁻¹¹ 6.72 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	9.24 x 10 ⁻⁰⁸ 4.98 x 10 ⁻⁰⁷	4.68 x 10 ⁻⁰⁸ 2.58 x 10 ⁻⁰⁶	4.29 x 10 ⁻⁰⁷ 2.70 x 10 ⁻⁰⁷	8.74 x 10 ⁻⁰⁷ 4.45 x 10 ⁻⁰⁵
	Radium-226 Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.69 x 10 ⁻⁰⁸	2.22 x 10 ⁻¹⁵	1.16 x 10 ⁻¹⁰	4.66 x 10 ⁻¹⁰	3.31 x 10 ⁻⁰⁸	3.05 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	4.00 x 10 ⁻⁰⁸	9.31 x 10 ⁻¹⁰	8.94 x 10 ⁻⁰⁸	1.81 x 10 ⁻⁰⁷
	Polonium-210	0.00 x 10 40 0.00 x 10+00	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	3.49 x 10 ⁻¹⁰	1.42 x 10 ⁻¹³	7.73 x 10 ⁻⁰⁸	1.22 x 10 ⁻¹³	0.00 x 10 ⁺⁰⁰	2.24 x 10 ⁻⁰⁸	1.12 x 10 ⁻⁰⁸	8.34 x 10 ⁻⁰⁷	9.46 x 10 ⁻⁰⁷
	Total by Pathway	0.00 x 10 0.00 x 10+00	0.00 x 10 ⁺⁰⁰	1.74 x 10 ⁻⁰⁶	8.71 x 10 ⁻¹¹	8.31 x 10 ⁻⁰⁸	3.99 x 10 ⁻⁰⁵	7.64 x 10 ⁻⁰⁷	7.40 x 10 ⁻⁰⁷	0.00 x 10	1.23 x 10 ⁻⁰⁶	2.89 x 10 ⁻⁰⁶	1.82 x 10 ⁻⁰⁶	4.92 x 10 ⁻⁰⁵
	Total by Facilita	0.00 X 10	0.00 X 10	1.7 TX 10	0.7 T X TO	0.5 T X TO		plication Case	7.10 X 10	0.00 X 10	1.23 X 10	2.03 X 10	1.02 X 10	1.52 X 10
	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.91 x 10 ⁻⁰⁴	1.45 x 10 ⁻⁰⁹	2.64 × 10 ⁻⁰⁸	2.93 × 10 ⁻⁰⁴	1.74 × 10 ⁻⁰⁶	1.85 × 10- ⁰⁵	0.00 x 10 ⁺⁰⁰	5.31 x 10 ⁻⁰⁴	1.18 × 10 ⁻⁰⁴	2.38 × 10 ⁻⁰⁴	1.39 × 10 ⁻⁰³
	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.08 x 10 ⁻⁰⁴	2.53 x 10 ⁻¹⁰	2.88 x 10 ⁻⁰⁸	1.48 x 10 ⁻⁰⁶	1.90 x 10 ⁻⁰⁶	6.75 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	5.81 x 10 ⁻⁰⁴	1.28 x 10 ⁻⁰⁴	2.60 x 10 ⁻⁰⁴	1.18 x 10 ⁻⁰³
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	5.00 x 10 ⁻⁰⁵	3.39 x 10 ⁻¹¹	7.33 x 10 ⁻⁰⁷	1.12 x 10 ⁻⁰⁵	5.48 x 10 ⁻⁰⁸	1.35 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	6.70 x 10 ⁻⁰⁵	6.29 x 10 ⁻⁰⁵	5.31 x 10 ⁻⁰⁴	7.23 x 10 ⁻⁰⁴
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.70 x 10 ⁻⁰⁴	7.61 x 10 ⁻⁰⁸	1.20 x 10 ⁻⁰⁶	3.10 x 10 ⁻⁰²	9.43 x 10 ⁻⁰⁷	1.13 x 10 ⁻⁰⁴	0.00 x 10 ⁺⁰⁰	4.55 x 10 ⁻⁰⁴	1.97 x 10 ⁻⁰³	1.87 x 10 ⁻⁰³	3.55 x 10 ⁻⁰²
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	6.15 x 10 ⁻⁰⁵	5.42 x 10 ⁻¹¹	1.93 x 10 ⁻⁰⁹	4.47 x 10 ⁻⁰⁷	2.07 x 10 ⁻⁰⁶	1.17 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	5.48 x 10 ⁻⁰⁴	4.23 x 10 ⁻⁰⁷	9.16 x 10 ⁻⁰⁴	1.53 x 10 ⁻⁰³
Permanent	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.08 x 10 ⁻⁰⁴	3.24 x 10 ⁻¹³	3.41 x 10 ⁻⁰⁹	1.01 x 10 ⁻¹⁰	3.66 x 10 ⁻⁰⁶	5.02 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	3.42 x 10 ⁻⁰³	3.61 x 10 ⁻⁰⁶	7.21 x 10 ⁻⁰³	1.07 x 10 ⁻⁰²
Resident (Patterson Lake	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	7.89 x 10 ⁻⁰⁴	7.79 x 10 ⁻⁰⁸	1.99 x 10 ⁻⁰⁶	3.13 x 10 ⁻⁰²	1.04 x 10 ⁻⁰⁵	1.32 x 10 ⁻⁰⁴	0.00 x 10 ⁺⁰⁰	5.61 x 10 ⁻⁰³	2.28 x 10 ⁻⁰³	1.10 x 10 ⁻⁰²	5.11 x 10 ⁻⁰²
North Arm -						1		Bound Scenario						02
West Basin)	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.91 x 10 ⁻⁰⁴	1.45 x 10 ⁻⁰⁹	2.64 x 10 ⁻⁰⁸	2.93 x 10 ⁻⁰⁴	1.74 x 10 ⁻⁰⁶	1.86 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	5.32 x 10 ⁻⁰⁴	1.18 x 10 ⁻⁰⁴	2.38 x 10 ⁻⁰⁴	1.39 x 10 ⁻⁰³
	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.09 x 10 ⁻⁰⁴	2.53 x 10 ⁻¹⁰	2.88 x 10 ⁻⁰⁸	1.48 x 10 ⁻⁰⁶	1.90 x 10 ⁻⁰⁶	6.76 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	5.81 x 10 ⁻⁰⁴	1.28 x 10 ⁻⁰⁴	2.60 x 10 ⁻⁰⁴	1.18 x 10 ⁻⁰³
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	5.00 x 10 ⁻⁰⁵	3.39 x 10 ⁻¹¹	7.33 x 10 ⁻⁰⁷	1.12 x 10 ⁻⁰⁵	5.48 x 10 ⁻⁰⁸	1.35 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	6.70 x 10 ⁻⁰⁵	6.29 x 10 ⁻⁰⁵	5.31 x 10 ⁻⁰⁴	7.23 x 10 ⁻⁰⁴
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.71 x 10 ⁻⁰⁴	7.63 x 10 ⁻⁰⁸	1.20 x 10 ⁻⁰⁶	3.10 x 10 ⁻⁰² 4.47 x 10 ⁻⁰⁷	9.46 x 10 ⁻⁰⁷	1.13 x 10 ⁻⁰⁴	0.00 x 10 ⁺⁰⁰	4.56 x 10 ⁻⁰⁴	1.97 x 10 ⁻⁰³ 4.23 x 10 ⁻⁰⁷	1.87 x 10 ⁻⁰³	3.56 x 10 ⁻⁰²
	Lead-210 Polonium-210	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	6.18 x 10 ⁻⁰⁵ 1.08 x 10 ⁻⁰⁴	5.44 x 10 ⁻¹¹ 3.25 x 10 ⁻¹³	1.93 x 10 ⁻⁰⁹ 3.41 x 10 ⁻⁰⁹	1.01 x 10 ⁻¹⁰	2.08 x 10 ⁻⁰⁶ 3.67 x 10 ⁻⁰⁶	1.17 x 10 ⁻⁰⁷ 5.04 x 10 ⁻¹⁰	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	5.50 x 10 ⁻⁰⁴ 3.44 x 10 ⁻⁰³	3.61 x 10 ⁻⁰⁶	9.20 x 10 ⁻⁰⁴ 7.24 x 10 ⁻⁰³	1.53 x 10 ⁻⁰³ 1.08 x 10 ⁻⁰²
	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	7.90 x 10 ⁻⁰⁴	7.81 x 10 ⁻⁰⁸	1.99 x 10 ⁻⁰⁶	3.13 x 10 ⁻⁰²	1.04 x 10 ⁻⁰⁵	1.32 x 10 ⁻⁰⁴	0.00 x 10 ⁺⁰⁰	5.63 x 10 ⁻⁰³	2.28 x 10 ⁻⁰³	1.11 x 10 ⁻⁰²	5.12 x 10 ⁻⁰²
	Total by Fathway	0.00 X 10	0.00 X 10	7.90 X 10	7.01 X 10	1.99 X 10		plication Case	1.52 X 10	0.00 X 10	3.03 X 10	2.20 X 10	1.11 x 10	3.12 X 10
	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.33 x 10 ⁻⁰⁴	3.58 x 10 ⁻¹⁰	1.07 x 10 ⁻⁰⁶	3.92 x 10 ⁻⁰⁴	7.07 x 10 ⁻⁰⁵	2.41 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	3.13 x 10 ⁻⁰⁴	1.44 x 10 ⁻⁰⁴	1.08 x 10 ⁻⁰⁴	1.19 x 10 ⁻⁰³
	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.44 x 10 ⁻⁰⁴	6.27 x 10 ⁻¹¹	1.16 x 10 ⁻⁰⁶	1.98 x 10 ⁻⁰⁶	7.69 x 10 ⁻⁰⁵	8.78 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	3.41 x 10 ⁻⁰⁴	1.56 x 10 ⁻⁰⁴	1.17 x 10 ⁻⁰⁴	8.38 x 10 ⁻⁰⁴
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.54 x 10 ⁻⁰⁵	8.39 x 10 ⁻¹²	2.18 x 10 ⁻⁰⁵	1.28 x 10 ⁻⁰⁵	1.63 x 10 ⁻⁰⁶	1.76 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	2.89 x 10 ⁻⁰⁵	5.65 x 10 ⁻⁰⁵	2.54 x 10 ⁻⁰⁴	4.01 x 10 ⁻⁰⁴
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.52 x 10 ⁻⁰⁴	1.88 x 10 ⁻⁰⁸	6.25 x 10 ⁻⁰⁵	4.14 x 10 ⁻⁰²	4.93 x 10 ⁻⁰⁵	1.47 x 10 ⁻⁰⁴	0.00 x 10 ⁺⁰⁰	3.45 x 10 ⁻⁰⁴	3.11 x 10 ⁻⁰³	7.21 x 10 ⁻⁰⁴	4.60 x 10 ⁻⁰²
Permanent	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	8.36 x 10 ⁻⁰⁵	1.10 x 10 ⁻¹¹	1.53 x 10 ⁻⁰⁷	5.98 x 10 ⁻⁰⁷	1.65 x 10 ⁻⁰⁴	1.52 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	6.32 x 10 ⁻⁰⁴	1.02 x 10 ⁻⁰⁶	1.09 x 10 ⁻⁰³	1.97 x 10 ⁻⁰³
Resident One-	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.06 x 10 ⁻⁰⁴	8.02 x 10 ⁻¹⁴	3.81 x 10 ⁻⁰⁷	1.32 x 10 ⁻¹⁰	4.09 x 10 ⁻⁰⁴	6.52 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	5.56 x 10 ⁻⁰³	1.22 x 10 ⁻⁰⁵	9.16 x 10 ⁻⁰³	1.53 x 10 ⁻⁰²
Year-Old	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	7.44 x 10 ⁻⁰⁴	1.93 x 10 ⁻⁰⁸	8.72 x 10 ⁻⁰⁵	4.19 x 10 ⁻⁰²	7.72 x 10 ⁻⁰⁴	1.72 x 10 ⁻⁰⁴	0.00 x 10 ⁺⁰⁰	7.22 x 10 ⁻⁰³	3.48 x 10 ⁻⁰³	1.14 x 10 ⁻⁰²	6.58 x 10 ⁻⁰²
(Patterson Lake								Bound Scenario						
North Arm -	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.33 x 10 ⁻⁰⁴	3.58 x 10 ⁻¹⁰	1.07 x 10 ⁻⁰⁶	3.92 x 10 ⁻⁰⁴	7.08 x 10 ⁻⁰⁵	2.41 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	3.14 x 10 ⁻⁰⁴	1.44 x 10 ⁻⁰⁴	1.08 x 10 ⁻⁰⁴	1.19 x 10 ⁻⁰³
West Basin)	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.44 x 10 ⁻⁰⁴	6.27 x 10 ⁻¹¹	1.16 x 10 ⁻⁰⁶	1.98 x 10 ⁻⁰⁶	7.70 x 10 ⁻⁰⁵	8.79 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	3.41 x 10 ⁻⁰⁴	1.56 x 10 ⁻⁰⁴	1.17 x 10 ⁻⁰⁴	8.39 x 10 ⁻⁰⁴
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.54 x 10 ⁻⁰⁵	8.39 x 10 ⁻¹²	2.18 x 10 ⁻⁰⁵	1.28 x 10 ⁻⁰⁵	1.63 x 10 ⁻⁰⁶	1.76 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	2.89 x 10 ⁻⁰⁵	5.65 x 10 ⁻⁰⁵	2.54 x 10 ⁻⁰⁴	4.01 x 10 ⁻⁰⁴
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.52 x 10 ⁻⁰⁴	1.89 x 10 ⁻⁰⁸	6.25 x 10 ⁻⁰⁵	4.14 x 10 ⁻⁰²	4.93 x 10 ⁻⁰⁵	1.48 x 10 ⁻⁰⁴	0.00 x 10 ⁺⁰⁰	3.46 x 10 ⁻⁰⁴	3.11 x 10 ⁻⁰³	7.21 x 10 ⁻⁰⁴	4.60 x 10 ⁻⁰²
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	8.40 x 10 ⁻⁰⁵	1.10 x 10 ⁻¹¹	1.53 x 10 ⁻⁰⁷	5.98 x 10 ⁻⁰⁷	1.65 x 10 ⁻⁰⁴	1.53 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	6.35 x 10 ⁻⁰⁴	1.02 x 10 ⁻⁰⁶	1.09 x 10 ⁻⁰³	1.98 x 10 ⁻⁰³
	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.07 x 10 ⁻⁰⁴	8.05 x 10 ⁻¹⁴	3.81 x 10 ⁻⁰⁷	1.32 x 10 ⁻¹⁰	4.11 x 10 ⁻⁰⁴	6.55 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	5.58 x 10 ⁻⁰³	1.22 x 10 ⁻⁰⁵	9.20 x 10 ⁻⁰³	1.54 x 10 ⁻⁰²
	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	7.46 x 10 ⁻⁰⁴	1.93 x 10 ⁻⁰⁸	8.72 x 10 ⁻⁰⁵	4.19 x 10 ⁻⁰²	7.75 x 10 ⁻⁰⁴	1.72 x 10 ⁻⁰⁴	0.00 x 10 ⁺⁰⁰	7.25 x 10 ⁻⁰³	3.48 x 10 ⁻⁰³	1.15 x 10 ⁻⁰²	6.58 x 10 ⁻⁰²

mSv/yr = millisieverts per year; n/a = not applicable or not assessed.

5.2.4.1.4 Radon Dose

The radon dose was calculated separately from the dose from other radionuclides and was estimated outside of IMPACT.

Project activities would result in the release of radon gas into the atmosphere. The atmospheric model for the Project used AERMOD to estimate radon concentrations at various locations based on radon source emissions (EIS Appendix 7A). The mine vent is the dominant source of radon emissions for the Project, representing approximately 98.6% of the total modelled radon emissions. The atmospheric model assumed that the maximum ore grade in year 1 of the Project (4.07%) occurs throughout the Project. The air quality dispersion model was developed as a reasonable upper bound due to the inherent conservatism associated with its configuration and therefore only the Application Case was run. For example, the air quality dispersion model makes the conservative assumption that emission sources from each piece of emitting equipment emit simultaneously and at their maximum possible emission rate. It is unlikely that all sources would emit at the same time; therefore, this assumption leads to higher than actual predicted deposition rates.

The camp worker would be exposed to radon through inhalation while at the camp site, located southwest of the mill terrace. The camp worker represents an adult who resides at the camp for 6 months of the year and away from the site for the remaining 6 months of the year. For exposure to radon, it has been conservatively assumed that the camp worker spends 100% of their time indoors when on site. The predicted radon concentration at the camp site, from AERMOD, is 44.5 Bq/m³ (41.6 Bq/m³ based on incremental radon, where background radon has been subtracted from the total).

The dose from radon in air considers ingrowth of radon decay progeny (polonium-218, lead-214, bismuth-214) during dispersion of radon gas from the source to receptor. Ingrowth was quantified in terms of the radon progeny equilibrium ratio, according to the methods outlined in Health Canada's federal guidance on contaminated site radiological risk assessment in Canada, Part VI (Health Canada 2010b). Radon dose is dependent on the radon equilibrium fraction as well as the exposure time for the receptor.

Consistent with recommendations in CSA N288.6-22 and Health Canada (2010b), the dose from radon in air was calculated according to the equation in Appendix A, Section 2.4.3, with input values shown in Table 5-11.

Indoor radon dose dominates over outdoor radon dose; and therefore, only indoor radon dose was quantified. However, the outdoor equilibrium fraction (F_{out}) was needed to estimate the indoor equilibrium fraction (F_{in}), which is needed to include short-lived progeny in the radon dose calculation.

Radon and radon progeny are expected to be in partial equilibrium at the mine exhaust vent. An initial time correction for estimating equilibrium fractions for radon and progeny from the mine ventilation was accounted for using guidance from USEPA (1985). A time correction factor of



11.79 minutes that corresponds to an equilibrium fraction of 0.2 was considered reasonable to account for the degree of equilibrium between radon and radon progeny in the mine exhaust vent (USEPA 1985). This initial time correction was added to the travel time.

Table 5-11: Summary of Input Parameters for Radon Dose Calculation

Parameter	Value	Source
C _{Rn} (background)	2.9 Bq/m ³	From baseline monitoring at site
C _{Rn} (camp site)	Max-year EA Case: 44.5 Bq/m ³	Atmospheric Model (EIS Section 7.2)
Distance from mine vent to camp	941 m	From AERMOD
Mean wind speed	212 m/min	From meteorological dataset
Initial time correction	11.79 min	USEPA 1985 – to account for partial equilibrium of radon and radon progeny in the mine vent
t (travel time to camp)	16.2 min	Calculated t = Initial time correction + (distance / wind speed)
Exposure time	4,380 h/yr	Assumption based on camp worker residency of 0.5 of the year
Fout	0.26	Calculated
Fin	0.44	Calculated

 Bq/m^3 = becquerels per cubic metre; EA = Environmental Assessment; F_{out} = outdoor equilibrium fraction; F_{in} = indoor equilibrium fraction; C_{Rn} = concentration of radon in air; USEPA = United States Environmental Protection Agency.

The incremental radon dose to the camp worker during Operations from the Application Case is expected to be 0.51 mSv/yr.

5.2.4.2 Estimated Dose to Human Receptors – Reasonably Foreseeable Development Case

5.2.4.2.1 Non-carcinogen Doses

The doses for cobalt, copper, molybdenum, and uranium to the camp worker, subsistence harvester, and seasonal resident are presented by exposure pathway for the RFD Case in Table 5-12.



Table 5-12: Estimated Non-carcinogen Doses to Human Receptors – Reasonably Foreseeable Development Case

				Dos	e by Pathway R	FD Case (mg/k	g/d)		
	СОРС	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
					Base Case				
	Cobalt	8.58 x 10 ⁻⁰⁶	9.04 x 10 ⁻⁰⁹	1.01 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	1.15 x 10 ⁻⁰⁵	7.21 x 10 ⁻⁰⁶	1.97 x 10 ⁻⁰⁴	2.25 x 10 ⁻⁰⁴
	Copper	1.24 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	3.81 x 10 ⁻⁰⁴	1.15 x 10 ⁻⁰⁴	1.66 x 10 ⁻⁰²	1.71 x 10 ⁻⁰²
	Molybdenum	3.04 x 10 ⁻⁰⁶	2.98 x 10 ⁻⁰⁹	4.54 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	3.39 x 10 ⁻⁰⁸	4.10 x 10 ⁻⁰⁶	2.30 x 10 ⁻⁰³	2.31 x 10 ⁻⁰³
	Uranium	1.48 x 10 ⁻⁰⁶	5.17 x 10 ⁻⁰⁹	1.35 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.06 x 10 ⁻⁰⁶	1.15 x 10 ⁻⁰⁵	4.37 x 10 ⁻⁰⁵	5.88 x 10 ⁻⁰⁵
				Pı	oject Lifespan				
Camp Worker	Cobalt	8.63 x 10 ⁻⁰⁶	9.06 x 10 ⁻⁰⁹	1.02 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	1.33 x 10 ⁻⁰⁵	7.29 x 10 ⁻⁰⁶	2.01 x 10 ⁻⁰⁴	2.30 x 10 ⁻⁰⁴
(Patterson Lake North Arm -	Copper	1.24 x 10 ⁻⁰⁵	1.28 x 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	4.18 x 10 ⁻⁰⁴	1.15 x 10 ⁻⁰⁴	1.66 x 10 ⁻⁰²	1.72 x 10 ⁻⁰²
West Basin)	Molybdenum	3.04 x 10 ⁻⁰⁶	3.03 x 10 ⁻⁰⁹	4.55 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	6.14 x 10 ⁻⁰⁸	4.17 x 10 ⁻⁰⁶	2.31 x 10 ⁻⁰³	2.31 x 10 ⁻⁰³
	Uranium	1.48 x 10 ⁻⁰⁶	1.68 x 10 ⁻⁰⁸	1.35 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	6.36 x 10 ⁻⁰⁶	2.60 x 10 ⁻⁰⁵	5.29 x 10 ⁻⁰⁵	8.68 x 10 ⁻⁰⁵
					Far-Future				
	Cobalt	n/a							
	Copper	n/a							
	Molybdenum	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Uranium	n/a							
					Base Case				
	Cobalt	8.58 x 10 ⁻⁰⁶	9.04 x 10 ⁻⁰⁹	1.01 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.30 x 10 ⁻⁰⁵	1.44 x 10 ⁻⁰⁵	2.44 x 10 ⁻⁰⁴	2.90 × 10 ⁻⁰⁴
	Copper	1.24 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	7.61 x 10 ⁻⁰⁴	2.29 x 10 ⁻⁰⁴	1.75 x 10 ⁻⁰²	1.86 × 10 ⁻⁰²
Subsistence	Molybdenum	3.04 x 10 ⁻⁰⁶	2.98 x 10 ⁻⁰⁹	4.54 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	6.77 x 10 ⁻⁰⁸	8.18 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰³	1.96 × 10 ⁻⁰³
Harvester Adult (Patterson Lake	Uranium	1.48 x 10 ⁻⁰⁶	5.17 x 10 ⁻⁰⁹	1.35 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	4.11 x 10 ⁻⁰⁶	2.30 x 10 ⁻⁰⁵	4.92 x 10 ⁻⁰⁵	7.78 × 10 ⁻⁰⁵
South Arm)				Pı	oject Lifespan				
	Cobalt	1.00 x 10 ⁻⁰⁵	9.05 x 10 ⁻⁰⁹	1.12 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.65 x 10 ⁻⁰⁵	1.46 x 10 ⁻⁰⁵	2.51 x 10 ⁻⁰⁴	3.02 x 10 ⁻⁰⁴
	Copper	1.36 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.86 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	8.34 x 10 ⁻⁰⁴	2.30 x 10 ⁻⁰⁴	1.77 x 10 ⁻⁰²	1.88 x 10 ⁻⁰²
	Molybdenum	5.72 x 10 ⁻⁰⁶	2.98 x 10 ⁻⁰⁹	7.11 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	1.23 x 10 ⁻⁰⁷	8.34 x 10 ⁻⁰⁶	1.96 x 10 ⁻⁰³	1.97 x 10 ⁻⁰³



				Dos	e by Pathway R	FD Case (mg/k	g/d)		
	СОРС	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
	Uranium	4.68 x 10 ⁻⁰⁶	6.94 x 10 ⁻⁰⁹	2.23 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	1.27 x 10 ⁻⁰⁵	5.20 x 10 ⁻⁰⁵	6.74 x 10 ⁻⁰⁵	1.37 x 10 ⁻⁰⁴
					Far-Future				
	Cobalt	1.23 x 10 ⁻⁰⁵	9.05 x 10 ⁻⁰⁹	1.46 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	3.30 x 10 ⁻⁰⁵	1.44 x 10 ⁻⁰⁵	2.73 x 10 ⁻⁰⁴	3.33 x 10 ⁻⁰⁴
	Copper	1.82 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	2.57 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	1.12 x 10 ⁻⁰³	2.30 x 10 ⁻⁰⁴	1.79 x 10 ⁻⁰²	1.92 x 10 ⁻⁰²
	Molybdenum	2.18 x 10 ⁻⁰⁵	2.98 x 10 ⁻⁰⁹	3.26 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	4.86 x 10 ⁻⁰⁷	8.18 x 10 ⁻⁰⁶	1.97 x 10 ⁻⁰³	2.01 × 10 ⁻⁰³
	Uranium	3.31 x 10 ⁻⁰⁶	5.36 x 10 ⁻⁰⁹	3.02 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	9.23 x 10 ⁻⁰⁶	2.39 x 10 ⁻⁰⁵	5.18 x 10 ⁻⁰⁵	8.83 x 10 ⁻⁰⁵
					Base Case				
	Cobalt	9.58 x 10 ⁻⁰⁶	5.91 x 10 ⁻⁰⁷	6.62 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	2.18 x 10 ⁻⁰⁵	2.84 x 10 ⁻⁰⁵	4.78 x 10 ⁻⁰⁴	5.39 x 10 ⁻⁰⁴
	Copper	1.38 x 10 ⁻⁰⁵	8.21 x 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	7.21 x 10 ⁻⁰⁴	4.52 x 10 ⁻⁰⁴	4.13 x 10 ⁻⁰²	4.25 x 10 ⁻⁰²
	Molybdenum	3.39 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰⁷	2.97 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	6.42 x 10 ⁻⁰⁸	1.61 x 10 ⁻⁰⁵	6.06 x 10 ⁻⁰³	6.08 x 10 ⁻⁰³
	Uranium	1.65 x 10 ⁻⁰⁶	3.38 x 10 ⁻⁰⁷	8.79 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	3.90 x 10 ⁻⁰⁶	4.54 x 10 ⁻⁰⁵	1.02 x 10 ⁻⁰⁴	1.54 x 10 ⁻⁰⁴
Subsistence				Pı	oject Lifespan				
Harvester One-	Cobalt	1.12 x 10 ⁻⁰⁵	5.91 x 10 ⁻⁰⁷	7.34 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	2.52 x 10 ⁻⁰⁵	2.86 x 10 ⁻⁰⁵	4.85 x 10 ⁻⁰⁴	5.51 x 10 ⁻⁰⁴
Year-Old	Copper	1.52 x 10 ⁻⁰⁵	8.22 x 10 ⁻⁰⁷	1.22 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	7.91 x 10 ⁻⁰⁴	4.53 x 10 ⁻⁰⁴	4.15 x 10 ⁻⁰²	4.27 x 10 ⁻⁰²
(Patterson Lake South Arm)	Molybdenum	6.39 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰⁷	4.65 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.16 x 10 ⁻⁰⁷	1.63 x 10 ⁻⁰⁵	6.07 x 10 ⁻⁰³	6.09 x 10 ⁻⁰³
Journ Arm,	Uranium	5.23 x 10 ⁻⁰⁶	4.54 x 10 ⁻⁰⁷	1.46 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	1.20 x 10 ⁻⁰⁵	8.90 x 10 ⁻⁰⁵	1.15 x 10 ⁻⁰⁴	2.24 x 10 ⁻⁰⁴
					Far-Future				
	Cobalt	1.38 x 10 ⁻⁰⁵	5.91 x 10 ⁻⁰⁷	9.54 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	3.13 x 10 ⁻⁰⁵	2.84 x 10 ⁻⁰⁵	5.08 x 10 ⁻⁰⁴	5.83 x 10 ⁻⁰⁴
	Copper	2.03 x 10 ⁻⁰⁵	8.21 x 10 ⁻⁰⁷	1.68 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	1.06 x 10 ⁻⁰³	4.52 x 10 ⁻⁰⁴	4.16 x 10 ⁻⁰²	4.32 x 10 ⁻⁰²
	Molybdenum	2.43 x 10 ⁻⁰⁵	1.95 x 10 ⁻⁰⁷	2.13 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	4.61 x 10 ⁻⁰⁷	1.61 x 10 ⁻⁰⁵	6.08 x 10 ⁻⁰³	6.13 x 10 ⁻⁰³
	Uranium	3.70 x 10 ⁻⁰⁶	3.50 x 10 ⁻⁰⁷	1.98 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	8.75 x 10 ⁻⁰⁶	4.70 x 10 ⁻⁰⁵	1.04 x 10 ⁻⁰⁴	1.66 x 10 ⁻⁰⁴
					Base Case				
Subsistence Harvester Adult	Cobalt	8.58 x 10 ⁻⁰⁶	9.04 x 10 ⁻⁰⁹	1.01 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.30 x 10 ⁻⁰⁵	1.44 x 10 ⁻⁰⁵	2.44 x 10 ⁻⁰⁴	2.90 x 10 ⁻⁰⁴
(Beet Lake)	Copper	1.24 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	7.61 x 10 ⁻⁰⁴	2.29 x 10 ⁻⁰⁴	1.75 x 10 ⁻⁰²	1.86 x 10 ⁻⁰²
	Molybdenum	3.04 x 10 ⁻⁰⁶	2.98 x 10 ⁻⁰⁹	4.54 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	6.77 x 10 ⁻⁰⁸	8.18 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰³	1.96 x 10 ⁻⁰³



				Dose	e by Pathway R	FD Case (mg/k	g/d)		
	СОРС	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
	Uranium	1.48 x 10 ⁻⁰⁶	5.17 x 10 ⁻⁰⁹	1.35 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	4.11 x 10 ⁻⁰⁶	2.30 x 10 ⁻⁰⁵	4.92 x 10 ⁻⁰⁵	7.78 x 10 ⁻⁰⁵
				Pr	oject Lifespan				
	Cobalt	9.22 x 10 ⁻⁰⁶	9.05 x 10 ⁻⁰⁹	1.07 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.45 x 10 ⁻⁰⁵	1.46 x 10 ⁻⁰⁵	2.47 x 10 ⁻⁰⁴	2.96 x 10 ⁻⁰⁴
	Copper	1.29 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.81 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	7.93 x 10 ⁻⁰⁴	2.30 x 10 ⁻⁰⁴	1.76 x 10 ⁻⁰²	1.86 x 10 ⁻⁰²
	Molybdenum	4.23 x 10 ⁻⁰⁶	2.98 x 10 ⁻⁰⁹	5.88 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	9.23 x 10 ⁻⁰⁸	8.34 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰³	1.97 x 10 ⁻⁰³
	Uranium	2.59 x 10 ⁻⁰⁶	6.16 x 10 ⁻⁰⁹	1.67 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	7.09 x 10 ⁻⁰⁶	3.92 x 10 ⁻⁰⁵	5.60 x 10 ⁻⁰⁵	1.05 x 10 ⁻⁰⁴
					Far-Future				
	Cobalt	1.06 x 10 ⁻⁰⁵	9.05 x 10 ⁻⁰⁹	1.25 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.83 x 10 ⁻⁰⁵	1.44 x 10 ⁻⁰⁵	2.54 x 10 ⁻⁰⁴	3.08 x 10 ⁻⁰⁴
	Copper	1.54 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	2.18 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	9.46 x 10 ⁻⁰⁴	2.29 x 10 ⁻⁰⁴	1.77 x 10 ⁻⁰²	1.89 x 10 ⁻⁰²
	Molybdenum	1.28 x 10 ⁻⁰⁵	2.98 x 10 ⁻⁰⁹	1.92 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.86 x 10 ⁻⁰⁷	8.18 x 10 ⁻⁰⁶	1.96 x 10 ⁻⁰³	1.98 x 10 ⁻⁰³
	Uranium	2.16 x 10 ⁻⁰⁶	5.28 x 10 ⁻⁰⁹	1.97 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	6.01 x 10 ⁻⁰⁶	2.35 x 10 ⁻⁰⁵	5.01 x 10 ⁻⁰⁵	8.18 x 10 ⁻⁰⁵
					Base Case				
	Cobalt	9.58 x 10 ⁻⁰⁶	5.91 x 10 ⁻⁰⁷	6.62 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	2.18 x 10 ⁻⁰⁵	2.84 x 10 ⁻⁰⁵	4.78 x 10 ⁻⁰⁴	5.39 x 10 ⁻⁰⁴
	Copper	1.38 x 10 ⁻⁰⁵	8.21 x 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	7.21 x 10 ⁻⁰⁴	4.52 x 10 ⁻⁰⁴	4.13 x 10 ⁻⁰²	4.25 x 10 ⁻⁰²
	Molybdenum	3.39 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰⁷	2.97 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	6.42 x 10 ⁻⁰⁸	1.61 x 10 ⁻⁰⁵	6.06 x 10 ⁻⁰³	6.08 x 10 ⁻⁰³
	Uranium	1.65 x 10 ⁻⁰⁶	3.38 x 10 ⁻⁰⁷	8.79 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	3.90 x 10 ⁻⁰⁶	4.54 x 10 ⁻⁰⁵	1.02 x 10 ⁻⁰⁴	1.54 x 10 ⁻⁰⁴
Subsistence				Pr	oject Lifespan				
Harvester One-	Cobalt	1.03 x 10 ⁻⁰⁵	5.91 x 10 ⁻⁰⁷	7.01 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	2.33 x 10 ⁻⁰⁵	2.86 x 10 ⁻⁰⁵	4.81 x 10 ⁻⁰⁴	5.45 x 10 ⁻⁰⁴
Year-Old (Beet	Copper	1.44 x 10 ⁻⁰⁵	8.22 x 10 ⁻⁰⁷	1.18 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	7.52 x 10 ⁻⁰⁴	4.53 x 10 ⁻⁰⁴	4.13 x 10 ⁻⁰²	4.26 x 10 ⁻⁰²
Lake)	Molybdenum	4.73 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰⁷	3.84 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	8.75 x 10 ⁻⁰⁸	1.63 x 10 ⁻⁰⁵	6.06 x 10 ⁻⁰³	6.08 x 10 ⁻⁰³
	Uranium	2.89 x 10 ⁻⁰⁶	4.03 x 10 ⁻⁰⁷	1.09 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	6.73 x 10 ⁻⁰⁶	6.98 x 10 ⁻⁰⁵	1.07 x 10 ⁻⁰⁴	1.88 x 10 ⁻⁰⁴
					Far-Future				
	Cobalt	1.18 x 10 ⁻⁰⁵	5.91 x 10 ⁻⁰⁷	8.17 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	2.68 x 10 ⁻⁰⁵	2.84 x 10 ⁻⁰⁵	4.89 x 10 ⁻⁰⁴	5.57 x 10 ⁻⁰⁴
	Copper	1.72 x 10 ⁻⁰⁵	8.21 x 10 ⁻⁰⁷	1.43 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	8.97 x 10 ⁻⁰⁴	4.52 x 10 ⁻⁰⁴	4.14 x 10 ⁻⁰²	4.28 x 10 ⁻⁰²
	Molybdenum	1.43 x 10 ⁻⁰⁵	1.95 x 10 ⁻⁰⁷	1.25 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	2.71 x 10 ⁻⁰⁷	1.61 x 10 ⁻⁰⁵	6.07 x 10 ⁻⁰³	6.10 x 10 ⁻⁰³



				Dos	e by Pathway R	FD Case (mg/k	g/d)		
	СОРС	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
	Uranium	2.41 x 10 ⁻⁰⁶	3.45 x 10 ⁻⁰⁷	1.29 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	5.70 x 10 ⁻⁰⁶	4.63 x 10 ⁻⁰⁵	1.03 x 10 ⁻⁰⁴	1.59 x 10 ⁻⁰⁴
					Base Case				
	Cobalt	8.58 x 10 ⁻⁰⁶	9.04 x 10 ⁻⁰⁹	1.01 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.30 x 10 ⁻⁰⁵	1.44 x 10 ⁻⁰⁵	2.44 x 10 ⁻⁰⁴	2.90 x 10 ⁻⁰⁴
	Copper	1.24 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	7.61 x 10 ⁻⁰⁴	2.29 x 10 ⁻⁰⁴	1.75 x 10 ⁻⁰²	1.86 x 10 ⁻⁰²
	Molybdenum	3.04 x 10 ⁻⁰⁶	2.98 x 10 ⁻⁰⁹	4.54 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	6.77 x 10 ⁻⁰⁸	8.18 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰³	1.96 x 10 ⁻⁰³
	Uranium	1.48 x 10 ⁻⁰⁶	5.17 x 10 ⁻⁰⁹	1.35 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	4.11 x 10 ⁻⁰⁶	2.30 x 10 ⁻⁰⁵	4.92 x 10 ⁻⁰⁵	7.78 x 10 ⁻⁰⁵
				Pr	oject Lifespan				
Subsistence	Cobalt	8.64 x 10 ⁻⁰⁶	9.05 x 10 ⁻⁰⁹	1.02 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.31 x 10 ⁻⁰⁵	1.46 x 10 ⁻⁰⁵	2.45 x 10 ⁻⁰⁴	2.91 x 10 ⁻⁰⁴
Harvester Adult	Copper	1.24 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	7.64 x 10 ⁻⁰⁴	2.30 x 10 ⁻⁰⁴	1.76 x 10 ⁻⁰²	1.86 x 10 ⁻⁰²
(Lloyd Lake)	Molybdenum	3.16 x 10 ⁻⁰⁶	2.98 x 10 ⁻⁰⁹	4.68 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	7.02 x 10 ⁻⁰⁸	8.34 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰³	1.97 x 10 ⁻⁰³
	Uranium	1.58 x 10 ⁻⁰⁶	5.19 x 10 ⁻⁰⁹	1.38 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	4.39 x 10 ⁻⁰⁶	2.32 x 10 ⁻⁰⁵	4.93 x 10 ⁻⁰⁵	7.85 x 10 ⁻⁰⁵
		,			Far-Future				
	Cobalt	8.78 x 10 ⁻⁰⁶	9.05 x 10 ⁻⁰⁹	1.04 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.35 x 10 ⁻⁰⁵	1.44 x 10 ⁻⁰⁵	2.45 x 10 ⁻⁰⁴	2.92 x 10 ⁻⁰⁴
	Copper	1.27 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.80 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	7.80 x 10 ⁻⁰⁴	2.29 x 10 ⁻⁰⁴	1.76 x 10 ⁻⁰²	1.86 x 10 ⁻⁰²
	Molybdenum	4.03 x 10 ⁻⁰⁶	2.98 x 10 ⁻⁰⁹	6.03 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	8.99 x 10 ⁻⁰⁸	8.18 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰³	1.97 x 10 ⁻⁰³
	Uranium	1.54 x 10 ⁻⁰⁶	5.18 x 10 ⁻⁰⁹	1.40 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	4.28 x 10 ⁻⁰⁶	2.30 x 10 ⁻⁰⁵	4.92 x 10 ⁻⁰⁵	7.81 x 10 ⁻⁰⁵
					Base Case				
	Cobalt	9.58 x 10 ⁻⁰⁶	5.91 x 10 ⁻⁰⁷	6.62 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	2.18 x 10 ⁻⁰⁵	2.84 x 10 ⁻⁰⁵	4.78 x 10 ⁻⁰⁴	5.39 x 10 ⁻⁰⁴
	Copper	1.38 x 10 ⁻⁰⁵	8.21 x 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	7.21 x 10 ⁻⁰⁴	4.52 x 10 ⁻⁰⁴	4.13 x 10 ⁻⁰²	4.25 x 10 ⁻⁰²
Subsistence	Molybdenum	3.39 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰⁷	2.97 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	6.42 x 10 ⁻⁰⁸	1.61 x 10 ⁻⁰⁵	6.06 x 10 ⁻⁰³	6.08 x 10 ⁻⁰³
Harvester One- Year-Old (Lloyd	Uranium	1.65 x 10 ⁻⁰⁶	3.38 x 10 ⁻⁰⁷	8.79 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	3.90 x 10 ⁻⁰⁶	4.54 x 10 ⁻⁰⁵	1.02 x 10 ⁻⁰⁴	1.54 x 10 ⁻⁰⁴
Lake)		,		Pr	oject Lifespan				
	Cobalt	9.65 x 10 ⁻⁰⁶	5.91 x 10 ⁻⁰⁷	6.66 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	2.19 x 10 ⁻⁰⁵	2.86 x 10 ⁻⁰⁵	4.79 x 10 ⁻⁰⁴	5.40 x 10 ⁻⁰⁴
	Copper	1.39 x 10 ⁻⁰⁵	8.22 x 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	7.24 x 10 ⁻⁰⁴	4.53 x 10 ⁻⁰⁴	4.13 x 10 ⁻⁰²	4.25 x 10 ⁻⁰²
	Molybdenum	3.52 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰⁷	3.06 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	6.65 x 10 ⁻⁰⁸	1.63 x 10 ⁻⁰⁵	6.06 x 10 ⁻⁰³	6.08 x 10 ⁻⁰³



				Dos	e by Pathway R	FD Case (mg/k	g/d)		
	СОРС	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
	Uranium	1.76 x 10 ⁻⁰⁶	3.39 x 10 ⁻⁰⁷	8.99 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	4.16 x 10 ⁻⁰⁶	4.57 x 10 ⁻⁰⁵	1.02 x 10 ⁻⁰⁴	1.55 x 10 ⁻⁰⁴
					Far-Future				
	Cobalt	9.80 x 10 ⁻⁰⁶	5.91 x 10 ⁻⁰⁷	6.78 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	2.23 x 10 ⁻⁰⁵	2.84 x 10 ⁻⁰⁵	4.79 x 10 ⁻⁰⁴	5.41 x 10 ⁻⁰⁴
	Copper	1.42 x 10 ⁻⁰⁵	8.21 x 10 ⁻⁰⁷	1.18 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	7.39 x 10 ⁻⁰⁴	4.52 x 10 ⁻⁰⁴	4.13 x 10 ⁻⁰²	4.25 x 10 ⁻⁰²
	Molybdenum	4.50 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰⁷	3.94 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	8.52 x 10 ⁻⁰⁸	1.61 x 10 ⁻⁰⁵	6.06 x 10 ⁻⁰³	6.08 x 10 ⁻⁰³
	Uranium	1.72 x 10 ⁻⁰⁶	3.38 x 10 ⁻⁰⁷	9.16 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	4.06 x 10 ⁻⁰⁶	4.54 x 10 ⁻⁰⁵	1.02 x 10 ⁻⁰⁴	1.55 x 10 ⁻⁰⁴
					Base Case				
	Cobalt	8.58 x 10 ⁻⁰⁶	9.04 x 10 ⁻⁰⁹	1.01 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	7.27 x 10 ⁻⁰⁶	1.13 x 10 ⁻⁰⁵	1.88 x 10 ⁻⁰⁴	2.15 x 10 ⁻⁰⁴
	Copper	1.24 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.41 x 10 ⁻⁰⁴	1.80 x 10 ⁻⁰⁴	1.65 x 10 ⁻⁰²	1.69 x 10 ⁻⁰²
	Molybdenum	3.04 x 10 ⁻⁰⁶	2.98 x 10 ⁻⁰⁹	4.54 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	2.14 x 10 ⁻⁰⁸	6.42 x 10 ⁻⁰⁶	2.41 x 10 ⁻⁰³	2.42 x 10 ⁻⁰³
	Uranium	1.48 x 10 ⁻⁰⁶	5.17 x 10 ⁻⁰⁹	1.35 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	1.30 x 10 ⁻⁰⁶	1.81 x 10 ⁻⁰⁵	4.08 x 10 ⁻⁰⁵	6.16 x 10 ⁻⁰⁵
				Pı	roject Lifespan				
Seasonal	Cobalt	9.45 x 10 ⁻⁰⁶	9.05 x 10 ⁻⁰⁹	1.08 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	7.96 x 10 ⁻⁰⁶	1.13 x 10 ⁻⁰⁵	1.89 x 10 ⁻⁰⁴	2.18 x 10 ⁻⁰⁴
Resident Adult (Patterson Lake	Copper	1.31 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.82 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.55 x 10 ⁻⁰⁴	1.80 x 10 ⁻⁰⁴	1.65 x 10 ⁻⁰²	1.69 x 10 ⁻⁰²
South Arm)	Molybdenum	4.65 x 10 ⁻⁰⁶	2.98 x 10 ⁻⁰⁹	6.08 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	3.19 x 10 ⁻⁰⁸	6.46 x 10 ⁻⁰⁶	2.41 x 10 ⁻⁰³	2.42 x 10 ⁻⁰³
	Uranium	3.40 x 10 ⁻⁰⁶	6.24 x 10 ⁻⁰⁹	1.88 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.94 x 10 ⁻⁰⁶	2.71 x 10 ⁻⁰⁵	4.42 x 10 ⁻⁰⁵	7.77 x 10 ⁻⁰⁵
					Far-Future				
	Cobalt	1.08 x 10 ⁻⁰⁵	9.04 x 10 ⁻⁰⁹	1.28 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	9.19 x 10 ⁻⁰⁶	1.13 x 10 ⁻⁰⁵	1.95 x 10 ⁻⁰⁴	2.26 x 10 ⁻⁰⁴
	Copper	1.59 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	2.25 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	3.08 x 10 ⁻⁰⁴	1.80 x 10 ⁻⁰⁴	1.65 x 10 ⁻⁰²	1.70 x 10 ⁻⁰²
	Molybdenum	1.43 x 10 ⁻⁰⁵	2.98 x 10 ⁻⁰⁹	2.14 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	1.01 x 10 ⁻⁰⁷	6.42 x 10 ⁻⁰⁶	2.41 x 10 ⁻⁰³	2.43 x 10 ⁻⁰³
	Uranium	2.58 x 10 ⁻⁰⁶	5.29 x 10 ⁻⁰⁹	2.35 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.28 x 10 ⁻⁰⁶	1.85 x 10 ⁻⁰⁵	4.14 x 10 ⁻⁰⁵	6.47 x 10 ⁻⁰⁵
Seasonal					Base Case				
Resident One- Year-Old	Cobalt	9.58 x 10 ⁻⁰⁶	5.91 x 10 ⁻⁰⁷	6.62 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	6.89 x 10 ⁻⁰⁶	2.47 x 10 ⁻⁰⁵	4.13 x 10 ⁻⁰⁴	4.56 x 10 ⁻⁰⁴
(Patterson Lake	Copper	1.38 x 10 ⁻⁰⁵	8.21 x 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	2.29 x 10 ⁻⁰⁴	3.93 x 10 ⁻⁰⁴	3.97 x 10 ⁻⁰²	4.03 x 10 ⁻⁰²
South Arm)	Molybdenum	3.39 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰⁷	2.97 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	2.03 x 10 ⁻⁰⁸	1.40 x 10 ⁻⁰⁵	6.38 x 10 ⁻⁰³	6.40 x 10 ⁻⁰³



				Dose	e by Pathway R	FD Case (mg/k	g/d)		
	СОРС	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
	Uranium	1.65 x 10 ⁻⁰⁶	3.38 x 10 ⁻⁰⁷	8.79 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.23 x 10 ⁻⁰⁶	3.95 x 10 ⁻⁰⁵	9.62 x 10 ⁻⁰⁵	1.40 x 10 ⁻⁰⁴
				Pr	oject Lifespan				
	Cobalt	1.06 x 10 ⁻⁰⁵	5.91 x 10 ⁻⁰⁷	7.06 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	7.54 x 10 ⁻⁰⁶	2.48 x 10 ⁻⁰⁵	4.15 x 10 ⁻⁰⁴	4.59 x 10 ⁻⁰⁴
	Copper	1.47 x 10 ⁻⁰⁵	8.22 x 10 ⁻⁰⁷	1.19 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	2.42 x 10 ⁻⁰⁴	3.93 x 10 ⁻⁰⁴	3.97 x 10 ⁻⁰²	4.04 x 10 ⁻⁰²
	Molybdenum	5.19 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰⁷	3.98 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	3.03 x 10 ⁻⁰⁸	1.41 x 10 ⁻⁰⁵	6.38 x 10 ⁻⁰³	6.40 x 10 ⁻⁰³
	Uranium	3.80 x 10 ⁻⁰⁶	4.07 x 10 ⁻⁰⁷	1.23 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	2.78 x 10 ⁻⁰⁶	5.57 x 10 ⁻⁰⁵	9.87 x 10 ⁻⁰⁵	1.63 x 10 ⁻⁰⁴
					Far-Future				
	Cobalt	1.21 x 10 ⁻⁰⁵	5.91 x 10 ⁻⁰⁷	8.37 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	8.71 x 10 ⁻⁰⁶	2.47 x 10 ⁻⁰⁵	4.21 x 10 ⁻⁰⁴	4.68 x 10 ⁻⁰⁴
	Copper	1.77 x 10 ⁻⁰⁵	8.21 x 10 ⁻⁰⁷	1.47 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	2.92 x 10 ⁻⁰⁴	3.93 x 10 ⁻⁰⁴	3.98 x 10 ⁻⁰²	4.05 x 10 ⁻⁰²
	Molybdenum	1.60 x 10 ⁻⁰⁵	1.95 x 10 ⁻⁰⁷	1.40 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	9.58 x 10 ⁻⁰⁸	1.40 x 10 ⁻⁰⁵	6.38 x 10 ⁻⁰³	6.41 x 10 ⁻⁰³
	Uranium	2.88 x 10 ⁻⁰⁶	3.45 x 10 ⁻⁰⁷	1.54 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	2.16 x 10 ⁻⁰⁶	4.03 x 10 ⁻⁰⁵	9.67 x 10 ⁻⁰⁵	1.44 x 10 ⁻⁰⁴
					Base Case				
	Cobalt	8.58 x 10 ⁻⁰⁶	9.04 x 10 ⁻⁰⁹	1.01 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	7.27 x 10 ⁻⁰⁶	1.13 x 10 ⁻⁰⁵	1.88 x 10 ⁻⁰⁴	2.15 x 10 ⁻⁰⁴
	Copper	1.24 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.41 x 10 ⁻⁰⁴	1.80 x 10 ⁻⁰⁴	1.65 x 10 ⁻⁰²	1.69 x 10 ⁻⁰²
	Molybdenum	3.04 x 10 ⁻⁰⁶	2.98 x 10 ⁻⁰⁹	4.54 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	2.14 x 10 ⁻⁰⁸	6.42 x 10 ⁻⁰⁶	2.41 x 10 ⁻⁰³	2.42 x 10 ⁻⁰³
	Uranium	1.48 x 10 ⁻⁰⁶	5.17 x 10 ⁻⁰⁹	1.35 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	1.30 x 10 ⁻⁰⁶	1.81 x 10 ⁻⁰⁵	4.08 x 10 ⁻⁰⁵	6.16 x 10 ⁻⁰⁵
				Pr	oject Lifespan				
Seasonal Resident Adult	Cobalt	8.62 x 10 ⁻⁰⁶	9.05 x 10 ⁻⁰⁹	1.02 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	7.33 x 10 ⁻⁰⁶	1.13 x 10 ⁻⁰⁵	1.88 x 10 ⁻⁰⁴	2.15 x 10 ⁻⁰⁴
(Lloyd Lake)	Copper	1.24 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	3.21 x 10 ⁻⁰⁴	1.80 x 10 ⁻⁰⁴	1.65 x 10 ⁻⁰²	1.70 x 10 ⁻⁰²
	Molybdenum	3.11 x 10 ⁻⁰⁶	2.98 x 10 ⁻⁰⁹	4.63 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	2.23 x 10 ⁻⁰⁸	6.46 x 10 ⁻⁰⁶	2.41 x 10 ⁻⁰³	2.42 x 10 ⁻⁰³
	Uranium	1.54 x 10 ⁻⁰⁶	5.18 x 10 ⁻⁰⁹	1.36 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	1.73 x 10 ⁻⁰⁶	1.81 x 10 ⁻⁰⁵	4.08 x 10 ⁻⁰⁵	6.22 x 10 ⁻⁰⁵
					Far-Future				
	Cobalt	8.70 x 10 ⁻⁰⁶	9.04 x 10 ⁻⁰⁹	1.03 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	7.45 x 10 ⁻⁰⁶	1.13 x 10 ⁻⁰⁵	1.88 x 10 ⁻⁰⁴	2.16 x 10 ⁻⁰⁴
	Copper	1.26 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.78 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	3.28 x 10 ⁻⁰⁴	1.80 x 10 ⁻⁰⁴	1.65 x 10 ⁻⁰²	1.70 x 10 ⁻⁰²
	Molybdenum	3.63 x 10 ⁻⁰⁶	2.98 x 10 ⁻⁰⁹	5.44 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	2.85 x 10 ⁻⁰⁸	6.42 x 10 ⁻⁰⁶	2.41 x 10 ⁻⁰³	2.42 x 10 ⁻⁰³



				Dos	e by Pathway R	FD Case (mg/k	g/d)						
	СОРС	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC				
	Uranium	1.51 x 10 ⁻⁰⁶	5.18 x 10 ⁻⁰⁹	1.38 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	1.68 x 10 ⁻⁰⁶	1.81 x 10 ⁻⁰⁵	4.08 x 10 ⁻⁰⁵	6.21 x 10 ⁻⁰⁵				
					Base Case								
	Cobalt	9.58 x 10 ⁻⁰⁶	5.91 x 10 ⁻⁰⁷	6.62 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	6.89 x 10 ⁻⁰⁶	2.47 x 10 ⁻⁰⁵	4.13 x 10 ⁻⁰⁴	4.56 x 10 ⁻⁰⁴				
	Copper	1.38 x 10 ⁻⁰⁵	8.21 x 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	2.29 x 10 ⁻⁰⁴	3.93 x 10 ⁻⁰⁴	3.97 x 10 ⁻⁰²	4.03 x 10 ⁻⁰²				
	Molybdenum	3.39 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰⁷	2.97 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	2.03 x 10 ⁻⁰⁸	1.40 x 10 ⁻⁰⁵	6.38 x 10 ⁻⁰³	6.40 x 10 ⁻⁰³				
Seasonal Resident One- Year-Old (Lloyd	Uranium	1.65 x 10 ⁻⁰⁶	3.38 x 10 ⁻⁰⁷	8.79 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.23 x 10 ⁻⁰⁶	3.95 x 10 ⁻⁰⁵	9.62 x 10 ⁻⁰⁵	1.40 x 10 ⁻⁰⁴				
	Project Lifespan												
	Cobalt	9.62 x 10 ⁻⁰⁶	5.91 x 10 ⁻⁰⁷	6.65 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	6.92 x 10 ⁻⁰⁶	2.48 x 10 ⁻⁰⁵	4.14 x 10 ⁻⁰⁴	4.56 x 10 ⁻⁰⁴				
	Copper	1.39 x 10 ⁻⁰⁵	8.22 x 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	2.29 x 10 ⁻⁰⁴	3.93 x 10 ⁻⁰⁴	3.97 x 10 ⁻⁰²	4.03 x 10 ⁻⁰²				
Lake)	Molybdenum	3.47 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰⁷	3.02 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	2.08 x 10 ⁻⁰⁸	1.41 x 10 ⁻⁰⁵	6.38 x 10 ⁻⁰³	6.40 x 10 ⁻⁰³				
	Uranium	1.72 x 10 ⁻⁰⁶	3.39 x 10 ⁻⁰⁷	8.91 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.28 x 10 ⁻⁰⁶	3.96 x 10 ⁻⁰⁵	9.63 x 10 ⁻⁰⁵	1.40 x 10 ⁻⁰⁴				
	Far-Future Far-Future												
	Cobalt	9.71 x 10 ⁻⁰⁶	5.91 x 10 ⁻⁰⁷	6.72 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	6.99 x 10 ⁻⁰⁶	2.47 x 10 ⁻⁰⁵	4.14 x 10 ⁻⁰⁴	4.56 x 10 ⁻⁰⁴				
	Copper	1.40 x 10 ⁻⁰⁵	8.21 x 10 ⁻⁰⁷	1.16 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	2.32 x 10 ⁻⁰⁴	3.93 x 10 ⁻⁰⁴	3.97 x 10 ⁻⁰²	4.03 x 10 ⁻⁰²				
	Molybdenum	4.06 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰⁷	3.55 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	2.43 x 10 ⁻⁰⁸	1.40 x 10 ⁻⁰⁵	6.38 x 10 ⁻⁰³	6.40 x 10 ⁻⁰³				
	Uranium	1.69 x 10 ⁻⁰⁶	3.38 x 10 ⁻⁰⁷	9.02 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.27 x 10 ⁻⁰⁶	3.95 x 10 ⁻⁰⁵	9.63 x 10 ⁻⁰⁵	1.40 x 10 ⁻⁰⁴				
				Ва	ase Case Adult								
	Cobalt	8.58 x 10 ⁻⁰⁶	9.04 x 10 ⁻⁰⁹	1.01 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.30 x 10 ⁻⁰⁵	1.44 x 10 ⁻⁰⁵	2.44 x 10 ⁻⁰⁴	1.24 x 10 ⁻⁰⁵				
Downsnort	Copper	1.24 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	1.76 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	7.61 x 10 ⁻⁰⁴	2.29 x 10 ⁻⁰⁴	1.75 x 10 ⁻⁰²	1.48 x 10 ⁻⁰⁶				
Permanent Resident	Molybdenum	3.04 x 10 ⁻⁰⁶	2.98 x 10 ⁻⁰⁹	4.54 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	6.77 x 10 ⁻⁰⁸	8.18 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰³	0.00 x 10 ⁺⁰⁰				
(Patterson Lake	Uranium	1.48 x 10 ⁻⁰⁶	5.17 x 10 ⁻⁰⁹	1.35 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	4.11 x 10 ⁻⁰⁶	2.30 x 10 ⁻⁰⁵	4.92 x 10 ⁻⁰⁵	3.04 x 10 ⁻⁰⁶				
North Arm - West Basin)				Base (Case One-Year (Old							
West Bushi)	Cobalt	9.58 x 10 ⁻⁰⁶	5.91 x 10 ⁻⁰⁷	6.62 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	2.18 x 10 ⁻⁰⁵	2.84 x 10 ⁻⁰⁵	4.78 x 10 ⁻⁰⁴	1.38 x 10 ⁻⁰⁵				
	Copper	1.38 x 10 ⁻⁰⁵	8.21 x 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	7.21 x 10 ⁻⁰⁴	4.52 x 10 ⁻⁰⁴	4.13 x 10 ⁻⁰²	1.65 x 10 ⁻⁰⁶				
	Molybdenum	3.39 x 10 ⁻⁰⁶	1.95 x 10 ⁻⁰⁷	2.97 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	6.42 x 10 ⁻⁰⁸	1.61 x 10 ⁻⁰⁵	6.06 x 10 ⁻⁰³	0.00 x 10 ⁺⁰⁰				



ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT HUMAN HEALTH RISK ASSESSMENT

			Dose	e by Pathway R	FD Case (mg/k	g/d)		
СОРС	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
Uranium	1.65 x 10 ⁻⁰⁶	3.38 x 10 ⁻⁰⁷	8.79 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	3.90 x 10 ⁻⁰⁶	4.54 x 10 ⁻⁰⁵	1.02 x 10 ⁻⁰⁴	3.39 x 10 ⁻¹
			Fa	r-Future Adult				
Cobalt	2.10 x 10 ⁻⁰⁵	9.05 x 10 ⁻⁰⁹	2.48 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	5.62 x 10 ⁻⁰⁵	1.44 x 10 ⁻⁰⁵	3.08 x 10 ⁻⁰⁴	3.21 x 10 ⁻¹
Copper	3.21 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁸	4.54 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	1.97 x 10 ⁻⁰³	2.30 x 10 ⁻⁰⁴	1.83 x 10 ⁻⁰²	1.48 x 10 ⁻⁰
Molybdenum	6.77 x 10 ⁻⁰⁵	2.98 x 10 ⁻⁰⁹	1.01 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.51 x 10 ⁻⁰⁶	8.18 x 10 ⁻⁰⁶	2.00 x 10 ⁻⁰³	0.00 x 10 ⁺⁰
Uranium	1.48 x 10 ⁻⁰⁵	7.50 x 10 ⁻⁰⁹	1.35 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	4.11 x 10 ⁻⁰⁵	3.34 x 10 ⁻⁰⁵	6.77 x 10 ⁻⁰⁵	6.77 x 10 ⁻⁰
			Far Fu	ture One-Year	Old			
Cobalt	2.34 x 10 ⁻⁰⁵	5.92 x 10 ⁻⁰⁷	1.62 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	5.33 x 10 ⁻⁰⁵	2.84 x 10 ⁻⁰⁵	5.43 x 10 ⁻⁰⁴	3.58 x 10 ⁻⁰
Copper	3.58 x 10 ⁻⁰⁵	8.23 x 10 ⁻⁰⁷	2.97 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	1.87 x 10 ⁻⁰³	4.53 x 10 ⁻⁰⁴	4.20 x 10 ⁻⁰²	1.65 x 10 ⁻⁰
Molybdenum	7.56 x 10 ⁻⁰⁵	1.95 x 10 ⁻⁰⁷	6.62 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	1.43 x 10 ⁻⁰⁶	1.61 x 10 ⁻⁰⁵	6.11 x 10 ⁻⁰³	0.00 x 10+
Uranium	1.65 x 10 ⁻⁰⁵	4.90 x 10 ⁻⁰⁷	8.80 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	3.90 x 10 ⁻⁰⁵	6.58 x 10 ⁻⁰⁵	1.16 x 10 ⁻⁰⁴	7.56 x 10 ⁻⁰

 $\overline{\text{COPC}}$ = constituents of potential concern; n/a = not applicable or not assessed; RFD = reasonably foreseeable development.



5.2.4.2.2 Carcinogen Doses

Arsenic was evaluated in the HHRA as a non-threshold carcinogen; therefore, the exposure would be averaged over the receptor's lifetime to estimate a lifetime average daily dose (Health Canada 2021a). For this assessment, the lifetime average daily dose was estimated for various age groups (toddler, child, teen, adult) to permit estimation of the lifetime risk to a composite receptor for each of the subsistence harvester, seasonal resident, and permanent resident. For the camp worker, an adult receptor was considered appropriate.

The estimated lifetime average daily dose for the RFD case for Operations and the far-future projection is presented in Table 5-13.



Table 5-13: Estimated Carcinogen Doses for Arsenic to Human Receptors – Reasonably Foreseeable Development Case – Operations and Far-Future Projection

	Lifetime Average Daily Dose (mg/kg/d)													
		Reasona	bly Foreseeable Dev	elopment Case – O	perations	Reasonably Foreseeable Development – Far-Future Projection								
Age Group	Age Group Camp Worker F		Subsistence Harvester - Beet Lake	Subsistence Seasonal Resident - Harvester - Lloyd Patterson Lake Lake South Arm		Seasonal Resident Permanent - Lloyd Lake Resident		Subsistence Harvester - Patterson Lake South Arm	Subsistence Subsistence Harvester - Beet Harvester - Lloy Lake Lake		Seasonal Resident - Patterson Lake South Arm	Seasonal Resident - Lloyd Lake		
1 year old	n/a	2.80 × 10 ⁻⁰⁶	5.63 × 10 ⁻⁰⁸	2.60 × 10 ⁻⁰⁸	5.80 × 10 ⁻⁰⁷	6.15 × 10 ⁻⁰⁹	1.25 × 10 ⁻⁰⁷	2.37 × 10 ⁻⁰⁸	8.31 × 10 ⁻¹⁰	4.79 × 10 ⁻¹¹	8.28 × 10 ⁻⁰⁹	1.60 × 10 ⁻¹¹		
Child	n/a	6.55 × 10 ⁻⁰⁶	1.28 × 10 ⁻⁰⁷	5.77 × 10 ⁻⁰⁸	1.35 × 10 ⁻⁰⁶	1.28 × 10 ⁻⁰⁸	2.90×10^{-07}	5.05 × 10 ⁻⁰⁸	1.92 × 10 ⁻⁰⁹	1.11 × 10 ⁻¹⁰	1.77 × 10 ⁻⁰⁸	3.70 × 10 ⁻¹¹		
Teen	n/a	4.86 × 10 ⁻⁰⁶	1.02 × 10 ⁻⁰⁷	4.79 × 10 ⁻⁰⁸	9.96 × 10 ⁻⁰⁷	9.95 × 10 ⁻⁰⁹	2.26×10^{-07}	3.94 × 10 ⁻⁰⁸	1.50 × 10 ⁻⁰⁹	8.62 × 10 ⁻¹¹	1.35 × 10 ⁻⁰⁸	2.84 × 10 ⁻¹¹		
Adult	8.70 × 10 ⁻⁰⁵	4.96 × 10 ⁻⁰⁵	1.03 × 10 ⁻⁰⁶	4.68 × 10 ⁻⁰⁷	1.01 × 10 ⁻⁰⁵	9.64 × 10 ⁻⁰⁸	2.35 × 10 ⁻⁰⁶	3.97 × 10 ⁻⁰⁷	1.56 × 10 ⁻⁰⁸	8.98 × 10 ⁻¹⁰	1.34 × 10 ⁻⁰⁷	2.91 × 10 ⁻¹⁰		

n/a = not applicable or not assessed.

5.2.4.2.3 Radiological Dose

The estimated radiation doses to human receptors for the RFD Case are shown in Table 5-14. The doses shown represent the maximum annual dose over the assessment period with the addition of the Fission Patterson Lake South Property. The dose is presented by radionuclide and exposure pathway, as well as for total dose. The radiation dose is presented as an incremental dose (i.e., only considering Project effects and effects from the Fission Patterson Lake South Property).

The maximum estimated radiation dose for the RFD Case would be 0.2 mSv/yr for the subsistence harvester (one-year-old) who eats Traditional Foods gathered at Patterson Lake South Arm. The main contribution to total dose would be from polonium-210 from eating terrestrial animals in the Traditional Food diet, including moose (meat and organs), beaver, grouse, and mallard.

In the far-future projection, the maximum estimated dose for the RFD Case was predicted to be 0.1 mSv/yr for the future permanent resident residing at the decommissioned and reclaimed Project site well beyond Closure.

5.2.4.2.4 Radon Dose

The camp worker would be exposed to radon through inhalation while at the camp site, located southwest of the mill terrace. The camp worker represents an adult who resides at the camp for 6 months of the year and away from the site for the remaining 6 months of the year. For exposure to radon, it has been conservatively assumed that the camp worker spends 100% of their time indoors when on site.

For the RFD Case, the camp worker would also be exposed to radon from the Fission Patterson Lake South Property. The predicted radon concentration at the camp site, from AERMOD, is 49.3 Bq/m³ (46.4 Bq/m³ based on incremental radon where background radon has been subtracted from the total).

The method to estimate radon dose to the camp worker for the RFD Case follows the equations in Appendix A, Section 2.4.3 and assumptions presented in Section 5.2.4.1.4, Radon Dose. The incremental radon dose to the camp worker during Operations from the RFD Case would be 0.57 mSv/yr.



Ref. 19-2574 APRIL 2024

Table 5-14: Estimated Radiation Doses to Human Receptors – Reasonably Foreseeable Development Case

							Incrementa	al Dose by Pathway R	FD Case (mSv/yr)						
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal) RFD Case	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by Radionuclide	
	Uranium-238	8.63 x 10 ⁻⁰⁴	2.37 x 10 ⁻¹²	2.20 x 10 ⁻⁰⁸	1.67 x 10 ⁻¹³	1.36 x 10 ⁻⁰⁷	1.51 x 10 ⁻⁰³	7.65 x 10 ⁻¹¹	8.15 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	6.22 x 10 ⁻⁰⁵	5.63 x 10 ⁻⁰⁵	7.24 x 10 ⁻⁰⁵	2.56 x 10 ⁻⁰³	
	Uranium-234	1.04 x 10 ⁻⁰³	5.79 x 10 ⁻¹²	2.40 x 10 ⁻⁰⁸	2.92 x 10 ⁻¹⁴	1.48 x 10 ⁻⁰⁷	7.60 x 10 ⁻⁰⁶	8.36 x 10 ⁻¹¹	2.97 x 10 ⁻¹²	0.00 x 10	6.80 x 10 ⁻⁰⁵	6.13 x 10 ⁻⁰⁵	7.88 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰³	
	Thorium-230	4.17 x 10 ⁻⁰³	1.40 x 10 ⁻¹¹	1.44 x 10 ⁻⁰⁶	9.77 x 10 ⁻¹³	7.10 x 10 ⁻⁰⁷	1.09 x 10 ⁻⁰⁵	1.03 x 10 ⁻⁰⁹	2.54 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	1.13 x 10 ⁻⁰³	6.12 x 10 ⁻⁰⁴	6.63 x 10 ⁻⁰³	1.26 x 10 ⁻⁰²	
	Radium-226	1.04 x 10 ⁻⁰³	2.69 x 10 ⁻¹⁰	6.77 x 10 ⁻⁰⁷	3.02 x 10 ⁻¹⁰	9.59 x 10 ⁻⁰⁷	2.48 x 10 ⁻⁰²	1.61 x 10 ⁻⁰⁹	1.93 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	3.71 x 10 ⁻⁰⁴	1.06 x 10 ⁻⁰³	8.00 x 10 ⁻⁰⁴	2.81 x 10 ⁻⁰²	
	Lead-210	3.27 x 10 ⁻⁰⁴	4.23 x 10 ⁻¹¹	2.22 x 10 ⁻⁰⁶	1.95 x 10 ⁻¹²	1.52 x 10 ⁻⁰⁶	3.54 x 10 ⁻⁰⁴	1.96 x 10 ⁻⁰⁸	1.11 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	1.59 x 10 ⁻⁰³	2.29 x 10 ⁻⁰³	6.51 × 10 ⁻⁰³	1.11 x 10 ⁻⁰²	
	Polonium-210	9.82 x 10 ⁻⁰⁴	3.69 x 10 ⁻¹³	1.58 x 10 ⁻⁰⁶	4.74 x 10 ⁻¹⁵	2.69 x 10 ⁻⁰⁶	7.98 x 10 ⁻⁰⁸	3.26 x 10 ⁻⁰⁸	4.48 x 10 ⁻¹²	0.00 x 10 ⁺⁰⁰	9.46 x 10 ⁻⁰³	3.65 x 10 ⁻⁰³	1.62 x 10 ⁻⁰²	3.03 x 10 ⁻⁰²	
Camp Worker	Total by Pathway	8.42 x 10 ⁻⁰³	3.34 x 10 ⁻¹⁰	5.96 x 10 ⁻⁰⁶	3.05 x 10 ⁻¹⁰	6.17 x 10 ⁻⁰⁶	2.67 x 10 ⁻⁰²	5.50 x 10 ⁻⁰⁸	1.95 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.27 x 10 ⁻⁰²	7.73 x 10 ⁻⁰³	3.02 x 10 ⁻⁰²	8.58 x 10 ⁻⁰²	
Patterson Lake orth Arm - West	Total by Fathway	0.42 X 10	3.54 X 10	3.30 X 10	3.03 X 10	0.17 X 10	2.07 X 10	Far-Future	1.55 X 10	0.00 X 10	1.27 X 10	7.75 X 10	5.02 X 10	0.30 X 10	
Basin)	Harrison 220	/-	/-	- /-	- 1-	/-	/-		/-	/s	/s	77./5	/a	/a	
	Uranium-238	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	
	Uranium-234	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	
	Thorium-230	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	
	Radium-226	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	
	Lead-210	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	
	Polonium-210	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	
	Total by Pathway	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	
								RFD Case							
	Uranium-238	1.57 x 10 ⁻⁰⁴	4.30 x 10 ⁻¹³	4.64 x 10 ⁻⁰⁵	3.52 x 10 ⁻¹⁰	2.46 x 10 ⁻⁰⁸	2.73 x 10 ⁻⁰⁴	1.28 x 10 ⁻⁰⁷	1.36 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	1.24 x 10 ⁻⁰⁴	1.12 x 10 ⁻⁰⁴	1.45 x 10 ⁻⁰⁴	8.59 x 10 ⁻⁰⁴	
	Uranium-234	1.89 x 10 ⁻⁰⁴	1.05 x 10 ⁻¹²	5.07 x 10 ⁻⁰⁵	6.16 x 10 ⁻¹¹	2.68 x 10 ⁻⁰⁸	1.38 x 10 ⁻⁰⁶	1.40 x 10 ⁻⁰⁷	4.96 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	1.36 x 10 ⁻⁰⁴	1.22 x 10 ⁻⁰⁴	1.58 x 10 ⁻⁰⁴	6.57 x 10 ⁻⁰⁴	
	Thorium-230	7.56 x 10 ⁻⁰⁴	2.55 x 10 ⁻¹²	1.83 x 10 ⁻⁰³	1.24 x 10 ⁻⁰⁹	1.29 x 10 ⁻⁰⁷	1.97 x 10 ⁻⁰⁶	1.30 x 10 ⁻⁰⁶	3.20 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.26 x 10 ⁻⁰³	1.22 x 10 ⁻⁰³	1.32 x 10 ⁻⁰²	1.93 x 10 ⁻⁰²	
	Radium-226	1.89 x 10 ⁻⁰⁴	4.88 x 10 ⁻¹¹	3.07 x 10 ⁻⁰⁴	1.37 x 10 ⁻⁰⁷	1.74 x 10 ⁻⁰⁷	4.50 x 10 ⁻⁰³	5.92 x 10 ⁻⁰⁷	7.10 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	7.41 x 10 ⁻⁰⁴	2.12 x 10 ⁻⁰³	1.60 x 10 ⁻⁰³	9.53 x 10 ⁻⁰³	
	Lead-210	5.94 x 10 ⁻⁰⁵	7.68 x 10 ⁻¹²	3.67 x 10 ⁻⁰⁴	3.24 x 10 ⁻¹⁰	2.76 x 10 ⁻⁰⁷	6.41 x 10 ⁻⁰⁵	5.06 x 10 ⁻⁰⁶	2.86 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	3.19 x 10 ⁻⁰³	4.57 x 10 ⁻⁰³	1.30 × 10 ⁻⁰²	2.12 x 10 ⁻⁰²	
Subsistence	Polonium-210	1.78 x 10 ⁻⁰⁴	6.70 x 10 ⁻¹⁴	6.25 x 10 ⁻⁰⁴	1.88 x 10 ⁻¹²	4.89 x 10 ⁻⁰⁷	1.45 x 10 ⁻⁰⁸	8.92 x 10 ⁻⁰⁶	1.22 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	1.89 x 10 ⁻⁰²	7.29 x 10 ⁻⁰³	3.24 x 10 ⁻⁰²	5.94 x 10 ⁻⁰²	
Harvester	Total by Pathway	1.53 x 10 ⁻⁰³	6.05 x 10 ⁻¹¹	3.23 x 10 ⁻⁰³	1.39 x 10 ⁻⁰⁷	1.12 x 10 ⁻⁰⁶	4.84 x 10 ⁻⁰³	1.61 x 10 ⁻⁰⁵	7.27 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	2.54 x 10 ⁻⁰²	1.54 x 10 ⁻⁰²	6.05 x 10 ⁻⁰²	1.11 x 10 ⁻⁰¹	
Patterson Lake	Far-Future Far-Future														
South Arm)	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.64 x 10 ⁻⁰⁵	2.00 x 10 ⁻¹⁰	2.59 x 10 ⁻⁰⁹	2.87 x 10 ⁻⁰⁵	2.41 x 10 ⁻⁰⁷	2.57 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	7.35 x 10 ⁻⁰⁵	1.15 x 10 ⁻⁰⁵	3.64 x 10 ⁻⁰⁵	1.79 x 10 ⁻⁰⁴	
	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.88 x 10 ⁻⁰⁵	3.50 x 10 ⁻¹¹	2.82 x 10 ⁻⁰⁹	1.45 x 10 ⁻⁰⁷	2.63 x 10 ⁻⁰⁷	9.34 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	8.03 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁵	3.97 x 10 ⁻⁰⁵	1.62 x 10 ⁻⁰⁴	
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.54 x 10 ⁻⁰⁵	1.04 x 10 ⁻¹¹	7.19 x 10 ⁻⁰⁸	1.10 x 10 ⁻⁰⁶	1.68 x 10 ⁻⁰⁸	4.16 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	2.06 x 10 ⁻⁰⁵	6.16 x 10 ⁻⁰⁶	1.94 x 10 ⁻⁰⁴	2.37 x 10 ⁻⁰⁴	
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	3.14 x 10 ⁻⁰⁵	1.40 x 10 ⁻⁰⁸	1.17 x 10 ⁻⁰⁷	3.03 x 10 ⁻⁰³	1.74 x 10 ⁻⁰⁷	2.08 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	8.37 x 10 ⁻⁰⁵	1.93 x 10 ⁻⁰⁴	4.01 x 10 ⁻⁰⁴	3.77 x 10 ⁻⁰³	
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.02 x 10 ⁻⁰⁶	1.78 x 10 ⁻¹²	1.90 x 10 ⁻¹⁰	4.38 x 10 ⁻⁰⁸	6.81 x 10 ⁻⁰⁸	3.85 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	1.80 x 10 ⁻⁰⁵	4.14 x 10 ⁻⁰⁸	4.02 x 10 ⁻⁰⁴	4.22 x 10 ⁻⁰⁴	
	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	3.55 x 10 ⁻⁰⁶	1.07 x 10 ⁻¹⁴	3.34 x 10 ⁻¹⁰	9.90 x 10 ⁻¹²	1.20 x 10 ⁻⁰⁷	1.65 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	1.13 x 10 ⁻⁰⁴	3.54 x 10 ⁻⁰⁷	3.23 x 10 ⁻⁰⁴	4.40 x 10 ⁻⁰⁴	
	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.08 x 10 ⁻⁰⁴	1.42 x 10 ⁻⁰⁸	1.95 x 10 ⁻⁰⁷	3.06 x 10 ⁻⁰³	8.82 x 10 ⁻⁰⁷	2.34 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	3.89 x 10 ⁻⁰⁴	2.24 x 10 ⁻⁰⁴	1.40 x 10 ⁻⁰³	5.21 x 10 ⁻⁰³	
								RFD Case							
	Uranium-238	1.56 x 10 ⁻⁰⁴	5.59 x 10 ⁻¹³	3.22 x 10 ⁻⁰⁵	8.70 x 10 ⁻¹¹	1.00 x 10 ⁻⁰⁶	3.66 x 10 ⁻⁰⁴	5.19 x 10 ⁻⁰⁶	1.77 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	1.39 x 10 ⁻⁰⁴	6.53 x 10 ⁻⁰⁵	8.39 x 10 ⁻⁰⁴	1.39 x 10 ⁻⁰⁴	
	Uranium-234	1.83 x 10 ⁻⁰⁴	1.37 x 10 ⁻¹²	3.51 x 10 ⁻⁰⁵	1.52 x 10 ⁻¹¹	1.09 x 10 ⁻⁰⁶	1.84 x 10 ⁻⁰⁶	5.65 x 10 ⁻⁰⁶	6.45 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	1.50 x 10 ⁻⁰⁴	7.07 x 10 ⁻⁰⁵	5.27 x 10 ⁻⁰⁴	1.50 x 10 ⁻⁰⁴	
61	Thorium-230	5.81 x 10 ⁻⁰⁴	3.31 x 10 ⁻¹²	9.31 x 10 ⁻⁰⁴	3.07 x 10 ⁻¹⁰	3.83 x 10 ⁻⁰⁶	2.24 x 10 ⁻⁰⁶	3.86 x 10 ⁻⁰⁵	4.17 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	6.69 x 10 ⁻⁰⁴	6.39 x 10 ⁻⁰³	9.59 x 10 ⁻⁰³	6.69 x 10 ⁻⁰⁴	
Subsistence larvester One-	Radium-226	1.83 x 10 ⁻⁰⁴	6.34 x 10 ⁻¹¹	2.74 x 10 ⁻⁰⁴	3.39 x 10 ⁻⁰⁸	9.10 x 10 ⁻⁰⁶	6.03 x 10 ⁻⁰³	3.10 x 10 ⁻⁰⁵	9.26 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	2.40 x 10 ⁻⁰³	7.12 x 10 ⁻⁰⁴	1.03 x 10 ⁻⁰²	2.40 x 10 ⁻⁰³	
Year-Old	Lead-210	6.14 x 10 ⁻⁰⁵	9.96 x 10 ⁻¹²	4.99 x 10 ⁻⁰⁴	6.57 x 10 ⁻¹¹	2.20 x 10 ⁻⁰⁵	8.58 x 10 ⁻⁰⁵	4.02 x 10 ⁻⁰⁴	3.72 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	7.46 x 10 ⁻⁰³	1.60 x 10 ⁻⁰²	2.82 x 10 ⁻⁰²	7.46 x 10 ⁻⁰³	
Patterson Lake	Polonium-210	1.83 x 10 ⁻⁰⁴	8.71 x 10 ⁻¹⁴	1.19 x 10 ⁻⁰³	4.65 x 10 ⁻¹³	5.46 x 10 ⁻⁰⁵	1.89 x 10 ⁻⁰⁸	9.98 x 10 ⁻⁰⁴	1.59 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	1.73 x 10 ⁻⁰²	3.65 x 10 ⁻⁰²	8.68 x 10 ⁻⁰²	1.73 x 10 ⁻⁰²	
South Arm)	Total by Pathway	1.35 x 10 ⁻⁰³	7.87 x 10 ⁻¹¹	2.97 x 10 ⁻⁰³	3.44 x 10 ⁻⁰⁸	9.16 x 10 ⁻⁰⁵	6.48 x 10 ⁻⁰³	1.48 x 10 ⁻⁰³	9.48 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	2.81 x 10 ⁻⁰²	5.97 x 10 ⁻⁰²	1.36 x 10 ⁻⁰¹	2.81 x 10 ⁻⁰²	
	2 2 2 7 7 2 2 2 7 2 2 7 2 7 2 7 2 7 2 7					22.%		Far-Future				2.2			
	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.83 x 10 ⁻⁰⁵	4.95 x 10 ⁻¹¹	1.05 x 10 ⁻⁰⁷	3.84 x 10 ⁻⁰⁵	9.78 x 10 ⁻⁰⁶	3.34 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	1.41 x 10 ⁻⁰⁵	1.67 x 10 ⁻⁰⁵	1.44 x 10 ⁻⁰⁴	1.41 x 10 ⁻⁰⁵	
		0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.83 x 10 ° 5 1.99 x 10 -05	8.67 x 10 ⁻¹²	1.05 x 10 ° 7	1.94 x 10 ⁻⁰⁷		1.21 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	1.41 x 10 ° 5 1.53 x 10 ° 0 5	1.81 x 10 ⁻⁰⁵	1.44 x 10 ° · · · · · · · · · · · · · · · · · ·	1.41 x 10 ° s	
	Uranium-234	U.UU X 10	U.UU X 103	ו א פפ.ו _ב	0.07 X 10 .2	1.14 X 10 -	1.54 X 10	1.06 x 10 ⁻⁰⁵	1.21 X 10	0.00 x 10	1.35 X 10	1.01 X 10	1.11 X 1U -	1.33 X 10 °	

							Incrementa	al Dose by Pathway	RFD Case (mSv/yr)						
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by Radionuclide	
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	7.83 x 10 ⁻⁰⁶	2.58 x 10 ⁻¹²	2.14 x 10 ⁻⁰⁶	1.25 x 10 ⁻⁰⁶	5.00 x 10 ⁻⁰⁷	5.40 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	5.53 x 10 ⁻⁰⁶	9.35 x 10 ⁻⁰⁵	1.20 x 10 ⁻⁰⁴	5.53 x 10 ⁻⁰⁶	
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.80 x 10 ⁻⁰⁵	3.47 x 10 ⁻⁰⁹	6.13 x 10 ⁻⁰⁶	4.06 x 10 ⁻⁰³	9.08 x 10 ⁻⁰⁶	2.71 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	3.05 x 10 ⁻⁰⁴	1.79 x 10 ⁻⁰⁴	4.68 x 10 ⁻⁰³	3.05 x 10 ⁻⁰⁴	
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.75 x 10 ⁻⁰⁶	3.62 x 10 ⁻¹³	1.50 x 10 ⁻⁰⁸	5.87 x 10 ⁻⁰⁸	5.42 x 10 ⁻⁰⁶	5.01 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	1.01 x 10 ⁻⁰⁷	5.02 x 10 ⁻⁰⁴	5.32 x 10 ⁻⁰⁴	1.01 x 10 ⁻⁰⁷	
	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	6.79 x 10 ⁻⁰⁶	2.64 x 10 ⁻¹⁵	3.73 x 10 ⁻⁰⁸	1.29 x 10 ⁻¹¹	1.35 x 10 ⁻⁰⁵	2.15 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	1.20 x 10 ⁻⁰⁶	3.77 x 10 ⁻⁰⁴	5.82 x 10 ⁻⁰⁴	1.20 x 10 ⁻⁰⁶	
	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	8.37 x 10 ⁻⁰⁵	3.53 x 10 ⁻⁰⁹	8.54 x 10 ⁻⁰⁶	4.10 x 10 ⁻⁰³	4.89 x 10 ⁻⁰⁵	3.05 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	3.41 x 10 ⁻⁰⁴	1.19 x 10 ⁻⁰³	6.17 x 10 ⁻⁰³	3.41 x 10 ⁻⁰⁴	
		<u>, </u>						RFD Case				·			
	Uranium-238	8.56 x 10 ⁻⁰⁵	2.35 x 10 ⁻¹³	1.61 x 10 ⁻⁰⁵	1.22 x 10 ⁻¹⁰	1.35 x 10 ⁻⁰⁸	1.49 x 10 ⁻⁰⁴	4.72 x 10 ⁻⁰⁸	5.03 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	6.15 x 10 ⁻⁰⁵	5.72 x 10 ⁻⁰⁵	4.14 x 10 ⁻⁰⁴	6.15 x 10 ⁻⁰⁵	
	Uranium-234	1.03 x 10 ⁻⁰⁴	5.75 x 10 ⁻¹³	1.76 x 10 ⁻⁰⁵	2.14 x 10 ⁻¹¹	1.47 x 10 ⁻⁰⁸	7.53 x 10 ⁻⁰⁷	5.16 x 10 ⁻⁰⁸	1.83 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	6.70 x 10 ⁻⁰⁵	6.23 x 10 ⁻⁰⁵	2.98 x 10 ⁻⁰⁴	6.70 x 10 ⁻⁰⁵	
Subsistence	Thorium-230	4.13 x 10 ⁻⁰⁴	1.39 x 10 ⁻¹²	8.23 x 10 ⁻⁰⁴	5.58 x 10 ⁻¹⁰	7.04 x 10 ⁻⁰⁸	1.08 x 10 ⁻⁰⁶	6.90 x 10 ⁻⁰⁷	1.70 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	6.69 x 10 ⁻⁰⁴	3.87 x 10 ⁻⁰³	6.80 x 10 ⁻⁰³	6.69 x 10 ⁻⁰⁴	
	Radium-226	1.03 x 10 ⁻⁰⁴	2.67 x 10 ⁻¹¹	1.18 x 10 ⁻⁰⁴	5.27 x 10 ⁻⁰⁸	9.51 x 10 ⁻⁰⁸	2.46 x 10 ⁻⁰³	2.57 x 10 ⁻⁰⁷	3.08 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	1.16 x 10 ⁻⁰³	4.11 x 10 ⁻⁰⁴	4.57 x 10 ⁻⁰³	1.16 x 10 ⁻⁰³	
	Lead-210	3.25 x 10 ⁻⁰⁵	4.20 x 10 ⁻¹²	5.33 x 10 ⁻⁰⁵	4.70 x 10 ⁻¹¹	1.51 x 10 ⁻⁰⁷	3.51 x 10 ⁻⁰⁵	7.33 x 10 ⁻⁰⁷	4.14 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.50 x 10 ⁻⁰³	1.75 x 10 ⁻⁰³	4.83 x 10 ⁻⁰³	2.50 x 10 ⁻⁰³	
	Polonium-210	9.74 x 10 ⁻⁰⁵	3.66 x 10 ⁻¹⁴	9.19 x 10 ⁻⁰⁵	2.76 x 10 ⁻¹³	2.67 x 10 ⁻⁰⁷	7.91 x 10 ⁻⁰⁹	1.29 x 10 ⁻⁰⁶	1.77 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	3.99 x 10 ⁻⁰³	7.70 x 10 ⁻⁰³	1.47 x 10 ⁻⁰²	3.99 x 10 ⁻⁰³	
	Total by Pathway	8.35 x 10 ⁻⁰⁴	3.31 x 10 ⁻¹¹	1.12 x 10 ⁻⁰³	5.34 x 10 ⁻⁰⁸	6.12 x 10 ⁻⁰⁷	2.65 x 10 ⁻⁰³	3.07 x 10 ⁻⁰⁶	3.14 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	8.44 x 10 ⁻⁰³	1.38 x 10 ⁻⁰²	3.16 x 10 ⁻⁰²	8.44 x 10 ⁻⁰³	
Harvester (Beet Lake)	Far-Future Convious														
Luncy	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	9.81 x 10 ⁻⁰⁶	7.44 x 10 ⁻¹¹	1.42 x 10 ⁻⁰⁹	1.57 x 10 ⁻⁰⁵	8.94 x 10 ⁻⁰⁸	9.54 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	6.30 x 10 ⁻⁰⁶	1.25 x 10 ⁻⁰⁵	7.26 x 10 ⁻⁰⁵	6.30 x 10 ⁻⁰⁶	
	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.07 x 10 ⁻⁰⁵	1.30 x 10 ⁻¹¹	1.54 x 10 ⁻⁰⁹	7.92 x 10 ⁻⁰⁸	9.78 x 10 ⁻⁰⁸	3.47 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	6.86 x 10 ⁻⁰⁶	1.36 x 10 ⁻⁰⁵	6.12 x 10 ⁻⁰⁵	6.86 x 10 ⁻⁰⁶	
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	8.19 x 10 ⁻⁰⁶	5.55 x 10 ⁻¹²	3.93 x 10 ⁻⁰⁸	6.02 x 10 ⁻⁰⁷	8.93 x 10 ⁻⁰⁹	2.21 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	3.37 x 10 ⁻⁰⁶	4.93 x 10 ⁻⁰⁵	7.25 x 10 ⁻⁰⁵	3.37 x 10 ⁻⁰⁶	
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.35 x 10 ⁻⁰⁵	6.02 x 10 ⁻⁰⁹	6.41 x 10 ⁻⁰⁸	1.66 x 10 ⁻⁰³	7.47 x 10 ⁻⁰⁸	8.95 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	1.06 x 10 ⁻⁰⁴	1.03 x 10 ⁻⁰⁴	1.93 x 10 ⁻⁰³	1.06 x 10 ⁻⁰⁴	
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.79 x 10 ⁻⁰⁷	2.46 x 10 ⁻¹³	1.03 x 10 ⁻¹⁰	2.37 x 10 ⁻⁰⁸	9.37 x 10 ⁻⁰⁹	5.30 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	2.26 x 10 ⁻⁰⁸	4.16 x 10 ⁻⁰⁶	6.98 x 10 ⁻⁰⁶	2.26 x 10 ⁻⁰⁸	
	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	4.75 x 10 ⁻⁰⁷	1.42 x 10 ⁻¹⁵	1.83 x 10 ⁻¹⁰	5.41 x 10 ⁻¹²	1.65 x 10 ⁻⁰⁸	2.27 x 10 ⁻¹²	0.00 x 10 ⁺⁰⁰	1.94 x 10 ⁻⁰⁷	3.28 x 10 ⁻⁰⁵	4.85 x 10 ⁻⁰⁵	1.94 x 10 ⁻⁰⁷	
	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	4.30 x 10 ⁻⁰⁵	6.11 x 10 ⁻⁰⁹	1.07 x 10 ⁻⁰⁷	1.68 x 10 ⁻⁰³	2.97 x 10 ⁻⁰⁷	9.91 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	1.22 x 10 ⁻⁰⁴	2.15 x 10 ⁻⁰⁴	2.19 x 10 ⁻⁰³	1.22 x 10 ⁻⁰⁴	
								RFD Case							
	Uranium-238	8.53 x 10 ⁻⁰⁵	3.05 x 10 ⁻¹³	1.12 x 10 ⁻⁰⁵	3.02 x 10 ⁻¹¹	5.48 x 10 ⁻⁰⁷	2.00 x 10 ⁻⁰⁴	1.92 x 10 ⁻⁰⁶	6.54 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	2.55 x 10 ⁻⁰⁵	7.59 x 10 ⁻⁰⁵	2.54 x 10 ⁻⁰⁵	4.26 x 10 ⁻⁰⁴	
	Uranium-234	9.99 x 10 ⁻⁰⁵	7.47 x 10 ⁻¹³	1.22 x 10 ⁻⁰⁵	5.30 x 10 ⁻¹²	5.94 x 10 ⁻⁰⁷	1.01 x 10 ⁻⁰⁶	2.09 x 10 ⁻⁰⁶	2.38 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	2.77 x 10 ⁻⁰⁵	8.22 x 10 ⁻⁰⁵	2.75 x 10 ⁻⁰⁵	2.53 x 10 ⁻⁰⁴	
	Thorium-230	3.18 x 10 ⁻⁰⁴	1.81 x 10 ⁻¹²	4.19 x 10 ⁻⁰⁴	1.38 x 10 ⁻¹⁰	2.10 x 10 ⁻⁰⁶	1.23 x 10 ⁻⁰⁶	2.05 x 10 ⁻⁰⁵	2.22 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	4.41 x 10 ⁻⁰⁴	3.66 x 10 ⁻⁰⁴	1.86 × 10 ⁻⁰³	3.43 x 10 ⁻⁰³	
	Radium-226	9.99 x 10 ⁻⁰⁵	3.46 x 10 ⁻¹¹	1.05 x 10 ⁻⁰⁴	1.30 x 10 ⁻⁰⁸	4.97 x 10 ⁻⁰⁶	3.30 x 10 ⁻⁰³	1.34 x 10 ⁻⁰⁵	4.02 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	2.15 x 10 ⁻⁰⁴	1.31 x 10 ⁻⁰³	1.58 x 10 ⁻⁰⁴	5.25 x 10 ⁻⁰³	
	Lead-210	3.36 x 10 ⁻⁰⁵	5.45 x 10 ⁻¹²	7.25 x 10 ⁻⁰⁵	9.53 x 10 ⁻¹²	1.20 x 10 ⁻⁰⁵	4.69 x 10 ⁻⁰⁵	5.83 x 10 ⁻⁰⁵	5.39 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	5.33 x 10 ⁻⁰⁴	4.08 x 10 ⁻⁰³	2.01 × 10 ⁻⁰³	6.85 x 10 ⁻⁰³	
Subsistence	Polonium-210	9.99 x 10 ⁻⁰⁵	4.76 x 10 ⁻¹⁴	1.76 x 10 ⁻⁰⁴	6.84 x 10 ⁻¹⁴	2.99 x 10 ⁻⁰⁵	1.03 x 10 ⁻⁰⁸	1.45 x 10 ⁻⁰⁴	2.31 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	4.51 x 10 ⁻⁰³	9.43 x 10 ⁻⁰³	8.38 x 10 ⁻⁰³	2.28 x 10 ⁻⁰²	
Harvester One-	Total by Pathway	7.36 x 10 ⁻⁰⁴	4.30 x 10 ⁻¹¹	7.96 x 10 ⁻⁰⁴	1.32 x 10 ⁻⁰⁸	5.01 x 10 ⁻⁰⁵	3.54 x 10 ⁻⁰³	2.41 x 10 ⁻⁰⁴	4.09 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	5.75 x 10 ⁻⁰³	1.54 x 10 ⁻⁰²	1.25 x 10 ⁻⁰²	3.90 x 10 ⁻⁰²	
Year-Old (Beet								Far-Future							
Lake)	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	6.81 x 10 ⁻⁰⁶	1.84 x 10 ⁻¹¹	5.76 x 10 ⁻⁰⁸	2.10 x 10 ⁻⁰⁵	3.64 x 10 ⁻⁰⁶	1.24 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	1.61 x 10 ⁻⁰⁵	7.73 x 10 ⁻⁰⁶	5.64 x 10 ⁻⁰⁶	6.23 x 10 ⁻⁰⁵	
	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	7.41 x 10 ⁻⁰⁶	3.22 x 10 ⁻¹²	6.24 x 10 ⁻⁰⁸	1.06 x 10 ⁻⁰⁷	3.96 x 10 ⁻⁰⁶	4.51 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	1.75 x 10 ⁻⁰⁵	8.38 x 10 ⁻⁰⁶	6.12 x 10 ⁻⁰⁶	4.36 x 10 ⁻⁰⁵	
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	4.17 x 10 ⁻⁰⁶	1.37 x 10 ⁻¹²	1.17 x 10 ⁻⁰⁶	6.85 x 10 ⁻⁰⁷	2.66 x 10 ⁻⁰⁷	2.87 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	4.74 x 10 ⁻⁰⁶	3.03 x 10 ⁻⁰⁶	2.37 x 10 ⁻⁰⁵	3.77 x 10 ⁻⁰⁵	
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.20 x 10 ⁻⁰⁵	1.49 x 10 ⁻⁰⁹	3.35 x 10 ⁻⁰⁶	2.22 x 10 ⁻⁰³	3.90 x 10 ⁻⁰⁶	1.17 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	2.73 x 10 ⁻⁰⁵	1.67 x 10 ⁻⁰⁴	3.97 x 10 ⁻⁰⁵	2.49 x 10 ⁻⁰³	
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	3.79 x 10 ⁻⁰⁷	4.98 x 10 ⁻¹⁴	8.15 x 10 ⁻⁰⁹	3.21 x 10 ⁻⁰⁸	7.46 x 10 ⁻⁰⁷	6.89 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	2.86 x 10 ⁻⁰⁶	5.49 x 10 ⁻⁰⁸	4.94 × 10 ⁻⁰⁶	9.02 x 10 ⁻⁰⁶	
	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	9.06 x 10 ⁻⁰⁷	3.53 x 10 ⁻¹⁶	2.04 x 10 ⁻⁰⁸	7.08 x 10 ⁻¹²	1.85 x 10 ⁻⁰⁶	2.95 x 10 ⁻¹²	0.00 x 10 ⁺⁰⁰	2.45 x 10 ⁻⁰⁵	6.54 x 10 ⁻⁰⁷	4.16 x 10 ⁻⁰⁵	6.95 x 10 ⁻⁰⁵	
	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	3.17 x 10 ⁻⁰⁵	1.51 x 10 ⁻⁰⁹	4.67 x 10 ⁻⁰⁶	2.24 x 10 ⁻⁰³	1.44 x 10 ⁻⁰⁵	1.29 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	9.31 x 10 ⁻⁰⁵	1.87 x 10 ⁻⁰⁴	1.22 x 10 ⁻⁰⁴	2.71 x 10 ⁻⁰³	
	, -,							RFD Case							
						1.0	7 12 10-06	4.32 x 10 ⁻⁰⁹	4.61 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	4.01 x 10 ⁻⁰⁶	2.93 x 10 ⁻⁰⁶	2.88 x 10 ⁻⁰⁶	2.26 x 10 ⁻⁰⁵	
	Uranium-238	4.08 x 10 ⁻⁰⁶	1.12 x 10 ⁻¹⁴	1.50 x 10 ⁻⁰⁶	1.14 x 10 ⁻¹¹	6.42 x 10 ⁻¹⁰	/. 2 X) **								
	Uranium-238 Uranium-234	4.08 x 10 ⁻⁰⁶ 4.92 x 10 ⁻⁰⁶	1.12 x 10 ⁻¹⁴ 2.74 x 10 ⁻¹⁴	1.50 x 10 ⁻⁰⁶	1.14 x 10 ⁻¹¹ 1.99 x 10 ⁻¹²	6.42 x 10 ⁻¹⁰	7.12 x 10 ⁻⁰⁶ 3.59 x 10 ⁻⁰⁸			+		†	+	1.73 x 10 ⁻⁰⁵	
Subsistence	Uranium-234	4.92 x 10 ⁻⁰⁶	2.74 x 10 ⁻¹⁴	1.64 x 10 ⁻⁰⁶	1.99 x 10 ⁻¹²	6.99 x 10 ⁻¹⁰	3.59 x 10 ⁻⁰⁸	4.72 x 10 ⁻⁰⁹	1.68 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	4.38 x 10 ⁻⁰⁶	3.19 x 10 ⁻⁰⁶	3.13 x 10 ⁻⁰⁶	1.73 x 10 ⁻⁰⁵ 5 95 x 10 ⁻⁰⁴	
larvester (Lloyd	Uranium-234 Thorium-230	4.92 x 10 ⁻⁰⁶ 1.97 x 10 ⁻⁰⁵	2.74 x 10 ⁻¹⁴ 6.64 x 10 ⁻¹⁴	1.64 x 10 ⁻⁰⁶ 8.37 x 10 ⁻⁰⁵	1.99 x 10 ⁻¹² 5.67 x 10 ⁻¹¹	6.99 x 10 ⁻¹⁰ 3.35 x 10 ⁻⁰⁹	3.59 x 10 ⁻⁰⁸ 5.14 x 10 ⁻⁰⁸	4.72 x 10 ⁻⁰⁹ 7.12 x 10 ⁻⁰⁸	1.68 x 10 ⁻¹⁰ 1.76 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰ 0.00 x 10 ⁺⁰⁰	4.38 x 10 ⁻⁰⁶ 1.04 x 10 ⁻⁰⁴	3.19 x 10 ⁻⁰⁶ 3.19 x 10 ⁻⁰⁵	3.13 x 10 ⁻⁰⁶ 3.56 × 10 ⁻⁰⁴	5.95 x 10 ⁻⁰⁴	
	Uranium-234 Thorium-230 Radium-226	4.92 x 10 ⁻⁰⁶ 1.97 x 10 ⁻⁰⁵ 4.92 x 10 ⁻⁰⁶	2.74 x 10 ⁻¹⁴ 6.64 x 10 ⁻¹⁴ 1.27 x 10 ⁻¹²	1.64 x 10 ⁻⁰⁶ 8.37 x 10 ⁻⁰⁵ 1.14 x 10 ⁻⁰⁵	1.99 x 10 ⁻¹² 5.67 x 10 ⁻¹¹ 5.09 x 10 ⁻⁰⁹	6.99 x 10 ⁻¹⁰ 3.35 x 10 ⁻⁰⁹ 4.53 x 10 ⁻⁰⁹	3.59 x 10 ⁻⁰⁸ 5.14 x 10 ⁻⁰⁸ 1.17 x 10 ⁻⁰⁴	4.72 x 10 ⁻⁰⁹ 7.12 x 10 ⁻⁰⁸ 2.49 x 10 ⁻⁰⁸	1.68 x 10 ⁻¹⁰ 1.76 x 10 ⁻⁰⁹ 2.98 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	4.38 x 10 ⁻⁰⁶ 1.04 x 10 ⁻⁰⁴ 2.74 x 10 ⁻⁰⁵	3.19 x 10 ⁻⁰⁶ 3.19 x 10 ⁻⁰⁵ 5.52 x 10 ⁻⁰⁵	3.13 x 10 ⁻⁰⁶ 3.56 × 10 ⁻⁰⁴ 2.09 x 10 ⁻⁰⁵	5.95 x 10 ⁻⁰⁴ 2.40 x 10 ⁻⁰⁴	
Harvester (Lloyd	Uranium-234 Thorium-230	4.92 x 10 ⁻⁰⁶ 1.97 x 10 ⁻⁰⁵	2.74 x 10 ⁻¹⁴ 6.64 x 10 ⁻¹⁴	1.64 x 10 ⁻⁰⁶ 8.37 x 10 ⁻⁰⁵	1.99 x 10 ⁻¹² 5.67 x 10 ⁻¹¹	6.99 x 10 ⁻¹⁰ 3.35 x 10 ⁻⁰⁹	3.59 x 10 ⁻⁰⁸ 5.14 x 10 ⁻⁰⁸	4.72 x 10 ⁻⁰⁹ 7.12 x 10 ⁻⁰⁸	1.68 x 10 ⁻¹⁰ 1.76 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰ 0.00 x 10 ⁺⁰⁰	4.38 x 10 ⁻⁰⁶ 1.04 x 10 ⁻⁰⁴	3.19 x 10 ⁻⁰⁶ 3.19 x 10 ⁻⁰⁵	3.13 x 10 ⁻⁰⁶ 3.56 × 10 ⁻⁰⁴	5.95 x 10 ⁻⁰⁴	

							Increment	al Dose by Pathway R	FD Case (mSv/yr)						
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by Radionuclide	
	Harrison, 220	0.00 10+00	0.00 x 10 ⁺⁰⁰	8.88 x 10 ⁻⁰⁷	C 74 10-12	C 7F 10-11	7.40 10-07	Far-Future	0.64 10-08	0.00 10+00	2.47 10-06	2.00 10-07	0.00 10-07	F 20 · · 10-06	
	Uranium-238 Uranium-234	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	0.00 x 10 ⁺⁰⁰	9.71 x 10 ⁻⁰⁷	6.74 x 10 ⁻¹²	6.75 x 10 ⁻¹¹ 7.35 x 10 ⁻¹¹	7.49 x 10 ⁻⁰⁷ 3.77 x 10 ⁻⁰⁹	8.10 x 10 ⁻⁰⁹ 8.85 x 10 ⁻⁰⁹	8.64 x 10 ⁻⁰⁸ 3.14 x 10 ⁻¹⁰	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	2.47 x 10 ⁻⁰⁶ 2.70 x 10 ⁻⁰⁶	3.00 x 10 ⁻⁰⁷ 3.27 x 10 ⁻⁰⁷	9.68 x 10 ⁻⁰⁷	5.39 x 10 ⁻⁰⁶ 4.98 x 10 ⁻⁰⁶	
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	8.40 x 10 ⁻⁰⁷	5.69 x 10 ⁻¹³	1.87 x 10 ⁻⁰⁹	2.87 x 10 ⁻⁰⁸	9.16 x 10 ⁻¹⁰	2.26 x 10 ⁻¹¹	0.00 x 10	1.12 x 10 ⁻⁰⁶	1.61 x 10 ⁻⁰⁷	4.08 x 10 ⁻⁰⁶	6.24 x 10 ⁻⁰⁶	
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.29 x 10 ⁻⁰⁶	5.76 x 10 ⁻¹⁰	3.05 x 10 ⁻⁰⁹	7.91 x 10 ⁻⁰⁵	7.15 x 10 ⁻⁰⁹	8.56 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	3.45 x 10 ⁻⁰⁶	5.04 x 10 ⁻⁰⁶	5.26 x 10 ⁻⁰⁶	9.50 x 10 ⁻⁰⁵	
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.04 x 10 ⁻⁰⁸	1.81 x 10 ⁻¹⁴	4.55 × 10 ⁻¹²	9.31 × 10 ⁻¹⁰	6.88 x 10 ⁻¹⁰	3.89 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	1.83 x 10 ⁻⁰⁷	1.16 x 10 ⁻⁰⁹	3.09 × 10 ⁻⁰⁷	5.15 x 10 ⁻⁰⁷	
	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	8.19 x 10 ⁻¹²	2.70 x 10 ⁻¹³	1.14 x 10 ⁻⁰⁹	1.56 x 10 ⁻¹³	0.00 x 10 ⁺⁰⁰	5.96 x 10 ⁻⁰⁸	9.31 x 10 ⁻⁰⁹	2.18 x 10 ⁻⁰⁶	2.25 x 10 ⁻⁰⁶	
	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	4.01 x 10 ⁻⁰⁶	5.84 x 10 ⁻¹⁰	5.08 x 10 ⁻⁰⁹	7.99 x 10 ⁻⁰⁵	2.68 x 10 ⁻⁰⁸	9.43 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	9.99 x 10 ⁻⁰⁶	5.84 x 10 ⁻⁰⁶	1.37 x 10 ⁻⁰⁵	1.14 x 10 ⁻⁰⁴	
								RFD Case							
	Uranium-238	4.07 x 10 ⁻⁰⁶	1.46 x 10 ⁻¹⁴	1.04 x 10 ⁻⁰⁶	2.81 x 10 ⁻¹²	2.61 x 10 ⁻⁰⁸	9.53 x 10 ⁻⁰⁶	1.76 x 10 ⁻⁰⁷	5.99 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.37 x 10 ⁻⁰⁶	3.62 x 10 ⁻⁰⁶	1.28 x 10 ⁻⁰⁶	2.22 x 10 ⁻⁰⁵	
	Uranium-234	4.76 x 10 ⁻⁰⁶	3.56 x 10 ⁻¹⁴	1.13 x 10 ⁻⁰⁶	4.93 x 10 ⁻¹³	2.83 x 10 ⁻⁰⁸	4.81 x 10 ⁻⁰⁸	1.91 x 10 ⁻⁰⁷	2.18 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	2.57 x 10 ⁻⁰⁶	3.92 x 10 ⁻⁰⁶	1.38 x 10 ⁻⁰⁶	1.40 x 10 ⁻⁰⁵	
	Thorium-230	1.51 x 10 ⁻⁰⁵	8.63 x 10 ⁻¹⁴	4.26 x 10 ⁻⁰⁵	1.40 x 10 ⁻¹¹	9.99 x 10 ⁻⁰⁸	5.84 x 10 ⁻⁰⁸	2.12 x 10 ⁻⁰⁶	2.29 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	4.51 x 10 ⁻⁰⁵	1.74 x 10 ⁻⁰⁵	1.71 × 10 ⁻⁰⁴	2.94 x 10 ⁻⁰⁴	
	Radium-226	4.76 x 10 ⁻⁰⁶	1.65 x 10 ⁻¹²	1.02 x 10 ⁻⁰⁵	1.26 x 10 ⁻⁰⁹	2.37 x 10 ⁻⁰⁷	1.57 x 10 ⁻⁰⁴	1.30 x 10 ⁻⁰⁶	3.89 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	2.08 x 10 ⁻⁰⁵	6.27 x 10 ⁻⁰⁵	8.11 x 10 ⁻⁰⁶	2.69 x 10 ⁻⁰⁴	
	Lead-210	1.60 x 10 ⁻⁰⁶	2.60 x 10 ⁻¹³	3.95 x 10 ⁻⁰⁶	5.19 x 10 ⁻¹³	5.73 x 10 ⁻⁰⁷	2.24 x 10 ⁻⁰⁶	3.01 x 10 ⁻⁰⁶	2.78 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	2.90 x 10 ⁻⁰⁵	1.94 x 10 ⁻⁰⁴	1.01 × 10 ⁻⁰⁴	3.36 x 10 ⁻⁰⁴	
Subsistence	Polonium-210	4.76 x 10 ⁻⁰⁶	2.27 x 10 ⁻¹⁵	9.48 x 10 ⁻⁰⁶	3.69 x 10 ⁻¹⁵	1.42 x 10 ⁻⁰⁶	4.93 x 10 ⁻¹⁰	7.46 x 10 ⁻⁰⁶	1.19 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	2.43 x 10 ⁻⁰⁴	4.50 x 10 ⁻⁰⁴	4.12 x 10 ⁻⁰⁴	1.13 x 10 ⁻⁰³	
Harvester One-	Total by Pathway	3.51 x 10 ⁻⁰⁵	2.05 x 10 ⁻¹²	6.84 x 10 ⁻⁰⁵	1.28 x 10 ⁻⁰⁹	2.39 x 10 ⁻⁰⁶	1.69 x 10 ⁻⁰⁴	1.43 x 10 ⁻⁰⁵	3.95 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	3.43 x 10 ⁻⁰⁴	7.32 x 10 ⁻⁰⁴	6.96 x 10 ⁻⁰⁴	2.06 x 10 ⁻⁰³	
Year-Old (Lloyd Lake)	Far-Future Converse														
	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	6.17 x 10 ⁻⁰⁷	1.67 x 10 ⁻¹²	2.75 x 10 ⁻⁰⁹	1.00 x 10 ⁻⁰⁶	3.29 x 10 ⁻⁰⁷	1.12 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.46 x 10 ⁻⁰⁶	3.69 x 10 ⁻⁰⁷	4.03 x 10 ⁻⁰⁷	4.29 x 10 ⁻⁰⁶	
	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	6.71 x 10 ⁻⁰⁷	2.92 x 10 ⁻¹³	2.97 x 10 ⁻⁰⁹	5.05 x 10 ⁻⁰⁹	3.58 x 10 ⁻⁰⁷	4.09 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	1.59 x 10 ⁻⁰⁶	3.99 x 10 ⁻⁰⁷	4.38 x 10 ⁻⁰⁷	3.46 x 10 ⁻⁰⁶	
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	4.28 x 10 ⁻⁰⁷	1.41 x 10 ⁻¹³	5.58 x 10 ⁻⁰⁸	3.26 x 10 ⁻⁰⁸	2.73 x 10 ⁻⁰⁸	2.94 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	4.86 x 10 ⁻⁰⁷	1.44 x 10 ⁻⁰⁷	1.96 x 10 ⁻⁰⁶	3.13 x 10 ⁻⁰⁶	
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.15 x 10 ⁻⁰⁶	1.43 x 10 ⁻¹⁰	1.60 x 10 ⁻⁰⁷	1.06 x 10 ⁻⁰⁴	3.74 x 10 ⁻⁰⁷	1.12 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	2.61 x 10 ⁻⁰⁶	7.94 x 10 ⁻⁰⁶	2.05 x 10 ⁻⁰⁶	1.21 x 10 ⁻⁰⁴	
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.79 x 10 ⁻⁰⁸	3.66 x 10 ⁻¹⁵	3.49 × 10 ⁻¹⁰	1.86 x 10 ⁻⁰⁹	5.49 x 10 ⁻⁰⁸	5.07 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	2.11 x 10 ⁻⁰⁷	2.79 × 10 ⁻⁰⁹	3.65 × 10 ⁻⁰⁷	6.64 x 10 ⁻⁰⁷	
	Polonium-210	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	0.00 x 10 ⁺⁰⁰ 0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰ 2.90 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	9.31 x 10 ⁻¹⁰ 2.23 x 10 ⁻⁰⁷	3.41 x 10 ⁻¹³ 1.07 x 10 ⁻⁰⁴	1.29 x 10 ⁻⁰⁷ 1.27 x 10 ⁻⁰⁶	2.03 x 10 ⁻¹³ 1.23 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	9.69 x 10 ⁻⁰⁸ 6.45 x 10 ⁻⁰⁶	3.17 x 10 ⁻⁰⁸ 8.89 x 10 ⁻⁰⁶	2.80 x 10 ⁻⁰⁶ 8.02 x 10 ⁻⁰⁶	3.06 x 10 ⁻⁰⁶ 1.36 x 10 ⁻⁰⁴	
	Total by Pathway	0.00 x 10	0.00 x 10	2.90 X 10 **	1.45 X 10	2.23 X 10 °	1.07 x 10	RFD Case	1.23 X 10 **	0.00 x 10	0.43 X 10 **	6.69 X 10	6.02 X 10	1.50 X 10 °	
	Uranium-238	9.39 x 10 ⁻⁰⁵	2.58 x 10 ⁻¹³	2.78 x 10 ⁻⁰⁵	2.11 x 10 ⁻¹⁰	1.48 x 10 ⁻⁰⁸	1.64 x 10 ⁻⁰⁴	7.66 x 10 ⁻⁰⁸	8.17 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	2.36 x 10 ⁻⁰⁵	5.34 x 10 ⁻⁰⁵	2.71 x 10 ⁻⁰⁵	3.91 x 10 ⁻⁰⁴	
	Uranium-234	1.13 x 10 ⁻⁰⁴	6.30 x 10 ⁻¹³	3.04 x 10 ⁻⁰⁵	3.69 x 10 ⁻¹¹	1.48 x 10 1.61 x 10 ⁻⁰⁸	8.27 x 10 ⁻⁰⁷	8.38 x 10 ⁻⁰⁸	2.97 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	2.58 x 10 ⁻⁰⁵	5.81 x 10 ⁻⁰⁵	2.96 x 10 ⁻⁰⁵	2.58 x 10 ⁻⁰⁴	
	Thorium-230	4.54 x 10 ⁻⁰⁴	1.53 x 10 ⁻¹²	1.10 x 10 ⁻⁰³	7.44 x 10 ⁻¹⁰	7.72 x 10 ⁻⁰⁸	1.18 x 10 ⁻⁰⁶	7.78 x 10 ⁻⁰⁷	1.92 x 10 ⁻⁰⁸	0.00 x 10	4.30 x 10 ⁻⁰⁴	2.57 x 10 ⁻⁰⁴	3.09 x 10 ⁻⁰³	5.33 x 10 ⁻⁰³	
	Radium-226	1.13 x 10 ⁻⁰⁴	2.93 x 10 ⁻¹¹	1.84 x 10 ⁻⁰⁴	8.22 x 10 ⁻⁰⁸	1.04 x 10 ⁻⁰⁷	2.70 x 10 ⁻⁰³	3.55 x 10 ⁻⁰⁷	4.26 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	1.41 x 10 ⁻⁰⁴	6.01 x 10 ⁻⁰⁴	3.08 x 10 ⁻⁰⁴	4.09 x 10 ⁻⁰³	
	Lead-210	3.56 x 10 ⁻⁰⁵	4.61 x 10 ⁻¹²	2.20 x 10 ⁻⁰⁴	1.94 x 10 ⁻¹⁰	1.66 x 10 ⁻⁰⁷	3.85 x 10 ⁻⁰⁵	3.03 x 10 ⁻⁰⁶	1.72 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	6.06 x 10 ⁻⁰⁴	1.18 x 10 ⁻⁰³	2.98 × 10 ⁻⁰³	5.06 x 10 ⁻⁰³	
	Polonium-210	1.07 x 10 ⁻⁰⁴	4.02 x 10 ⁻¹⁴	3.75 x 10 ⁻⁰⁴	1.13 x 10 ⁻¹²	2.93 x 10 ⁻⁰⁷	8.69 x 10 ⁻⁰⁹	5.35 x 10 ⁻⁰⁶	7.34 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	3.60 x 10 ⁻⁰³	1.98 x 10 ⁻⁰³	7.92 x 10 ⁻⁰³	1.40 x 10 ⁻⁰²	
Seasonal Resident	Total by Pathway	9.17 x 10 ⁻⁰⁴	3.63 x 10 ⁻¹¹	1.94 x 10 ⁻⁰³	8.33 x 10 ⁻⁰⁸	6.71 x 10 ⁻⁰⁷	2.91 x 10 ⁻⁰³	9.68 x 10 ⁻⁰⁶	4.36 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	4.82 x 10 ⁻⁰³	4.13 x 10 ⁻⁰³	1.44 x 10 ⁻⁰²	2.91 x 10 ⁻⁰²	
(Patterson Lake South Arm)	Far-Future														
	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.58 x 10 ⁻⁰⁵	1.20 x 10 ⁻¹⁰	1.55 x 10 ⁻⁰⁹	1.72 x 10 ⁻⁰⁵	1.44 x 10 ⁻⁰⁷	1.54 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	1.40 x 10 ⁻⁰⁵	5.42 x 10 ⁻⁰⁶	8.71 x 10 ⁻⁰⁶	6.29 x 10 ⁻⁰⁵	
	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.73 x 10 ⁻⁰⁵	2.10 x 10 ⁻¹¹	1.69 x 10 ⁻⁰⁹	8.69 x 10 ⁻⁰⁸	1.58 x 10 ⁻⁰⁷	5.60 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	1.53 x 10 ⁻⁰⁵	5.91 x 10 ⁻⁰⁶	9.51 x 10 ⁻⁰⁶	4.82 x 10 ⁻⁰⁵	
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	9.24 x 10 ⁻⁰⁶	6.26 x 10 ⁻¹²	4.31 x 10 ⁻⁰⁸	6.61 x 10 ⁻⁰⁷	1.01 x 10 ⁻⁰⁸	2.49 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	3.92 x 10 ⁻⁰⁶	2.90 x 10 ⁻⁰⁶	4.52 x 10 ⁻⁰⁵	6.20 x 10 ⁻⁰⁵	
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.88 x 10 ⁻⁰⁵	8.40 x 10 ⁻⁰⁹	7.03 x 10 ⁻⁰⁸	1.82 x 10 ⁻⁰³	1.04 x 10 ⁻⁰⁷	1.25 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	1.59 x 10 ⁻⁰⁵	9.09 x 10 ⁻⁰⁵	7.73 x 10 ⁻⁰⁵	2.04 x 10 ⁻⁰³	
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.21 x 10 ⁻⁰⁶	1.07 x 10 ⁻¹²	1.13 x 10 ⁻¹⁰	2.63 x 10 ⁻⁰⁸	4.08 x 10 ⁻⁰⁸	2.31 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	3.43 x 10 ⁻⁰⁶	1.93 x 10 ⁻⁰⁸	9.27 x 10 ⁻⁰⁵	9.74 x 10 ⁻⁰⁵	
	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.13 x 10 ⁻⁰⁶	6.40 x 10 ⁻¹⁵	2.00 x 10 ⁻¹⁰	5.95 x 10 ⁻¹²	7.22 x 10 ⁻⁰⁸	9.91 x 10 ⁻¹²	0.00 x 10 ⁺⁰⁰	2.15 x 10 ⁻⁰⁵	1.67 x 10 ⁻⁰⁷	8.72 x 10 ⁻⁰⁵	1.11 x 10 ⁻⁰⁴	
	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	6.45 x 10 ⁻⁰⁵	8.55 x 10 ⁻⁰⁹	1.17 x 10 ⁻⁰⁷	1.84 x 10 ⁻⁰³	5.29 x 10 ⁻⁰⁷	1.40 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	7.40 x 10 ⁻⁰⁵	1.05 x 10 ⁻⁰⁴	3.21 x 10 ⁻⁰⁴	2.42 x 10 ⁻⁰³	
Seasonal Resident		0.07	0.05 (5.12	1.00	5.00 15.11	1.10 15.00	1.64 04	RFD Case	0.47	0.00 15:00	0.05	504 (505	0.74	2.24 15.04	
One-Year-Old (Patterson Lake	Uranium-238	9.37 x 10 ⁻⁰⁵	3.35 x 10 ⁻¹³	1.93 x 10 ⁻⁰⁵	5.22 x 10 ⁻¹¹	1.48 x 10 ⁻⁰⁸	1.64 x 10 ⁻⁰⁴	7.66 x 10 ⁻⁰⁸	8.17 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	2.36 x 10 ⁻⁰⁵	5.34 x 10 ⁻⁰⁵	2.71 x 10 ⁻⁰⁵	3.91 x 10 ⁻⁰⁴	
South Arm)	Uranium-234	1.10 x 10 ⁻⁰⁴	8.20 x 10 ⁻¹³	2.10 x 10 ⁻⁰⁵	9.15 x 10 ⁻¹²	1.61 x 10 ⁻⁰⁸	8.27 x 10 ⁻⁰⁷	8.38 x 10 ⁻⁰⁸	2.97 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	2.58 x 10 ⁻⁰⁵	5.81 x 10 ⁻⁰⁵	2.96 x 10 ⁻⁰⁵	2.58 x 10 ⁻⁰⁴	
	Thorium-230	3.49 x 10 ⁻⁰⁴	1.99 x 10 ⁻¹²	5.59 x 10 ⁻⁰⁴	1.84 x 10 ⁻¹⁰	7.72 x 10 ⁻⁰⁸	1.18 x 10 ⁻⁰⁶	7.78 x 10 ⁻⁰⁷	1.92 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	4.30 x 10 ⁻⁰⁴	2.57 x 10 ⁻⁰⁴	3.09 x 10 ⁻⁰³	5.33 x 10 ⁻⁰³	

							Incrementa	l Dose by Pathway	RFD Case (mSv/yr)					
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by Radionuclide
	Radium-226	1.10 x 10 ⁻⁰⁴	3.80 x 10 ⁻¹¹	1.64 x 10 ⁻⁰⁴	2.03 x 10 ⁻⁰⁸	1.04 x 10 ⁻⁰⁷	2.70 x 10 ⁻⁰³	3.55 x 10 ⁻⁰⁷	4.26 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	1.41 x 10 ⁻⁰⁴	6.01 x 10 ⁻⁰⁴	3.08 x 10 ⁻⁰⁴	4.09 x 10 ⁻⁰³
	Lead-210	3.69 x 10 ⁻⁰⁵	5.98 x 10 ⁻¹²	3.00 x 10 ⁻⁰⁴	3.94 x 10 ⁻¹¹	1.66 x 10 ⁻⁰⁷	3.85 x 10 ⁻⁰⁵	3.03 x 10 ⁻⁰⁶	1.72 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	6.06 x 10 ⁻⁰⁴	1.18 x 10 ⁻⁰³	2.98 x 10 ⁻⁰³	5.06 x 10 ⁻⁰³
	Polonium-210	1.10 x 10 ⁻⁰⁴	5.23 x 10 ⁻¹⁴	7.16 x 10 ⁻⁰⁴	2.79 x 10 ⁻¹³	2.93 x 10 ⁻⁰⁷	8.69 x 10 ⁻⁰⁹	5.35 x 10 ⁻⁰⁶	7.34 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	3.60 x 10 ⁻⁰³	1.98 x 10 ⁻⁰³	7.92 x 10 ⁻⁰³	1.40 x 10 ⁻⁰²
	Total by Pathway	8.08 x 10 ⁻⁰⁴	4.72 x 10 ⁻¹¹	1.78 x 10 ⁻⁰³	2.06 x 10 ⁻⁰⁸	6.71 x 10 ⁻⁰⁷	2.91 x 10 ⁻⁰³	9.68 x 10 ⁻⁰⁶	4.36 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	4.82 x 10 ⁻⁰³	4.13 x 10 ⁻⁰³	1.44 x 10 ⁻⁰²	2.91 x 10 ⁻⁰²
								Far-Future						
	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.10 x 10 ⁻⁰⁵	2.97 x 10 ⁻¹¹	1.55 x 10 ⁻⁰⁹	1.72 x 10 ⁻⁰⁵	1.44 x 10 ⁻⁰⁷	1.54 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	8.24 x 10 ⁻⁰⁶	5.42 x 10 ⁻⁰⁶	8.71 x 10 ⁻⁰⁶	6.29 x 10 ⁻⁰⁵
	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.20 x 10 ⁻⁰⁵	5.20 x 10 ⁻¹²	1.69 x 10 ⁻⁰⁹	8.69 x 10 ⁻⁰⁸	1.58 x 10 ⁻⁰⁷	5.60 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	8.96 x 10 ⁻⁰⁶	5.91 x 10 ⁻⁰⁶	9.51 x 10 ⁻⁰⁶	4.82 x 10 ⁻⁰⁵
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	4.70 x 10 ⁻⁰⁶	1.55 x 10 ⁻¹²	4.31 x 10 ⁻⁰⁸	6.61 x 10 ⁻⁰⁷	1.01 x 10 ⁻⁰⁸	2.49 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	1.69 x 10 ⁻⁰⁶	2.90 x 10 ⁻⁰⁶	4.52 x 10 ⁻⁰⁵	6.20 x 10 ⁻⁰⁵
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.68 x 10 ⁻⁰⁵	2.08 x 10 ⁻⁰⁹	7.03 x 10 ⁻⁰⁸	1.82 x 10 ⁻⁰³	1.04 x 10 ⁻⁰⁷	1.25 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	1.21 x 10 ⁻⁰⁵	9.09 x 10 ⁻⁰⁵	7.73 x 10 ⁻⁰⁵	2.04 x 10 ⁻⁰³
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.65 x 10 ⁻⁰⁶	2.17 x 10 ⁻¹³	1.13 x 10 ⁻¹⁰	2.63 x 10 ⁻⁰⁸	4.08 x 10 ⁻⁰⁸	2.31 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	3.95 x 10 ⁻⁰⁶	1.93 x 10 ⁻⁰⁸	9.27 x 10 ⁻⁰⁵	9.74 x 10 ⁻⁰⁵
	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	4.07 x 10 ⁻⁰⁶	1.59 x 10 ⁻¹⁵	2.00 x 10 ⁻¹⁰	5.95 x 10 ⁻¹²	7.22 x 10 ⁻⁰⁸	9.91 x 10 ⁻¹²	0.00 x 10 ⁺⁰⁰	3.48 x 10 ⁻⁰⁵	1.67 x 10 ⁻⁰⁷	8.72 x 10 ⁻⁰⁵	1.11 x 10 ⁻⁰⁴
	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	5.02 x 10 ⁻⁰⁵	2.12 x 10 ⁻⁰⁹	1.17 x 10 ⁻⁰⁷	1.84 x 10 ⁻⁰³	5.29 x 10 ⁻⁰⁷	1.40 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	6.98 x 10 ⁻⁰⁵	1.05 x 10 ⁻⁰⁴	3.21 x 10 ⁻⁰⁴	2.42 x 10 ⁻⁰³
								RFD Case						
	Uranium-238	2.45 x 10 ⁻⁰⁶	6.72 x 10 ⁻¹⁵	9.00 x 10 ⁻⁰⁷	6.83 x 10 ⁻¹²	3.85 x 10 ⁻¹⁰	4.27 x 10 ⁻⁰⁶	2.59 x 10 ⁻⁰⁹	2.76 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.90 x 10 ⁻⁰⁶	1.39 x 10 ⁻⁰⁶	5.27 x 10 ⁻⁰⁷	1.25 x 10 ⁻⁰⁵
	Uranium-234	2.95 x 10 ⁻⁰⁶	1.64 x 10 ⁻¹⁴	9.83 x 10 ⁻⁰⁷	1.19 x 10 ⁻¹²	4.20 x 10 ⁻¹⁰	2.15 x 10 ⁻⁰⁸	2.83 x 10 ⁻⁰⁹	1.01 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	3.17 x 10 ⁻⁰⁶	1.51 x 10 ⁻⁰⁶	5.74 x 10 ⁻⁰⁷	9.22 x 10 ⁻⁰⁶
	Thorium-230	1.18 x 10 ⁻⁰⁵	3.98 x 10 ⁻¹⁴	5.02 x 10 ⁻⁰⁵	3.40 x 10 ⁻¹¹	2.01 x 10 ⁻⁰⁹	3.08 x 10 ⁻⁰⁸	4.27 x 10 ⁻⁰⁸	1.06 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	3.41 x 10 ⁻⁰⁵	6.70 x 10 ⁻⁰⁶	8.29 × 10 ⁻⁰⁵	1.86 x 10 ⁻⁰⁴
	Radium-226	2.95 x 10 ⁻⁰⁶	7.62 x 10 ⁻¹³	6.84 x 10 ⁻⁰⁶	3.05 x 10 ⁻⁰⁹	2.72 x 10 ⁻⁰⁹	7.04 x 10 ⁻⁰⁵	1.49 x 10 ⁻⁰⁸	1.79 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	9.75 x 10 ⁻⁰⁶	1.57 x 10 ⁻⁰⁵	3.93 x 10 ⁻⁰⁶	1.11 x 10 ⁻⁰⁴
	Lead-210	9.29 x 10 ⁻⁰⁷	1.20 x 10 ⁻¹³	1.74 x 10 ⁻⁰⁶	1.53 x 10 ⁻¹²	4.32 x 10 ⁻⁰⁹	1.00 x 10 ⁻⁰⁶	2.20 x 10 ⁻⁰⁸	1.24 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	7.65 x 10 ⁻⁰⁴	3.07 x 10 ⁻⁰⁵	1.97 × 10 ⁻⁰⁵	8.19 x 10 ⁻⁰⁴
	Polonium-210	2.79 x 10 ⁻⁰⁶	1.05 x 10 ⁻¹⁵	2.97 x 10 ⁻⁰⁶	8.94 x 10 ⁻¹⁵	7.64 x 10 ⁻⁰⁹	2.26 x 10 ⁻¹⁰	3.88 x 10 ⁻⁰⁸	5.32 x 10 ⁻¹²	0.00 x 10 ⁺⁰⁰	5.59 x 10 ⁻⁰⁴	5.16 x 10 ⁻⁰⁵	8.13 x 10 ⁻⁰⁵	6.98 x 10 ⁻⁰⁴
easonal Resident	Total by Pathway	2.39 x 10 ⁻⁰⁵	9.47 x 10 ⁻¹³	6.37 x 10 ⁻⁰⁵	3.10 x 10 ⁻⁰⁹	1.75 x 10 ⁻⁰⁸	7.57 x 10 ⁻⁰⁵	1.24 x 10 ⁻⁰⁷	1.82 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	1.37 x 10 ⁻⁰³	1.08 x 10 ⁻⁰⁴	1.89 x 10 ⁻⁰⁴	1.84 x 10 ⁻⁰³
(Lloyd Lake)	Far-Future Far-Future													
	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	5.33 x 10 ⁻⁰⁷	4.04 x 10 ⁻¹²	4.05 x 10 ⁻¹¹	4.49 x 10 ⁻⁰⁷	4.86 x 10 ⁻⁰⁹	5.18 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.27 x 10 ⁻⁰⁶	1.41 x 10 ⁻⁰⁷	2.26 x 10 ⁻⁰⁷	3.67 x 10 ⁻⁰⁶
	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	5.82 x 10 ⁻⁰⁷	7.07 x 10 ⁻¹³	4.41 x 10 ⁻¹¹	2.26 x 10 ⁻⁰⁹	5.31 x 10 ⁻⁰⁹	1.89 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	2.47 x 10 ⁻⁰⁶	1.54 x 10 ⁻⁰⁷	2.46 x 10 ⁻⁰⁷	3.46 x 10 ⁻⁰⁶
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	5.04 x 10 ⁻⁰⁷	3.42 x 10 ⁻¹³	1.12 x 10 ⁻⁰⁹	1.72 x 10 ⁻⁰⁸	5.50 x 10 ⁻¹⁰	1.36 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	4.47 x 10 ⁻⁰⁷	7.56 x 10 ⁻⁰⁸	9.52 x 10 ⁻⁰⁷	2.00 x 10 ⁻⁰⁶
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	7.74 x 10 ⁻⁰⁷	3.46 x 10 ⁻¹⁰	1.83 x 10 ⁻⁰⁹	4.75 x 10 ⁻⁰⁵	4.29 x 10 ⁻⁰⁹	5.14 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.77 x 10 ⁻⁰⁶	2.37 x 10 ⁻⁰⁶	9.88 x 10 ⁻⁰⁷	5.39 x 10 ⁻⁰⁵
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.22 x 10 ⁻⁰⁸	1.09 x 10 ⁻¹⁴	2.73 × 10 ⁻¹²	6.98 x 10 ⁻¹⁰	4.15 x 10 ⁻¹⁰	2.32 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	7.56 x 10 ⁻⁰⁴	2.33 × 10 ⁻¹⁰	7.08 × 10 ⁻⁰⁸	7.56 x 10 ⁻⁰⁴
	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	5.00 x 10 ⁻¹²	1.56 x 10 ⁻¹³	6.84 x 10 ⁻¹⁰	9.33 x 10 ⁻¹⁴	0.00 x 10 ⁺⁰⁰	5.10 x 10 ⁻⁰⁴	4.66 x 10 ⁻⁰⁹	6.56 x 10 ⁻⁰⁷	5.10 x 10 ⁻⁰⁴
	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.41 x 10 ⁻⁰⁶	3.51 x 10 ⁻¹⁰	3.05 x 10 ⁻⁰⁹	4.79 x 10 ⁻⁰⁵	1.61 x 10 ⁻⁰⁸	5.66 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.27 x 10 ⁻⁰³	2.75 x 10 ⁻⁰⁶	3.14 x 10 ⁻⁰⁶	1.33 x 10 ⁻⁰³
								RFD Case						
	Uranium-238	2.44 x 10 ⁻⁰⁶	8.73 x 10 ⁻¹⁵	6.25 x 10 ⁻⁰⁷	1.69 x 10 ⁻¹²	1.57 x 10 ⁻⁰⁸	5.72 x 10 ⁻⁰⁶	1.05 x 10 ⁻⁰⁷	3.59 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	4.50 x 10 ⁻⁰⁷	1.90 x 10 ⁻⁰⁶	2.31 x 10 ⁻⁰⁷	1.15 x 10 ⁻⁰⁵
	Uranium-234	2.86 x 10 ⁻⁰⁶	2.14 x 10 ⁻¹⁴	6.80 x 10 ⁻⁰⁷	2.96 x 10 ⁻¹³	1.70 x 10 ⁻⁰⁸	2.88 x 10 ⁻⁰⁸	1.15 x 10 ⁻⁰⁷	1.31 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	4.89 x 10 ⁻⁰⁷	2.06 x 10 ⁻⁰⁶	2.50 x 10 ⁻⁰⁷	6.49 x 10 ⁻⁰⁶
	Thorium-230	9.09 x 10 ⁻⁰⁶	5.18 x 10 ⁻¹⁴	2.55 x 10 ⁻⁰⁵	8.43 x 10 ⁻¹²	5.99 x 10 ⁻⁰⁸	3.51 x 10 ⁻⁰⁸	1.27 x 10 ⁻⁰⁶	1.37 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	8.57 x 10 ⁻⁰⁶	3.71 x 10 ⁻⁰⁶	4.03 × 10 ⁻⁰⁵	8.86 x 10 ⁻⁰⁵
	Radium-226	2.86 x 10 ⁻⁰⁶	9.91 x 10 ⁻¹³	6.11 x 10 ⁻⁰⁶	7.56 x 10 ⁻¹⁰	1.42 x 10 ⁻⁰⁷	9.42 x 10 ⁻⁰⁵	7.80 x 10 ⁻⁰⁷	2.33 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	3.95 x 10 ⁻⁰⁶	2.09 x 10 ⁻⁰⁵	1.56 x 10 ⁻⁰⁶	1.33 x 10 ⁻⁰⁴
	Lead-210	9.61 x 10 ⁻⁰⁷	1.56 x 10 ⁻¹³	2.37 x 10 ⁻⁰⁶	3.11 x 10 ⁻¹³	3.44 x 10 ⁻⁰⁷	1.34 x 10 ⁻⁰⁶	1.75 x 10 ⁻⁰⁶	1.62 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	5.52 x 10 ⁻⁰⁶	5.74 x 10 ⁻⁰⁵	2.34 × 10 ⁻⁰⁵	9.31 x 10 ⁻⁰⁵
Sananal Basil	Polonium-210	2.86 x 10 ⁻⁰⁶	1.36 x 10 ⁻¹⁵	5.68 x 10 ⁻⁰⁶	2.21 x 10 ⁻¹⁵	8.54 x 10 ⁻⁰⁷	2.96 x 10 ⁻¹⁰	4.34 x 10 ⁻⁰⁶	6.91 x 10 ⁻¹²	0.00 x 10 ⁺⁰⁰	4.62 x 10 ⁻⁰⁵	1.42 x 10 ⁻⁰⁴	8.88 x 10 ⁻⁰⁵	2.91 x 10 ⁻⁰⁴
Seasonal Resident One-Year-Old	Total by Pathway	2.11 x 10 ⁻⁰⁵	1.23 x 10 ⁻¹²	4.10 x 10 ⁻⁰⁵	7.67 x 10 ⁻¹⁰	1.43 x 10 ⁻⁰⁶	1.01 x 10 ⁻⁰⁴	8.36 x 10 ⁻⁰⁶	2.37 x 10 ⁻⁰⁶	0.00 x 10 ⁺⁰⁰	6.51 x 10 ⁻⁰⁵	2.28 x 10 ⁻⁰⁴	1.55 x 10 ⁻⁰⁴	6.24 x 10 ⁻⁰⁴
(Lloyd Lake)														
	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	3.70 x 10 ⁻⁰⁷	9.99 x 10 ⁻¹³	1.65 x 10 ⁻⁰⁹	6.01 x 10 ⁻⁰⁷	1.98 x 10 ⁻⁰⁷	6.74 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	2.77 x 10 ⁻⁰⁷	1.92 x 10 ⁻⁰⁷	1.01 x 10 ⁻⁰⁷	1.81 x 10 ⁻⁰⁶
	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	4.03 x 10 ⁻⁰⁷	1.75 x 10 ⁻¹³	1.78 x 10 ⁻⁰⁹	3.03 x 10 ⁻⁰⁹	2.15 x 10 ⁻⁰⁷	2.45 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	3.02 x 10 ⁻⁰⁷	2.08 x 10 ⁻⁰⁷	1.09 x 10 ⁻⁰⁷	1.24 x 10 ⁻⁰⁶
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.57 x 10 ⁻⁰⁷	8.46 x 10 ⁻¹⁴	3.35 x 10 ⁻⁰⁸	1.96 x 10 ⁻⁰⁸	1.64 x 10 ⁻⁰⁸	1.77 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	9.24 x 10 ⁻⁰⁸	7.52 x 10 ⁻⁰⁸	4.62 x 10 ⁻⁰⁷	9.56 x 10 ⁻⁰⁷
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	6.92 x 10 ⁻⁰⁷	8.56 x 10 ⁻¹¹	9.58 x 10 ⁻⁰⁸	6.35 x 10 ⁻⁰⁵	2.24 x 10 ⁻⁰⁷	6.70 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	4.97 x 10 ⁻⁰⁷	4.14 x 10 ⁻⁰⁶	3.95 x 10 ⁻⁰⁷	7.02 x 10 ⁻⁰⁵
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.66 x 10 ⁻⁰⁸	2.22 x 10 ⁻¹⁵	2.33 × 10 ⁻¹⁰	9.31 × 10 ⁻¹⁰	3.28 x 10 ⁻⁰⁸	3.05 x 10 ⁻¹¹	0.00 x 10 ⁺⁰⁰	4.00 x 10 ⁻⁰⁸	1.40 × 10 ⁻⁰⁹	8.94 × 10 ⁻⁰⁸	1.82 x 10 ⁻⁰⁷
	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	5.82 x 10 ⁻¹⁰	1.99 x 10 ⁻¹³	7.73 x 10 ⁻⁰⁸	1.22 x 10 ⁻¹³	0.00 x 10 ⁺⁰⁰	2.24 x 10 ⁻⁰⁸	1.68 x 10 ⁻⁰⁸	8.34 x 10 ⁻⁰⁷	9.51 x 10 ⁻⁰⁷
	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.74 x 10 ⁻⁰⁶	8.68 x 10 ⁻¹¹	1.34 x 10 ⁻⁰⁷	6.41 x 10 ⁻⁰⁵	7.63 x 10 ⁻⁰⁷	7.38 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	1.23 x 10 ⁻⁰⁶	4.64 x 10 ⁻⁰⁶	1.99 x 10 ⁻⁰⁶	7.54 x 10 ⁻⁰⁵
								Far-Future Adı	ılt					

							Increment	al Dose by Pathway	RFD Case (mSv/yr)					
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by Radionuclide
	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.91 x 10 ⁻⁰⁴	1.45 x 10 ⁻⁰⁹	2.69 x 10 ⁻⁰⁸	2.98 x 10 ⁻⁰⁴	1.74 x 10 ⁻⁰⁶	1.85 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	5.31 x 10 ⁻⁰⁴	1.20 x 10 ⁻⁰⁴	2.40 x 10 ⁻⁰⁴	1.40 x 10 ⁻⁰³
	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.08 x 10 ⁻⁰⁴	2.53 x 10 ⁻¹⁰	2.93 x 10 ⁻⁰⁸	1.50 x 10 ⁻⁰⁶	1.90 x 10 ⁻⁰⁶	6.75 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	5.81 x 10 ⁻⁰⁴	1.30 x 10 ⁻⁰⁴	2.62 x 10 ⁻⁰⁴	1.18 x 10 ⁻⁰³
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	5.00 x 10 ⁻⁰⁵	3.39 x 10 ⁻¹¹	7.47 x 10 ⁻⁰⁷	1.14 x 10 ⁻⁰⁵	5.48 x 10 ⁻⁰⁸	1.35 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	6.70 x 10 ⁻⁰⁵	6.40 x 10 ⁻⁰⁵	5.37 x 10 ⁻⁰⁴	7.30 x 10 ⁻⁰⁴
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.70 x 10 ⁻⁰⁴	7.61 x 10 ⁻⁰⁸	1.22 x 10 ⁻⁰⁶	3.15 x 10 ⁻⁰²	9.43 x 10 ⁻⁰⁷	1.13 x 10 ⁻⁰⁴	0.00 x 10 ⁺⁰⁰	4.55 x 10 ⁻⁰⁴	2.01 x 10 ⁻⁰³	1.90 x 10 ⁻⁰³	3.62 x 10 ⁻⁰²
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	6.15 x 10 ⁻⁰⁵	5.42 x 10 ⁻¹¹	1.96 x 10 ⁻⁰⁹	4.55 x 10 ⁻⁰⁷	2.07 x 10 ⁻⁰⁶	1.17 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	5.48 x 10 ⁻⁰⁴	4.31 x 10 ⁻⁰⁷	9.16 x 10 ⁻⁰⁴	1.53 x 10 ⁻⁰³
	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.08 x 10 ⁻⁰⁴	3.24 x 10 ⁻¹³	3.47 x 10 ⁻⁰⁹	1.03 x 10 ⁻¹⁰	3.66 x 10 ⁻⁰⁶	5.02 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	3.42 x 10 ⁻⁰³	3.68 x 10 ⁻⁰⁶	7.21 x 10 ⁻⁰³	1.07 x 10 ⁻⁰²
Permanent	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	7.89 x 10 ⁻⁰⁴	7.79 x 10 ⁻⁰⁸	2.03 x 10 ⁻⁰⁶	3.18 x 10 ⁻⁰²	1.04 x 10 ⁻⁰⁵	1.32 x 10 ⁻⁰⁴	0.00 x 10 ⁺⁰⁰	5.61 x 10 ⁻⁰³	2.33 x 10 ⁻⁰³	1.11 x 10 ⁻⁰²	5.18 x 10 ⁻⁰²
Resident (Pattersor Lake North Arm -								Far-Future One-Ye	ar-Old					
West Basin)	Uranium-238	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.33 x 10 ⁻⁰⁴	3.58 x 10 ⁻¹⁰	1.09 x 10 ⁻⁰⁶	3.99 x 10 ⁻⁰⁴	7.07 x 10 ⁻⁰⁵	2.41 x 10 ⁻⁰⁵	0.00 x 10 ⁺⁰⁰	3.13 x 10 ⁻⁰⁴	1.47 x 10 ⁻⁰⁴	1.09 x 10 ⁻⁰⁴	1.20 x 10 ⁻⁰³
	Uranium-234	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.44 x 10 ⁻⁰⁴	6.27 x 10 ⁻¹¹	1.19 x 10 ⁻⁰⁶	2.01 x 10 ⁻⁰⁶	7.69 x 10 ⁻⁰⁵	8.78 x 10 ⁻⁰⁸	0.00 x 10 ⁺⁰⁰	3.41 x 10 ⁻⁰⁴	1.59 x 10 ⁻⁰⁴	1.18 x 10 ⁻⁰⁴	8.42 x 10 ⁻⁰⁴
	Thorium-230	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.54 x 10 ⁻⁰⁵	8.39 x 10 ⁻¹²	2.22 x 10 ⁻⁰⁵	1.30 x 10 ⁻⁰⁵	1.63 x 10 ⁻⁰⁶	1.76 x 10 ⁻⁰⁹	0.00 x 10 ⁺⁰⁰	2.89 x 10 ⁻⁰⁵	5.75 x 10 ⁻⁰⁵	2.57 x 10 ⁻⁰⁴	4.06 x 10 ⁻⁰⁴
	Radium-226	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	1.52 x 10 ⁻⁰⁴	1.88 x 10 ⁻⁰⁸	6.37 x 10 ⁻⁰⁵	4.22 x 10 ⁻⁰²	4.93 x 10 ⁻⁰⁵	1.47 x 10 ⁻⁰⁴	0.00 x 10 ⁺⁰⁰	3.45 x 10 ⁻⁰⁴	3.17 x 10 ⁻⁰³	7.33 x 10 ⁻⁰⁴	4.69 x 10 ⁻⁰²
	Lead-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	8.36 x 10 ⁻⁰⁵	1.10 x 10 ⁻¹¹	1.56 x 10 ⁻⁰⁷	6.09 x 10 ⁻⁰⁷	1.65 x 10 ⁻⁰⁴	1.52 x 10 ⁻⁰⁷	0.00 x 10 ⁺⁰⁰	6.32 x 10 ⁻⁰⁴	1.03 x 10 ⁻⁰⁶	1.09 x 10 ⁻⁰³	1.97 x 10 ⁻⁰³
	Polonium-210	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	2.06 x 10 ⁻⁰⁴	8.02 x 10 ⁻¹⁴	3.88 x 10 ⁻⁰⁷	1.34 x 10 ⁻¹⁰	4.09 x 10 ⁻⁰⁴	6.52 x 10 ⁻¹⁰	0.00 x 10 ⁺⁰⁰	5.56 x 10 ⁻⁰³	1.24 x 10 ⁻⁰⁵	9.16 x 10 ⁻⁰³	1.53 x 10 ⁻⁰²
	Total by Pathway	0.00 x 10 ⁺⁰⁰	0.00 x 10 ⁺⁰⁰	7.44 x 10 ⁻⁰⁴	1.93 x 10 ⁻⁰⁸	8.87 x 10 ⁻⁰⁵	4.26 x 10 ⁻⁰²	7.72 x 10 ⁻⁰⁴	1.72 x 10 ⁻⁰⁴	0.00 x 10 ⁺⁰⁰	7.22 x 10 ⁻⁰³	3.54 x 10 ⁻⁰³	1.15 x 10 ⁻⁰²	6.66 x 10 ⁻⁰²

mSv/yr = millisieverts per year; n/a = not applicable or not assessed; RFD = reasonably foreseeable development.

5.2.5 Uncertainty in Exposure Assessment

The exposure assessment followed CSA, Health Canada and USEPA guidance. Key uncertainties in the human health exposure assumptions and how they are addressed in the HHRA are summarized in Table 5-15.

Concentrations of COPCs in environmental media including water, sediment, air, soil, and Traditional Food items were estimated based on the assumption that human and ecological receptors are exposed to the maximum exposure concentrations at their location for each model scenario and Project phase. The duration of this exposure was assumed to be sufficient for each receptor to be in equilibrium with their environment. This results in conservatively high predicted uptakes of COPCs by ecological receptors and exposures to human health receptors.

The assumptions to address uncertainties in the exposure assessment were anticipated to produce conservative exposure estimates for human health receptors. Therefore, the risk that the exposure assessment underestimated potential exposure of human health receptors to COPCs from the Project is low.



Table 5-15: How Uncertainties in the Human Health Exposure Assessment Are Addressed

Area of Concern	Uncertainties	Description of How the Uncertainties Have Been Addressed
Receptor selection	There are no permanent residents in the RSA, but the area is known to be used for subsistence harvesting including fishing, hunting, and gathering, and there are cabins and outfitters in the LSA that can be accessed by road. There are uncertainties on how potential receptors would realistically use the LSA and RSA (i.e., locations and residency times).	 Based residency and location assumptions on current understanding of how people use the project RSA Sought feedback from local Indigenous Groups Assumed reasonably conservative residency times for receptors that conservatively represent receptors with shorter residency times, such as a recreational visitor Located HH receptors in the LSA and RSA at locations known to be in the area of cabins, camps, outfitters Included two receptors located at the Project site to cover all Project phases: camp worker during all phases and a family permanently residing at the decommissioned and reclaimed Project site well beyond Closure (i.e., far-future projection)
Traditional Foods diet	Detailed site-specific Traditional Foods dietary information is not currently available for the Rook I study area.	 Assumed all receptors consume Traditional Foods. Receptors included a high consumer and an average consumer of Traditional Foods Assumed that all Traditional Foods in a receptor's diet is from the LSA or RSA Used recognized Traditional Foods survey results for relevant Saskatchewan ecozones (FNFNES) as a starting point Sought feedback from local Indigenous Groups and regulatory bodies. Based the total food intake for male and female HH receptors on an adult male diet (N288.1-20 central tendency) Based the Traditional Foods diet average and high ingestion rates on the Boreal Shield ecozone, which has higher ingestion rates than the Boreal Plains ecozone Adjusted the Traditional Foods diet to account for a higher ingestion rate for fruits and berries than considered in the FNFNES data
Selection of representative ecological receptors for the IMPACT model to represent Traditional Foods receptors	Where possible, there is interest to simplify the environmental pathways model used to estimate potential HH risks without leading to an underestimate of potential risk.	 Selected representative foods from the top three Traditional Foods items in the Boreal Shield diet Selected representative foods from the Traditional Foods items known to be used by local Indigenous Groups Applied feedback from local Indigenous Groups and regulatory bodies Representative foods with linkages to the aquatic environment were preferred over terrestrial receptors from the same location because they have the potential to be more exposed to Project-related COPCs through atmospheric and aquatic pathways

COPC = constituent of potential concern; FNFNES = First Nations Food, Nutrition, and Environmental Study; HH = human health; LSA = local study area; RSA = regional study area.



5.3 Toxicity Assessment

The human health toxicity assessment considers potential adverse health effects from non-radiological and radiological exposures via two internal exposure pathways: ingestion and inhalation. In the case of chemical toxicity from exposure to radiological COPCs, the assessment also considers radiological dose from these exposures.

For assessment of non-radiological COPCs, a toxicity reference value (TRV) is used. A TRV is a toxicological index, associating specific health effects with a level of exposure to a chemical. TRVs may include slope factors and unit risks for carcinogens, and reference doses, tolerable daily intakes, or acceptable daily intakes for non-carcinogens.

Toxicity reference values are used in the risk characterization to determine incremental lifetime cancer risks (ILCRs) for carcinogens and hazard quotients (HQs) for non-carcinogens.

For assessment of radiological COPCs, a radiation dose benchmark that combines external and internal radiological doses from all radiological COPCs is used.

5.3.1 Toxicity Reference Values

No COPCs in air were identified for further evaluation of potential risks for human health; therefore, toxicity via inhalation was not included in the toxicity assessment. Separate toxicity benchmarks for direct contact effects from dermal exposure are not available. Although some of the COPCs present in soil may cause direct contact dermatitis, information is not available to suggest that such effects can occur at environmental levels (CSA 2022). A summary of the TRVs used in the HHRA is shown in Table 5-16.

Chloride and sulphate are COPCs identified in Section 4.4, Final List of Constituents of Potential Concern for the Environmental Risk Assessment. These COPCs are associated with water ingestion. Chloride does not have a drinking water standard and is not considered to present a risk to human health at concentrations found in drinking water. Sulphate in drinking water is associated with adverse physiological effects such as diarrhoea or dehydration at concentrations above 500 mg/L. The upper bound concentration of sulphate at the edge of the treated effluent mixing zone was 291 mg/L, as shown in Table 4-2, and therefore concentrations at exposure points farther downstream would be less than those associated with adverse physiological effects. For these reasons, chloride and sulphate were not assessed further in the HHRA.

Arsenic, cobalt, copper, molybdenum, and uranium were retained for further evaluation in the HHRA because they were predicted to exceed water or sediment quality screening benchmarks (Section 4.2.5, Water Quality Constituents for Further Evaluation in the Environmental Risk Assessment).

The relevant non-cancer TRVs are expressed as a quantity of a chemical per unit body weight per unit time (mg/kg/d) for oral exposure and have generally been derived for sensitive individuals in the public based on sensitive endpoints. Additionally, these factors typically involve the incorporation of uncertainty factors by regulatory agencies to account for uncertainties inherent



in the underlying studies or their applicability for protection of members of the public. Carcinogenic effects TRVs are generally referred to as slope factors or unit risks and are used to estimate upper-bound lifetime probabilities of an individual developing cancer as a result of exposure to a particular level of a potential carcinogen. The carcinogen slope factor or unit risk is, therefore, the lifetime cancer risk per unit of dose or concentration. The slope factor is expressed as risk per mg/kg/d, or (mg/kg/d)⁻¹, for oral exposure. Arsenic was the only Project-related COPC evaluated as a carcinogen.

In the selection of TRVs, preference was given to toxicological benchmarks derived by Health Canada, the USEPA Integrated Risk Information System database, the Agency for Toxic Substances and Disease Registry (ATSDR) and the WHO. The supporting documentation for each toxicity benchmark was reviewed and professional judgment was used to evaluate the appropriateness of the benchmark value.

The human health TRVs for arsenic, copper, molybdenum, and uranium were obtained from Health Canada's TRV Guidance (Health Canada 2021b; 2010c). Since Health Canada does not have a published TRV for cobalt, the cobalt TRV was obtained from the ATSDR (ATSDR 2004).

Table 5-16: Human Health Oral Exposure Toxicity Reference Values

СОРС	Benchmark Value	Unit	Reference
Arsenic (cancer)	1.8	(mg/kg/d) ⁻¹	Health Canada 2021b
Cobalt	0.01	mg/kg/d	ATSDR 2004
Copper	0.426	mg/kg/d	Health Canada 2021b
Molybdenum	0.023 (infant) 0.028 (adult)	mg/kg/d	Health Canada 2010c
Uranium	0.0006	mg/kg/d	Health Canada 2021b

COPC = constituent of potential concern.

5.3.1.1 Arsenic

Arsenic is classified as a Class I carcinogen to humans (Environment Canada and Health Canada 1993). Health Canada (2021b) recommends 1.8 (mg/kg/d)⁻¹ as the oral slope factor for arsenic. It was originally developed by Health Canada when the agency was deriving a Guideline for Canadian Drinking Water Quality (Health Canada 2006). The TRV is based on the risk of bladder, lung, and liver cancer in people exposed to arsenic in their drinking water (Morales et al. 2000; Chen et al. 1985; Wu et al. 1989).

5.3.1.2 Cobalt

Cobalt is a trace element that is essential to human health (Health Canada 2021b). Cobalt is not classified as a human carcinogen. Health Canada (Health Canada 2021b) does not provide a threshold oral TRV for cobalt. The listed TRV of 0.01 mg/kg/d is recommended by ATSDR (2004). The ATSDR TRV is an intermediate Minimal Risk Level, and is based on a study by Davis and Fields (Davis and Fields 1958), in which human males ingested a 2% cobalt chloride solution (in water or milk) for up to 22 days. The critical endpoint was hematological effects (increased levels of



erythrocytes). The ATSDR took the lowest observed adverse effect level (LOAEL) of 1 mg/kg/d and applied a total uncertainty factor of 100 to arrive at their intermediate minimal risk level. The ATSDR did not derive a chronic minimal risk level for cobalt due to a lack of relevant animal and human studies.

5.3.1.3 Copper

Copper is a trace element that is essential for human health (Health Canada 2021b). Copper is not classified as a human carcinogen. Health Canada recommends 0.426 mg/kg/d as the threshold oral TRV for copper for all age groups. The TRV was originally developed by Health Canada when the agency was deriving a Guideline for Canadian Drinking Water Quality (Health Canada 2019a). The TRV is based on a critical health effect of gastrointestinal toxicity and liver function (hepatoxicity) in human infants exposed to copper in drinking water (Olivares et al. 1998). The TRV is based directly (no uncertainty factors applied) on the upper bound of the 95th confidence interval for a no observed adverse effect level (NOAEL) of 2 mg/L copper in drinking water (0.318 mg/kg/d).

5.3.1.4 Molybdenum

Molybdenum is considered to be an essential trace element for human health (Health Canada 2021b). However, Health Canada recommends that potential health risks to human receptors be characterized if molybdenum is identified as a COPC (Health Canada 2021b). The TRVs for essential trace elements are tolerable upper intake levels (ULs), which are considered to be the highest average daily nutrient intake levels that are likely to pose no risk of adverse health effects to almost all individuals in the general population. Health Canada (2010c) recommends age-specific ULs for molybdenum that are based on a NOAEL value derived for adults (IOM 2001) from sub-chronic developmental and reproductive effects on rats consuming molybdate in drinking water. An uncertainty factor of 30 was applied (10 for interspecies variability and 3 for intraspecies variability) to the NOAEL value of 0.9 mg/kg-d. The adult UL was weight adjusted to derive age-based TRVs. As with other essential trace elements, Health Canada (2010c) recommends that adjustments for relative bioavailability of molybdenum may be necessary when considering oral exposures from different pathways.

5.3.1.5 Uranium

Health Canada (2021b) recommends 6.0×10^{-4} mg/kg/d as the threshold oral TRV for uranium for non-radiological effects for all age groups. Uranium (non-radiological) is not classified as a human carcinogen. The TRV was originally developed by Health Canada when the agency was deriving a Guideline for Canadian Drinking Water Quality and it has since been re-affirmed (Health Canada 2019b). The TRV is based on the critical health effect of kidney toxicity in rats exposed to uranium in drinking water (Gilman et al. 1998). The TRV is based on a NOAEL of 0.06 mg/kg/d and a total uncertainty factor of 100.



5.3.2 Radiation Dose Limits and Targets

Radiological dose from radiological COPCs was retained for further evaluation because screening values were not available for individual radionuclides emitted to water and air.

Potential effects from radiation were compared to an effective dose limit (Table 5-17). The effective dose is defined as the sum of all tissue equivalent doses multiplied by the appropriate tissue weighting factors associated with each respective tissue (Health Canada 2010b). The limit is incremental and is exclusive of natural background, such as natural levels of radon, and medical exposures. The public dose limit for radiation protection is 1 mSv/yr, as described in the Radiation Protection Regulations under the *Nuclear Safety and Control Act*, and as recommended in CSA N288.6-22. A higher incremental dose than the effective dose limit is considered unacceptable.

Incremental dose from the Project can also be compared to a dose constraint. A dose constraint is a conservative value for the annual increment dose applied to a single operation that is considered protective without further demonstration in situations where multiple sources may contribute to incremental dose (Health Canada 2011b). Application of a dose constraint is meant to ensure that the combined doses from multiple sources do not exceed the public dose limit of 1 mSv/yr. A dose constraint of 0.3 mSv/yr is used in the ERA, as recommended by Health Canada (Health Canada 2011b). The dose constraint represents a dose, lower than the public dose limit that ensures that the combined dose from multiple sources does not result in exceedance of the public dose limit. Exceedance of the dose constraint does not indicate that adverse effects would occur, but instead indicates that the assumptions used in the calculation of exposure estimates for the operation should be examined in more detail.

Table 5-17: Radiation Dose Limits for Human Health

СОРС	Benchmark Value	Unit	Reference
Effective dose limit: public and non-nuclear energy worker	1	mSv/yr	Radiation Protection Regulations. SOR/2000-203
Dose constraint	0.3	mSv/yr	Health Canada 2014; ICRP 2007; IAEA 2014

mSv/yr = millisieverts per year.

5.3.3 Uncertainty in the Toxicity Assessment

In general, TRV are usually based on limited toxicological data. For this reason, a margin of safety is built into TRV estimates, by use of uncertainty factors or conservative confidence levels, and actual risks are lower than those estimated. In this risk assessment, TRV recommended by Health Canada were used when available to reduce uncertainty that potential health risks for human receptors would be underestimated in the risk evaluation.

The two major areas of uncertainty introduced in this toxicity assessment are animal to human extrapolation for Health Canada's recommend TRV for uranium, and use of an intermediate-duration TRV from another regulatory agency than Health Canada for cobalt. In both cases, uncertainty factors were applied in the derivation of the TRVs. For uranium, the chronic TRV was based on a no observed adverse effects level for rats and a total uncertainty factor of 100. For



cobalt, the intermediate (sub-chronic) TRV was based on a LOAEL for humans and a total uncertainty factor of 100. As a result, overestimation of the potential for adverse effects on humans is more likely than underestimation for similar exposure scenarios.

5.4 Risk Characterization

Risk assessment is the process of estimating the likelihood of undesirable effects on human health resulting from exposure to chemical contaminants. Three components must be present for risks to human health to exist:

- The COPC must be present at concentrations sufficient to cause a possible adverse effect.
- A receptor (theoretical human health receptor) must be present.
- There must be a complete exposure pathway by which the receptor can come into contact with the chemical.

5.4.1 Risk Estimation

The potential for adverse effects on human receptors was determined in the risk assessment through the risk characterization step, where risk estimates were calculated to determine the potential for effects on the human receptors identified. The risk estimate was determined by comparing the predicted exposures, in terms of doses, with exposures that are known to be protective based on effects data (i.e., TRVs or radiation dose limits).

The methods of non-radiological risk estimation used for the HHRA are:

- HQs for non-carcinogens; and
- ILCR for carcinogens.

Hazard quotients are calculated in IMPACT as the ratio of the exposure concentration or intake rate divided by the benchmark value, as shown below:

$$HQ = \frac{Exposure (Dose) Estimate}{TRV}$$

The HQs were compared to threshold values. Non-carcinogenic constituents are not expected to cause any adverse health effects below the toxicity reference or threshold value. The HQs can be compared to a benchmark value of one (1) if all exposure pathways (exposures from all pathways including background) are accounted for.

To account for uncertainty in pathways beyond Project activities (i.e., exposure to background sources unrelated to the Project), it was determined that to be protective a benchmark HQ value of 0.2 per medium (e.g., water, soil, food, air) would be considered acceptable for the assessment. This approach is consistent with the approach taken by Health Canada in its guidance on human health preliminary quantitative risk assessment (Health Canada 2021a).



For carcinogens (e.g., arsenic), the incremental risk (i.e., total risk minus background risk) of developing cancer over a lifetime was estimated by multiplying the predicted dose above background by the cancer slope factor, as shown below:

$$ILCR = \sum LADD_i \times SF \times ADAF_i$$

where,

ILCR = incremental lifetime cancer risk (unitless)

LADD_i = dose received during lifestage i averaged over a lifetime (mg/kg/d)

SF = adult cancer slope factor (per mg/kg/d)

ADAF_i = age-dependent adjustment factors for lifestage i

Health Canada (2013) recommends that for carcinogens where the mode of action is unknown or the burden of proof for a threshold mode of action is not met, that the assessment should follow the non-threshold approach (i.e., a linear dose-response relationship). The Canadian drinking water guideline technical document for arsenic indicates that there is limited data on the mode of action for arsenic and that the use of a non-linear relationship may overestimate cancer risks of internal organs (Health Canada 2006). Therefore, for this assessment, a linear approach for arsenic was used, and the age-dependent adjustment factors for all life stages were set at 1.

Incremental lifetime cancer risks were compared to de minimis risk levels that are considered essentially negligible compared to background cancer risks. Cancer risks that are considered acceptable can range from 1 in 10,000 to 1 in 1,000,000 in different jurisdictions. Health Canada (2021a) considers an increase in lifetime cancer risk of 1 in 100,000 (or 0.00001) to be essentially negligible compared to the background cancer risk level in North America of approximately 5 in 10 (or 0.5).

Total radiation doses due to radionuclides in the uranium-238 decay series were predicted. Incremental radiation doses were compared to the regulatory public dose limit and dose limit for a non-nuclear energy worker of 1 mSv/yr and a dose constraint of 0.3 mSv/yr, as described in Section 5.3.2, Radiation Dose Limits and Targets. Radon dose was calculated separately from the dose due to other radionuclides and was also compared to the dose limit of 1 mSv/yr.

5.4.1.1 Estimated Risk to Human Receptors – Application Case

This subsection presents the estimated risks to human receptors due to releases from the Project during all phases including the far-future projection once groundwater solutes have been released to Patterson Lake. Results are presented for the Application Case and the Upper Bound sensitivity scenario.

5.4.1.1.1 Non-carcinogen Risk

The estimated incremental HQs for all human receptors at all assessed locations during Operations and for the far-future projection are shown in Table 5-18. The HQs represent the



ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT HUMAN HEALTH RISK ASSESSMENT

maximum HQ over the Project phase for the COPCs of interest, which is a conservative representation as exposure varies over the different Project phases. HQs were evaluated for the adult and the one-year-old; however, for assessment of non-carcinogens, the one-year-old is typically considered the most sensitive receptor (Health Canada 2010a).

The maximum predicted Project HQ was for uranium for the subsistence harvester (one-year-old) at the Patterson Lake South Arm. This maximum HQ would be primarily from ingestion of terrestrial animals (e.g., beaver, grouse, mallard, moose, moose organs), which is attributed to slightly elevated concentrations of uranium in soil resulting from atmospheric deposition, which then bioaccumulates in terrestrial plants. The subsistence harvester would also eat a high proportion of locally grown and harvested blueberries and Labrador Tea.

For molybdenum, HQs for the base case (without the Project) slightly exceeded 0.2 for the terrestrial animal ingestion pathway for the reference subsistence harvester (one-year-old); however, considering the Project only, HQs remain below 0.2 for all receptors during Operations and for the far-future projection for both the Application Case and the reasonable upper bound sensitivity scenario. The terrestrial animal ingestion pathway for the one-year-old includes ingestion of moose meat and organs, as well as beaver. For molybdenum, the addition of the Project effects does not significantly change the existing base case HQ predictions indicating that the Project has a minimal contribution to existing risk from molybdenum in Traditional Foods.

No significant adverse effect would be likely on any human receptors as a result of releases from the Project during Operations for both the Application Case and reasonable upper bound sensitivity scenario. All estimated incremental Project HQs for all non-carcinogenic COPCs (i.e., cobalt, copper, molybdenum, and uranium) would remain below the acceptable risk level of 0.2 per pathway for the one-year-old and adult for all human receptors over the Project lifespan and in the far future.



Table 5-18: Estimated Non-carcinogen Risk to Human Receptors – Project Lifespan and Far-Future – Application Case and Upper Bound Scenario

Table 5-16. Estillat				,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	Project Life			- 1- 1					Far Fut	ure HQs			
	СОРС	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
								Ва	se Case								
	Cobalt	8.58 × 10 ⁻⁰⁴	9.04 × 10 ⁻⁰⁷	1.01 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	1.15 × 10 ⁻⁰³	7.21 × 10 ⁻⁰⁴	1.97 × 10 ⁻⁰²	2.25 x 10 ⁻⁰²	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Copper	2.91 × 10 ⁻⁰⁵	2.95 × 10 ⁻⁰⁸	4.12 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	8.94 × 10 ⁻⁰⁴	2.69 × 10 ⁻⁰⁴	3.89 × 10 ⁻⁰²	4.01 x 10 ⁻⁰²	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Molybdenum	1.08 × 10 ⁻⁰⁴	1.06 × 10 ⁻⁰⁷	1.62 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	1.21 × 10 ⁻⁰⁶	1.46 × 10 ⁻⁰⁴	8.23 × 10 ⁻⁰²	8.25 x 10 ⁻⁰²	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Uranium	2.46 × 10 ⁻⁰³	8.62 × 10 ⁻⁰⁶	2.24 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	3.43×10^{-03}	1.92 × 10 ⁻⁰²	7.29 × 10 ⁻⁰²	9.80 x 10 ⁻⁰²	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
Campa Mankan		06				05			Incremental Pro	1	l	I	1			ı	
Camp Worker (Patterson Lake	Cobalt	5.31 × 10 ⁻⁰⁶	3.03 x 10 ⁻⁰⁹	3.99 x 10 ⁻⁰⁹	0.00 × 10 ⁺⁰⁰	9.83 × 10 ⁻⁰⁵	8.04 x 10 ⁻⁰⁶	3.37 x 10 ⁻⁰⁴	4.49 x 10 ⁻⁰⁴	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
North Arm – West	Copper	2.44 × 10 ⁻⁰⁸	4.79 x 10 ⁻¹⁰	2.10 x 10 ⁻¹¹	0.00 × 10 ⁺⁰⁰	4.22 × 10 ⁻⁰⁵	1.22 x 10 ⁻⁰⁶	1.44 x 10 ⁻⁰⁴	1.87 x 10 ⁻⁰⁴	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
Basin)	Molybdenum	1.84 × 10 ⁻⁰⁷	1.92 x 10 ⁻⁰⁹	1.64 x 10 ⁻¹⁰	0.00 × 10 ⁺⁰⁰	5.75 × 10 ⁻⁰⁷	5.42 x 10 ⁻⁰⁶	8.58 x 10 ⁻⁰⁵	9.20 x 10 ⁻⁰⁵	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Uranium	2.55 × 10 ⁻⁰⁶	1.91 x 10 ⁻⁰⁵	8.88 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	1.28 × 10 ⁻⁰³	2.14 x 10 ⁻⁰²	1.39 x 10 ⁻⁰²	3.66 x 10 ⁻⁰²	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Calcalt	F 30 10-06	2.02 v 10-09	4.02 v 10-09	0.00 × 10 ⁺⁰⁰	1.02 × 10 ⁻⁰⁴			- Incremental I		n/a	n /a	n /a	n /a	n/a	n/a	2/2
	Cobalt	5.38×10^{-06} 2.55×10^{-08}	3.03 x 10 ⁻⁰⁹ 4.79 x 10 ⁻¹⁰	4.03 x 10 ⁻⁰⁹ 2.20 x 10 ⁻¹¹	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	1.02×10^{-05} 4.34×10^{-05}	8.04 x 10 ⁻⁰⁶ 1.22 x 10 ⁻⁰⁶	3.50 x 10 ⁻⁰⁴ 1.45 x 10 ⁻⁰⁴	4.66 x 10 ⁻⁰⁴ 1.90 x 10 ⁻⁰⁴	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Copper	1.88 × 10 ⁻⁰⁷	1.92 x 10 ⁻⁰⁹	1.67 x 10 ⁻¹⁰	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	4.34 × 10 ⁻⁰⁶	5.42 x 10 ⁻⁰⁶	1.45 x 10 ° · · · · · · · · · · · · · · · · · ·	1.54 x 10 ⁻⁰⁴	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Molybdenum	1.06 × 10 ⁻⁰⁵	1.92 x 10 ⁻⁰⁵	3.72 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	3.53 × 10 ⁻⁰³	2.14 x 10 ⁻⁰²	1.47 x 10 ⁻⁰²	3.95 x 10 ⁻⁰²	n/a	n/a n/a	n/a n/a	n/a n/a	n/a n/a	n/a n/a	n/a n/a	n/a
	Uranium	1.00 × 10 **	1.91 X 10 ··	3.72 X 10 **	0.00 × 10 **	3.33 × 10 ···	2.14 X 10		se Case	n/a	II/ a	П/а	II/a	11/ a	TI/a	Ti/a	n/a
	Cobalt	8.58 × 10 ⁻⁰⁴	9.04 × 10 ⁻⁰⁷	1.01 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.30 × 10 ⁻⁰³	1.44 × 10 ⁻⁰³	2.44 × 10 ⁻⁰²	2.90 x 10 ⁻⁰²	8.58 × 10 ⁻⁰⁴	9.04 × 10 ⁻⁰⁷	1.01 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.30 × 10 ⁻⁰³	1.44 × 10 ⁻⁰³	2.44 × 10 ⁻⁰²	2.90 x 10 ⁻⁰²
	Copper	2.91 × 10 ⁻⁰⁵	2.95 × 10 ⁻⁰⁸	4.12 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	1.79 × 10 ⁻⁰³	5.38 × 10 ⁻⁰⁴	4.12 × 10 ⁻⁰²	4.35 x 10 ⁻⁰²	2.91 × 10 ⁻⁰⁵	2.95 × 10 ⁻⁰⁸	4.12 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	1.79 × 10 ⁻⁰³	5.38 × 10 ⁻⁰⁴	4.12 × 10 ⁻⁰²	4.35 x 10 ⁻⁰²
	Molybdenum	1.08 × 10 ⁻⁰⁴	1.06 × 10 ⁻⁰⁷	1.62 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.42 × 10 ⁻⁰⁶	2.92 × 10 ⁻⁰⁴	6.98 × 10 ⁻⁰²	7.02 x 10 ⁻⁰²	1.08 × 10 ⁻⁰⁴	1.06 × 10 ⁻⁰⁷	1.62 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.42 × 10 ⁻⁰⁶	2.92 × 10 ⁻⁰⁴	6.98 × 10 ⁻⁰²	7.02 x 10 ⁻⁰²
	Uranium	2.46 × 10 ⁻⁰³	8.62 × 10 ⁻⁰⁶	2.24 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	6.85 × 10 ⁻⁰³	3.84 × 10 ⁻⁰²	8.20 × 10 ⁻⁰²	1.30 x 10 ⁻⁰¹	2.46 × 10 ⁻⁰³	8.62 × 10 ⁻⁰⁶	2.24 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	6.85 × 10 ⁻⁰³	3.84 × 10 ⁻⁰²	8.20 × 10 ⁻⁰²	1.30 x 10 ⁻⁰¹
			3,02					L	Incremental Pro	·						0.20	
Subsistence	Cobalt	7.48 × 10 ⁻⁰⁵	1.31 x 10 ⁻⁰⁹	7.93 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	1.96 × 10 ⁻⁰⁴	1.61 x 10 ⁻⁰⁵	6.73 x 10 ⁻⁰⁴	9.60 x 10 ⁻⁰⁴	3.77 × 10 ⁻⁰⁴	7.08 x 10 ⁻¹⁰	4.46 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	1.01 × 10 ⁻⁰³	1.13 x 10 ⁻⁰⁶	2.93 x 10 ⁻⁰³	4.31 x 10 ⁻⁰³
Harvester	Copper	1.38 × 10 ⁻⁰⁶	7.99 x 10 ⁻¹¹	1.71 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	8.43 × 10 ⁻⁰⁵	2.43 x 10 ⁻⁰⁶	2.87 x 10 ⁻⁰⁴	3.75 x 10 ⁻⁰⁴	1.36 × 10 ⁻⁰⁵	5.34 x 10 ⁻¹²	1.92 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	8.32 × 10 ⁻⁰⁴	9.74 x 10 ⁻⁰⁸	7.56 x 10 ⁻⁰⁴	1.60 x 10 ⁻⁰³
(Patterson Lake South Arm)	Molybdenum	5.26 × 10 ⁻⁰⁵	3.03 x 10 ⁻¹⁰	6.99 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	1.15 × 10 ⁻⁰⁶	1.08 x 10 ⁻⁰⁵	1.71 x 10 ⁻⁰⁴	2.36 x 10 ⁻⁰⁴	6.70 × 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	1.00 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	1.50 × 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	7.56 × 10 ⁻⁰⁴	1.44 x 10 ⁻⁰³
South Aim,	Uranium	9.33 × 10 ⁻⁰⁴	2.61 x 10 ⁻⁰⁶	6.77 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.57 × 10 ⁻⁰³	4.28 x 10 ⁻⁰²	2.78 x 10 ⁻⁰²	7.41 x 10 ⁻⁰²	3.06 × 10 ⁻⁰³	2.74 x 10 ⁻⁰⁷	2.79 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	8.53 × 10 ⁻⁰³	1.22 x 10 ⁻⁰³	4.26 x 10 ⁻⁰³	1.71 x 10 ⁻⁰²
							Upper I	Bound Scenario	o – Incremental I	Project Risk							
	Cobalt	7.78 × 10 ⁻⁰⁵	1.31 x 10 ⁻⁰⁹	8.26 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	2.04×10^{-04}	1.61 x 10 ⁻⁰⁵	6.98 x 10 ⁻⁰⁴	9.97 x 10 ⁻⁰⁴	5.77 × 10 ⁻⁰⁴	7.08 x 10 ⁻¹⁰	6.83 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	1.55 × 10 ⁻⁰³	1.13 x 10 ⁻⁰⁶	4.49 x 10 ⁻⁰³	6.62 x 10 ⁻⁰³
	Copper	1.42 × 10 ⁻⁰⁶	7.99 x 10 ⁻¹¹	1.77 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	8.68 × 10 ⁻⁰⁵	2.43 x 10 ⁻⁰⁶	2.89 x 10 ⁻⁰⁴	3.80 x 10 ⁻⁰⁴	2.04 × 10 ⁻⁰⁵	5.34 x 10 ⁻¹²	2.89 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	1.25 × 10 ⁻⁰³	9.74 x 10 ⁻⁰⁸	1.14 x 10 ⁻⁰³	2.41 x 10 ⁻⁰³
	Molybdenum	1.60 × 10 ⁻⁰⁴	3.03 x 10 ⁻¹⁰	2.17 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	3.51 × 10 ⁻⁰⁶	1.08 x 10 ⁻⁰⁵	2.93 x 10 ⁻⁰⁴	4.68 x 10 ⁻⁰⁴	2.57 × 10 ⁻⁰³	$0.00 \times 10^{+00}$	3.85 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	5.74 × 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.92 × 10 ⁻⁰³	5.55 x 10 ⁻⁰³
	Uranium	2.56 × 10 ⁻⁰³	2.61 x 10 ⁻⁰⁶	1.89 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	7.06×10^{-03}	4.28 x 10 ⁻⁰²	2.89 x 10 ⁻⁰²	8.14 x 10 ⁻⁰²	3.07 × 10 ⁻⁰³	2.74 x 10 ⁻⁰⁷	2.80 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	8.54 × 10 ⁻⁰³	1.22 x 10 ⁻⁰³	4.26 x 10 ⁻⁰³	1.71 x 10 ⁻⁰²
			1			1	1		se Case		1		1	1		1	
	Cobalt	9.58 × 10 ⁻⁰⁴	5.91 × 10 ⁻⁰⁵		$0.00 \times 10^{+00}$											4.78 × 10 ⁻⁰²	
	Copper	3.25 × 10 ⁻⁰⁵	1.93 × 10 ⁻⁰⁶	2.69 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	1.69 × 10 ⁻⁰³	1.06 × 10 ⁻⁰³	9.70 × 10 ⁻⁰²	9.98 x 10 ⁻⁰²	3.25 × 10 ⁻⁰⁵	1.93 × 10 ⁻⁰⁶	2.69 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	1.69 × 10 ⁻⁰³	1.06 × 10 ⁻⁰³	9.70 × 10 ⁻⁰²	9.98 x 10 ⁻⁰²
	Molybdenum	1.47 × 10 ⁻⁰⁴	8.46 × 10 ⁻⁰⁶	1.29 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.79 × 10 ⁻⁰⁶	7.01 × 10 ⁻⁰⁴	2.64 × 10 ⁻⁰¹	2.64 x 10 ⁻⁰¹	1.47 × 10 ⁻⁰⁴	8.46 × 10 ⁻⁰⁶	1.29 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.79 × 10 ⁻⁰⁶	7.01 × 10 ⁻⁰⁴	2.64 × 10 ⁻⁰¹	2.64 x 10 ⁻⁰¹
	Uranium	2.75 × 10 ⁻⁰³	5.64 × 10 ⁻⁰⁴	1.47 x 10 ⁻⁰³	$0.00 \times 10^{+00}$	6.49×10^{-03}	7.57 × 10 ⁻⁰²	1.70 × 10 ⁻⁰¹	2.57 x 10 ⁻⁰¹	2.75 × 10 ⁻⁰³	5.64 × 10 ⁻⁰⁴	1.47 x 10 ⁻⁰³	$0.00 \times 10^{+00}$	6.49 × 10 ⁻⁰³	7.57 × 10 ⁻⁰²	1.70 × 10 ⁻⁰¹	2.57 x 10 ⁻⁰¹
Subsistence				06					Incremental Pro	-							
Harvester One-	Cobalt	8.35 × 10 ⁻⁰⁵	8.54 x 10 ⁻⁰⁸	5.18 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	1.86 × 10 ⁻⁰⁴	2.27 x 10 ⁻⁰⁵	6.97 x 10 ⁻⁰⁴	9.95 x 10 ⁻⁰⁴	4.21 × 10 ⁻⁰⁴	4.63 x 10 ⁻⁰⁸	2.91 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	9.56 × 10 ⁻⁰⁴	2.22 x 10 ⁻⁰⁶	3.02 x 10 ⁻⁰³	4.43 x 10 ⁻⁰³
Year-Old	Copper	1.54 × 10 ⁻⁰⁶	5.22 x 10 ⁻⁰⁹	1.12 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	7.99 × 10 ⁻⁰⁵	4.17 x 10 ⁻⁰⁶	2.89 x 10 ⁻⁰⁴	3.74 x 10 ⁻⁰⁴	1.51 × 10 ⁻⁰⁵	3.49 x 10 ⁻¹⁰	1.25 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	7.89 × 10 ⁻⁰⁴	1.92 x 10 ⁻⁰⁷	7.25 x 10 ⁻⁰⁴	1.53 x 10 ⁻⁰³
(Patterson Lake South Arm)	Molybdenum	7.16×10^{-05}	2.41 x 10 ⁻⁰⁸	5.56 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	1.33 × 10 ⁻⁰⁶	1.82 x 10 ⁻⁰⁵	2.17 x 10 ⁻⁰⁴	3.14 x 10 ⁻⁰⁴	9.11 × 10 ⁻⁰⁴	0.00 × 10 ⁺⁰⁰	7.98 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	1.73 × 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	9.41 × 10 ⁻⁰⁴	1.95 x 10 ⁻⁰³
South Alling	Uranium	1.04 × 10 ⁻⁰³	1.71 x 10 ⁻⁰⁴	4.42 x 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	2.43 × 10 ⁻⁰³	6.44 x 10 ⁻⁰²	2.02 x 10 ⁻⁰²	8.87 x 10 ⁻⁰²	3.42 × 10 ⁻⁰³	1.79 x 10 ⁻⁰⁵	1.83 x 10 ⁻⁰³	$0.00 \times 10^{+00}$	8.09 × 10 ⁻⁰³	2.41 x 10 ⁻⁰³	3.14 x 10 ⁻⁰³	1.89 x 10 ⁻⁰²
	Cabalt	0.60 10-05	0.54 10-08	F 40 x 10-06	0.00 10+00	1.94 × 10 ⁻⁰⁴		7.24 x 10 ⁻⁰⁴	- Incremental I	-	4.62 .: 10-08	4.46 10-05	0.00 .: 10+00	1 47 10-03	2 22 10-06	4.6.4 -: 10-03	6.00 1: 10-03
	Copper	8.69 × 10 ⁻⁰⁵	8.54 x 10 ⁻⁰⁸	5.40 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	-	2.27 x 10 ⁻⁰⁵		1.03 x 10 ⁻⁰³	6.45×10^{-04}	4.63 x 10 ⁻⁰⁸ 3.49 x 10 ⁻¹⁰	4.46 x 10 ⁻⁰⁵ 1.89 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	1.47×10^{-03}	2.22 x 10 ⁻⁰⁶	4.64 x 10 ⁻⁰³	6.80 x 10 ⁻⁰³
	Copper	1.58×10^{-06} 2.17×10^{-04}	5.22 x 10 ⁻⁰⁹ 2.41 x 10 ⁻⁰⁸	1.15 x 10 ⁻⁰⁷ 1.73 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	8.23×10^{-05} 4.05×10^{-06}	4.17 x 10 ⁻⁰⁶ 1.82 x 10 ⁻⁰⁵	2.91 x 10 ⁻⁰⁴ 3.70 x 10 ⁻⁰⁴	3.79 x 10 ⁻⁰⁴ 6.27 x 10 ⁻⁰⁴	2.27×10^{-05} 3.50×10^{-03}	$3.49 \times 10^{+00}$ $0.00 \times 10^{+00}$	3.06 x 10 ⁻⁰⁴	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	1.19×10^{-03} 6.63×10^{-05}	1.92×10^{-07} $0.00 \times 10^{+00}$	1.09×10^{-03} 3.63×10^{-03}	2.30 x 10 ⁻⁰³
	Molybdenum	2.17 × 10 ° 4 2.86 × 10 ° 03	2.41 x 10 °° 1.71 x 10 °°	1.73 x 10 ° 3 1.24 x 10 ° 3	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	4.05×10^{-03} 6.69×10^{-03}	6.44 x 10 ⁻⁰²	2.10 x 10 ⁻⁰²	9.64 x 10 ⁻⁰²	3.50×10^{-03} 3.43×10^{-03}	0.00×10^{-05} 1.79×10^{-05}	1.83 x 10 ⁻⁰³	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	8.10×10^{-03}	2.41 x 10 ⁻⁰³	3.63 × 10 ° 3 3.14 x 10 ° 3	7.50 x 10 ⁻⁰³ 1.89 x 10 ⁻⁰²
	Uranium	2.00 × 10 ³³	1.71 X 10 ° 4	1.24 X 10 °	0.00 × 10 · 30	0.03 × 10 °3	0.44 X 10 °2	2.10 X 10 °2	9.04 X 10 °	3.43 × 10 °	1.79 × 10 °	1.05 X 10 °3	0.00 × 10.50	0.10 × 10 3	2.41 X 10 °3	5.14 X 10 °	1.09 X 10 °-

					Project Life	span HQs							Far Futi	ure HQs			
	СОРС	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
	C L II	0.50 40.04	0.04 4.0-07	1.04 10-06	0.00 40+00	2.20 40-02	4.44.40-02		se Case	0.50 40-04	0.04 40-07	1.04 10.06	0.00 40+00	2.20 40-02	4.4.4.0-03	2.44 40-02	2.00 40:02
	Cobalt	8.58 × 10 ⁻⁰⁴	9.04 × 10 ⁻⁰⁷	1.01 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	2.30 × 10 ⁻⁰³	1.44 × 10 ⁻⁰³	2.44 × 10 ⁻⁰²	2.90 x 10 ⁻⁰²	8.58 × 10 ⁻⁰⁴	9.04 × 10 ⁻⁰⁷	1.01 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	2.30 × 10 ⁻⁰³	1.44 × 10 ⁻⁰³	2.44 × 10 ⁻⁰²	2.90 x 10 ⁻⁰²
	Copper	2.91 × 10 ⁻⁰⁵ 1.08 × 10 ⁻⁰⁴	2.95×10^{-08} 1.06×10^{-07}	4.12 x 10 ⁻⁰⁸ 1.62 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	1.79×10^{-03} 2.42×10^{-06}	5.38×10^{-04} 2.92×10^{-04}	4.12×10^{-02} 6.98×10^{-02}	4.35 x 10 ⁻⁰² 7.02 x 10 ⁻⁰²	2.91×10^{-05} 1.08×10^{-04}	2.95×10^{-08} 1.06×10^{-07}	4.12 x 10 ⁻⁰⁸ 1.62 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	1.79×10^{-03} 2.42×10^{-06}	5.38×10^{-04} 2.92×10^{-04}	4.12×10^{-02} 6.98×10^{-02}	4.35 x 10 ⁻⁰² 7.02 x 10 ⁻⁰²
	Molybdenum Uranium	2.46 × 10 ⁻⁰³	8.62 × 10 ⁻⁰⁶	2.24 x 10 ⁻⁰⁵	$0.00 \times 10^{-0.00}$	6.85 × 10 ⁻⁰³	3.84 × 10 ⁻⁰²	8.20 × 10 ⁻⁰²	1.30 x 10 ⁻⁰¹	2.46×10^{-03}	8.62 × 10 ⁻⁰⁶	2.24 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	6.85 × 10 ⁻⁰³	3.84 × 10 ⁻⁰²	8.20 × 10 ⁻⁰²	1.30 x 10 ⁻⁰¹
	Oramium	2.40 × 10	8.02 × 10	2.24 X 10	0.00 × 10	0.03 × 10			Incremental Pro	4	8.02 × 10	2.24 X 10	0.00 × 10	0.83 × 10	3.04 × 10	8.20 × 10	1.30 X 10
Cubaiatanaa	Cobalt	3.89 × 10 ⁻⁰⁵	1.31 x 10 ⁻⁰⁹	4.13 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	1.02 × 10 ⁻⁰⁴	1.61 x 10 ⁻⁰⁵	2.28 x 10 ⁻⁰⁴	3.85 x 10 ⁻⁰⁴	1.99 × 10 ⁻⁰⁴	7.08 x 10 ⁻¹⁰	2.36 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	5.34 × 10 ⁻⁰⁴	1.13 x 10 ⁻⁰⁶	1.02 x 10 ⁻⁰³	1.76 x 10 ⁻⁰³
Subsistence Harvester (Beet	Copper	7.07 × 10 ⁻⁰⁷	5.33 x 10 ⁻¹¹	8.78 x 10 ⁻¹⁰	0.00 × 10 ⁺⁰⁰	4.32 × 10 ⁻⁰⁵	1.62 x 10 ⁻⁰⁶	5.49 x 10 ⁻⁰⁵	1.00 x 10 ⁻⁰⁴	7.09 × 10 ⁻⁰⁶	3.56 x 10 ⁻¹²	1.00 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	4.35 × 10 ⁻⁰⁴	6.50 x 10 ⁻⁰⁸	2.76 x 10 ⁻⁰⁴	7.18 x 10 ⁻⁰⁴
Lake)	Molybdenum	2.70 × 10 ⁻⁰⁵	1.52 x 10 ⁻¹⁰	3.58 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	5.89 × 10 ⁻⁰⁷	5.41 x 10 ⁻⁰⁶	2.82 x 10 ⁻⁰⁵	6.12 x 10 ⁻⁰⁵	3.49 × 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	5.22 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	7.79 × 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.71 × 10 ⁻⁰⁴	6.29 x 10 ⁻⁰⁴
	Uranium	3.43 × 10 ⁻⁰⁴	1.54 x 10 ⁻⁰⁶	2.48 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	9.44 × 10 ⁻⁰⁴	2.52 x 10 ⁻⁰²	1.05 x 10 ⁻⁰²	3.70 x 10 ⁻⁰²	1.14 × 10 ⁻⁰³	1.62 x 10 ⁻⁰⁷	1.04 x 10 ⁻⁰⁵	0.00 × 10 ⁺⁰⁰	3.17 × 10 ⁻⁰³	7.20 x 10 ⁻⁰⁴	1.44 x 10 ⁻⁰³	6.48 x 10 ⁻⁰³
									– Incremental I								
	Cobalt	4.05 × 10 ⁻⁰⁵	1.31 x 10 ⁻⁰⁹	4.30 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	1.06 × 10 ⁻⁰⁴	1.61 x 10 ⁻⁰⁵	2.36 x 10 ⁻⁰⁴	3.99 x 10 ⁻⁰⁴	3.06 × 10 ⁻⁰⁴	7.08 x 10 ⁻¹⁰	3.61 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	8.18 × 10 ⁻⁰⁴	1.13 x 10 ⁻⁰⁶	1.57 x 10 ⁻⁰³	2.69 x 10 ⁻⁰³
	Copper	7.28 × 10 ⁻⁰⁷	5.33 x 10 ⁻¹¹	9.05 x 10 ⁻¹⁰	0.00 × 10 ⁺⁰⁰	4.45×10^{-05}	1.62 x 10 ⁻⁰⁶	5.56 x 10 ⁻⁰⁵	1.02 x 10 ⁻⁰⁴	1.06 × 10 ⁻⁰⁵	3.56 x 10 ⁻¹²	1.51 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	6.54 × 10 ⁻⁰⁴	6.50 x 10 ⁻⁰⁸	4.14 x 10 ⁻⁰⁴	1.08 x 10 ⁻⁰³
	Molybdenum	8.25 × 10 ⁻⁰⁵	1.52 x 10 ⁻¹⁰	1.11 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	1.81 × 10 ⁻⁰⁶	5.41 x 10 ⁻⁰⁶	6.50 x 10 ⁻⁰⁵	1.55 x 10 ⁻⁰⁴	1.34 × 10 ⁻⁰³	$0.00 \times 10^{+00}$	2.01 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.99 × 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	1.04 × 10 ⁻⁰³	2.41 x 10 ⁻⁰³
	Uranium	9.43 × 10 ⁻⁰⁴	1.54 x 10 ⁻⁰⁶	6.95 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	2.59 × 10 ⁻⁰³	2.52 x 10 ⁻⁰²	1.08 x 10 ⁻⁰²	3.96 x 10 ⁻⁰²	1.14 × 10 ⁻⁰³	1.62 x 10 ⁻⁰⁷	1.04 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	3.18 × 10 ⁻⁰³	7.20 x 10 ⁻⁰⁴	1.44 x 10 ⁻⁰³	6.48 x 10 ⁻⁰³
								Bas	se Case	_							
	Cobalt	9.58 × 10 ⁻⁰⁴	5.91 × 10 ⁻⁰⁵	6.62 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.18 × 10 ⁻⁰³	2.84 × 10 ⁻⁰³	4.78 × 10 ⁻⁰²	5.39 x 10 ⁻⁰²	9.58 × 10 ⁻⁰⁴	5.91 × 10 ⁻⁰⁵	6.62 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.18 × 10 ⁻⁰³	2.84×10^{-03}	4.78 × 10 ⁻⁰²	5.39 x 10 ⁻⁰²
	Copper	3.25 × 10 ⁻⁰⁵	1.93 × 10 ⁻⁰⁶	2.69 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	1.69×10^{-03}	1.06 × 10 ⁻⁰³	9.70 × 10 ⁻⁰²	9.98 x 10 ⁻⁰²	3.25 × 10 ⁻⁰⁵	1.93×10^{-06}	2.69 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	1.69 × 10 ⁻⁰³	1.06×10^{-03}	9.70 × 10 ⁻⁰²	9.98 x 10 ⁻⁰²
	Molybdenum	1.47 × 10 ⁻⁰⁴	8.46×10^{-06}	1.29 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.79×10^{-06}	7.01 × 10 ⁻⁰⁴	2.64 × 10 ⁻⁰¹	2.64 x 10 ⁻⁰¹	1.47 × 10 ⁻⁰⁴	8.46 × 10 ⁻⁰⁶	1.29 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.79 × 10 ⁻⁰⁶	7.01×10^{-04}	2.64 × 10 ⁻⁰¹	2.64 x 10 ⁻⁰¹
	Uranium	2.75 × 10 ⁻⁰³	5.64 × 10 ⁻⁰⁴	1.47 x 10 ⁻⁰³	$0.00 \times 10^{+00}$	6.49×10^{-03}	7.57 × 10 ⁻⁰²	1.70×10^{-01}	2.57 x 10 ⁻⁰¹	2.75 × 10 ⁻⁰³	5.64 × 10 ⁻⁰⁴	1.47 x 10 ⁻⁰³	$0.00 \times 10^{+00}$	6.49 × 10 ⁻⁰³	7.57×10^{-02}	1.70 × 10 ⁻⁰¹	2.57 x 10 ⁻⁰¹
								ication Case -	Incremental Pro	•							
Subsistence Harvester One-	Cobalt	4.34 × 10 ⁻⁰⁵	8.54 x 10 ⁻⁰⁸	2.70 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	9.69 × 10 ⁻⁰⁵	2.27 x 10 ⁻⁰⁵	2.32 x 10 ⁻⁰⁴	3.98 x 10 ⁻⁰⁴	2.23 × 10 ⁻⁰⁴	4.63 x 10 ⁻⁰⁸	1.54 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	5.06 × 10 ⁻⁰⁴	2.22 x 10 ⁻⁰⁶	1.04 x 10 ⁻⁰³	1.79 x 10 ⁻⁰³
Year-Old (Beet	Copper	7.89 × 10 ⁻⁰⁷	3.48 x 10 ⁻⁰⁹	5.74 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	4.10×10^{-05}	2.78 x 10 ⁻⁰⁶	5.16 x 10 ⁻⁰⁵	9.62 x 10 ⁻⁰⁵	7.91 × 10 ⁻⁰⁶	2.33 x 10 ⁻¹⁰	6.56 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	4.12 × 10 ⁻⁰⁴	1.28 x 10 ⁻⁰⁷	2.54 x 10 ⁻⁰⁴	6.75 x 10 ⁻⁰⁴
Lake)	Molybdenum	3.67 × 10 ⁻⁰⁵	1.21 x 10 ⁻⁰⁸	2.85 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	6.80×10^{-07}	9.10 x 10 ⁻⁰⁶	3.46 x 10 ⁻⁰⁵	8.39 x 10 ⁻⁰⁵	4.74 × 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	4.16 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	8.99 × 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	3.32 × 10 ⁻⁰⁴	8.57 x 10 ⁻⁰⁴
	Uranium	3.83 × 10 ⁻⁰⁴	1.01 x 10 ⁻⁰⁴	1.62 x 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	8.95 × 10 ⁻⁰⁴	3.80 x 10 ⁻⁰²	7.40 x 10 ⁻⁰³	4.69 x 10 ⁻⁰²	1.27×10^{-03}	1.06 x 10 ⁻⁰⁵	6.79 x 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	3.01 × 10 ⁻⁰³	1.42 x 10 ⁻⁰³	1.04 x 10 ⁻⁰³	7.43 x 10 ⁻⁰³
		05							- Incremental I	_				0.4	06		
	Cobalt	4.52 × 10 ⁻⁰⁵	8.54 x 10 ⁻⁰⁸	2.81 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	1.01 × 10 ⁻⁰⁴	2.27 x 10 ⁻⁰⁵	2.40 x 10 ⁻⁰⁴	4.12 x 10 ⁻⁰⁴	3.41 × 10 ⁻⁰⁴	4.63 x 10 ⁻⁰⁸	2.36 x 10 ⁻⁰⁵	0.00 × 10 ⁺⁰⁰	7.75 × 10 ⁻⁰⁴	2.22 x 10 ⁻⁰⁶	1.59 x 10 ⁻⁰³	2.73 x 10 ⁻⁰³
	Copper	8.13 × 10 ⁻⁰⁷	3.48 x 10 ⁻⁰⁹	5.91 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	4.22 × 10 ⁻⁰⁵	2.78 x 10 ⁻⁰⁶	5.23 x 10 ⁻⁰⁵	9.81 x 10 ⁻⁰⁵	1.19 × 10 ⁻⁰⁵	2.33 x 10 ⁻¹⁰	9.86 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	6.20 × 10 ⁻⁰⁴	1.28 x 10 ⁻⁰⁷	3.82 x 10 ⁻⁰⁴	1.01 x 10 ⁻⁰³
	Molybdenum	1.12 × 10 ⁻⁰⁴	1.21 x 10 ⁻⁰⁸	8.86 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	2.09 × 10 ⁻⁰⁶	9.10 x 10 ⁻⁰⁶	7.97 x 10 ⁻⁰⁵	2.12 x 10 ⁻⁰⁴	1.82 × 10 ⁻⁰³	0.00 × 10 ⁺⁰⁰	1.60 x 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	3.45 × 10 ⁻⁰⁵	0.00 × 10 ⁺⁰⁰	1.28 × 10 ⁻⁰³	3.29 x 10 ⁻⁰³
	Uranium	1.05 × 10 ⁻⁰³	1.01 x 10 ⁻⁰⁴	4.54 x 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	2.46 × 10 ⁻⁰³	3.80 x 10 ⁻⁰²	7.65 x 10 ⁻⁰³	4.97 x 10 ⁻⁰²	1.27×10^{-03}	1.06 x 10 ⁻⁰⁵	6.79 x 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	3.01 × 10 ⁻⁰³	1.42 x 10 ⁻⁰³	1.04 x 10 ⁻⁰³	7.44 x 10 ⁻⁰³
	Cobolt	8.58 × 10 ⁻⁰⁴	0.04 .: 10-07	1.01 x 10 ⁻⁰⁶	0.00 .: 10+00	2.30 × 10 ⁻⁰³	1.44 . 10-03		se Case	0.00 10-04	9.04 × 10 ⁻⁰⁷	1.01 10-06	0.00 .: 10+00	2.30 × 10 ⁻⁰³	1.44 10-03	2.44 × 10 ⁻⁰²	2.90 x 10 ⁻⁰²
	Copper	8.58 × 10 ⁻⁰⁴ 2.91 × 10 ⁻⁰⁵	9.04×10^{-07} 2.95×10^{-08}	4.12 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	2.30 × 10 ⁻⁰³	1.44×10^{-03} 5.38×10^{-04}	2.44×10^{-02} 4.12×10^{-02}	4.35 x 10 ⁻⁰²	8.58 × 10 ⁻⁰⁴ 2.91 × 10 ⁻⁰⁵	9.04×10^{-07} 2.95×10^{-08}	4.12 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	2.30×10^{-03} 1.79×10^{-03}	1.44×10^{-03} 5.38×10^{-04}	2.44×10^{-02} 4.12×10^{-02}	4.35 x 10 ⁻⁰²
	Copper	1.08 × 10 ⁻⁰⁴	1.06 × 10 ⁻⁰⁷	+	$0.00 \times 10^{-0.00}$	2.42 × 10 ⁻⁰⁶	2.92 × 10 ⁻⁰⁴	6.98 × 10 ⁻⁰²		1.08×10^{-04}	1.06 × 10 ⁻⁰⁷	1.62 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	2.42 × 10 ⁻⁰⁶	2.92 × 10 ⁻⁰⁴	6.98 × 10 ⁻⁰²	
	Molybdenum		8.62 × 10 ⁻⁰⁶	1.62 x 10 ⁻⁰⁷ 2.24 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	6.85×10^{-03}	3.84 × 10 ⁻⁰²	8.20 × 10 ⁻⁰²	7.02 x 10 ⁻⁰²	2.46×10^{-03}	8.62 × 10 ⁻⁰⁶		$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	6.85 × 10 ⁻⁰³		8.20 × 10 ⁻⁰²	7.02 x 10 ⁻⁰²
	Uranium	2.46 × 10 ⁻⁰³	8.62 × 10 °°	2.24 X 10 ° 3	0.00 × 10 · · · ·	6.85 × 10 ° 3			1.30 x 10 ⁻⁰¹ Incremental Pro		8.62 × 10 °°	2.24 x 10 ⁻⁰⁵	0.00 × 10***	6.85 × 10 °	3.84 × 10 °	8.20 × 10 °2	1.30 x 10 ⁻⁰¹
	Cobalt	4.05 × 10 ⁻⁰⁶	1.31 x 10 ⁻⁰⁹	4.26 x 10 ⁻⁰⁹	0.00 × 10 ⁺⁰⁰	1.06 × 10 ⁻⁰⁵	1.61 x 10 ⁻⁰⁵	5.65 x 10 ⁻⁰⁵	8.73 x 10 ⁻⁰⁵	2.04 × 10 ⁻⁰⁵	7.08 x 10 ⁻¹⁰	2.42 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	5.47 × 10 ⁻⁰⁵	1.13 x 10 ⁻⁰⁶	1.10 x 10 ⁻⁰⁴	1.87 x 10 ⁻⁰⁴
Subsistence Harvester (Lloyd	Copper	7.33 × 10 ⁻⁰⁸	2.66 x 10 ⁻¹¹	9.02 x 10 ⁻¹¹	$0.00 \times 10^{+00}$	4.48 × 10 ⁻⁰⁶	8.11 x 10 ⁻⁰⁷	1.73 x 10 ⁻⁰⁵	2.27 x 10 ⁻⁰⁵	7.23×10^{-07}	1.78 x 10 ⁻¹²	1.02 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	4.44 × 10 ⁻⁰⁵	3.25 x 10 ⁻⁰⁸	2.84 x 10 ⁻⁰⁵	7.36 x 10 ⁻⁰⁵
Lake)	Molybdenum	2.79 × 10 ⁻⁰⁶	1.52 x 10 ⁻¹⁰	3.67 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	6.08 × 10 ⁻⁰⁸	5.41 x 10 ⁻⁰⁶	1.73 x 10 1.22 x 10 ⁻⁰⁵	2.27 x 10 2.05 x 10 ⁻⁰⁵	3.56×10^{-05}	$0.00 \times 10^{+00}$	5.33 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	7.94 × 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.77 × 10 ⁻⁰⁵	6.41 x 10 ⁻⁰⁵
	Uranium	3.13 × 10 ⁻⁰⁵	5.85 x 10 ⁻⁰⁸	2.25 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	8.61 × 10 ⁻⁰⁵	9.58 x 10 ⁻⁰⁴	4.07 x 10 ⁻⁰⁴	1.48 x 10 ⁻⁰³	1.03 × 10 ⁻⁰⁴	6.15 x 10 ⁻⁰⁹	9.41 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.87×10^{-04}	2.74 x 10 ⁻⁰⁵	9.62 x 10 ⁻⁰⁵	5.15 x 10 ⁻⁰⁴
	Oraniani	3.13 × 10	3.03 × 10	2.23 X 10	0.00 % 10	0.01 × 10			- Incremental F		0.13 X 10	J. 71 X 10	0.00 × 10	2.07 × 10	2.73 10	3.02 X 10	3.13 X 10
	Cobalt	4.21 × 10 ⁻⁰⁶	1.31 x 10 ⁻⁰⁹	4.44 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	1.10 × 10 ⁻⁰⁵	1.61 x 10 ⁻⁰⁵	5.73 x 10 ⁻⁰⁵	8.86 x 10 ⁻⁰⁵	3.13 × 10 ⁻⁰⁵	7.08 x 10 ⁻¹⁰	3.70 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	8.38 × 10 ⁻⁰⁵	1.13 x 10 ⁻⁰⁶	1.66 x 10 ⁻⁰⁴	2.82 x 10 ⁻⁰⁴
	Copper	7.54 × 10 ⁻⁰⁸	2.66 x 10 ⁻¹¹	9.29 x 10 ⁻¹¹	$0.00 \times 10^{+00}$	4.61 × 10 ⁻⁰⁶	8.11 x 10 ⁻⁰⁷	1.74 x 10 ⁻⁰⁵	2.29 x 10 ⁻⁰⁵	1.09 × 10 ⁻⁰⁶	1.78 x 10 ⁻¹²	1.54 x 10 ⁻⁰⁹	0.00 × 10 ⁺⁰⁰	6.67 × 10 ⁻⁰⁵	3.25 x 10 ⁻⁰⁸	4.25 x 10 ⁻⁰⁵	1.10 x 10 ⁻⁰⁴
	Molybdenum	8.51 × 10 ⁻⁰⁶	1.52 x 10 ⁻¹⁰	1.14 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	1.86 × 10 ⁻⁰⁷	5.41 x 10 ⁻⁰⁶	1.60 x 10 ⁻⁰⁵	3.01 x 10 ⁻⁰⁵	1.37 × 10 ⁻⁰⁴	0.00 × 10 ⁺⁰⁰	2.05 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	3.05 × 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	1.06 × 10 ⁻⁰⁴	2.46 x 10 ⁻⁰⁴
	Uranium	8.67 × 10 ⁻⁰⁵	5.85 x 10 ⁻⁰⁸	6.31 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	2.38 × 10 ⁻⁰⁴	9.58 x 10 ⁻⁰⁴	4.37 x 10 ⁻⁰⁴	1.72 x 10 ⁻⁰³	1.03 × 10 ⁻⁰⁴	6.15 x 10 ⁻⁰⁹	9.41 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	2.88 × 10 ⁻⁰⁴	2.74 x 10 ⁻⁰⁵	9.62 x 10 ⁻⁰⁵	5.15 x 10 ⁻⁰⁴
		1 2.2	1				1	1	1 = 3	1	1	1	1	1	1	1	1

Coper Molybdenum Uranium	9.58 × 10 ⁻⁰⁴ 3.25 × 10 ⁻⁰⁵ 1.47 × 10 ⁻⁰⁴ 2.75 × 10 ⁻⁰³	Soil (Internal) 5.91 × 10 ⁻⁰⁵ 1.93 × 10 ⁻⁰⁶	Sediment (Internal)	Project Lifes Aquatic Plants	Aquatic Animals	Terrestrial	Terrestrial	Total by	Water	Soil	Sediment	Aquatic	Aquatic	Terrestrial	Terrestrial	Table
Copper Molybdenum	9.58 × 10 ⁻⁰⁴ 3.25 × 10 ⁻⁰⁵ 1.47 × 10 ⁻⁰⁴	5.91 × 10 ⁻⁰⁵ 1.93 × 10 ⁻⁰⁶		Plants	Animals					00		Aquatic	Aquatic	i ci i coti iai	i ei i esti iai	Total by
Copper Molybdenum	3.25×10^{-05} 1.47×10^{-04}	1.93 × 10 ⁻⁰⁶	6.62 x 10 ⁻⁰⁵			Plants	Animals	СОРС	(Internal)	(Internal)	(Internal)	Plants	Animals	Plants	Animals	СОРС
Copper Molybdenum	3.25×10^{-05} 1.47×10^{-04}	1.93 × 10 ⁻⁰⁶	0.02 X 10	$0.00 \times 10^{+00}$	2.18 × 10 ⁻⁰³	2.84 × 10 ⁻⁰³	4.78 × 10 ⁻⁰²	5.39 x 10 ⁻⁰²	9.58 × 10 ⁻⁰⁴	5.91 × 10 ⁻⁰⁵	6.62 x 10 ⁻⁰⁵	0.00 × 10 ⁺⁰⁰	2.18 × 10 ⁻⁰³	2.84 × 10 ⁻⁰³	4.78 × 10 ⁻⁰²	5.39 x 10 ⁻⁰²
Molybdenum	1.47 × 10 ⁻⁰⁴	+	2.69 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	1.69 × 10 ⁻⁰³	1.06 × 10 ⁻⁰³	9.70 × 10 ⁻⁰²	9.98 x 10 ⁻⁰²	3.25 × 10 ⁻⁰⁵	1.93 × 10 ⁻⁰⁶	2.69 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	1.69 × 10 ⁻⁰³	1.06 × 10 ⁻⁰³	9.70 × 10 ⁻⁰²	9.98 x 10 ⁻⁰²
	+	8.46×10^{-06}	1.29 x 10 ⁻⁰⁵	0.00 × 10 ⁺⁰⁰	2.79 × 10 ⁻⁰⁶	7.01 × 10 ⁻⁰⁴	2.64 × 10 ⁻⁰¹	2.64 x 10 ⁻⁰¹	1.47×10^{-04}	8.46 × 10 ⁻⁰⁶	1.29 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.79 × 10 ⁻⁰⁶	7.00×10^{-04}	2.64 × 10 ⁻⁰¹	2.64 x 10 ⁻⁰¹
Clanian	E.75 × 10	5.64 × 10 ⁻⁰⁴	1.47 x 10 ⁻⁰³	0.00 × 10 ⁺⁰⁰	6.49 × 10 ⁻⁰³	7.57×10^{-02}	1.70 × 10 ⁻⁰¹	2.57 x 10 ⁻⁰¹	2.75×10^{-03}	5.64 × 10 ⁻⁰⁴	1.47 x 10 ⁻⁰³	$0.00 \times 10^{+00}$	6.49×10^{-03}	7.57×10^{-02}	1.70 × 10 ⁻⁰¹	2.57 x 10 ⁻⁰¹
		3.04 × 10	1.47 X 10	0.00 × 10	0.43 × 10		ication Case – I			3.04 × 10	1.47 X 10	0.00 × 10	0.43 × 10	7.57 × 10	1.70 × 10	2.37 X 10
Subsistence Cobalt	4.52 × 10 ⁻⁰⁶	8.54 x 10 ⁻⁰⁸	2.78 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	1.01 × 10 ⁻⁰⁵	2.27 x 10 ⁻⁰⁵	5.72 x 10 ⁻⁰⁵	9.49 x 10 ⁻⁰⁵	2.28 × 10 ⁻⁰⁵	4.63 x 10 ⁻⁰⁸	1.58 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	5.19 × 10 ⁻⁰⁵	2.22 x 10 ⁻⁰⁶	1.12 x 10 ⁻⁰⁴	1.91 x 10 ⁻⁰⁴
Harvester One-	8.18 × 10 ⁻⁰⁸	1.74 x 10 ⁻⁰⁹	5.89 x 10 ⁻⁰⁹	0.00 × 10 ⁺⁰⁰	4.25 × 10 ⁻⁰⁶	1.39 x 10 ⁻⁰⁶	1.63 x 10 ⁻⁰⁵	2.21 x 10 ⁻⁰⁵	8.08 × 10 ⁻⁰⁷	1.16 x 10 ⁻¹⁰	6.70 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	4.21 × 10 ⁻⁰⁵	6.40 x 10 ⁻⁰⁸	2.62 x 10 ⁻⁰⁵	6.93 x 10 ⁻⁰⁵
Year-Old (Lloyd Lake) Molybdenum	3.79 × 10 ⁻⁰⁶	1.21 x 10 ⁻⁰⁸	2.92 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	7.01 × 10 ⁻⁰⁸	9.10 x 10 ⁻⁰⁶	1.50 x 10 ⁻⁰⁵	2.83 x 10 ⁻⁰⁵	4.84×10^{-05}	0.00 × 10 ⁺⁰⁰	4.24 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	9.16 × 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	3.39 × 10 ⁻⁰⁵	8.74 x 10 ⁻⁰⁵
Uranium	3.50 × 10 ⁻⁰⁵	3.82 x 10 ⁻⁰⁶	1.47 x 10 ⁻⁰⁵	0.00 × 10 ⁺⁰⁰	8.16 × 10 ⁻⁰⁵	1.44 x 10 ⁻⁰³	2.89 x 10 ⁻⁰⁴	1.87 x 10 ⁻⁰³	1.15 × 10 ⁻⁰⁴	4.02 x 10 ⁻⁰⁷	6.15 x 10 ⁻⁰⁵	0.00 × 10 ⁺⁰⁰	2.72 × 10 ⁻⁰⁴	5.39 x 10 ⁻⁰⁵	7.05 x 10 ⁻⁰⁵	5.74 x 10 ⁻⁰⁴
	<u> </u>	<u>'</u>				Upper E	Sound Scenario	- Incremental F	Project Risk							
Cobalt	4.70 × 10 ⁻⁰⁶	8.54 x 10 ⁻⁰⁸	2.90 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	1.05 × 10 ⁻⁰⁵	2.27 x 10 ⁻⁰⁵	5.80 x 10 ⁻⁰⁵	9.63 x 10 ⁻⁰⁵	3.50×10^{-05}	4.63 x 10 ⁻⁰⁸	2.42 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	7.94 × 10 ⁻⁰⁵	2.22 x 10 ⁻⁰⁶	1.68 x 10 ⁻⁰⁴	2.87 x 10 ⁻⁰⁴
Copper	8.42 × 10 ⁻⁰⁸	1.74 x 10 ⁻⁰⁹	6.07 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	4.37 × 10 ⁻⁰⁶	1.39 x 10 ⁻⁰⁶	1.64 x 10 ⁻⁰⁵	2.23 x 10 ⁻⁰⁵	1.21×10^{-06}	1.16 x 10 ⁻¹⁰	1.01 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	6.33 × 10 ⁻⁰⁵	6.40 x 10 ⁻⁰⁸	3.92 x 10 ⁻⁰⁵	1.04 x 10 ⁻⁰⁴
Molybdenum	1.16 × 10 ⁻⁰⁵	1.21 x 10 ⁻⁰⁸	9.09 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.15 × 10 ⁻⁰⁷	9.10 x 10 ⁻⁰⁶	1.96 x 10 ⁻⁰⁵	4.14 x 10 ⁻⁰⁵	1.86×10^{-04}	$0.00 \times 10^{+00}$	1.63 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	3.52×10^{-06}	$0.00 \times 10^{+00}$	1.30×10^{-04}	3.36 x 10 ⁻⁰⁴
Uranium	9.68 × 10 ⁻⁰⁵	3.82 x 10 ⁻⁰⁶	4.12 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.26 × 10 ⁻⁰⁴	1.44 x 10 ⁻⁰³	3.11 x 10 ⁻⁰⁴	2.12 x 10 ⁻⁰³	1.15×10^{-04}	4.02 x 10 ⁻⁰⁷	6.15 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.73×10^{-04}	5.39 x 10 ⁻⁰⁵	7.05 x 10 ⁻⁰⁵	5.74 x 10 ⁻⁰⁴
							Bas	e Case	_							
Cobalt	8.58 × 10 ⁻⁰⁴	9.04 × 10 ⁻⁰⁷	1.01×10^{-1}	$0.00 \times 10^{+00}$	7.27 × 10 ⁻⁰⁴	1.13×10^{-03}	1.88 × 10 ⁻⁰²	2.15 x 10 ⁻⁰²	8.58 × 10 ⁻⁰⁴	9.04 × 10 ⁻⁰⁷	1.01 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	7.27 × 10 ⁻⁰⁴	1.13 × 10 ⁻⁰³	1.88 × 10 ⁻⁰²	2.15 x 10 ⁻⁰²
Copper	2.91 × 10 ⁻⁰⁵	2.95 × 10 ⁻⁰⁸	4.12 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	5.66 × 10 ⁻⁰⁴	4.22 × 10 ⁻⁰⁴	3.86 × 10 ⁻⁰²	3.96 x 10 ⁻⁰²	2.91 × 10 ⁻⁰⁵	2.95 × 10 ⁻⁰⁸	4.12 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	5.66 × 10 ⁻⁰⁴	4.22 × 10 ⁻⁰⁴	3.86 × 10 ⁻⁰²	3.96 x 10 ⁻⁰²
Molybdenum	1.08 × 10 ⁻⁰⁴	1.06 × 10 ⁻⁰⁷	1.62 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	7.66 × 10 ⁻⁰⁷	2.29 × 10 ⁻⁰⁴	8.60 × 10 ⁻⁰²	8.63 x 10 ⁻⁰²	1.08 × 10 ⁻⁰⁴	1.06 × 10 ⁻⁰⁷	1.62 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	7.66 × 10 ⁻⁰⁷	2.29 × 10 ⁻⁰⁴	8.60 × 10 ⁻⁰²	8.63 x 10 ⁻⁰²
Uranium	2.46 × 10 ⁻⁰³	8.62 × 10 ⁻⁰⁶	2.24 x 10 ⁻⁰⁵	0.00 × 10 ⁺⁰⁰	2.17 × 10 ⁻⁰³	3.01 × 10 ⁻⁰²	6.79 × 10 ⁻⁰²	1.03 x 10 ⁻⁰¹	2.46×10^{-03}	8.62 × 10 ⁻⁰⁶	2.24 x 10 ⁻⁰⁵	0.00 × 10 ⁺⁰⁰	2.17 × 10 ⁻⁰³	3.01 × 10 ⁻⁰²	6.79 × 10 ⁻⁰²	1.03 x 10 ⁻⁰¹
						Appl	ication Case – I	ncremental Pro	ject Risk							
Seasonal Resident Cobalt	4.49 × 10 ⁻⁰⁵	7.84 x 10 ⁻¹⁰	4.76 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	3.74 × 10 ⁻⁰⁵	4.53 x 10 ⁻⁰⁶	1.60 x 10 ⁻⁰⁴	2.47 x 10 ⁻⁰⁴	2.26 × 10 ⁻⁰⁴	4.25 x 10 ⁻¹⁰	2.67 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	1.92 × 10 ⁻⁰⁴	5.31 x 10 ⁻⁰⁷	7.01 x 10 ⁻⁰⁴	1.12 x 10 ⁻⁰³
(Patterson Lake South Arm) Copper	8.27 × 10 ⁻⁰⁷	4.80 x 10 ⁻¹¹	1.03 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	1.60 × 10 ⁻⁰⁵	9.32 x 10 ⁻⁰⁷	6.70 x 10 ⁻⁰⁵	8.48 x 10 ⁻⁰⁵	8.13 × 10 ⁻⁰⁶	3.20 x 10 ⁻¹²	1.15 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	1.58 × 10 ⁻⁰⁴	4.58 x 10 ⁻⁰⁸	1.87 x 10 ⁻⁰⁴	3.53 x 10 ⁻⁰⁴
Molybdenum	3.16 × 10 ⁻⁰⁵	1.82 x 10 ⁻¹⁰	4.20 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	2.19 × 10 ⁻⁰⁷	2.94 x 10 ⁻⁰⁶	4.01 x 10 ⁻⁰⁵	7.48 x 10 ⁻⁰⁵	4.02×10^{-04}	$0.00 \times 10^{+00}$	6.02 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.84 × 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	1.80 × 10 ⁻⁰⁴	5.86 x 10 ⁻⁰⁴
Uranium	5.60 × 10 ⁻⁰⁴	1.57 x 10 ⁻⁰⁶	4.06 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	4.88 × 10 ⁻⁰⁴	1.34 x 10 ⁻⁰²	5.17 x 10 ⁻⁰³	1.96 x 10 ⁻⁰²	1.84×10^{-03}	1.65 x 10 ⁻⁰⁷	1.68 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	1.62 × 10 ⁻⁰³	5.75 x 10 ⁻⁰⁴	1.02 x 10 ⁻⁰³	5.08 x 10 ⁻⁰³
			ı				Bound Scenario		1 -			ı				I
Cobalt	4.67 × 10 ⁻⁰⁵	7.84 x 10 ⁻¹⁰	4.96 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	3.89 × 10 ⁻⁰⁵	4.53 x 10 ⁻⁰⁶	1.66 x 10 ⁻⁰⁴	2.56 x 10 ⁻⁰⁴	3.46 × 10 ⁻⁰⁴	4.25 x 10 ⁻¹⁰	4.10 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	2.94 × 10 ⁻⁰⁴	5.31 x 10 ⁻⁰⁷	1.08 x 10 ⁻⁰³	1.72 x 10 ⁻⁰³
Copper	8.51 × 10 ⁻⁰⁷	4.80 x 10 ⁻¹¹	1.06 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	1.65 × 10 ⁻⁰⁵	9.32 x 10 ⁻⁰⁷	6.76 x 10 ⁻⁰⁵	8.59 x 10 ⁻⁰⁵	1.22 × 10 ⁻⁰⁵	3.20 x 10 ⁻¹²	1.73 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	2.38 × 10 ⁻⁰⁴	4.58 x 10 ⁻⁰⁸	2.80 x 10 ⁻⁰⁴	5.30 x 10 ⁻⁰⁴
Molybdenum	9.59 × 10 ⁻⁰⁵	1.82 x 10 ⁻¹⁰	1.30 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	6.67 × 10 ⁻⁰⁷	2.94 x 10 ⁻⁰⁶	6.90 x 10 ⁻⁰⁵	1.69 x 10 ⁻⁰⁴	1.54 × 10 ⁻⁰³	0.00 × 10 ⁺⁰⁰	2.31 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	1.09 × 10 ⁻⁰⁵	0.00 × 10 ⁺⁰⁰	6.94 × 10 ⁻⁰⁴	2.25 x 10 ⁻⁰³
Uranium	1.54 × 10 ⁻⁰³	1.57 x 10 ⁻⁰⁶	1.14 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	1.34 × 10 ⁻⁰³	1.34 x 10 ⁻⁰²	5.49 x 10 ⁻⁰³	2.18 x 10 ⁻⁰²	1.84 × 10 ⁻⁰³	1.65 x 10 ⁻⁰⁷	1.68 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	1.62 × 10 ⁻⁰³	5.75 x 10 ⁻⁰⁴	1.02 x 10 ⁻⁰³	5.08 x 10 ⁻⁰³
Cobalt	9.58 × 10 ⁻⁰⁴	5.91 × 10 ⁻⁰⁵	6.62 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	6.89 × 10 ⁻⁰⁴	2.47 × 10 ⁻⁰³	4.13 × 10 ⁻⁰²	4.56 x 10 ⁻⁰²	9.58 × 10 ⁻⁰⁴	5.91 × 10 ⁻⁰⁵	6.62 x 10 ⁻⁰⁵	0.00 × 10 ⁺⁰⁰	6.89 × 10 ⁻⁰⁴	2.47 × 10 ⁻⁰³	4.13 × 10 ⁻⁰²	4.56 x 10 ⁻⁰²
Cobalt						9.22 × 10 ⁻⁰⁴	+			1.93 × 10 ⁻⁰⁶	2.69 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$			9.32 × 10 ⁻⁰²	
Copper	3.25×10^{-05}	1.93×10^{-06} 8.46×10^{-06}	2.69 x 10 ⁻⁰⁶ 1.29 x 10 ⁻⁰⁵	0.00 × 10 ⁺⁰⁰	5.36 × 10 ⁻⁰⁴		9.32 × 10 ⁻⁰² 2.77 × 10⁻⁰¹	9.47 x 10 ⁻⁰²	3.25×10^{-05} 1.47×10^{-04}			$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	5.36×10^{-04} 8.84×10^{-07}	9.22 × 10 ⁻⁰⁴		9.47 x 10 ⁻⁰²
Molybdenum	1.47×10^{-04} 2.75×10^{-03}	5.64 × 10 ⁻⁰⁴	1.47 x 10 ⁻⁰³	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	8.84×10^{-07} 2.06×10^{-03}	6.10×10^{-04} 6.58×10^{-02}	1.60 × 10 ⁻⁰¹	2.78 x 10 ⁻⁰¹ 2.33 x 10 ⁻⁰¹	1.47×10^{-03} 2.75×10^{-03}	8.46×10^{-06} 5.64×10^{-04}	1.29 x 10 ⁻⁰⁵ 1.47 x 10 ⁻⁰³	$0.00 \times 10^{+00}$ $0.00 \times 10^{+00}$	2.06 × 10 ⁻⁰³	6.10×10^{-04} 6.58×10^{-02}	2.77×10^{-01} 1.60×10^{-01}	2.78 x 10 ⁻⁰¹ 2.33 x 10 ⁻⁰¹
Uranium	2.75 × 10 ···	5.64 × 10 °	1.47 x 10 ···	0.00 × 10 · · ·	2.06 × 10 ···		ication Case – I			5.64 × 10 °	1.47 x 10 °°	0.00 × 10***	2.06 × 10 °°	6.58 × 10 °-	1.60 × 10 °	2.33 X 10 **
Seasonal Resident Cobalt	5.01 × 10 ⁻⁰⁵	5.12 x 10 ⁻⁰⁸	3.11 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	3.54 × 10 ⁻⁰⁵	7.54 x 10 ⁻⁰⁶	1.66 x 10 ⁻⁰⁴	2.62 x 10 ⁻⁰⁴	2.52 × 10 ⁻⁰⁴	2.78 x 10 ⁻⁰⁸	1.75 x 10 ⁻⁰⁵	0.00 × 10 ⁺⁰⁰	1.82 × 10 ⁻⁰⁴	1.16 x 10 ⁻⁰⁶	7.23 x 10 ⁻⁰⁴	1.18 x 10 ⁻⁰³
One-Year-Old Copper	9.24 × 10 ⁻⁰⁷	3.13 x 10 ⁻⁰⁹	6.72 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	1.52 × 10 ⁻⁰⁵	1.87 x 10 ⁻⁰⁶	6.82 x 10 ⁻⁰⁵	8.62 x 10 ⁻⁰⁵	9.08 × 10 ⁻⁰⁶	2.09 x 10 ⁻¹⁰	7.53 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	1.50 × 10 ⁻⁰⁴	1.00 x 10 ⁻⁰⁷	1.77 x 10 ⁻⁰⁴	3.37 x 10 ⁻⁰⁴
(Patterson Lake	4.29 × 10 ⁻⁰⁵	1.45 x 10 ⁻⁰⁸	3.34 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.52 × 10 ⁻⁰⁷	5.75 x 10 ⁻⁰⁶	5.12 x 10 ⁻⁰⁵	1.04 x 10 ⁻⁰⁴	5.47 × 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	4.79 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	3.28 × 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	2.24 × 10 ⁻⁰⁴	8.22 x 10 ⁻⁰⁴
South Arm) Uranium	6.25 × 10 ⁻⁰⁴	1.02 x 10 ⁻⁰⁴	2.65 x 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	4.62 × 10 ⁻⁰⁴	2.40 x 10 ⁻⁰²	3.81 x 10 ⁻⁰³	2.92 x 10 ⁻⁰²	2.05 × 10 ⁻⁰³	1.08 x 10 ⁻⁰⁵	1.10 x 10 ⁻⁰³	$0.00 \times 10^{+00}$	1.54 × 10 ⁻⁰³	1.26 x 10 ⁻⁰³	7.49 x 10 ⁻⁰⁴	6.70 x 10 ⁻⁰³
							Bound Scenario									
Cobalt	5.21 × 10 ⁻⁰⁵	5.12 x 10 ⁻⁰⁸	3.24 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	3.68 × 10 ⁻⁰⁵	7.54 x 10 ⁻⁰⁶	1.72 x 10 ⁻⁰⁴	2.72 x 10 ⁻⁰⁴	3.87 × 10 ⁻⁰⁴	2.78 x 10 ⁻⁰⁸	2.68 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.79 × 10 ⁻⁰⁴	1.16 x 10 ⁻⁰⁶	1.11 x 10 ⁻⁰³	1.80 x 10 ⁻⁰³
Copper	9.51 × 10 ⁻⁰⁷	3.13 x 10 ⁻⁰⁹	6.92 x 10 ⁻⁰⁸	0.00 × 10 ⁺⁰⁰	1.56 × 10 ⁻⁰⁵	1.87 x 10 ⁻⁰⁶	6.87 x 10 ⁻⁰⁵	8.73 x 10 ⁻⁰⁵	1.36 × 10 ⁻⁰⁵	2.09 x 10 ⁻¹⁰	1.13 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.25 × 10 ⁻⁰⁴	1.00 x 10 ⁻⁰⁷	2.66 x 10 ⁻⁰⁴	5.06 x 10 ⁻⁰⁴
Molybdenum	1.30 × 10 ⁻⁰⁴	1.45 x 10 ⁻⁰⁸	1.04 x 10 ⁻⁰⁵	0.00 × 10 ⁺⁰⁰	7.70 × 10 ⁻⁰⁷	5.75 x 10 ⁻⁰⁶	8.76 x 10 ⁻⁰⁵	2.35 x 10 ⁻⁰⁴	2.10 × 10 ⁻⁰³	$0.00 \times 10^{+00}$	1.84 x 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	1.26 × 10 ⁻⁰⁵	0.00 × 10 ⁺⁰⁰	8.65 × 10 ⁻⁰⁴	3.16 x 10 ⁻⁰³
Uranium	1.72 × 10 ⁻⁰³	1.02 x 10 ⁻⁰⁴	7.43 x 10 ⁻⁰⁴	0.00 × 10 ⁺⁰⁰	1.27 × 10 ⁻⁰³	2.40 x 10 ⁻⁰²	4.05 x 10 ⁻⁰³	3.18 x 10 ⁻⁰²	2.06 × 10 ⁻⁰³	1.08 x 10 ⁻⁰⁵	1.10 x 10 ⁻⁰³	$0.00 \times 10^{+00}$	1.54 × 10 ⁻⁰³	1.26 x 10 ⁻⁰³	7.49 x 10 ⁻⁰⁴	6.71 x 10 ⁻⁰³

					Project Life	span HQs							Far F <u>ut</u> ı	ure HQs			
	СОРС	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
	Cobalt	8.58 × 10 ⁻⁰⁴	9.04 × 10 ⁻⁰⁷	1.01 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	7.27 × 10 ⁻⁰⁴	1.13 × 10 ⁻⁰³	1.88 × 10 ⁻⁰²	se Case 2.15 x 10 ⁻⁰²	8.58 × 10 ⁻⁰⁴	9.04 × 10 ⁻⁰⁷	1.01 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	7.27 × 10 ⁻⁰⁴	1.13 × 10 ⁻⁰³	1.88 × 10 ⁻⁰²	2.15 x 10 ⁻⁰²
	Copper	2.91 × 10 ⁻⁰⁵	2.95 × 10 ⁻⁰⁸	4.12 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	5.66 × 10 ⁻⁰⁴	4.22 × 10 ⁻⁰⁴	3.86 × 10 ⁻⁰²	3.96 x 10 ⁻⁰²	2.91 × 10 ⁻⁰⁵	2.95 × 10 ⁻⁰⁸	4.12 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	5.66 × 10 ⁻⁰⁴	4.22 × 10 ⁻⁰⁴	3.86 × 10 ⁻⁰²	3.96 x 10 ⁻⁰²
	Molybdenum	1.08 × 10 ⁻⁰⁴	1.06 × 10 ⁻⁰⁷	1.62 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	7.66 × 10 ⁻⁰⁷	2.29 × 10 ⁻⁰⁴	8.60 × 10 ⁻⁰²	8.63 x 10 ⁻⁰²	1.08 × 10 ⁻⁰⁴	1.06 × 10 ⁻⁰⁷	1.62 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	7.66×10^{-07}	2.29 × 10 ⁻⁰⁴	8.60 × 10 ⁻⁰²	8.63 x 10 ⁻⁰²
	Uranium	2.46 × 10 ⁻⁰³	8.62 × 10 ⁻⁰⁶	2.24 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.17 × 10 ⁻⁰³	3.01 × 10 ⁻⁰²	6.79 × 10 ⁻⁰²	1.03 x 10 ⁻⁰¹	2.46 × 10 ⁻⁰³	8.62 × 10 ⁻⁰⁶	2.24 x 10 ⁻⁰⁵	0.00 × 10 ⁺⁰⁰	2.17 × 10 ⁻⁰³	3.01 × 10 ⁻⁰²	6.79 × 10 ⁻⁰²	1.03 x 10 ⁻⁰¹
	- Cramam	2.70	0.02	2,2 1 % 10	0.00	2			Incremental Pro		0.02	2,2 1 % 10	0.00		3.61	0.73	1100 % 10
	Cobalt	2.43 × 10 ⁻⁰⁶	7.84 x 10 ⁻¹⁰	2.55 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	3.62 × 10 ⁻⁰⁶	4.53 x 10 ⁻⁰⁶	1.30 x 10 ⁻⁰⁵	2.36 x 10 ⁻⁰⁵	1.23 × 10 ⁻⁰⁵	4.25 x 10 ⁻¹⁰	1.45 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	1.76 × 10 ⁻⁰⁵	5.31 x 10 ⁻⁰⁷	2.67 x 10 ⁻⁰⁵	5.70 x 10 ⁻⁰⁵
Seasonal Resident	Copper	4.40 × 10 ⁻⁰⁸	1.60 x 10 ⁻¹¹	5.41 x 10 ⁻¹¹	$0.00 \times 10^{+00}$	1.87 × 10 ⁻⁰⁴	3.11 x 10 ⁻⁰⁷	3.94 x 10 ⁻⁰⁶	1.91 x 10 ⁻⁰⁴	4.34 × 10 ⁻⁰⁷	1.07 x 10 ⁻¹²	6.15 x 10 ⁻¹⁰	$0.00 \times 10^{+00}$	2.04 × 10 ⁻⁰⁴	1.53 x 10 ⁻⁰⁸	7.19 x 10 ⁻⁰⁶	2.12 x 10 ⁻⁰⁴
(Lloyd Lake)	Molybdenum	1.67 × 10 ⁻⁰⁶	9.10 x 10 ⁻¹¹	2.20 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	1.97 × 10 ⁻⁰⁸	1.47 x 10 ⁻⁰⁶	2.81 x 10 ⁻⁰⁶	5.97 x 10 ⁻⁰⁶	2.14 × 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	3.20 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	2.52 × 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	6.65 × 10 ⁻⁰⁶	2.83 x 10 ⁻⁰⁵
	Uranium	1.88 × 10 ⁻⁰⁵	3.51 x 10 ⁻⁰⁸	1.35 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	5.58 × 10 ⁻⁰⁴	3.00 x 10 ⁻⁰⁴	7.14 x 10 ⁻⁰⁵	9.48 x 10 ⁻⁰⁴	6.19 × 10 ⁻⁰⁵	3.69 x 10 ⁻⁰⁹	5.64 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	6.37 × 10 ⁻⁰⁴	1.29 x 10 ⁻⁰⁵	2.51 x 10 ⁻⁰⁵	7.37 x 10 ⁻⁰⁴
								Bound Scenario	– Incremental I	Project Risk					,		
	Cobalt	2.53 × 10 ⁻⁰⁶	7.84 x 10 ⁻¹⁰	2.66 x 10 ⁰⁹	$0.00 \times 10^{+00}$	3.76×10^{-06}	4.53 x 10 ⁻⁰⁶	1.32 x 10 ⁻⁰⁵	2.40 x 10 ⁻⁰⁵	1.88 × 10 ⁻⁰⁵	4.25 x 10 ⁻¹⁰	2.22 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	2.68 × 10 ⁻⁰⁵	5.31 x 10 ⁻⁰⁷	4.01 x 10 ⁻⁰⁵	8.63 x 10 ⁻⁰⁵
	Copper	4.53 × 10 ⁻⁰⁸	1.60 x 10 ⁻¹¹	5.57 x 10 ⁻¹¹	$0.00 \times 10^{+00}$	1.87 × 10 ⁻⁰⁴	3.11 x 10 ⁻⁰⁷	3.96 x 10 ⁻⁰⁶	1.92 x 10 ⁻⁰⁴	6.52 × 10 ⁻⁰⁷	1.07 x 10 ⁻¹²	9.24 x 10 ⁻¹⁰	$0.00 \times 10^{+00}$	2.13 × 10 ⁻⁰⁴	1.53 x 10 ⁻⁰⁸	1.08 x 10 ⁻⁰⁵	2.25 x 10 ⁻⁰⁴
	Molybdenum	5.11 × 10 ⁻⁰⁶	9.10 x 10 ⁻¹¹	6.85 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	5.98 × 10 ⁻⁰⁸	1.47 x 10 ⁻⁰⁶	3.70 x 10 ⁻⁰⁶	1.03 x 10 ⁻⁰⁵	8.20 × 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	1.23 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	9.66 × 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.55 × 10 ⁻⁰⁵	1.09 x 10 ⁻⁰⁴
	Uranium	5.20 × 10 ⁻⁰⁵	3.51 x 10 ⁻⁰⁸	3.79 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	6.18×10^{-04}	3.00 x 10 ⁻⁰⁴	8.02 x 10 ⁻⁰⁵	1.05 x 10 ⁻⁰³	6.19 × 10 ⁻⁰⁵	3.69 x 10 ⁻⁰⁹	5.65 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	6.37×10^{-04}	1.29 x 10 ⁻⁰⁵	2.51 x 10 ⁻⁰⁵	7.37 x 10 ⁻⁰⁴
		1		-		1			se Case		1	1		1			00
	Cobalt	9.58 × 10 ⁻⁰⁴	5.91 × 10 ⁻⁰⁵	6.62 x 10 ⁻⁰⁵	0.00 × 10 ⁺⁰⁰	6.89 × 10 ⁻⁰⁴	2.47 × 10 ⁻⁰³	4.13 × 10 ⁻⁰²	4.56 x 10 ⁻⁰²	9.58 × 10 ⁻⁰⁴	5.91 × 10 ⁻⁰⁵	6.62 x 10 ⁻⁰⁵	0.00 × 10 ⁺⁰⁰	6.89 × 10 ⁻⁰⁴	2.47 × 10 ⁻⁰³	4.13 × 10 ⁻⁰²	4.56 x 10 ⁻⁰²
	Copper	3.25 × 10 ⁻⁰⁵	1.93 × 10 ⁻⁰⁶	2.69 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	5.36 × 10 ⁻⁰⁴	9.22 × 10 ⁻⁰⁴	9.32 × 10 ⁻⁰²	9.47 x 10 ⁻⁰²	3.25 × 10 ⁻⁰⁵	1.93 × 10 ⁻⁰⁶	2.69 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	5.36 × 10 ⁻⁰⁴	9.22 × 10 ⁻⁰⁴	9.32 × 10 ⁻⁰²	9.47 x 10 ⁻⁰²
	Molybdenum	1.47 × 10 ⁻⁰⁴	8.46 × 10 ⁻⁰⁶	1.29 x 10 ⁻⁰⁵	0.00 × 10 ⁺⁰⁰	8.84 × 10 ⁻⁰⁷	6.10 × 10 ⁻⁰⁴	2.77 × 10 ⁻⁰¹	2.78 x 10 ⁻⁰¹	1.47 × 10 ⁻⁰⁴	8.46 × 10 ⁻⁰⁶	1.29 x 10 ⁻⁰⁵	0.00 × 10 ⁺⁰⁰	8.84 × 10 ⁻⁰⁷	6.10 × 10 ⁻⁰⁴	2.77 × 10 ⁻⁰¹	2.78 x 10 ⁻⁰¹
	Uranium	2.75 × 10 ⁻⁰³	5.64 × 10 ⁻⁰⁴	1.47 x 10 ⁻⁰³	$0.00 \times 10^{+00}$	2.06 × 10 ⁻⁰³	6.58 × 10 ⁻⁰²	1.60 × 10 ⁻⁰¹	2.33 x 10 ⁻⁰¹	2.75 × 10 ⁻⁰³	5.64 × 10 ⁻⁰⁴	1.47 x 10 ⁻⁰³	$0.00 \times 10^{+00}$	2.06 × 10 ⁻⁰³	6.58 × 10 ⁻⁰²	1.60 × 10 ⁻⁰¹	2.33 x 10 ⁻⁰¹
	C. L. II	2.71 10-06	F 12 10-08	1.67 10:07	0.00 10+00	1.01 10-06			Incremental Pro	•	2.70 10-08	0.47 10-07	0.00 10+00	0.06 10-06	1.16 10-06	2.70 10-05	F 26 10-05
Seasonal Resident	Cobalt	2.71 × 10 ⁻⁰⁶	5.12 x 10 ⁻⁰⁸	1.67 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	1.91 × 10 ⁻⁰⁶	7.54 x 10 ⁻⁰⁶	1.33 x 10 ⁻⁰⁵	2.57 x 10 ⁻⁰⁵	1.37 × 10 ⁻⁰⁵	2.78 x 10 ⁻⁰⁸	9.47 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	9.86 × 10 ⁻⁰⁶	1.16 x 10 ⁻⁰⁶	2.70 x 10 ⁻⁰⁵	5.26 x 10 ⁻⁰⁵
One-Year-Old (Lloyd Lake)	Copper	4.91 × 10 ⁻⁰⁸	1.04 x 10 ⁻⁰⁹	3.54 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	8.07 × 10 ⁻⁰⁷	6.24 x 10 ⁻⁰⁷	3.81 x 10 ⁻⁰⁶	5.29 x 10 ⁻⁰⁶	4.85 × 10 ⁻⁰⁷	6.98 x 10 ⁻¹¹	4.02 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	8.00 × 10 ⁻⁰⁶	3.34 x 10 ⁻⁰⁸	6.53 x 10 ⁻⁰⁶	1.51 x 10 ⁻⁰⁵
(Lioyu Lake)	Molybdenum	2.27 × 10 ⁻⁰⁶	7.24 x 10 ⁻⁰⁹	1.75 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	1.33 × 10 ⁻⁰⁸	2.88 x 10 ⁻⁰⁶	3.52 x 10 ⁻⁰⁶	8.86 x 10 ⁻⁰⁶	2.90 × 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.54 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	1.74 × 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	8.14 × 10 ⁻⁰⁶	3.99 x 10 ⁻⁰⁵
	Uranium	2.10 × 10 ⁻⁰⁵	2.29 x 10 ⁻⁰⁶	8.84 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	1.55 × 10 ⁻⁰⁵	5.37 x 10 ⁻⁰⁴	4.99 x 10 ⁻⁰⁵	6.34 x 10 ⁻⁰⁴ - Incremental I	6.91 × 10 ⁻⁰⁵	2.41 x 10 ⁻⁰⁷	3.69 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	5.18 × 10 ⁻⁰⁵	2.81 x 10 ⁻⁰⁵	1.80 x 10 ⁻⁰⁵	2.04 x 10 ⁻⁰⁴
	Cobalt	2.82 × 10 ⁻⁰⁶	5.12 x 10 ⁻⁰⁸	1.74 x 10 ⁻⁰⁷	0.00 × 10 ⁺⁰⁰	1.99 × 10 ⁻⁰⁶	7.54 x 10 ⁻⁰⁶	1.35 x 10 ⁻⁰⁵	2.61 x 10 ⁻⁰⁵	2.10 × 10 ⁻⁰⁵	2.78 x 10 ⁻⁰⁸	1.45 x 10 ⁻⁰⁶	0.00 × 10 ⁺⁰⁰	1.51 × 10 ⁻⁰⁵	1.16 x 10 ⁻⁰⁶	4.06 x 10 ⁻⁰⁵	7.93 x 10 ⁻⁰⁵
	Copper	5.05 × 10 ⁻⁰⁸	1.04 x 10 ⁻⁰⁹	3.64 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	8.31 × 10 ⁻⁰⁷	6.24 x 10 ⁻⁰⁷	3.83 x 10 ⁻⁰⁶	5.34 x 10 ⁻⁰⁶	7.28×10^{-07}	6.98 x 10 ⁻¹¹	6.04 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	1.20 × 10 ⁻⁰⁵	3.34 x 10 ⁻⁰⁸	9.78 x 10 ⁻⁰⁶	2.26 x 10 ⁻⁰⁵
	Molybdenum	6.94 × 10 ⁻⁰⁶	7.24 x 10 ⁻⁰⁹	5.45 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	4.09 × 10 ⁻⁰⁸	2.88 x 10 ⁻⁰⁶	4.62 x 10 ⁻⁰⁶	1.50 x 10 ⁻⁰⁵	1.11 × 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	9.76 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	6.69 × 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	3.12 × 10 ⁻⁰⁵	1.53 x 10 ⁻⁰⁴
	Uranium	5.81 × 10 ⁻⁰⁵	2.29 x 10 ⁻⁰⁶	2.47 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	4.30 × 10 ⁻⁰⁵	5.37 x 10 ⁻⁰⁴	5.63 x 10 ⁻⁰⁵	7.21 x 10 ⁻⁰⁴	6.92 × 10 ⁻⁰⁵	2.41 x 10 ⁻⁰⁷	3.69 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	5.18 × 10 ⁻⁰⁵	2.81 x 10 ⁻⁰⁵	1.80 x 10 ⁻⁰⁵	2.04 x 10 ⁻⁰⁴
	Oraniani	3.01 × 10	2.23 X 10	2.47 X 10	0.00 × 10	4.50 × 10	3.37 X 10		se Case	0.52 × 10	2.41 X 10	3.03 X 10	0.00 × 10	3.10 × 10	2.01 X 10	1.00 X 10	2.04 X 10
	Cobalt	n/a	8.58 × 10 ⁻⁰⁴	9.04 × 10 ⁻⁰⁷	1.01 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.30 × 10 ⁻⁰³	1.44 × 10 ⁻⁰³	2.44 × 10 ⁻⁰²	2.90 x 10 ⁻⁰²							
	Copper	n/a	2.91 × 10 ⁻⁰⁵	2.95 × 10 ⁻⁰⁸	4.12 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	1.79 × 10 ⁻⁰³	5.38 × 10 ⁻⁰⁴	4.12 × 10 ⁻⁰²	4.35 x 10 ⁻⁰²							
	Molybdenum	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	1.08 × 10 ⁻⁰⁴	1.06 × 10 ⁻⁰⁷	1.62 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.42 × 10 ⁻⁰⁶	2.92 × 10 ⁻⁰⁴	6.98 × 10 ⁻⁰²	7.02 x 10 ⁻⁰²
	Uranium	n/a	2.46 × 10 ⁻⁰³	8.62 × 10 ⁻⁰⁶	2.24 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	6.85 × 10 ⁻⁰³	3.84 × 10 ⁻⁰²	8.20 × 10 ⁻⁰²	1.30 x 10 ⁻⁰¹							
					·	,			Incremental Pro								
Permanent	Cobalt	n/a	1.24 × 10 ⁻⁰³	2.83 x 10 ⁻⁰⁹	1.47 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	3.32 × 10 ⁻⁰³	4.51 x 10 ⁻⁰⁶	6.37 x 10 ⁻⁰³	1.09 x 10 ⁻⁰²							
Resident (Patterson Lake North Arm –	Copper	n/a	4.62 × 10 ⁻⁰⁵	6.05 x 10 ⁻¹¹	6.54 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	2.84 × 10 ⁻⁰³	1.10 x 10 ⁻⁰⁶	1.81 x 10 ⁻⁰³	4.69 x 10 ⁻⁰³							
West Basin)	Molybdenum	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	2.31 × 10 ⁻⁰³	$0.00 \times 10^{+00}$	3.46 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	5.15 × 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	1.80 × 10 ⁻⁰³	4.16 x 10 ⁻⁰³
	Uranium	n/a	2.21 × 10 ⁻⁰²	3.84 x 10 ⁻⁰⁶	2.02 x 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	6.17 × 10 ⁻⁰²	1.71 x 10 ⁻⁰²	3.07 x 10 ⁻⁰²	1.32 x 10 ⁻⁰¹							
							Upper E	Bound Scenario	– Incremental I		1	<u> </u>		1		<u> </u>	
	Cobalt	n/a	1.90 × 10 ⁻⁰³	2.83 x 10 ⁻⁰⁹	2.25 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	5.09 × 10 ⁻⁰³	4.51 x 10 ⁻⁰⁶	9.75 x 10 ⁻⁰³	1.67 x 10 ⁻⁰²							
	Copper	n/a	6.94×10^{-05}	6.05 x 10 ⁻¹¹	9.83 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	4.26 × 10 ⁻⁰³	1.10 x 10 ⁻⁰⁶	2.71 x 10 ⁻⁰³	7.04 x 10 ⁻⁰³							
	Molybdenum	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	8.87 × 10 ⁻⁰³	$0.00 \times 10^{+00}$	1.33 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	1.98 × 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	6.90 × 10 ⁻⁰³	1.60 x 10 ⁻⁰²
	Uranium	n/a	2.22 × 10 ⁻⁰²	3.84 x 10 ⁻⁰⁶	2.02 x 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	6.17 × 10 ⁻⁰²	1.71 x 10 ⁻⁰²	3.07 x 10 ⁻⁰²	1.32 x 10 ⁻⁰¹							

ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT

HUMAN HEALTH RISK ASSESSMENT

					Project Life	span HQs							Far Fut	ıre HQs			
	СОРС	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
								Bas	se Case								
	Cobalt	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	9.58 × 10 ⁻⁰⁴	5.91 × 10 ⁻⁰⁵	6.62 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.18×10^{-03}	2.84×10^{-03}	4.78 × 10 ⁻⁰²	5.39 x 10 ⁻⁰²
	Copper	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	3.25×10^{-05}	1.93 × 10 ⁻⁰⁶	2.69 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	1.69×10^{-03}	1.06×10^{-03}	9.70 × 10 ⁻⁰²	9.98 x 10 ⁻⁰²
	Molybdenum	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	1.47×10^{-04}	8.46 × 10 ⁻⁰⁶	1.29 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.79×10^{-06}	7.01×10^{-04}	2.64 × 10 ⁻⁰¹	2.64 x 10 ⁻⁰¹
	Uranium	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	2.75×10^{-03}	5.64 × 10 ⁻⁰⁴	1.47 x 10 ⁻⁰³	$0.00 \times 10^{+00}$	6.49×10^{-03}	7.57×10^{-02}	1.70×10^{-01}	2.57 x 10 ⁻⁰¹
Permanent							Appl	ication Case -	ncremental Pro	oject Risk							
Resident One-Year-	Cobalt	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	1.39×10^{-03}	1.85 x 10 ⁻⁰⁷	9.59 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	3.15×10^{-03}	8.90 x 10 ⁻⁰⁶	6.47 x 10 ⁻⁰³	1.11 x 10 ⁻⁰²
Old (Patterson	Copper	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	5.16 × 10 ⁻⁰⁵	3.95 x 10 ⁻⁰⁹	4.28 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.69×10^{-03}	2.18 x 10 ⁻⁰⁶	1.67 x 10 ⁻⁰³	4.41 x 10 ⁻⁰³
Lake North Arm –	Molybdenum	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	3.14×10^{-03}	$0.00 \times 10^{+00}$	2.75 x 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	5.95×10^{-05}	$0.00 \times 10^{+00}$	2.20×10^{-03}	5.67 x 10 ⁻⁰³
West Basin)	Uranium	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	2.47×10^{-02}	2.51 x 10 ⁻⁰⁴	1.32 x 10 ⁻⁰²	$0.00 \times 10^{+00}$	5.85×10^{-02}	3.37 x 10 ⁻⁰²	2.23 x 10 ⁻⁰²	1.53 x 10 ⁻⁰¹
							Upper E	Bound Scenario	- Incremental	Project Risk							
	Cobalt	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	2.12×10^{-03}	1.85 x 10 ⁻⁰⁷	1.47 x 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	4.83×10^{-03}	8.90 x 10 ⁻⁰⁶	9.90 x 10 ⁻⁰³	1.70 x 10 ⁻⁰²
	Copper	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	7.75×10^{-05}	3.95 x 10 ⁻⁰⁹	6.42 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	4.04×10^{-03}	2.18 x 10 ⁻⁰⁶	2.50 x 10 ⁻⁰³	6.62 x 10 ⁻⁰³
	Molybdenum	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	1.21 × 10 ⁻⁰²	$0.00 \times 10^{+00}$	1.06 x 10 ⁻⁰³	$0.00 \times 10^{+00}$	2.28×10^{-04}	$0.00 \times 10^{+00}$	8.44 × 10 ⁻⁰³	2.18 x 10 ⁻⁰²
	Uranium	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	2.48 × 10 ⁻⁰²	2.51 x 10 ⁻⁰⁴	1.32 x 10 ⁻⁰²	$0.00 \times 10^{+00}$	5.85 × 10 ⁻⁰²	3.37 x 10 ⁻⁰²	2.23 x 10 ⁻⁰²	1.53 x 10 ⁻⁰¹

Note: **Bold** indicates exceedance of the HQ of 0.2.

HQ = hazard quotient; COPC = constituent of potential concern; n/a = receptor not assessed in that phase.

5.4.1.1.2 Carcinogen Risk

The arsenic ILCR is expected to exceed the negligible cancer risk level of 1 in 100,000 for the subsistence harvester (i.e., composite receptor throughout all life stages) harvesting Traditional Foods from Patterson Lake South Arm during the Project lifespan for both the Application Case and the reasonable upper bound sensitivity scenario (Table 5-19). The arsenic ILCR is below the negligible cancer risk level of 1 in 100,000 for all other human receptors during the Project lifespan and in the far-future scenario.

Incremental lifetime cancer risks from arsenic for the subsistence harvester at Patterson Lake South Arm were predicted to be 4 in 100,000 in both the Application Case and reasonable upper bound sensitivity scenario. In comparison, baseline cancer risks from arsenic for the reference subsistence harvester were predicted to be 69 in 100,000 for the selected regional background conditions in the IMPACT model. The reference subsistence harvester represents an adult subsistence harvester who is exposed to baseline concentrations (i.e., existing conditions) of arsenic in foodstuffs only, without any Project-related additions. The ILCR for the human receptors from arsenic from the Project is a small portion of the existing baseline cancer risks for these receptors.

The main ingestion exposure pathway for arsenic for the adult camp worker, subsistence harvester and seasonal resident was consumption of local terrestrial animals including beaver, grouse, mallard, moose, and moose organs, as well as locally caught fish represented in the HHRA by lake whitefish and northern pike. The ILCRs for the Application Case and reasonable upper bound sensitivity scenario by food type are provided in Table 5-20 and Table 5-21 for the seasonal resident and subsistence harvester, respectively. The greatest contributor to the predicted ILCRs for the Application Case for both the seasonal resident and subsistence harvester at Patterson Lake South Arm during the Project lifespan is consumption of moose meat and organs.

Arsenic uptake by fish and their maximum arsenic concentrations in flesh were predicted as part of the environmental pathways assessment. For the Patterson Lake South Arm, the maximum predicted arsenic concentration in lake whitefish (0.043 mg/kg fw) was within the range of background variability observed in the baseline LSA and RSA (0.011 mg/kg to 0.11 mg/kg fw; Annex V.1, Aquatic Environment Baseline Report), and within the range of tissue concentrations from lake whitefish samples as part of the Eastern Athabasca Region community monitoring program for 2011 to 2020 (0.01 mg/kg to 0.52 mg/kg fw; CanNorth 2018). The maximum predicted concentration of arsenic in northern pike (0.045 mg/kg fw) was also within the range of background variability measured in northern pike during baseline studies (0.011 mg/kg to 0.36 mg/kg fw; Annex V.1), and below the range of tissue concentrations from northern pike samples as part of the Eastern Athabasca Region community monitoring program for 2011 to 2020 (0.09 mg/kg to 0.15 mg/kg fw; CanNorth 2018), and the FNFNES sampling program for 2015 (maximum concentration of 0.1 mg/kg fw; Chan et al. 2018).



The potential for arsenic to represent health risks for consumers of Traditional Foods was assessed for the Eastern Athabasca Region and for the Boreal Shield region of Saskatchewan by CanNorth (2018) and Chan et al. (2018), respectively. Each of these HHRAs concluded that arsenic did not pose a significant risk to consumers of Traditional Foods. Since predicted incremental cancer risks from fish consumption are based on predicted arsenic concentrations in fish that are within the range of concentrations assessed for other regions of Saskatchewan, arsenic originating from the Project is not considered to represent a significant health risk.

Additionally, the main contribution to the arsenic cancer risk for the human receptors is from ingestion of moose and moose organs. The ingestion rates for moose and moose organs in the Traditional Foods diet (Table 5-5) are considered conservative and were based on the ingestion rates provided in the FNFNES (Chan et al. 2018). The diet is conservative in that it was based on the higher ingestion rates in the male diet and the high consumer of Traditional Foods.

In the far-future projection, for the future permanent resident at the camp site, the arsenic ILCR was lower than for the camp worker during Operations and below the negligible cancer risk level (Table 5-19). The risk would be lower during the far-future projection because the main source of arsenic to the environment is during Operations when treated effluent is being released to Patterson Lake North Arm – West Basin.

Table 5-19: Estimated Incremental Lifetime Cancer Risk from Arsenic to Human Receptors – Application Case and Upper Bound Scenario

Receptor		se Cancer Risk 00,000)		cenario Cancer 100,000)
	Operations	Far-Future	Operations	Far-Future
Camp worker (adult)	1	n/a	1	n/a
Subsistence harvester - Patterson Lake South Arm (composite)	4	0.1	4	0.1
Subsistence harvester - Beet Lake (composite)	0.2	0.004	0.2	0.005
Subsistence harvester - Lloyd Lake (composite)	0.1	0.0002	0.1	0.0003
Seasonal resident - Patterson Lake South Arm (composite)	0.8	0.02	0.8	0.03
Seasonal resident - Lloyd Lake (composite)	0.02	0.00004	0.02	0.00007
Permanent resident - camp site (composite)	n/a	0.5	n/a	0.8

Bold indicates exceedance of the negligible cancer risk level of 1 in 100,000.

n/a = not applicable, receptor was not assessed.



Table 5-20: Estimated Carcinogen Risks for Arsenic to Seasonal Residents - Operations - Application Case and Upper Bound Scenario

Food Type		Application Cas	se – Operations			Upper Bound Sce	nario – Operation	S
Food Type		dent – Patterson outh Arm	Seasonal Resid	lent – Lloyd Lake		dent – Patterson outh Arm	Seasonal Resid	ent – Lloyd Lake
	ILCR	% Contribution	ILCR	% Contribution	ILCR	% Contribution	ILCR	% Contribution
Labrador tea	5.04 × 10 ⁻⁰⁸	1%	5.04 × 10 ⁻⁰⁸	22%	5.04 × 10 ⁻⁰⁸	1%	5.04 × 10 ⁻⁰⁸	22%
Blueberry	2.08×10^{-08}	0.3%	2.08 × 10 ⁻⁰⁸	9%	2.08 × 10 ⁻⁰⁸	0.3%	2.08×10^{-08}	9%
Northern pike	2.21 × 10 ⁻⁰⁷	3%	2.10 × 10 ⁻⁰⁹	1%	2.27 × 10 ⁻⁰⁷	3%	2.15 × 10 ⁻⁰⁹	1%
Lake whitefish	2.03 × 10 ⁻⁰⁷	3%	1.92 × 10 ⁻⁰⁹	1%	2.08 × 10 ⁻⁰⁷	3%	1.96 × 10 ⁻⁰⁹	1%
Store- bought food	$0.00 \times 10^{+00}$	0%	0.00 × 10 ⁺⁰⁰	0%	0.00 × 10 ⁺⁰⁰	0%	$0.00 \times 10^{+00}$	0%
Moose	2.38 × 10 ⁻⁰⁶	30%	1.82 × 10 ⁻⁰⁸	8%	2.44 × 10 ⁻⁰⁶	30%	1.83 × 10 ⁻⁰⁸	8%
Moose organs	4.13 × 10 ⁻⁰⁶	53%	3.16 × 10 ⁻⁰⁸	14%	4.23 × 10 ⁻⁰⁶	53%	3.17 × 10 ⁻⁰⁸	14%
Beaver	4.97 × 10 ⁻⁰⁷	6%	8.59 × 10 ⁻⁰⁸	38%	5.08 × 10 ⁻⁰⁷	6%	8.60 × 10 ⁻⁰⁸	38%
Mallard	3.42 × 10 ⁻⁰⁷	4%	3.21 × 10 ⁻⁰⁹	1%	3.50 × 10 ⁻⁰⁷	4%	3.30 × 10 ⁻⁰⁹	1%
Spruce grouse	1.08 × 10 ⁻⁰⁸	0.1%	1.06 × 10 ⁻⁰⁸	5%	1.08 × 10 ⁻⁰⁸	0.1%	1.06 × 10 ⁻⁰⁸	5%
Total	7.86 × 10 ⁻⁰⁶	100%	2.25 × 10 ⁻⁰⁷	100%	8.05 × 10 ⁻⁰⁶	100%	2.25 × 10 ⁻⁰⁷	100%

Note: Total percentages may not sum to 100% due to rounding. ILCR = incremental lifetime cancer risk.



Table 5-21: Estimated Carcinogen Risks for Arsenic to Subsistence Harvesters – Operations – Application Case and Upper Bound Scenario

		A	pplication Ca	se – Operatior	ıs			Upp	er Bound Sce	nario - Operat	tions	
Food		Harvester – ke South Arm	Subsistence Harvester – Beet Lake		Subsistence Harvester – Lloyd Lake			Harvester – ke South Arm		Harvester –		Harvester – I Lake
Type	ILCR	% Contribution	ILCR	% Contribution	ILCR	% Contribution	ILCR	% Contribution	ILCR	% Contribution	ILCR	% Contribution
Labrador tea	2.44 × 10 ⁻⁰⁷	1%	2.44 × 10 ⁻⁰⁷	11%	2.44 × 10 ⁻⁰⁷	23%	2.44 × 10 ⁻⁰⁷	1%	2.44 × 10 ⁻⁰⁷	10%	2.44 × 10 ⁻⁰⁷	23%
Blueberry	3.46 × 10 ⁻⁰⁸	0.1%	3.46 × 10 ⁻⁰⁸	1%	3.46 × 10 ⁻⁰⁸	3%	3.46 × 10 ⁻⁰⁸	0.1%	3.46×10^{-08}	1%	3.46 × 10 ⁻⁰⁸	3%
Northern pike	1.16 × 10 ⁻⁰⁶	3%	1.89 × 10 ⁻⁰⁷	8%	1.10 × 10 ⁻⁰⁸	1%	1.19 × 10 ⁻⁰⁶	3%	1.94 × 10 ⁻⁰⁷	8%	1.13 × 10 ⁻⁰⁸	1%
Lake whitefish	1.07 × 10 ⁻⁰⁶	3%	1.74 × 10 ⁻⁰⁷	8%	1.01 × 10 ⁻⁰⁸	1%	1.09 × 10 ⁻⁰⁶	3%	1.78 × 10 ⁻⁰⁷	8%	1.03 × 10 ⁻⁰⁸	1%
Store- bought food	0.00 × 10 ⁺⁰⁰	0%	0.00 × 10 ⁺⁰⁰	0%	0.00 × 10 ⁺⁰⁰	0%	0.00 × 10 ⁺⁰⁰	0%	0.00 × 10 ⁺⁰⁰	0%	0.00 × 10 ⁺⁰⁰	0%
Moose	1.26 × 10 ⁻⁰⁵	34%	2.50 × 10 ⁻⁰⁷	11%	9.61 × 10 ⁻⁰⁸	9%	1.29 × 10 ⁻⁰⁵	34%	2.54 × 10 ⁻⁰⁷	11%	9.63 × 10 ⁻⁰⁸	9%
Moose Organs	1.76 × 10 ⁻⁰⁵	48%	3.50 × 10 ⁻⁰⁷	15%	1.34 × 10 ⁻⁰⁷	12%	1.80 × 10 ⁻⁰⁵	48%	3.56 × 10 ⁻⁰⁷	15%	1.35 × 10 ⁻⁰⁷	13%
Beaver	2.72 × 10 ⁻⁰⁶	7%	8.14 × 10 ⁻⁰⁷	35%	4.69 × 10 ⁻⁰⁷	44%	2.78 × 10 ⁻⁰⁶	7%	8.24 × 10 ⁻⁰⁷	35%	4.70 × 10 ⁻⁰⁷	44%
Mallard	1.16 × 10 ⁻⁰⁶	3%	1.88 × 10 ⁻⁰⁷	8%	1.09 × 10 ⁻⁰⁸	1%	1.19 × 10 ⁻⁰⁶	3%	1.93 × 10 ⁻⁰⁷	8%	1.12 × 10 ⁻⁰⁸	1%
Spruce grouse	6.74 × 10 ⁻⁰⁸	0.2%	6.63 × 10 ⁻⁰⁸	3%	6.61 × 10 ⁻⁰⁸	6%	6.74 × 10 ⁻⁰⁸	0.2%	6.63 × 10 ⁻⁰⁸	3%	6.61 × 10 ⁻⁰⁸	6%
Total	3.66 × 10 ⁻⁰⁵	100%	2.31 × 10 ⁻⁰⁶	100%	1.08 × 10 ⁻⁰⁶	100%	3.75 × 10 ⁻⁰⁵	100%	2.34 × 10 ⁻⁰⁶	100%	1.08 × 10 ⁻⁰⁶	100%

Note: Total percentages may not sum to 100% due to rounding. ILCR = incremental lifetime cancer risk.



5.4.1.1.3 Radiological Risk

The incremental radiation dose to all human receptors during all Project phases and the far-future projection would be below the public regulatory dose limit of 1 mSv/yr for both the Application Case and the reasonable upper bound sensitivity scenario, as shown in Table 5-22. The maximum dose in the Application Case was predicted for the subsistence harvester (one-year-old) who eats Traditional Foods gathered at Patterson Lake South Arm during Operations. The main contribution to the total dose is from polonium-210 from eating terrestrial animals in the Traditional Food diet, including moose (e.g., meat, organs), beaver, grouse, and mallard.

If a dose constraint of 0.3 mSv/yr is applied, the dose to the subsistence harvester (one-year-old) would be less than the dose constraint for the Application Case and for the reasonable upper bound sensitivity scenario, and well below the regulatory public dose limit.

In the far-future, a future permanent resident, residing at the decommissioned and reclaimed Project site well beyond Closure could receive a dose up to 0.07 mSv/yr, which is well below both the regulatory public dose limit and the dose constraint.

Overall, since the radiation dose estimates were below the public dose limit, no discernable health effects would be anticipated due to exposure of these receptors to radioactive releases from the Project. The presence and concentrations of radionuclides in the receiving environment would be monitored and the associated radiation dose estimates would be periodically reassessed in accordance with the processes outlined in the Environmental Protection Program.

Table 5-22: Summary of All Radiation Doses to Human Receptors – Application Case and Upper Bound Scenario

		Applica	tion Case	Upper	-Bound
Receptor	Location	Project Lifespan	Far-Future	Project Lifespan	Far-Future
			Dose (n	nSv/yr)	
Camp worker	Permanent Camp	6.50 x 10 ⁻⁰²	n/a	8.24 x 10 ⁻⁰²	n/a
Harvester	Patterson Lake South Arm	6.80 x 10 ⁻⁰²	4.62 x 10 ⁻⁰³	1.03 x 10 ⁻⁰¹	4.62 x 10 ⁻⁰³
Harvester_1y	Patterson Lake South Arm	7.45 x 10 ⁻⁰²	5.41 x 10 ⁻⁰³	1.19 x 10 ⁻⁰¹	5.41 x 10 ⁻⁰³
Harvester	Beet Lake	2.18 x 10 ⁻⁰²	1.97 x 10 ⁻⁰³	2.38 x 10 ⁻⁰²	1.97 x 10 ⁻⁰³
Harvester_1y	Beet Lake	2.68 x 10 ⁻⁰²	2.43 x 10 ⁻⁰³	2.94 x 10 ⁻⁰²	2.43 x 10 ⁻⁰³
Harvester	Lloyd Lake	9.53 x 10 ⁻⁰⁴	7.98 x 10 ⁻⁰⁵	1.06 x 10 ⁻⁰³	7.98 x 10 ⁻⁰⁵
Harvester_1y	Lloyd Lake	1.00 x 10 ⁻⁰³	9.12 x 10 ⁻⁰⁵	1.14 x 10 ⁻⁰³	9.12 x 10 ⁻⁰⁵
Seasonal Resident	Patterson Lake South Arm	1.77 x 10 ⁻⁰²	2.09 x 10 ⁻⁰³	2.60 x 10 ⁻⁰²	2.09 x 10 ⁻⁰³
Seasonal Resident_1y	Patterson Lake South Arm	2.11 x 10 ⁻⁰²	2.65 x 10 ⁻⁰³	3.20 x 10 ⁻⁰²	2.65 x 10 ⁻⁰³
Seasonal Resident	Lloyd Lake	1.55 x 10 ⁻⁰³	1.31 x 10 ⁻⁰³	1.59 x 10 ⁻⁰³	1.31 x 10 ⁻⁰³
Seasonal Resident_1y	Lloyd Lake	3.13 x 10 ⁻⁰⁴	4.91 x 10 ⁻⁰⁵	3.49 x 10 ⁻⁰⁴	4.92 x 10 ⁻⁰⁵
Permanent Resident	Permanent Camp	n/a	5.11 x 10 ⁻⁰²	n/a	5.12 x 10 ⁻⁰²
Permanent Resident_1y	Permanent Camp	n/a	6.58 x 10 ⁻⁰²	n/a	6.58 x 10 ⁻⁰²

n/a = not applicable, receptor was not assessed.



5.4.1.1.4 Radon Risk

The incremental radon dose to the camp worker was estimated to be 0.51 mSv/yr. The incremental radon dose was based on atmospheric modelling of the radon released during the maximum ore grade year (Year 1 of the Project). This is a conservative assumption, as radon is expected to be released at a much lower rate over the duration of the Project. Additionally, the assessment is conservative in that it assumes that the camp worker spends 100% of the time indoors.

The total incremental dose to the camp worker from all radionuclides in the U-238 decay chain including radon would be 0.58 mSv/yr for the Application Case and 0.59 mSv/yr for the Upper Bound scenario, both of which are below the dose limit for a non-nuclear energy worker of 1 mSv/yr (Table 5-23). The estimate of total dose including radon is conservative based on the following assumptions:

- The camp worker spends 100% of their time indoors when on site for exposure to radon (Section 5.2.4.1.4).
- Receptors are exposed to the maximum exposure concentrations at their location for each model scenario and Project phase (Section 5.2.5).
- For radionuclides in the U-238 decay chain (other than radon), the camp worker is also exposed to radionuclides through ingestion (water and food) pathways resulting in a conservative dose when also factoring in the dose from radon indoors.

Table 5-23: Total Radiation Dose to Camp Worker from all Radionuclides including Radon Progeny – Project Lifespan

Human Receptor	Location	Scenario	Maximum Total Incremental Dose – U-238 Decay Chain (mSv/yr)	Maximum Radon Dose (mSv/yr)	Maximum Total Dose (mSv/yr)	% of Dose Limit for Non-Nuclear Energy Worker
Camp Worker	Camp	Application Case	6.50 x 10 ⁻⁰²	5.10 x 10 ⁻⁰¹	5.75 x 10 ⁻⁰¹	58%
		Upper Bound	8.24 x 10 ⁻⁰²	5.10 x 10 ⁻⁰¹	5.92 x 10 ⁻⁰¹	59%

5.4.1.2 Estimated Risk to Human Receptors – Reasonably Foreseeable Development Case 5.4.1.2.1 Non-carcinogen Risk

With the addition of releases from the Fission Patterson Lake South Property, the estimated incremental HQs for all human receptors at all assessed locations were predicted to increase slightly under the RFD Case; however, they would remain below the acceptable risk level of 0.2 per pathway for the one-year-old and adult for all human receptors (Table 5-24). For assessment of non-carcinogens, the one-year-old is typically considered the most sensitive receptor (Health Canada 2010a). Table 5-24 shows the Project only risk, for the Project lifespan and for the far future, as well as the Base Case risk, all for the RFD Case. These results indicate that the Project only risks would be minimal.



Table 5-24: Estimated Non-carcinogen Risk to Human Receptors – Operations – Reasonably Foreseeable Development Case

				орогалоно	RFD Case Haza		Nopmont Gao		
	СОРС	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
					Base Case				
	Cobalt	8.58 × 10 ⁻⁰⁴	9.04×10^{-07}	1.01x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	1.15×10^{-03}	7.21×10^{-04}	1.97 × 10 ⁻⁰²	2.25 x 10 ⁻⁰²
	Copper	2.91 × 10 ⁻⁰⁵	2.95 × 10 ⁻⁰⁸	4.12 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	8.94×10^{-04}	2.69×10^{-04}	3.89×10^{-02}	4.01 x 10 ⁻⁰²
	Molybdenum	1.08 × 10 ⁻⁰⁴	1.06 × 10 ⁻⁰⁷	1.62 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	1.21×10^{-06}	1.46 × 10 ⁻⁰⁴	8.23 × 10 ⁻⁰²	8.25 x 10 ⁻⁰²
	Uranium	2.46 × 10 ⁻⁰³	8.62 × 10 ⁻⁰⁶	2.24 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	3.43×10^{-03}	1.92 × 10 ⁻⁰²	7.29 × 10 ⁻⁰²	9.80 x 10 ⁻⁰²
		,		Project Lifespa	ın - Incremental P	roject Risk		1	
Camp worker	Cobalt	5.31 × 10 ⁻⁰⁶	1.72 x 10 ⁻⁰⁹	3.99 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	1.79 × 10 ⁻⁰⁴	8.04 x 10 ⁻⁰⁶	3.32 x 10 ⁻⁰⁴	5.25 x 10 ⁻⁰⁴
(Patterson Lake North	Copper	2.44 × 10 ⁻⁰⁸	4.67 x 10 ⁻¹⁰	2.10 x 10 ⁻¹¹	$0.00 \times 10^{+00}$	8.64×10^{-05}	4.06 x 10 ⁻⁰⁷	1.47 x 10 ⁻⁰⁴	2.34 x 10 ⁻⁰⁴
Arm – West Basin)	Molybdenum	1.84 × 10 ⁻⁰⁷	1.91 x 10 ⁻⁰⁹	1.64 x 10 ⁻¹⁰	$0.00 \times 10^{+00}$	9.84×10^{-07}	2.71 x 10 ⁻⁰⁶	8.76 x 10 ⁻⁰⁵	9.15 x 10 ⁻⁰⁵
	Uranium	2.55 × 10 ⁻⁰⁶	1.94 x 10 ⁻⁰⁵	8.88 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	7.17×10^{-03}	2.42 x 10 ⁻⁰²	1.52 x 10 ⁻⁰²	4.66 x 10 ⁻⁰²
				Far-Future -	Incremental Proj	ect Risk			
	Cobalt	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Copper	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Molybdenum	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Uranium	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
					Base Case				
	Cobalt	8.58 × 10 ⁻⁰⁴	9.04 × 10 ⁻⁰⁷	1.01 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.30 × 10 ⁻⁰³	1.44 × 10 ⁻⁰³	2.44 × 10 ⁻⁰²	2.90 x 10 ⁻⁰²
	Copper	2.91 × 10 ⁻⁰⁵	2.95 × 10 ⁻⁰⁸	4.12 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	1.79 × 10 ⁻⁰³	5.38 × 10 ⁻⁰⁴	4.12 × 10 ⁻⁰²	4.35 x 10 ⁻⁰²
	Molybdenum	1.08 × 10 ⁻⁰⁴	1.06 × 10 ⁻⁰⁷	1.62 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.42×10^{-06}	2.92 × 10 ⁻⁰⁴	6.98 × 10 ⁻⁰²	7.02 x 10 ⁻⁰²
	Uranium	2.46 × 10 ⁻⁰³	8.62 × 10 ⁻⁰⁶	2.24 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	6.85×10^{-03}	3.84 × 10 ⁻⁰²	8.20 × 10 ⁻⁰²	1.30 x 10 ⁻⁰¹
				Proje	ct Lifespan - Incre	emental Project I	Risk	1	
Subsistence Harvester	Cobalt	1.46 × 10 ⁻⁰⁴	1.31 x 10 ⁻⁰⁹	1.10 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	3.59 × 10 ⁻⁰⁴	1.61 x 10 ⁻⁰⁵	6.63 x 10 ⁻⁰⁴	1.18 x 10 ⁻⁰³
(Patterson Lake South	Copper	2.86 × 10 ⁻⁰⁶	2.66 x 10 ⁻¹¹	2.49 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	1.73 × 10 ⁻⁰⁴	8.11 x 10 ⁻⁰⁷	2.93 x 10 ⁻⁰⁴	4.70 x 10 ⁻⁰⁴
Arm)	Molybdenum	9.60 × 10 ⁻⁰⁵	1.52 x 10 ⁻¹⁰	9.18 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	1.97 × 10 ⁻⁰⁶	5.41 x 10 ⁻⁰⁶	1.75 x 10 ⁻⁰⁴	2.78 x 10 ⁻⁰⁴
	Uranium	5.35 × 10 ⁻⁰³	2.95 x 10 ⁻⁰⁶	1.47 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	1.43 × 10 ⁻⁰²	4.83 x 10 ⁻⁰²	3.04 x 10 ⁻⁰²	9.84 x 10 ⁻⁰²
					Incremental Proj			1	
	Cobalt	3.77 × 10 ⁻⁰⁴	7.08 x 10 ⁻¹⁰	4.46 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	1.01 × 10 ⁻⁰³	1.13 x 10 ⁻⁰⁶	2.92 x 10 ⁻⁰³	4.31 x 10 ⁻⁰³
	Copper	1.36 × 10 ⁻⁰⁵	1.78 x 10 ⁻¹²	1.92 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	8.32×10^{-04}	3.25 x 10 ⁻⁰⁸	7.56 x 10 ⁻⁰⁴	1.60 x 10 ⁻⁰³
	Molybdenum	6.70 × 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	1.00 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	1.50×10^{-05}	$0.00 \times 10^{+00}$	7.56 × 10 ⁻⁰⁴	1.44 x 10 ⁻⁰³
	Uranium	3.06×10^{-03}	3.10 x 10 ⁻⁰⁷	2.79 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	8.53 × 10 ⁻⁰³	1.38 x 10 ⁻⁰³	4.39 x 10 ⁻⁰³	1.74 x 10 ⁻⁰²
Subsistence Harvester					Base Case				
One-Year-Old	Cobalt	9.58 × 10 ⁻⁰⁴	5.91 × 10 ⁻⁰⁵	6.62 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.18×10^{-03}	2.84×10^{-03}	4.78 × 10 ⁻⁰²	5.39 x 10 ⁻⁰²



					RFD Case Haza	rd Quotients			
	СОРС	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
(Patterson Lake South	Copper	3.25 × 10 ⁻⁰⁵	1.93 × 10 ⁻⁰⁶	2.69 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	1.69 × 10 ⁻⁰³	1.06 × 10 ⁻⁰³	9.70 × 10 ⁻⁰²	9.98 x 10 ⁻⁰²
Arm)	Molybdenum	1.47 × 10 ⁻⁰⁴	8.46 × 10 ⁻⁰⁶	1.29 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.79 × 10 ⁻⁰⁶	7.01 × 10 ⁻⁰⁴	2.64 × 10 ⁻⁰¹	2.64 x 10 ⁻⁰¹
	Uranium	2.75 × 10 ⁻⁰³	5.64 × 10 ⁻⁰⁴	1.47 x 10 ⁻⁰³	$0.00 \times 10^{+00}$	6.49 × 10 ⁻⁰³	7.57 × 10 ⁻⁰²	1.70 × 10 ⁻⁰¹	2.57 x 10 ⁻⁰¹
				Project Lifesp	an - Incremental F	Project Risk			
	Cobalt	1.63 × 10 ⁻⁰⁴	8.54 x 10 ⁻⁰⁸	7.19 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	3.40×10^{-04}	2.27 x 10 ⁻⁰⁵	6.80 x 10 ⁻⁰⁴	1.21 x 10 ⁻⁰³
	Copper	3.19 × 10 ⁻⁰⁶	1.74 x 10 ⁻⁰⁹	1.63 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	1.64×10^{-04}	1.39 x 10 ⁻⁰⁶	2.95 x 10 ⁻⁰⁴	4.64 x 10 ⁻⁰⁴
	Molybdenum	1.30 × 10 ⁻⁰⁴	1.21 x 10 ⁻⁰⁸	7.30 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.27 × 10 ⁻⁰⁶	9.10 x 10 ⁻⁰⁶	2.21 x 10 ⁻⁰⁴	3.71 x 10 ⁻⁰⁴
	Uranium	5.97 × 10 ⁻⁰³	1.93 x 10 ⁻⁰⁴	9.61 x 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	1.36 × 10 ⁻⁰²	7.27 x 10 ⁻⁰²	2.20 x 10 ⁻⁰²	1.15 x 10 ⁻⁰¹
				Far-Future	- Incremental Pro	ject Risk			
	Cobalt	4.21 × 10 ⁻⁰⁴	4.63 x 10 ⁻⁰⁸	2.91 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	9.56×10^{-04}	2.22 x 10 ⁻⁰⁶	3.01 x 10 ⁻⁰³	4.42 x 10 ⁻⁰³
	Copper	1.51 × 10 ⁻⁰⁵	1.16 x 10 ⁻¹⁰	1.25 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	7.89×10^{-04}	6.40 x 10 ⁻⁰⁸	7.25 x 10 ⁻⁰⁴	1.53 x 10 ⁻⁰³
	Molybdenum	9.11 × 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	7.98 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	1.73×10^{-05}	$0.00 \times 10^{+00}$	9.41 × 10 ⁻⁰⁴	1.95 x 10 ⁻⁰³
	Uranium	3.42 × 10 ⁻⁰³	2.03 x 10 ⁻⁰⁵	1.83 x 10 ⁻⁰³	$0.00 \times 10^{+00}$	8.09 × 10 ⁻⁰³	2.72 x 10 ⁻⁰³	3.23 x 10 ⁻⁰³	1.93 x 10 ⁻⁰²
					Base Case				
	Cobalt	8.58 × 10 ⁻⁰⁴	9.04 × 10 ⁻⁰⁷	1.01 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.30×10^{-03}	1.44 × 10 ⁻⁰³	2.44 × 10 ⁻⁰²	2.90 x 10 ⁻⁰²
	Copper	2.91 × 10 ⁻⁰⁵	2.95 × 10 ⁻⁰⁸	4.12 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	1.79×10^{-03}	5.38 × 10 ⁻⁰⁴	4.12×10^{-02}	4.35 x 10 ⁻⁰²
	Molybdenum	1.08 × 10 ⁻⁰⁴	1.06 × 10 ⁻⁰⁷	1.62 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.42×10^{-06}	2.92 × 10 ⁻⁰⁴	6.98 × 10 ⁻⁰²	7.02 x 10 ⁻⁰²
	Uranium	2.46 × 10 ⁻⁰³	8.62 × 10 ⁻⁰⁶	2.24 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	6.85×10^{-03}	3.84 × 10 ⁻⁰²	8.20 × 10 ⁻⁰²	1.30 x 10 ⁻⁰¹
				Project Lifesp	an - Incremental F	Project Risk			
	Cobalt	6.47 × 10 ⁻⁰⁵	1.31 x 10 ⁻⁰⁹	5.83 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	1.59 × 10 ⁻⁰⁴	1.61 x 10 ⁻⁰⁵	3.21 x 10 ⁻⁰⁴	5.61 x 10 ⁻⁰⁴
Subsistence Harvester (Beet Lake)	Copper	1.24 × 10 ⁻⁰⁶	2.66 x 10 ⁻¹¹	1.30 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	7.51 × 10 ⁻⁰⁵	8.11 x 10 ⁻⁰⁷	5.12 x 10 ⁻⁰⁵	1.28 x 10 ⁻⁰⁴
(Deet Lake)	Molybdenum	4.28 × 10 ⁻⁰⁵	1.52 x 10 ⁻¹⁰	4.78 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	8.77 × 10 ⁻⁰⁷	5.41 x 10 ⁻⁰⁶	3.50 x 10 ⁻⁰⁵	8.42 x 10 ⁻⁰⁵
	Uranium	1.86 × 10 ⁻⁰³	1.65 x 10 ⁻⁰⁶	5.43 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	4.97×10^{-03}	2.70 x 10 ⁻⁰²	1.14 x 10 ⁻⁰²	4.52 x 10 ⁻⁰²
				Far-Future	- Incremental Pro	ject Risk			
	Cobalt	1.99 × 10 ⁻⁰⁴	7.08 x 10 ⁻¹⁰	2.36 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	5.34 × 10 ⁻⁰⁴	1.13 x 10 ⁻⁰⁶	1.02 x 10 ⁻⁰³	1.76 x 10 ⁻⁰³
	Copper	7.09 × 10 ⁻⁰⁶	1.78 x 10 ⁻¹²	1.00 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	4.35 × 10 ⁻⁰⁴	3.25 x 10 ⁻⁰⁸	2.75 x 10 ⁻⁰⁴	7.18 x 10 ⁻⁰⁴
	Molybdenum	3.49 × 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	5.22 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	7.79 × 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.71 × 10 ⁻⁰⁴	6.29 x 10 ⁻⁰⁴
	Uranium	1.14 × 10 ⁻⁰³	1.73 x 10 ⁻⁰⁷	1.04 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	3.17×10^{-03}	7.71 x 10 ⁻⁰⁴	1.48 x 10 ⁻⁰³	6.58 x 10 ⁻⁰³
Subsistence Harvester					Base Case				
One-Year-Old (Beet	Cobalt	9.58 × 10 ⁻⁰⁴	5.91 × 10 ⁻⁰⁵	6.62 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.18 × 10 ⁻⁰³	2.84 × 10 ⁻⁰³	4.78×10^{-02}	5.39 x 10 ⁻⁰²
Lake)	Copper	3.25 × 10 ⁻⁰⁵	1.93 × 10 ⁻⁰⁶	2.69 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	1.69 × 10 ⁻⁰³	1.06×10^{-03}	9.70×10^{-02}	9.98 x 10 ⁻⁰²



		RFD Case Hazard Quotients									
	СОРС	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC		
	Molybdenum	1.47 × 10 ⁻⁰⁴	8.46 × 10 ⁻⁰⁶	1.29 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.79 × 10 ⁻⁰⁶	7.01 × 10 ⁻⁰⁴	2.64 × 10 ⁻⁰¹	2.64 x 10 ⁻⁰¹		
	Uranium	2.75×10^{-03}	5.64×10^{-04}	1.47 x 10 ⁻⁰³	$0.00 \times 10^{+00}$	6.49×10^{-03}	7.57 × 10 ⁻⁰²	1.70×10^{-01}	2.57 x 10 ⁻⁰¹		
				Project Lifesp	an - Incremental F	Project Risk					
	Cobalt	7.22 × 10 ⁻⁰⁵	8.54 x 10 ⁻⁰⁸	3.81 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	1.51 × 10 ⁻⁰⁴	2.27 x 10 ⁻⁰⁵	3.31 × 10 ⁻⁰⁴	5.81 x 10 ⁻⁰⁴		
	Copper	1.39 × 10 ⁻⁰⁶	1.74 x 10 ⁻⁰⁹	8.50 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	7.12×10^{-05}	1.39 x 10 ⁻⁰⁶	4.76 x 10 ⁻⁰⁵	1.22 x 10 ⁻⁰⁴		
	Molybdenum	5.82 × 10 ⁻⁰⁵	1.21 x 10 ⁻⁰⁸	3.80 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	1.01×10^{-06}	9.10 x 10 ⁻⁰⁶	4.30 x 10 ⁻⁰⁵	1.15 x 10 ⁻⁰⁴		
	Uranium	2.07×10^{-03}	1.08 x 10 ⁻⁰⁴	3.55 x 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	4.71×10^{-03}	4.06 x 10 ⁻⁰²	8.06 x 10 ⁻⁰³	5.59 x 10 ⁻⁰²		
				Far-Future	- Incremental Pro	ect Risk					
	Cobalt	2.23 × 10 ⁻⁰⁴	4.63 x 10 ⁻⁰⁸	1.54 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	5.06 × 10 ⁻⁰⁴	2.22 x 10 ⁻⁰⁶	1.04 x 10 ⁻⁰³	1.79 x 10 ⁻⁰³		
	Copper	7.91 × 10 ⁻⁰⁶	1.16 x 10 ⁻¹⁰	6.56 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	4.12×10^{-04}	6.40 x 10 ⁻⁰⁸	2.54 x 10 ⁻⁰⁴	6.75 x 10 ⁻⁰⁴		
	Molybdenum	4.74×10^{-04}	$0.00 \times 10^{+00}$	4.16 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	8.99×10^{-06}	$0.00 \times 10^{+00}$	3.32 × 10 ⁻⁰⁴	8.57 x 10 ⁻⁰⁴		
	Uranium	1.27 × 10 ⁻⁰³	1.13 x 10 ⁻⁰⁵	6.79 x 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	3.01×10^{-03}	1.52 x 10 ⁻⁰³	1.08 x 10 ⁻⁰³	7.57 x 10 ⁻⁰³		
					Base (Case					
	Cobalt	8.58 × 10 ⁻⁰⁴	9.04×10^{-07}	1.01 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.30×10^{-03}	1.44×10^{-03}	2.44 × 10 ⁻⁰²	2.90 x 10 ⁻⁰²		
	Copper	2.91 × 10 ⁻⁰⁵	2.95×10^{-08}	4.12 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	1.79×10^{-03}	5.38 × 10 ⁻⁰⁴	4.12 × 10 ⁻⁰²	4.35 x 10 ⁻⁰²		
	Molybdenum	1.08 × 10 ⁻⁰⁴	1.06×10^{-07}	1.62 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.42×10^{-06}	2.92 × 10 ⁻⁰⁴	6.98 × 10 ⁻⁰²	7.02 x 10 ⁻⁰²		
	Uranium	2.46 × 10 ⁻⁰³	8.62 × 10 ⁻⁰⁶	2.24 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	6.85×10^{-03}	3.84 × 10 ⁻⁰²	8.20 × 10 ⁻⁰²	1.30 x 10 ⁻⁰¹		
	Project Lifespan - Incremental Project Risk										
	Cobalt	6.56 × 10 ⁻⁰⁶	1.31 x 10 ⁻⁰⁹	6.04 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	1.63×10^{-05}	1.61 x 10 ⁻⁰⁵	6.36 x 10 ⁻⁰⁵	1.02 x 10 ⁻⁰⁴		
Subsistence Harvester (Lloyd Lake)	Copper	1.25 × 10 ⁻⁰⁷	2.66 x 10 ⁻¹¹	1.34 x 10 ⁻¹⁰	$0.00 \times 10^{+00}$	7.57×10^{-06}	8.11 x 10 ⁻⁰⁷	1.85 x 10 ⁻⁰⁵	2.70 x 10 ⁻⁰⁵		
(Lioyu Lake)	Molybdenum	4.30 × 10 ⁻⁰⁶	1.52 x 10 ⁻¹⁰	4.92 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	8.86×10^{-08}	5.41 x 10 ⁻⁰⁶	1.29 x 10 ⁻⁰⁵	2.27 x 10 ⁻⁰⁵		
	Uranium	1.73 × 10 ⁻⁰⁴	1.95 x 10 ⁻⁰⁸	4.97 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	4.62 × 10 ⁻⁰⁴	3.19 x 10 ⁻⁰⁴	1.66 x 10 ⁻⁰⁴	1.12 x 10 ⁻⁰³		
				Far-Future	- Incremental Pro	ject Risk					
	Cobalt	2.04 × 10 ⁻⁰⁵	7.08 x 10 ⁻¹⁰	2.42 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	5.47×10^{-05}	1.13 x 10 ⁻⁰⁶	1.10 x 10 ⁻⁰⁴	1.87 x 10 ⁻⁰⁴		
	Copper	7.23×10^{-07}	1.78 x 10 ⁻¹²	1.02 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	4.44×10^{-05}	3.25 x 10 ⁻⁰⁸	2.84 x 10 ⁻⁰⁵	7.36 x 10 ⁻⁰⁵		
	Molybdenum	3.56 × 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	5.33 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	7.94×10^{-07}	$0.00 \times 10^{+00}$	2.77×10^{-05}	6.41 x 10 ⁻⁰⁵		
	Uranium	1.03 × 10 ⁻⁰⁴	2.05 x 10 ⁻⁰⁹	9.41 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.87 × 10 ⁻⁰⁴	9.12 x 10 ⁻⁰⁶	7.98 x 10 ⁻⁰⁵	4.80 x 10 ⁻⁰⁴		
					Base Case						
	Cobalt	9.58 × 10 ⁻⁰⁴	5.91 × 10 ⁻⁰⁵	6.62 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.18×10^{-03}	2.84 × 10 ⁻⁰³	4.78 × 10 ⁻⁰²	5.39 x 10 ⁻⁰²		
Subsistence Harvester One-Year-Old (Lloyd	Copper	3.25 × 10 ⁻⁰⁵	1.93 × 10 ⁻⁰⁶	2.69 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	1.69×10^{-03}	1.06 × 10 ⁻⁰³	9.70 × 10 ⁻⁰²	9.98 x 10 ⁻⁰²		
Lake)	Molybdenum	1.47 × 10 ⁻⁰⁴	8.46 × 10 ⁻⁰⁶	1.29 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.79×10^{-06}	7.01 × 10 ⁻⁰⁴	2.64 × 10 ⁻⁰¹	2.64 x 10 ⁻⁰¹		
<u> </u>	Uranium	2.75 × 10 ⁻⁰³	5.64 × 10 ⁻⁰⁴	1.47 x 10 ⁻⁰³	$0.00 \times 10^{+00}$	6.49 × 10 ⁻⁰³	7.57 × 10 ⁻⁰²	1.70 × 10 ⁻⁰¹	2.57 x 10 ⁻⁰¹		
				Project Lifesp	an - Incremental F	Project R <u>isk</u>					



			RFD Case Hazard Quotients									
	СОРС	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC			
	Cobalt	7.32 × 10 ⁻⁰⁶	8.54 x 10 ⁻⁰⁸	3.95 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	1.54×10^{-05}	2.27 x 10 ⁻⁰⁵	6.44 x 10 ⁻⁰⁵	1.10 x 10 ⁻⁰⁴			
	Copper	1.40×10^{-07}	1.74 x 10 ⁻⁰⁹	8.76 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	7.18×10^{-06}	1.39 x 10 ⁻⁰⁶	1.74 x 10 ⁻⁰⁵	2.61 x 10 ⁻⁰⁵			
	Molybdenum	5.84 × 10 ⁻⁰⁶	1.21 x 10 ⁻⁰⁸	3.91 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	1.02×10^{-07}	9.10 x 10 ⁻⁰⁶	1.59 x 10 ⁻⁰⁵	3.13 x 10 ⁻⁰⁵			
	Uranium	1.93 × 10 ⁻⁰⁴	1.27 x 10 ⁻⁰⁶	3.25 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	4.38×10^{-04}	4.80 x 10 ⁻⁰⁴	1.18 x 10 ⁻⁰⁴	1.26 x 10 ⁻⁰³			
					- Incremental Proj	ect Risk						
	Cobalt	2.28 × 10 ⁻⁰⁵	4.63 x 10 ⁻⁰⁸	1.58 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	5.19 × 10 ⁻⁰⁵	2.22 x 10 ⁻⁰⁶	1.12 x 10 ⁻⁰⁴	1.91 x 10 ⁻⁰⁴			
	Copper	8.08×10^{-07}	1.16 x 10 ⁻¹⁰	6.70 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	4.21×10^{-05}	6.40 x 10 ⁻⁰⁸	2.62 x 10 ⁻⁰⁵	6.93 x 10 ⁻⁰⁵			
	Molybdenum	4.84×10^{-05}	$0.00 \times 10^{+00}$	4.24 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	9.16×10^{-07}	$0.00 \times 10^{+00}$	3.39×10^{-05}	8.74 x 10 ⁻⁰⁵			
	Uranium	1.15 × 10 ⁻⁰⁴	1.34 x 10 ⁻⁰⁷	6.15 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.72×10^{-04}	1.80 x 10 ⁻⁰⁵	5.89 x 10 ⁻⁰⁵	5.26 x 10 ⁻⁰⁴			
					Base Case							
	Cobalt	8.58 × 10 ⁻⁰⁴	9.04×10^{-07}	1.01 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	7.27×10^{-04}	1.13 × 10 ⁻⁰³	1.88 × 10 ⁻⁰²	2.15 x 10 ⁻⁰²			
	Copper	2.91 × 10 ⁻⁰⁵	2.95×10^{-08}	4.12 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	5.66×10^{-04}	4.22×10^{-04}	3.86 × 10 ⁻⁰²	3.96 x 10 ⁻⁰²			
	Molybdenum	1.08 × 10 ⁻⁰⁴	1.06 × 10 ⁻⁰⁷	1.62 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	7.66×10^{-07}	2.29×10^{-04}	8.60 × 10 ⁻⁰²	8.63 x 10 ⁻⁰²			
	Uranium	2.46 × 10 ⁻⁰³	8.62 × 10 ⁻⁰⁶	2.24 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.17×10^{-03}	3.01 × 10 ⁻⁰²	6.79 × 10 ⁻⁰²	1.03×10^{-01}			
				Project Lifespa	an - Incremental P	roject Risk						
Seasonal Resident	Cobalt	8.76 × 10 ⁻⁰⁵	7.84 x 10 ⁻¹⁰	6.61 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	6.82×10^{-05}	4.53 x 10 ⁻⁰⁶	1.59 x 10 ⁻⁰⁴	3.19 x 10 ⁻⁰⁴			
(Patterson Lake South	Copper	1.72 × 10 ⁻⁰⁶	1.60 x 10 ⁻¹¹	1.49 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	3.28×10^{-05}	3.11 x 10 ⁻⁰⁷	6.93 x 10 ⁻⁰⁵	1.04 x 10 ⁻⁰⁴			
Arm)	Molybdenum	5.76 × 10 ⁻⁰⁵	9.10 x 10 ⁻¹¹	5.51 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	3.74×10^{-07}	1.47 x 10 ⁻⁰⁶	4.10 x 10 ⁻⁰⁵	1.01 x 10 ⁻⁰⁴			
	Uranium	3.21 × 10 ⁻⁰³	1.77 x 10 ⁻⁰⁶	8.83 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.72×10^{-03}	1.51 x 10 ⁻⁰²	5.67 x 10 ⁻⁰³	2.67 x 10 ⁻⁰²			
				Far-Future	- Incremental Proj	ect Risk						
	Cobalt	2.26 × 10 ⁻⁰⁴	4.25 x 10 ⁻¹⁰	2.67 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	1.92×10^{-04}	5.31 x 10 ⁻⁰⁷	7.00 x 10 ⁻⁰⁴	1.12 x 10 ⁻⁰³			
	Copper	8.13 × 10 ⁻⁰⁶	1.07 x 10 ⁻¹²	1.15 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	1.58×10^{-04}	1.53 x 10 ⁻⁰⁸	1.87 x 10 ⁻⁰⁴	3.53 x 10 ⁻⁰⁴			
	Molybdenum	4.02×10^{-04}	$0.00 \times 10^{+00}$	6.02 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.84×10^{-06}	$0.00 \times 10^{+00}$	1.80×10^{-04}	5.86 x 10 ⁻⁰⁴			
	Uranium	1.84 × 10 ⁻⁰³	1.86 x 10 ⁻⁰⁷	1.68 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	1.62×10^{-03}	6.49 x 10 ⁻⁰⁴	1.05 x 10 ⁻⁰³	5.17 x 10 ⁻⁰³			
					Base Case							
	Cobalt	9.58 × 10 ⁻⁰⁴	5.91 × 10 ⁻⁰⁵	6.62 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	6.89×10^{-04}	2.47 × 10 ⁻⁰³	4.13 × 10 ⁻⁰²	4.56 x 10 ⁻⁰²			
	Copper	3.25 × 10 ⁻⁰⁵	1.93 × 10 ⁻⁰⁶	2.69 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	5.36 × 10 ⁻⁰⁴	9.22 × 10 ⁻⁰⁴	9.32 × 10 ⁻⁰²	9.47 x 10 ⁻⁰²			
Seasonal Resident One-	Molybdenum	1.47×10^{-04}	8.46×10^{-06}	1.29 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	8.84×10^{-07}	6.10 × 10 ⁻⁰⁴	2.77 × 10 ⁻⁰¹	2.78 x 10 ⁻⁰¹			
Year-Old (Patterson	Uranium	2.75 × 10 ⁻⁰³	5.64 × 10 ⁻⁰⁴	1.47 x 10 ⁻⁰³	$0.00 \times 10^{+00}$	2.06×10^{-03}	6.58 × 10 ⁻⁰²	1.60 × 10 ⁻⁰¹	2.33 x 10 ⁻⁰¹			
Lake South Arm)				Project Lifespa	an - Incremental P	roject Risk						
	Cobalt	9.78 × 10 ⁻⁰⁵	5.12 x 10 ⁻⁰⁸	4.32 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	6.46×10^{-05}	7.54 x 10 ⁻⁰⁶	1.63 x 10 ⁻⁰⁴	3.37 x 10 ⁻⁰⁴			
	Copper	1.92 × 10 ⁻⁰⁶	1.04 x 10 ⁻⁰⁹	9.77 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	3.11×10^{-05}	6.24 x 10 ⁻⁰⁷	7.02 x 10 ⁻⁰⁵	1.04 x 10 ⁻⁰⁴			
	Molybdenum	7.83 × 10 ⁻⁰⁵	7.24 x 10 ⁻⁰⁹	4.38 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	4.31×10^{-07}	2.88 x 10 ⁻⁰⁶	5.24 x 10 ⁻⁰⁵	1.38 x 10 ⁻⁰⁴			



	RFD Case Hazard Quotients									
	СОРС	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC	
	Uranium	3.58 × 10 ⁻⁰³	1.16 x 10 ⁻⁰⁴	5.77 x 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	2.58 × 10 ⁻⁰³	2.71 x 10 ⁻⁰²	4.17 x 10 ⁻⁰³	3.81 x 10 ⁻⁰²	
				Far-Future	- Incremental Proj	ect Risk				
	Cobalt	2.52 × 10 ⁻⁰⁴	2.78 x 10 ⁻⁰⁸	1.75 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	1.82 × 10 ⁻⁰⁴	1.16 x 10 ⁻⁰⁶	7.22 x 10 ⁻⁰⁴	1.17 x 10 ⁻⁰³	
	Copper	9.08 × 10 ⁻⁰⁶	6.98 x 10 ⁻¹¹	7.53 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	1.50 × 10 ⁻⁰⁴	3.34 x 10 ⁻⁰⁸	1.77 x 10 ⁻⁰⁴	3.37 x 10 ⁻⁰⁴	
	Molybdenum	5.47 × 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	4.79 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	3.28×10^{-06}	$0.00 \times 10^{+00}$	2.24×10^{-04}	8.22 x 10 ⁻⁰⁴	
	Uranium	2.05 × 10 ⁻⁰³	1.22 x 10 ⁻⁰⁵	1.10 x 10 ⁻⁰³	$0.00 \times 10^{+00}$	1.54×10^{-03}	1.42 x 10 ⁻⁰³	7.65 x 10 ⁻⁰⁴	6.88 x 10 ⁻⁰³	
					Base Case					
	Cobalt	8.58 × 10 ⁻⁰⁴	9.04 × 10 ⁻⁰⁷	1.01 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	7.27×10^{-04}	1.13 × 10 ⁻⁰³	1.88 × 10 ⁻⁰²	2.15 x 10 ⁻⁰²	
	Copper	2.91 × 10 ⁻⁰⁵	2.95 × 10 ⁻⁰⁸	4.12 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	5.66 × 10 ⁻⁰⁴	4.22 × 10 ⁻⁰⁴	3.86 × 10 ⁻⁰²	3.96 x 10 ⁻⁰²	
	Molybdenum	1.08 × 10 ⁻⁰⁴	1.06 × 10 ⁻⁰⁷	1.62 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	7.66 × 10 ⁻⁰⁷	2.29 × 10 ⁻⁰⁴	8.60 × 10 ⁻⁰²	8.63 x 10 ⁻⁰²	
	Uranium	2.46 × 10 ⁻⁰³	8.62 × 10 ⁻⁰⁶	2.24 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.17×10^{-03}	3.01 × 10 ⁻⁰²	6.79 × 10 ⁻⁰²	1.03 x 10 ⁻⁰¹	
				Project Lifesp	an - Incremental P	Project Risk				
C	Cobalt	3.93 × 10 ⁻⁰⁶	7.84 x 10 ⁻¹⁰	3.62 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	5.54 × 10 ⁻⁰⁶	4.53 x 10 ⁻⁰⁶	1.47 x 10 ⁻⁰⁵	2.87 x 10 ⁻⁰⁵	
Seasonal Resident (Lloyd Lake)	Copper	7.51 × 10 ⁻⁰⁸	1.60 x 10 ⁻¹¹	8.05 x 10 ⁻¹¹	$0.00 \times 10^{+00}$	1.88×10^{-04}	3.11 x 10 ⁻⁰⁷	4.24 x 10 ⁻⁰⁶	1.93 x 10 ⁻⁰⁴	
(Lioya Lake)	Molybdenum	2.58 × 10 ⁻⁰⁶	9.10 x 10 ⁻¹¹	2.95 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	2.93×10^{-08}	1.47 x 10 ⁻⁰⁶	2.98 x 10 ⁻⁰⁶	7.06 x 10 ⁻⁰⁶	
	Uranium	1.04 × 10 ⁻⁰⁴	1.17 x 10 ⁻⁰⁸	2.98 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	7.09×10^{-04}	9.99 x 10 ⁻⁰⁵	3.26 x 10 ⁻⁰⁵	9.46 x 10 ⁻⁰⁴	
					- Incremental Proj					
	Cobalt	1.23 × 10 ⁻⁰⁵	4.25 x 10 ⁻¹⁰	1.45 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	1.76 × 10 ⁻⁰⁵	5.31 x 10 ⁻⁰⁷	2.67 x 10 ⁻⁰⁵	5.70 x 10 ⁻⁰⁵	
	Copper	4.34 × 10 ⁻⁰⁷	1.07 x 10 ⁻¹²	6.15 x 10 ⁻¹⁰	$0.00 \times 10^{+00}$	2.04 × 10 ⁻⁰⁴	1.53 x 10 ⁻⁰⁸	7.19 x 10 ⁻⁰⁶	2.12 x 10 ⁻⁰⁴	
	Molybdenum	2.14 × 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	3.20 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	2.52 × 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	6.65 × 10 ⁻⁰⁶	2.83 x 10 ⁻⁰⁵	
	Uranium	6.19 × 10 ⁻⁰⁵	1.23 x 10 ⁻⁰⁹	5.64 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	6.37×10^{-04}	4.29 x 10 ⁻⁰⁶	2.26 x 10 ⁻⁰⁵	7.26 x 10 ⁻⁰⁴	
					Base Case					
	Cobalt	9.58 × 10 ⁻⁰⁴	5.91 × 10 ⁻⁰⁵	6.62 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	6.89 × 10 ⁻⁰⁴	2.47 × 10 ⁻⁰³	4.13 × 10 ⁻⁰²	4.56 x 10 ⁻⁰²	
	Copper	3.25 × 10 ⁻⁰⁵	1.93 × 10 ⁻⁰⁶	2.69 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	5.36 × 10 ⁻⁰⁴	9.22 × 10 ⁻⁰⁴	9.32 × 10 ⁻⁰²	9.47 x 10 ⁻⁰²	
	Molybdenum	1.47 × 10 ⁻⁰⁴	8.46 × 10 ⁻⁰⁶	1.29 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	8.84 × 10 ⁻⁰⁷	6.10 × 10 ⁻⁰⁴	2.77 × 10 ⁻⁰¹	2.78 x 10 ⁻⁰¹	
	Uranium	2.75 × 10 ⁻⁰³	5.64 × 10 ⁻⁰⁴	1.47 x 10 ⁻⁰³	$0.00 \times 10^{+00}$	2.06 × 10 ⁻⁰³	6.58 × 10 ⁻⁰²	1.60 × 10 ⁻⁰¹	2.33 x 10 ⁻⁰¹	
Seasonal Resident One		•	1	Project Lifesp	an - Incremental P	Project Risk				
Year-Old (Lloyd Lake)	Cobalt	4.39 × 10 ⁻⁰⁶	5.12 x 10 ⁻⁰⁸	2.37 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.93 × 10 ⁻⁰⁶	7.54 x 10 ⁻⁰⁶	1.50 x 10 ⁻⁰⁵	3.02 x 10 ⁻⁰⁵	
	Copper	8.39 × 10 ⁻⁰⁸	1.04 x 10 ⁻⁰⁹	5.26 x 10 ⁻⁰⁹	$0.00 \times 10^{+00}$	1.36 × 10 ⁻⁰⁶	6.24 x 10 ⁻⁰⁷	4.08 x 10 ⁻⁰⁶	6.15 x 10 ⁻⁰⁶	
	Molybdenum	3.51 × 10 ⁻⁰⁶	7.24 x 10 ⁻⁰⁹	2.35 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	1.94 × 10 ⁻⁰⁸	2.88 x 10 ⁻⁰⁶	3.73 x 10 ⁻⁰⁶	1.04 x 10 ⁻⁰⁵	
	Uranium	1.16 × 10 ⁻⁰⁴	7.64 x 10 ⁻⁰⁷	1.95 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	8.33 × 10 ⁻⁰⁵	1.79 x 10 ⁻⁰⁴	2.30 x 10 ⁻⁰⁵	4.21 x 10 ⁻⁰⁴	
			· · ·		- Incremental Proj					
	Cobalt	1.37 × 10 ⁻⁰⁵	2.78 x 10 ⁻⁰⁸	9.47 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	9.86 × 10 ⁻⁰⁶	1.16 x 10 ⁻⁰⁶	2.70 x 10 ⁻⁰⁵	5.26 x 10 ⁻⁰⁵	



					RFD Case Haza	rd Quotients					
	СОРС	Water (Internal)	Soil (Internal)	Sediment (Internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC		
	Copper	4.85 × 10 ⁻⁰⁷	6.98 x 10 ⁻¹¹	4.02 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	8.00 × 10 ⁻⁰⁶	3.34 x 10 ⁻⁰⁸	6.53 x 10 ⁻⁰⁶	1.51 x 10 ⁻⁰⁵		
	Molybdenum	2.90 × 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.54 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	1.74 × 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	8.14 × 10 ⁻⁰⁶	3.99 x 10 ⁻⁰⁵		
	Uranium	6.91 × 10 ⁻⁰⁵	8.03 x 10 ⁻⁰⁸	3.69 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	5.18 × 10 ⁻⁰⁵	9.37 x 10 ⁻⁰⁶	1.62 x 10 ⁻⁰⁵	1.83 x 10 ⁻⁰⁴		
					Base Case						
	Cobalt	8.58 × 10 ⁻⁰⁴	9.04×10^{-07}	1.01 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.30×10^{-03}	1.44×10^{-03}	2.44 × 10 ⁻⁰²	2.90 x 10 ⁻⁰²		
	Copper	2.91 × 10 ⁻⁰⁵	2.95×10^{-08}	4.12 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	1.79×10^{-03}	5.38 × 10 ⁻⁰⁴	4.12 × 10 ⁻⁰²	4.35 x 10 ⁻⁰²		
	Molybdenum	1.08 × 10 ⁻⁰⁴	1.06×10^{-07}	1.62 x 10 ⁻⁰⁷	$0.00 \times 10^{+00}$	2.42×10^{-06}	2.92 × 10 ⁻⁰⁴	6.98 × 10 ⁻⁰²	7.02 x 10 ⁻⁰²		
	Uranium	2.46 × 10 ⁻⁰³	8.62 × 10 ⁻⁰⁶	2.24 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	6.85×10^{-03}	3.84 × 10 ⁻⁰²	8.20 × 10 ⁻⁰²	1.30 x 10 ⁻⁰¹		
				Base	Case One-Year-O	ld					
	Cobalt	9.58 × 10 ⁻⁰⁴	5.91 × 10 ⁻⁰⁵	6.62 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.18×10^{-03}	2.84 × 10 ⁻⁰³	4.78 × 10 ⁻⁰²	5.39 x 10 ⁻⁰²		
	Copper	3.25 × 10 ⁻⁰⁵	1.93×10^{-06}	2.69 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	1.69×10^{-03}	1.06 × 10 ⁻⁰³	9.70 × 10 ⁻⁰²	9.98 x 10 ⁻⁰²		
	Molybdenum	1.47×10^{-04}	8.46×10^{-06}	1.29 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.79×10^{-06}	7.01×10^{-04}	2.64 × 10 ⁻⁰¹	2.64 x 10 ⁻⁰¹		
Permanent Resident	Uranium	2.75×10^{-03}	5.64 × 10 ⁻⁰⁴	1.47 x 10 ⁻⁰³	$0.00 \times 10^{+00}$	6.49×10^{-03}	7.57 × 10 ⁻⁰²	1.70×10^{-01}	2.57 x 10 ⁻⁰¹		
(Patterson Lake North	Far-Future Adult - Incremental Project Risk										
Arm - West Basin)	Cobalt	1.24 × 10 ⁻⁰³	1.42 x 10 ⁻⁰⁹	1.47 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	3.32×10^{-03}	2.26 x 10 ⁻⁰⁶	6.36 x 10 ⁻⁰³	1.09 x 10 ⁻⁰²		
	Copper	4.62 × 10 ⁻⁰⁵	6.13 x 10 ⁻¹¹	6.54 x 10 ⁻⁰⁸	$0.00 \times 10^{+00}$	2.84×10^{-03}	1.12 x 10 ⁻⁰⁶	1.81 x 10 ⁻⁰³	4.69 x 10 ⁻⁰³		
	Molybdenum	2.31 × 10 ⁻⁰³	$0.00 \times 10^{+00}$	3.46 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	5.15×10^{-05}	$0.00 \times 10^{+00}$	1.80×10^{-03}	4.16 x 10 ⁻⁰³		
	Uranium	2.21 × 10 ⁻⁰²	3.88 x 10 ⁻⁰⁶	2.02 x 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	6.17×10^{-02}	1.73 x 10 ⁻⁰²	3.09 x 10 ⁻⁰²	1.32 x 10 ⁻⁰¹		
			Fai	r-Future One-Ye	ar-Old - Incremen	tal Project Risk					
	Cobalt	1.39 × 10 ⁻⁰³	9.25 x 10 ⁻⁰⁸	9.59 x 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	3.15 × 10 ⁻⁰³	4.45 x 10 ⁻⁰⁶	6.46 x 10 ⁻⁰³	1.11 x 10 ⁻⁰²		
	Copper	5.16 × 10 ⁻⁰⁵	4.00 x 10 ⁻⁰⁹	4.28 x 10 ⁻⁰⁶	$0.00 \times 10^{+00}$	2.69 × 10 ⁻⁰³	2.20 x 10 ⁻⁰⁶	1.67 x 10 ⁻⁰³	4.41 x 10 ⁻⁰³		
	Molybdenum	3.14 × 10 ⁻⁰³	$0.00 \times 10^{+00}$	2.75 x 10 ⁻⁰⁴	$0.00 \times 10^{+00}$	5.95 × 10 ⁻⁰⁵	$0.00 \times 10^{+00}$	2.20 × 10 ⁻⁰³	5.67 x 10 ⁻⁰³		
	Uranium	2.47 × 10 ⁻⁰²	2.54 x 10 ⁻⁰⁴	1.32 x 10 ⁻⁰²	$0.00 \times 10^{+00}$	5.85×10^{-02}	3.41 x 10 ⁻⁰²	2.24 x 10 ⁻⁰²	1.53 x 10 ⁻⁰¹		

Note: **Bold** indicates exceedance of the HQ of 0.2.

HQ = hazard quotient; COPC = constituent of potential concern; RFD = reasonably foreseeable development; n/a = receptor not applicable.



5.4.1.2.2 Carcinogen Risk

The arsenic ILCR is expected to exceed the cancer risk level of 1 in 100,000 for the adult camp worker, subsistence harvester, and seasonal resident harvesting Traditional Foods from Patterson Lake South Arm during Operations for the RFD Case (Table 5-25). The arsenic ILCR would be below the cancer risk level of 1 in 100,000 for all other human receptors during Operations.

The predicted cancer risk (ILCR) for the RFD Case was slightly higher than the Application Case mainly due to the additional effluent assumed to be released to Patterson Lake South Arm from the Fission Patterson Lake South Property. The camp worker, subsistence harvester, and seasonal resident were assumed to consume Traditional Foods from Patterson Lake South Arm.

The greatest contributor to the predicted ILCRs for the RFD Case during the Project lifespan for the seasonal resident and subsistence harvester at Patterson Lake South Arm is consumption of northern pike and beaver (Table 5-26).

In the far-future projection, the cancer risk for the future permanent resident at the camp site and subsistence harvester would be similar between the RFD Case and the Application Case. There are no expected inputs to the environment in the far-future projection from the Fission Patterson Lake South Property.

The interpretation of cancer risk is discussed in Section 5.4.1.1.2, Carcinogen Risk.

Table 5-25: Estimated Incremental Lifetime Cancer Risk from Arsenic to Human Receptors – Reasonably Foreseeable Development Case

Basantan	RFD Case Cancer Risk (per 100,000)			
Receptor	Project Lifespan	Far-Future		
Camp worker (adult)	3	n/a		
Subsistence harvester - Patterson Lake South Arm (composite)	11	0.1		
Subsistence harvester - Beet Lake (composite)	0.2	0.004		
Subsistence harvester - Lloyd Lake (composite)	0.1	0.0002		
Seasonal resident - Patterson Lake South Arm (composite)	2	0.03		
Seasonal resident - Lloyd Lake (composite)	0.02	0.00007		
Permanent resident - camp (composite)	n/a	0.5		

Bold indicates exceedance of the negligible cancer risk level of 1 in 100,000.

n/a = not applicable, receptor was not assessed; RFD = reasonably foreseeable development.



Table 5-26: Estimated Carcinogen Risks for Arsenic to Subsistence Harvesters and Seasonal Resident – Operations – Reasonably Foreseeable Development

	•								
			istence Harvester – Subsistence Harvester – Seasonal Resident – Se			easonal Resident – Lloyd Lake			
ILCR	% Contribution	ILCR	% Contribution	ILCR	% Contribution	ILCR	% Contribution	ILCR	% Contribution
2.44 × 10 ⁻⁰⁷	0.2%	2.44 × 10 ⁻⁰⁷	10%	2.44 × 10 ⁻⁰⁷	23%	5.04 × 10 ⁻⁰⁸	0.2%	5.04 × 10 ⁻⁰⁸	22%
3.46 × 10 ⁻⁰⁸	0.03%	3.46 × 10 ⁻⁰⁸	1%	3.46 × 10 ⁻⁰⁸	3%	2.08 × 10 ⁻⁰⁸	0.1%	2.08 × 10 ⁻⁰⁸	9%
2.57 × 10 ⁻⁰⁵	22%	1.94 × 10 ⁻⁰⁷	8%	1.13 × 10 ⁻⁰⁸	1%	4.88 × 10 ⁻⁰⁶	21%	2.15 × 10 ⁻⁰⁹	1%
2.06 × 10 ⁻⁰⁵	18%	1.78 × 10 ⁻⁰⁷	8%	1.03 × 10 ⁻⁰⁸	1%	3.91 × 10 ⁻⁰⁶	17%	1.96 × 10 ⁻⁰⁹	1%
0.00 × 10 ⁺⁰⁰	0%	0.00 × 10 ⁺⁰⁰	0%	0.00 × 10 ⁺⁰⁰	0%	0.00 × 10 ⁺⁰⁰	0%	0.00 × 10 ⁺⁰⁰	0%
1.26 × 10 ⁻⁰⁵	11%	2.54 × 10 ⁻⁰⁷	11%	9.63 × 10 ⁻⁰⁸	9%	2.38 × 10 ⁻⁰⁶	10%	1.83 × 10 ⁻⁰⁸	8%
1.76 × 10 ⁻⁰⁵	15%	3.56 × 10 ⁻⁰⁷	15%	1.35 × 10 ⁻⁰⁷	13%	4.13 × 10 ⁻⁰⁶	18%	3.17 × 10 ⁻⁰⁸	14%
2.88 × 10 ⁻⁰⁵	25%	8.24 × 10 ⁻⁰⁷	35%	4.70 × 10 ⁻⁰⁷	44%	5.27 × 10 ⁻⁰⁶	23%	8.60 × 10 ⁻⁰⁸	38%
9.04 × 10 ⁻⁰⁶	8%	1.93 × 10 ⁻⁰⁷	8%	1.12 × 10 ⁻⁰⁸	1%	2.67 × 10 ⁻⁰⁶	11%	3.30 × 10 ⁻⁰⁹	1%
9.56 × 10 ⁻⁰⁸	0.1%	6.63 × 10 ⁻⁰⁸	3%	6.61 × 10 ⁻⁰⁸	6%	1.53 × 10 ⁻⁰⁸	0.1%	1.06 × 10 ⁻⁰⁸	5%
1.15 × 10 ⁻⁰⁴	100%	2.34 × 10 ⁻⁰⁶	100%	1.08 × 10 ⁻⁰⁶	100%	2.33 × 10 ⁻⁰⁵	100%	2.25 × 10 ⁻⁰⁷	100%
	Patterson La	ILCR Contribution 2.44×10^{-07} 0.2% 3.46×10^{-08} 0.03% 2.57×10^{-05} 22% 2.06×10^{-05} 18% $0.00 \times 10^{+00}$ 0% 1.26×10^{-05} 11% 1.76×10^{-05} 15% 2.88×10^{-05} 25% 9.04×10^{-06} 8% 9.56×10^{-08} 0.1%	Patterson Lake South Arm % Contribution ILCR 2.44 × 10 ⁻⁰⁷ 0.2% 2.44 × 10 ⁻⁰⁷ 3.46 × 10 ⁻⁰⁸ 0.03% 3.46 × 10 ⁻⁰⁸ 2.57 × 10 ⁻⁰⁵ 22% 1.94 × 10 ⁻⁰⁷ 2.06 × 10 ⁻⁰⁵ 18% 1.78 × 10 ⁻⁰⁷ 0.00 × 10 ⁺⁰⁰ 0% 0.00 × 10 ⁺⁰⁰ 1.26 × 10 ⁻⁰⁵ 11% 2.54 × 10 ⁻⁰⁷ 1.76 × 10 ⁻⁰⁵ 15% 3.56 × 10 ⁻⁰⁷ 2.88 × 10 ⁻⁰⁵ 25% 8.24 × 10 ⁻⁰⁷ 9.04 × 10 ⁻⁰⁶ 8% 1.93 × 10 ⁻⁰⁷ 9.56 × 10 ⁻⁰⁸ 0.1% 6.63 × 10 ⁻⁰⁸	Patterson Lake South Arm Beet Lake ILCR % Contribution ILCR % Contribution 2.44 × 10 ⁻⁰⁷ 0.2% 2.44 × 10 ⁻⁰⁷ 10% 3.46 × 10 ⁻⁰⁸ 0.03% 3.46 × 10 ⁻⁰⁸ 1% 2.57 × 10 ⁻⁰⁵ 22% 1.94 × 10 ⁻⁰⁷ 8% 2.06 × 10 ⁻⁰⁵ 18% 1.78 × 10 ⁻⁰⁷ 8% 0.00 × 10 ⁺⁰⁰ 0% 0.00 × 10 ⁺⁰⁰ 0% 1.26 × 10 ⁻⁰⁵ 11% 2.54 × 10 ⁻⁰⁷ 11% 1.76 × 10 ⁻⁰⁵ 15% 3.56 × 10 ⁻⁰⁷ 15% 2.88 × 10 ⁻⁰⁵ 25% 8.24 × 10 ⁻⁰⁷ 35% 9.04 × 10 ⁻⁰⁶ 8% 1.93 × 10 ⁻⁰⁷ 8% 9.56 × 10 ⁻⁰⁸ 0.1% 6.63 × 10 ⁻⁰⁸ 3%	ILCR % Contribution ILCR % Contribution ILCR % Contribution ILCR ILCR	ILCR % Contribution ILCR % Contribution ILCR % Contribution ILCR % Contribution 2.44×10^{-07} 0.2% 2.44×10^{-07} 10% 2.44×10^{-07} 23% 3.46×10^{-08} 0.03% 3.46×10^{-08} 1% 3.46×10^{-08} 3% 2.57×10^{-05} 22% 1.94×10^{-07} 8% 1.13×10^{-08} 1% 2.06×10^{-05} 18% 1.78×10^{-07} 8% 1.03×10^{-08} 1% $0.00 \times 10^{+00}$ 0% $0.00 \times 10^{+00}$ 0% $0.00 \times 10^{+00}$ 0% 1.26×10^{-05} 11% 2.54×10^{-07} 11% 9.63×10^{-08} 9% 1.76×10^{-05} 15% 3.56×10^{-07} 15% 1.35×10^{-07} 13% 2.88×10^{-05} 25% 8.24×10^{-07} 35% 4.70×10^{-07} 44% 9.04×10^{-06} 8% 1.93×10^{-07} 8% 1.12×10^{-08} 1% 9.56×10^{-08} 0.1%	Registron Lake South Arm Registration Lick South Arm Registration Lick South Arm Registration Registrati	Patterson Lake South Arm Beet Lake Lloyd Lake Patterson Lake South Arm	Patterson Lake South Arm Beet Lake Lloyd Lake Patterson Lake South Arm LCR % Contribution LCR Contribution
Note: Total percentages may not sum to 100% due to rounding.

ILCR = incremental lifetime cancer risk.



5.4.1.2.3 Radiological Risk

With the addition of releases from the Fission Patterson Lake South Property, the incremental radiation dose to the subsistence harvester (one-year-old) who eats Traditional Foods gathered at Patterson Lake South Arm would increase from 0.07 mSv/yr (Application Case) to 0.14 mSv/yr (RFD Case). For the RFD Case, the dose to all human receptors would remain below the public regulatory dose limit of 1 mSv/yr, and below the dose constraint of 0.3 mSv/yr (Table 5-27, Table 5-28).

In the far-future projection, a future permanent resident, residing at the decommissioned and reclaimed Project site well beyond Closure could receive a dose up to 0.1 mSv/yr, well below both the regulatory public dose limit and the dose constraint.

Overall, since the radiation dose estimates were below the public dose limit, no discernable health effects would be anticipated due to exposure of these receptors to cumulative radioactive releases from both the Project and the Fission Patterson Lake South Property.

Table 5-27: Summary of All Radiation Doses to Human Receptors – Reasonably Foreseeable Development Case

		RFD (Case
Receptor	Location	Project Lifespan	Far-Future
·		Dose (m	ıSv/yr)
Camp worker	Permanent Camp	8.58 x 10 ⁻⁰²	n/a
Harvester	Patterson Lake South Arm	1.11 x 10 ⁻⁰¹	5.21 x 10 ⁻⁰³
Harvester_1y	Patterson Lake South Arm	1.36 x 10 ⁻⁰¹	6.17 x 10 ⁻⁰³
Harvester	Beet Lake	3.16 x 10 ⁻⁰²	2.19 x 10 ⁻⁰³
Harvester_1y	Beet Lake	3.90 x 10 ⁻⁰²	2.71 x 10 ⁻⁰³
Harvester	Lloyd Lake	1.84 x 10 ⁻⁰³	1.14 x 10 ⁻⁰⁴
Harvester_1y	Lloyd Lake	2.06 x 10 ⁻⁰³	1.36 x 10 ⁻⁰⁴
Seasonal Resident	Patterson Lake South Arm	2.91 x 10 ⁻⁰²	2.42 x 10 ⁻⁰³
Seasonal Resident_1y	Patterson Lake South Arm	3.75 x 10 ⁻⁰²	3.10 x 10 ⁻⁰³
Seasonal Resident	Lloyd Lake	1.84 x 10 ⁻⁰³	1.33 x 10 ⁻⁰³
Seasonal Resident_1y	Lloyd Lake	6.24 x 10 ⁻⁰⁴	7.54 x 10 ⁻⁰⁵
Permanent Resident	Permanent Camp	n/a	5.18 x 10 ⁻⁰²
Permanent Resident_1y	Permanent Camp	n/a	6.66 x 10 ⁻⁰²

 $n/a = not \ applicable, \ receptor \ was \ not \ assessed; \ mSv/yr = millisieverts \ per \ year; \ RFD = reasonably \ foreseeable \ development.$



Table 5-28: Summary of Radiation Doses to Limiting Human Receptors

	Maximum Dose Receptor (mSv/yr)		Location	Largest Contributor to Dose	% of Dose Limit
		Proj	ject Lifespan		
Application Case	0.07	Subsistence Harvester one-year- old	Patterson Lake South Arm	Polonium-210 terrestrial animals	7%
Upper Bound Scenario	0.12	Subsistence Harvester one-year- old	Patterson Lake South Arm	Polonium-210 terrestrial animals	12%
RFD Case	0.14	Subsistence Harvester one-year- old	Patterson Lake South Arm	Polonium-210 terrestrial animals	14%
		F	ar-Future		
Application Case	0.07	Permanent resident one-year-old	Patterson Lake North Arm – West Basin	Polonium-210 terrestrial animals	7%
Upper Bound Scenario	0.07	Permanent resident one-year-old	Patterson Lake North Arm – West Basin	Polonium-210 terrestrial animals	7%
RFD Case	0.07	Permanent resident one-year-old	Patterson Lake North Arm – West Basin	Polonium-210 terrestrial animals	7%

RFD = reasonably foreseeable development; mSv/yr = millisieverts per year.

5.4.1.2.4 Radon Risk

For the RFD Case, the incremental radon dose to the camp worker was estimated to be 0.57 mSv/yr. The incremental radon dose was based on atmospheric modelling of radon released during the maximum ore grade year (Year 1 of the Project) as well as radon releases from the Fission Patterson Lake South Property. This is a conservative assumption, as radon is expected to be released at a much lower rate over the duration of the Project. Additionally, the assessment is conservative in that it assumes that the camp worker spends 100% of the time indoors.

The total incremental dose to the camp worker from all radionuclides in the U-238 decay chain including radon would be 0.66 mSv/yr, which is below the dose limit for a non-nuclear energy worker of 1 mSv/yr (Table 5-29). The estimate of total dose including radon is conservative based on the following assumptions:

- The camp worker spends 100% of their time indoors when on site for exposure to radon (Section 5.2.4.2.4).
- Receptors are exposed to the maximum exposure concentrations at their location for each model scenario and Project phase (Section 5.2.5).



• For radionuclides in the U-238 decay chain (other than radon), the camp worker is also exposed to radionuclides through ingestion (water and food) pathways resulting in a conservative dose when also factoring in the dose from radon indoors.

Table 5-29: Total Radiation Dose to Camp Worker from all Radionuclides including Radon Progeny – Project Lifespan (RFD Case)

Human Receptor	Location	Maximum Total Incremental Dose – U-238 Decay Chain (mSv/yr)	Maximum Radon Dose (mSv/yr)	Maximum Total Dose (mSv/yr)	% of Dose Limit for Non-Nuclear Energy Worker
Camp Worker	Camp	8.58 x 10 ⁻⁰²	5.70 x 10 ⁻⁰¹	6.56 x 10 ⁻⁰¹	65.6%

5.4.2 Uncertainty in the Risk Characterization

The problem formulation and toxicity and exposure information are combined in the risk characterization step to estimate the potential for human health effects. The uncertainties associated with each of the previous steps of the HHRA are discussed in Section 5.1.5, Uncertainty in Problem Formulation, Section 5.2.5, Uncertainty in Exposure Assessment, and Section 5.3.3, Uncertainty in the Toxicity Assessment. In each step of the HHRA, conservative assumptions were used to address uncertainties. The use of this approach is far more likely to overestimate potential risk than to underestimate risk.

6.0 ECOLOGICAL RISK ASSESSMENT

6.1 Problem Formulation

The intent of the problem formulation for an EcoRA is to define the goals of the risk assessment, develop an understanding of site conditions, and develop working hypotheses as to how potential exposure of ecological receptors to contaminants may result in potential ecological risks.

The problem formulation for this EcoRA:

- identifies COPCs for ecological health;
- identifies and characterizes the selected ecological receptors for the Project; and
- identifies the complete exposure pathways by which the COPCs may affect the receptors in a conceptual site model.

The conceptual site model for the EcoRA summarizes the links between contaminant sources, exposure pathways, and receptors of concern.

6.1.1 Selected Ecological Receptors

It is generally an impractical task to assess the effect of radiological and non-radiological emissions on all the species within a natural ecosystem (CSA 2022), and specifically within the ecosystem around the Project. Therefore, a representative group of organisms, referred to as ecological receptors, were chosen for the effects assessment and risk analysis. The ecological receptors selected for the ERA are a subset of valued components identified for the EA so that the results from the ERA could be integrated into the effects assessments for fish, vegetation, wildlife, human health, Indigenous land and resource use, and other land and resource uses.

The receptor selection process considered some of the same criteria as for VCs (EIS Section 6.3.1, Valued Components), such as: they are known to exist at the site and in the LSA, are representative of major taxonomic groups or exposure pathways, are listed federally and/or provincially, have a special importance or value to people including Indigenous communities, and/or meet other receptor selection criteria as further described in this subsection. In addition, the selection of ecological receptors considered the identification of potential exposure pathways to COPCs and availability of data from chemical analyses for radiological and non-radiological parameters for the species selected. As recognized in CSA N288.6-22, protection of the selected species should provide reasonable assurance that all species are protected.

6.1.1.1 Selection of Ecological Receptors

A preliminary list of ecological valued components for the Project was compiled from the species identified in EIS Section 6, Environmental Assessment Approach and Methods, as well as the following baseline reports:

Annex V.1, Aquatic Environment Baseline Report;



- Annex VII.2, Vegetation Baseline Report 2 (Inventory, Rare Plants, and Wetlands);
- Annex VIII.2, Wildlife Baseline Report 2 (Amphibians, Birds, and Bats);
- Annex VIII.3, Wildlife Baseline Report 3 (Bird Migration and Bats);
- Annex VII.1, Vegetation Baseline Report 1 (Mapping); and
- Annex VIII.1 Wildlife Baseline Report 1 (Mammals, Waterfowl, and Raptors).

The list was reviewed again once data from the 2020 updated aquatic and terrestrial baseline studies were available (Annex VIII.2; Annex VIII.3; Annex VII.1; Annex VIII.1) to confirm that the compiled list was still appropriate. Additional information from avian migration and acoustic bat studies were also considered (Annex VIII.3). Species were included in the preliminary list if they were quantified and/or incidentally observed through respective survey methods. Species that were identified to be located within the LSA from searches of the Hunting, Angling and Biodiversity Information of Saskatchewan (HABISask) database, but were not observed in the baseline surveys, were excluded from further consideration.

A representative subset of organisms was selected from each major plant or animal group to be carried forward as ecological receptors. Several factors were considered in the selection process, following the criteria provided in Table 7.1 of CSA N288.6-22:

- Major plant or animal group:
 - Availability of scientific information for the receptor that can be used in risk analysis. For example, the Canadian toad was not included on the ecological receptor list on the basis of limited scientific information for amphibian risk assessment. However, the assessment of a fish receptor is considered to be protective of the most sensitive, tadpole life stage.
- Presence in the site, LSA, and/or RSA:
 - Abundance of the species in the study area relative to other species from the same organism group. For example, the most frequently observed songbird species for the area may be carried forward to represent other songbirds.
 - Availability of baseline data from chemical analyses for radiological and non-radiological parameters for the species. For example, the southern red-backed vole was selected on this basis.
- Socio-economic or ecological value and value to Indigenous communities:
 - Value or importance to Indigenous communities, based on information from the draft Traditional Land Use Desktop Study (InterGroup 2018a), the IKTLU Studies (TSD II: BNDN; TSD III: BRDN; TSD IV: MN-S; TSD V.1: CRDN; TSD V.2: CRDN; TSD VI:



ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT ECOLOGICAL RISK ASSESSMENT

YNLR) and feedback during subsequent community information sessions and JWG meetings.

- Value or importance to other land and resource uses in the area, based on the Other Land and Resource Use Memorandum (InterGroup 2018b). This report identified northern pike and lake whitefish as targets of commercial fishing in Patterson Lake, Lloyd Lake, and Lake La Loche.
- Classification as endangered, threatened, or species of special concern identified by the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) and listed under the federal *Species at Risk Act* (SARA) (e.g., woodland caribou).
- Exposed to and/or sensitive to stressors:
 - Representing a potential exposure pathway to COPCs through releases to the environment.

The major plant and animal groups were defined based on taxonomy and ecology so as to represent the different possible pathways of exposure to COPCs. The organisms selected to represent each major group were either individual species of vertebrate organisms, or generic types of plants or invertebrates such as aquatic macrophytes or zooplankton.

Species at Risk (SAR) often lack the information needed for risk assessment because they are difficult to study. It is acceptable practice in ERA to use a surrogate species approach to assess potential risk to SAR, where SAR are represented by other more common species that have similar diets and exposure pathways. For example, the rusty blackbird was selected to represent other aerial insectivores, which include the following SAR: common nighthawk, olive-sided flycatcher, and barn swallow.

While a surrogate species may be selected to represent a SAR, the species listed in the Government of Canada's SARA or its regulations, or in corresponding Saskatchewan statues and legislation were assessed at the individual level in the ERA, as effects on even a single individual would not be acceptable (CSA 2022).

The selection of ecological receptors was informed by IKTLU Studies, information obtained during community information sessions, and JWG meetings.

- Terrestrial vegetation (lichen, blueberry, Labrador tea):
 - The CRDN IKTLU identified the use of berries (blueberries, cranberries [bog, low bush], gooseberries, Saskatoon, cloudberry, strawberries), medicines (kinnikinnick, Labrador tea, mint, spruce gum, sweet flag [rat root], mushrooms), shrubs (dogwood [red willow], willows [spp.]), trees (birch, jack pine, poplar, spruce [spp.], tamarack) and other vegetal matter (barks, mosses, roots, punk [rotten wood]; TSD V.1: CRDN).



- The MN-S identified the use of berries, such as blueberries, strawberries and raspberries, as important Traditional Foods since they are full of antioxidants and promote heart and brain health, and identified the Patterson Lake area as having a bounty for harvest (TSD IV: MN-S).
- The Athabasca Denesųłiné IKTLU identified specific plants harvested including Ts'aili Teli (Frog Tail), Labrador Tea, and blueberries (TSD VI: YNLR).

Fish (northern pike, lake whitefish):

- The BNDN IKTLU identified fishing in Patterson Lake whitefish, jackfish, suckers, ling cod (mariah; TSD II: BNDN).
- The CRDN IKTLU identified use of jackfish, herring, lake trout, mariah, pickerel and suckers (spp.; TSD V.1: CRDN).
- The MN-S IKTLU identified that fish is integral to the traditional diet, including whitefish, jackfish, pickerel, suckers, and catfish (TSD IV: MN-S).
- The Athabasca Denesųłiné IKTLU identified lake trout, northern pike, and whitefish (TSD VI: YNLR).

Woodland caribou:

- The BNDN IKTLU identified caribou as a species of importance and identified as hunted but increasingly less available (TSD III: BNDN).
- o In a JWG meeting, the CRDN Band Manager said "We've already had caribou taken off our menu." (CRDN-JWG 2020).

Moose:

- The BNDN IKTLU identified moose as an important resource (TSD II: BNDN).
- In a JWG meeting, MN-S identified moose as their main dietary animal (MN-S-JWG 2020).
- The Athabasca Denesųłiné IKTLU identified caribou, moose, black bear, and white-tailed deer as big game harvested in the area (TSD VI: YNLR).

Black bear:

o In a JWG meeting, the BNDN identified bear as of importance (BNDN-JWG 2019b).

• Spruce grouse:

o The BNDN IKTLU identified grouse as hunted (TSD II: BNDN).



- The Athabasca Denesųłiné IKTLU identified wild chicken (spruce grouse, ruffed grouse), ptarmigan, ducks, geese, owls, ravens, crows, and seagulls, as being harvested in the area (TSD VI: YNLR).
- In a JWG meeting, one of the action items was to consider adding ptarmigan and chicken (spruce grouse) to the list of VCs. Subsequently, grouse was added for assessment in the EcoRA (CRDN-JWG 2020).

Mallard:

- In a JWG meeting, MN-S said that they considered mallard to be a garbage eater and felt the goldeneye was more important (MN-S-JWG 2019b).
- o In a JWG meeting, the BRDN identified mallard as commonly eaten and does not eat goldeneye (BRDN-JWG 2019b).
- The CRDN IKTLU has identified its right to hunt ducks and harvest duck eggs within its traditional territory (TSD V.1: CRDN).

Table 6-1 provides the rationale for the selected ecological receptors and Table 6-2 provides a summary of the selected ecological receptors. A short description of each of the selected ecological receptors including their habitats and feeding habits is provided in Appendix B, Ecological Receptor Profiles.



Table 6-1: Rationale for Selection of Ecological Receptors for the Rook I Project Ecological Risk Assessment

		1	2	3	4	
Organism Category	Representative Species	Major Plant or Animal Group	Presence in the Site, LSA, or RSA	Socioeconomic or Ecological Value and Value to Indigenous Communities	Exposed to and/or Sensitive to Stressor	Other Species Represented ^(a)
Terrestrial invertebrates	Earthworm (Lumbricus terrestris)	Soil-dwelling detritivore	Ubiquitous in the environment.	Food source for other ecological receptors.	Exposed to atmospheric release through soil.	Assessed as a group.
	Lichen	Lichen	Observed in the LSA and RSA. Samples collected for analysis of metals and radionuclides.	Primary winter food source for woodland caribou. Some lichen species are provincially rare.		Represents all terrestrial and arboreal lichen species.
Terrestrial vegetation	Blueberry (Vaccinium myrtilloides)	Shrub	Observed in the study areas. Fruit, leaves, and stems collected and analyzed for metals and radionuclides.	Food source for other ecological receptors. Harvested by people for consumption.	Exposed to atmospheric release through soil and atmospheric deposition.	Represents all terrestrial vegetation species, including jackpine, black spruce, and common ericaceous shrub species (e.g., bearberry, Labrador tea, blueberry, mountain cranberry, green alder).
	Labrador tea (Rhododendron groenlandicum)	Shrub	Present in study areas.	Food source for other ecological receptors. Medicinal plant harvested by Indigenous Peoples.		
	Zooplankton (general category)	Zooplankton	Present in lakes / potential discharge locations.	Food source for other ecological receptors.	Exposed to aquatic release through surface water.	Assessed as a group.
Aquatic invertebrates	Benthic invertebrates (i.e., Chironomidae)	Benthic invertebrates (infaunal)	Most abundant taxon found in areas surveyed.	Food source for other ecological receptors.	Exposed to aquatic release through sediment.	Assessed as a group.
	Benthic invertebrates (i.e., Ephemeroptera)	Benthic invertebrates (epifaunal)	Represents relatively large fraction of biomass in Patterson Lake and other areas.	Food source for other ecological receptors.	Exposed to aquatic release through surface water and sediment.	Assessed as a group.
Aquatic vegetation	Macrophyte (e.g., <i>Carex</i> sp.)	Aquatic macrophyte	Present in downstream surface waterbodies. Shoot, root, and sediment samples collected at Lloyd Lake for analysis of metals and radionuclides.	Food source for other ecological receptors. Provides spawning substrate for some fish species (e.g., Northern Pike).	Exposed via surface water and sediment.	Assessed as a group.
	Phytoplankton	Phytoplankton	Present in lakes / potential discharge locations.	Food source for other ecological receptors.	Exposed through aquatic release to surface water.	Assessed as a group.



		1	2	3	4	
Organism Category	Representative Species	Major Plant or Animal Group	Presence in the Site, LSA, or RSA	Socioeconomic or Ecological Value and Value to Indigenous Communities	Exposed to and/or Sensitive to Stressor	Other Species Represented ^(a)
	Northern pike (Esox lucius)	Pelagic predator fish	Present in Patterson Lake and surrounding surface waterbodies. Retained for chemical analysis and age determination.	Food source for other ecological receptors. Commercial fishing documented at Patterson Lake, Lloyd Lake, and Lac La Loche.	Exposed to aquatic release through surface water and consumption of prey (fish and/or aquatic invertebrates).	Burbot, lake trout, walleye.
Fish	Lake Whitefish (Coregonus clupeaformis)	Benthopelagic fish	Present in Patterson Lake and surrounding surface waterbodies. Retained for chemical analysis and age determination.	Food source for other ecological receptors. Commercial fishing documented at Patterson Lake, Lloyd Lake, and Lac La Loche.	Exposed to aquatic release through water, sediment, and consumption of prey (planktonic crustaceans, benthic invertebrates, fish eggs).	Arctic grayling, lake herring (cisco), Johnny darter, lake chub, longnose sucker, ninespine stickleback, slimy sculpin, spottail shiner, trout perch, white sucker, yellow perch.
	Woodland caribou (Rangifer tarandus caribou)	Terrestrial herbivore (large)	Present in study areas.	Threatened status under COSEWIC and SARA. Trapped for fur/meat and hunted by Indigenous Peoples.	Exposed to atmospheric release through consumption of food (terrestrial and arboreal lichens); exposed to aquatic release through drinking surface water.	None.
	Snowshoe hare (Lepus americanus)	Terrestrial herbivore (small)	Present in study areas.	Food source for other ecological receptors. Trapped for fur/meat and hunted by Indigenous Peoples.	Exposed to atmospheric release through food (plants) and soil.	None.
Terrestrial mammals	Moose (Alces americanus)	Terrestrial herbivore (large)	Present in study areas.	Surrogate for other mammals such as white-tailed deer. Game animal for Indigenous Peoples.	Exposed to atmospheric release through food (leaves, upland plants, water plants) and soil; exposed to aquatic release through drinking surface water.	White-tailed deer.
	Little brown myotis (<i>Myotis lucifugus</i>)	Terrestrial insectivore	Present in study areas.	Threatened status under COSEWIC and SARA.	Exposed to atmospheric release through food (insects) and soil.	Northern myotis.
	Southern red-backed vole (Myodes gapperi)	Terrestrial omnivore (small)	Most abundant small mammal captured. Baseline tissue analysis collected (Aug 2018).	Food source for other ecological receptors selected.	Exposed to atmospheric release through food (plants, seeds, roots, insects, snails, berries) and soil.	Meadow jumping mouse, meadow vole, porcupine, deer mouse.
	Black bear (Ursus americanus)	Terrestrial omnivore (large)	Present in study areas.	Trapped for fur/meat and hunted by Indigenous Peoples.	Exposed to atmospheric release through food (berries, nuts, small mammals, fish, birds) and soil; exposed to aquatic release through drinking surface water.	None.



Organism Category		1	2	3	4	
	Representative Species	Major Plant or Animal Group	Presence in the Site, LSA, or RSA	Socioeconomic or Ecological Value and Value to Indigenous Communities	Exposed to and/or Sensitive to Stressor	Other Species Represented ^(a)
	Red fox (Vulpes vulpes)	Terrestrial carnivore	Present in study areas.	Trapped for fur/meat and hunted by Indigenous Peoples.	Exposed to atmospheric release through food (small mammals) and soil.	American marten, Canada lynx, least weasel, masked shrew, coyote, fisher, red squirrel.
Terrestrial mammals	Grey wolf (Canis lupus)	Terrestrial carnivore	Present in study areas.	Trapped for fur/meat.	Exposed to atmospheric release through food (small and large mammals) and soil.	None.
	Muskrat (Ondatra zibethicus)	Riparian herbivore	Present in study areas.	Trapped for fur/meat and hunted by Indigenous Peoples.	Exposed to aquatic release through water, food (aquatic vegetation), and sediment.	None.
Riparian mammals	North American beaver (Castor canadensis)	Riparian herbivore	Present in study areas.	Trapped for fur/meat and hunted by Indigenous Peoples.	Exposed to aquatic release through water, food (terrestrial and aquatic vegetation), and sediment.	None.
	American mink (<i>Neovison</i> vison)	Riparian carnivore	Present in study areas.	Trapped for fur/meat and hunted by Indigenous Peoples.	Exposed to aquatic release through water, food (small mammals, fish, amphibians, insects), and sediment, and atmospheric release through food and soil.	Common water shrew, short-tailed weasel (ermine), river otter.



			Selectio	n Criteria		
Organism Category	Representative Species	1 Major Plant or Animal Group	2 Presence in the Site, LSA, or RSA	3 Socioeconomic or Ecological Value and Value to Indigenous	4 Exposed to and/or Sensitive to Stressor	Other Species Represented ^(a)
	Canada goose (Branta canadensis)	Ground feeding herbivore	Commonly observed in study area.	Communities Hunted by Indigenous Peoples.	Exposed to atmospheric release through food (grasses) and soil.	None
	Spruce grouse (Falcipennis canadensis)	Ground feeding herbivore	Observed in study areas.	Hunted by Indigenous Peoples.	Exposed to atmospheric release through food (seeds and berries) and soil.	Bay-breasted warbler, blackpoll warbler, blue-headed vireo, dedar waxwing, chipping sparrow, clay-coloured sparrow, dark-eyed junco, hermit thrush, horned lark, orange-crowned warbler, palm warbler, Philadelphia vireo, pine grosbeak, pine siskin, red crossbill, red-eyed vireo, wainson's thrush, Tennessee warbler, white-winged crossbill, willow ptarmigan, winter wren, yellow warbler, yellow-rumped warbler.
Terrestrial birds	Rusty blackbird (Euphagus carolinus)	Ground/aquatic feeding omnivore	Present in study areas.	Special concern status (COSEWIC and SARA). Surrogate for other birds and at-risk species.	Exposed to aquatic release through food (aquatic invertebrates), and atmospheric release through food (seeds and fruit).	LeConte's sparrow, American pipit, upland sandpiper, Wilson's warbler, common nighthawk, northern flicker, olive-sided flycatcher, western wood-pewee, American crow, common raven, alder flycatcher, yellow-bellied flycatcher, least flycatcher, mourning warbler, barn swallow, Lincoln's sparrow, song sparrow, black-and-white warbler, Nashville warbler, fox sparrow, Canada jay, ciff swallow, black-backed woodpecker, hairy woodpecker, Boreal chickadee, ruby-crowned kinglet, goldencrowned kinglet, ovenbird, magnolia warbler, American robin, white-throated sparrow, white-crowned sparrow, Wilson's snipe, swamp sparrow, northern waterthrush, tree swallow.



			Selectio	n Criteria		
		1	2	3	4	
Organism Category	Representative Species	Major Plant or Animal Group	Presence in the Site, LSA, or RSA	Socioeconomic or Ecological Value and Value to Indigenous Communities	Exposed to and/or Sensitive to Stressor	Other Species Represented ^(a)
Riparian birds	Mallard (Anas platyrhynchos)	Riparian herbivore	Present in study areas.	Surrogate for other duck species (e.g., ring-necked duck). Hunted by Indigenous Peoples.	Exposed to aquatic release through water, food (aquatic plants and invertebrates), and sediment.	Green-winged teal, snow goose, sandhill crane, lesser scaup, redhead, ring-necked duck, bufflehead, common goldeneye, tundra swan, herring gull, ringbilled gull, solitary sandpiper.
	Common loon (Gavia immer)	Piscivore	Present in study areas.	Surrogate for other fish-eating birds (e.g., gull, bald eagle, osprey).	Exposed to aquatic release through water and aquatic food (fish).	Bald eagle, spotted sandpiper, Bonaparte's gull, California gull, belted kingfisher, common merganser, osprey, American white pelican, red-necked grebe, lesser yellowlegs, greater yellowlegs.

⁽a) Based on presence of species in the terrestrial and aquatic baseline studies. Species identified in this column are not an extensive list of all species identified; however, species were removed from list for either having an observation of 1, or no direct observations at all unless very strong evidence existed (e.g. lots of pellets, droppings, nests, eggs, seen on game cams).



COSEWIC = Committee on the Status of Endangered Wildlife in Canada; SARA = Species at Risk Act; LSA = local study area; RSA = regional study area.

Table 6-2: Selected Ecological Receptors for the Rook I Project Ecological Risk Assessment

Organism Category	Representative Species			
Terrestrial invertebrates	Earthworm (Lumbricus terrestris)			
	Lichen			
Terrestrial vegetation	Blueberry (Vaccinium myrtilloides)			
errestrial vegetation	Labrador tea (Rhododendron groenlandicum)			
	Zooplankton (general category)			
Aquatic invertebrates	Benthic invertebrates (i.e., Chironomidae)			
	Benthic invertebrates (i.e., Ephemeroptera)			
Agustic vogetation	Macrophyte (e.g., <i>Carex</i> sp.)			
Aquatic vegetation	Phytoplankton			
Fish	Northern pike (Esox lucius)			
FISH	Lake whitefish (Coregonus clupeaformis)			
	Woodland caribou (Rangifer tarandus caribou)			
	Snowshoe hare (Lepus americanus)			
	Moose (Alces americanus)			
Terrestrial mammals	Little brown myotis (Myotis lucifugus)			
Terrestriai mammais	Southern red-backed vole (Myodes gapperi)			
	Black bear (Ursus americanus)			
	Red fox (Vulpes vulpes)			
	Grey wolf (Canis lupus)			
	Muskrat (Ondatra zibethicus)			
Riparian mammals	North American beaver (Castor canadensis)			
	American mink (Neovison vison)			
	Canada goose (Branta canadensis)			
Terrestrial birds	Spruce grouse (Falcipennis canadensis)			
	Rusty blackbird (Euphagus carolinus)			
Pinarian hirde	Mallard (Anas platyrhynchos)			
Riparian birds	Common loon (Gavia immer)			

6.1.1.2 Consideration of Species at Risk

Table 6-3 lists federally listed species of conservation concern that may potentially interact with the Project as compiled partly based on terrestrial baseline studies (Annex VII.1; Annex VIII.1). The list was compiled through a combination of regional and provincial references, field data, and professional experience. The table includes species with endangered, threatened, or special concern status by COSEWIC or under SARA. The Saskatchewan Conservation Data Centre rankings are shown in the table and are intended to provide support in conservation planning and



ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT ECOLOGICAL RISK ASSESSMENT

monitoring of SAR; however, protection of species on the Saskatchewan Conservation Data Centre is not regulated. There are no provincially threatened or endangered species listed in the Wild Species at Risk Regulations, Chapter W-13.11 Reg. 1 of the *Wildlife Act*.

The list of species at risk compiled by Omnia during baseline studies (Annex VIII.1) was refined by considering the presence of the species on the HABISask database in the study area, and detections of the species during the 2018 baseline surveys. No additional species were detected during subsequent surveys that were not previously identified in 2018 (Annex VIII.1; Annex VIII.3; Annex V.1).

Table 6-4 lists the surrogate species selected for identified species at risk. While not all SAR are selected as ecological receptors to be assessed, the selected ecological receptors include species with similar feeding habits, so that surrogate species from the ecological receptor list can be used to infer dose and risk for the SAR species that are present.

Woodland caribou, little brown myotis, and rusty blackbird were selected as ecological receptors; therefore, surrogate species were not needed to represent them.



Table 6-3: List of Identified Species with Conservation Status in the Vicinity of the Project

Species	COSEWIC Status	SARA Status	SKCDC Ranking	Listed in HABISask in Study Area	2018 to 2020 Field Survey Observations
Amphibians					
Northern leopard frog (<i>Lithobates pipiens</i>)	Special Concern	Special Concern	S3	Yes	Not observed in targeted acoustic surveys.
Birds	1	1	1	T	
Common nighthawk (Chordeiles minor)	Special Concern	Threatened	S4B, S4M	Yes	Common and widespread; identified at 44 locations.
Olive-sided flycatcher (Contopus cooperi)	Special Concern	Threatened	S4B, S4M	Yes	Suitable habitat common and widespread. Identified at 12 locations.
Rusty blackbird (Euphagus carolinus)	Special Concern	Special Concern	S3B, S3M	Yes	Suitable habitat common and widespread. Identified at three locations.
Barn swallow (<i>Hirundo rustica</i>)	Threatened	Threatened	S4B, S4M	Yes	Limited breeding habitat. Identified at ten locations.
Horned grebe (Podiceps auritus)	Special Concern	Special Concern	S5B, S5M	No	Not observed.
Peregrine falcon (Falco peregrinus)	Not at Risk	Special Concern	S1B	No	Not observed.
Yellow rail (Coturnicops noveboracensis)	Special Concern	Special Concern	S3B, S3M	No	Not observed in targeted surveys.
Whooping crane (<i>Grus americana</i>)	Endangered	Endangered	S1M	No	Not observed.
Hudsonian godwit (Limosa haemastica)	Threatened	No status	S4M	No	Not observed.
Buff-breasted sandpiper (Calidris subruficollis)	Special Concern	Special Concern	S4M	No	Not observed.
Eskimo curlew (Numenius borealis)	Endangered	Endangered	SXB, SXM	No	Not observed.
Red-necked phalarope (Phalaropus lobatus)	Special Concern	No status	S4B, S3M	No	Not observed.
Short-eared owl (Asio flammeus)	Special Concern	Special Concern	S3B, S2N, S3M	No	Not observed.
Bank Swallow (<i>Riparia riparia</i>)	Threatened	Threatened	S4B, S5M	No	Not observed.
Canada Warbler (Cardellina canadensis)	Threatened	Threatened	S4B, S3M	No	Not observed.
Harris' Sparrow (Zonotrichia leucophrys)	Special Concern	No status	S5M	No	Not observed.



ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT ECOLOGICAL RISK ASSESSMENT

Species	COSEWIC Status	SARA Status	SKCDC Ranking	Listed in HABISask in Study Area	2018 to 2020 Field Survey Observations
Mammals					
Little brown myotis (Myotis lucifugus)	Endangered	Endangered	S4	Yes	Detected with high certainty.
Northern myotis (Myotis septentrionalis)	Endangered	Endangered	S3	Yes	Likely present but in less abundance relative to the little brown myotis.
Woodland caribou (Rangifer tarandus)	Threatened	Threatened	S3	Yes	Moderate amount of suitable habitat. Incidentally observed at two locations.
Wolverine (Gulo gulo)	Special Concern	Special Concern	S2	No	Not observed.

SARA = Species at Risk Act; COSEWIC = Committee on the Status of Endangered Wildlife in Canada; SKCDC = Saskatchewan Conservation Data Centre; HABISask = Hunting, Angling and Biodiversity Information of Saskatchewan.



ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT ECOLOGICAL RISK ASSESSMENT

Several species in Table 6-3 were identified by Omnia (Annex VIII.1) to be potentially present in the Project study area but were not observed in the 2018 field surveys, including northern leopard frog, wolverine, and 12 species of birds. With exception of northern leopard frog, these species do not appear to have ranges overlapping with the Project study area (Annex V.1; Annex VII.2; Annex VIII.2; Annex VIII.3) and thus these species were not considered to be present on site; therefore, they were not considered as ecological receptors requiring surrogate receptors for the assessment.

Northern leopard frog was targeted in nocturnal amphibian surveys during the 2018 season at 16 survey sites, situated in areas with high habitat potential for breeding amphibians (Annex V.1; Annex VII.2; Annex VIII.2; Annex VIII.3). Northern leopard frogs were not detected or incidentally observed in the field, or identified through automated species-specific recognitions at recorded survey sites (Annex V.1; Annex VII.2; Annex VIII.2; Annex VIII.3). Because the potential incidence of these species appears to be low, they were not selected for assessment in the EcoRA. While amphibians were not specifically selected as receptors, they are sufficiently represented by the fish model. The sensitive life stages of frogs (i.e., egg and tadpole) are aquatic and similar to the sensitive life stages of a fish. For example, during the tadpole stage, tadpoles and fish have similar exposure pathways (e.g., absorption through skin and gills). In addition, exposure factor and toxicity data for amphibians are limited. Therefore, the fish assessment model is considered to be appropriate for frogs, including the northern leopard frog, during their sensitive life stages.

The rusty blackbird has a varied diet, but feeds mostly on aquatic invertebrates such as insect larvae, crustaceans, and snails. Its diet is supplemented by seeds and small fruit in the fall and winter. Breeding habitat for the rusty blackbird include forest wetlands, peat bogs, sedge meadows, marshes and swamps, beaver ponds, and pasture edges within the boreal forest (COSEWIC 2006). Suitable habitat for rusty blackbirds was noted to be common and widespread in baseline surveys (Annex V.1; Annex VII.2; Annex VIII.2; Annex VIII.3). Rusty blackbird was selected as an ecological receptor in the EcoRA; therefore, a surrogate to represent it is not needed.

Common nighthawk, olive-sided flycatcher, and barn swallow are aerial insectivores, consuming flying insects over both land and water. Breeding habitat for the common nighthawk is varied and may include open habitats such as sand dunes and beaches, recently burned or logged areas, open forests, peatbogs, marshes, lakeshores, river banks, or rocky outcrops (COSEWIC 2007). Most of the olive-sided flycatcher observations within the study area were near large waterbodies (i.e., Patterson Lake and Forrest Lake), or along creeks and bogs with sparse vegetation (Annex V.1; Annex VII.2; Annex VIII.2; Annex VIII.3). Barn swallows nest almost exclusively on man-made structures such as barns, out-buildings, houses, bridges, and road culverts (COSEWIC 2011). In the study area, barn swallows were observed near bridge crossings. The rusty blackbird was considered a suitable surrogate species for common nighthawk, olive-sided flycatcher, and barn swallow. The rusty blackbird's diet is primarily composed of aquatic invertebrates and is modelled in the EcoRA as eating benthic invertebrates, which are anticipated to be more directly exposed to COPCs than flying insects. Therefore, the selection of this surrogate is conservative.



Little brown myotis can be found roosting in trees and buildings; whereas northern myotis prefer to roost in the cavities of trees and under loose bark and are associated with the boreal forest. Both species prefer to forage over water features and along forest edges. In the vicinity of the site, the most bat detections were noted near the Patterson Creek Bridge and near the south side of the Rook I site at the edge of Patterson Lake south (Annex V.1; Annex VII.2; Annex VIII.2; Annex VIII.3). Little brown myotis was selected as an ecological receptor in the EcoRA, as it was noted to be in higher abundance relative to northern myotis in baseline surveys. Based on the similarities in diet between the two species of bat, the assessment of one is considered to be representative of the other.

Woodland caribou have a relatively large home range and seek mature jack pine and black spruce-dominated forest, black spruce bog, open muskeg habitats common to the Boreal Shield (McLoughlin 2016), and inaccessible rough terrain such as muskeg or islands on lakes for calving. The food source for the woodland caribou in the winter is terrestrial or arboreal lichens; terrestrial and aquatic vegetation are also food sources in the remainder of the year. Although the current suitability of the site for woodland caribou is relatively low due to an abundance of recently burned or harvested forest in the vicinity of the Project, the habitat potential over the timeframe of the Project is moderate to high (Annex VII.1; Annex VIII.1). Based on the importance of the species and its unique diet, the woodland caribou was selected as an ecological receptor in the EcoRA.

Omnia (Annex VII.1; Annex VIII.1) also identified 28 provincially sensitive and at-risk plant species within the baseline monitoring boundary for the Project during 2018 field surveys. None of the species are listed under COSEWIC or SARA or as SAR under the Saskatchewan *Wildlife Act*. Therefore, these species were not considered to require individual assessment as ecological receptors.

Table 6-4: Surrogate Species for Identified Species at Risk

Species at Risk	Surrogate Species
Common nighthawk (Chordeiles minor)	Rusty blackbird
Olive-sided flycatcher (Contopus cooperi)	Rusty blackbird
Barn swallow (Hirundo rustica)	Rusty blackbird
Northern myotis (Myotis septentrionalis)	Little brown myotis

6.1.2 Constituents of Potential Concern Retained for Further Evaluation in the Ecological Risk Assessment

The selection of COPCs retained for further evaluation in the EcoRA is presented in Section 4.0, Model Integration and Evaluation of Sources. Chloride and sulphate were identified as COPCs in the aquatic environment, but not in the terrestrial environment.



ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT ECOLOGICAL RISK ASSESSMENT

The selection of chemical stressors to evaluate in the EcoRA followed a tiered approach to reduce the risk of overlooking any COPCs relevant to ecological health. The selection of COPCs in water was based on predicted upper bound concentrations for treated effluent at end of pipe, in the receiving environment at the boundary of the mixing zone, and in runoff. Also, if predicted maximum concentrations of a COPC exceeded a benchmark value in sediments, it was retained for further evaluation in the ERA. The selection of COPCs in air was based on maximum-year air quality for Construction or Operations. This approach assumes maximum emissions from the various Project sources would be occurring at the same time, and under worst-case meteorological conditions. Also, if predicted maximum concentrations of a COPC exceeded a benchmark value in soil, it was retained for further evaluation in the EcoRA.

6.1.3 Assessment and Measurement Endpoints

Assessment endpoints for the EcoRA are explicit expressions of the environmental values that are to be protected (FCSAP 2012a). Assessment endpoints for the EcoRA should include the ecological receptor and the attribute of the ecological receptor that is to be protected (e.g., abundance, viability of the population) (FCSAP 2012a). The EcoRA assessment endpoints to be evaluated are presented in Table 6-5.

Measurement endpoints for the EcoRA are conceptually related to assessment endpoints and are defined as the specific measures that would be used to judge potential for effect on the attribute of an assessment endpoint (e.g., if we predict an effect on organism growth or reproduction, we can infer a potential for effect on abundance). Measurement endpoints for the EcoRA may include endpoints such as survival, growth, or reproduction. Measurement endpoints for the EcoRA are the foundation for the lines of evidence that are used to estimate risks to ecological receptors (FCSAP 2012a).

In this EcoRA, the assessment endpoints are at the population or community level; however, for species at risk, the assessment endpoint is at the individual level. While exposure and risk estimates always pertain to individuals, for most receptors, when effects on individuals are predicted from constituent levels in a certain location, further discussion of population or community effects (or lack thereof) is appropriate. For species at risk, it is considered that effects on even a single individual represent an effect on the population.



Table 6-5: Assessment Endpoints, Measurement Endpoints, and Lines of Evidence

	Level of		Assessment	Lines of Evidence		
Ecological Receptor	Protection	Protection Goal	Endpoint	Line of Evidence	Measurement Endpoints and their Interpretation	
Bottom feeding fish	Population	Maintenance of bottom feeding fish populations in Patterson Lake, Beet Lake, and Naomi Lake as a source of food for piscivorous fish and wildlife.	Viability of bottom-feeding fish populations.	Water chemistry	COPC concentrations in water. Compare to toxicological reference values (low-effect threshold concentrations) for effect on survival, growth, or reproduction.	
(lake whitefish)	Population			Radiological dose	Compare estimated doses of COPCs to growth, survival, and reproduction benchmark values (low-effect threshold doses) relevant to the assessment endpoint.	
Pelagic fish	Population	Maintenance of pelagic fish populations in Patterson Lake, Beet Lake, and Naomi Lake as a source of food for piscivorous fish and wildlife.	Viability of pelagic fish populations.	Water chemistry	COPC concentrations in water. Compare to toxicological reference values (low-effect threshold concentrations) for effect on survival, growth, or reproduction.	
(northern pike)				Radiological dose	Compare estimated doses of COPCs to growth, survival, and reproduction benchmark values (low-effect threshold doses) relevant to the assessment endpoint.	
Aquatic vegetation	Population in Lal	Maintenance of aquatic plant populations in Patterson Lake, Beet Lake, and Naomi Lake as a source of food and cover for wildlife.	Viability of aquatic plant populations.	Water chemistry	COPC concentrations in water. Compare to toxicological reference values (low-effect threshold concentrations) for aquatic plants for effect on survival, growth, or reproduction.	
(macrophytes, phytoplankton)				Radiological dose	Compare estimated doses of COPCs to growth, survival, and reproduction benchmark values (low-effect threshold doses) relevant to the assessment endpoint.	



	Laural of		A = = = = = = = = = = = = = = = = = = =	Lines	Lines of Evidence		
Ecological Receptor	Level of Protection	Protection Goal	Assessment Endpoint	Line of Evidence	Measurement Endpoints and their Interpretation		
Zooplankton	Community	Maintenance of a diverse zooplankton community in Patterson Lake, Beet Lake, and Naomi Lake as a source of food for	Density, richness, and diversity of	Water chemistry	COPC concentrations in water. Compare to toxicological reference values (low-effect threshold concentrations) for zooplankton for effect on survival, growth, or reproduction.		
Zoopiankon	,	fish.	zooplankton community.	Radiological dose	Compare estimated doses of COPCs to growth, survival, and reproduction benchmark values (low-effect threshold doses) relevant to the assessment endpoint.		
				Water chemistry	Compare COPC concentrations to water quality guidelines.		
B 41: 1		Maintenance of a diverse aquatic and benthic invertebrate community in	Richness, diversity, abundance of benthic invertebrates.	Sediment chemistry	Compare COPC concentrations to sediment quality guidelines.		
Benthic invertebrates	Community	Patterson Lake, Beet Lake, and Naomi Lake as a source of food for fish and wildlife.		Radiological Dose	Compare estimated doses of COPCs to growth, survival, and reproduction benchmark values (low-effect threshold doses) relevant to the assessment endpoint.		
Riparian birds (mallard, common loon)	Population	Maintenance of riparian bird populations along the shorelines of Patterson Lake, Beet Lake, and Naomi Lake as a source of food for predatory wildlife.	Viability of aquatic riparian bird populations.	- Radiological and toxicological	Compare estimated doses of COPCs to growth, survival, and		
Riparian mammals (muskrat, beaver, American mink)	Population	Maintenance of riparian mammal population along the shorelines of Patterson Lake, Beet Lake, and Naomi Lake as a source of food for predatory wildlife.	Viability of aquatic riparian mammal populations.	doses	reproduction benchmark values (low-effect threshold doses) relevant to the assessment endpoint.		
Terrestrial		Maintanance of torrectrial investebrate	Viability of torrectrial	Soil chemistry	COPC concentrations in soil. Compare COPC concentrations to soil quality guidelines.		
invertebrates (earthworm)	Population Maintenance of terrestrial invertebrate population in the RSA as a source of food for wildlife.	Viability of terrestrial invertebrate populations.	Radiological dose	Compare estimated doses to growth, survival, and reproduction benchmark values (low-effect threshold doses) relevant to the assessment endpoint.			



ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT ECOLOGICAL RISK ASSESSMENT

	Level of			Lines of Evidence		
Ecological Receptor	Protection	Protection Goal	Assessment Endpoint	Line of Evidence	Measurement Endpoints and their Interpretation	
Terrestrial plants		Maintanance of towartrial plant		Soil chemistry	COPC concentrations in soil. Compare COPC concentrations to soil quality guidelines.	
.	Population	Maintenance of terrestrial plant population in the RSA as a source of food for wildlife.		Radiological dose	Compare estimated doses to growth, survival, and reproduction benchmark values (low-effect threshold doses) relevant to the assessment endpoint.	
Terrestrial birds (Canada goose, grouse, rusty blackbird ^(a))	Population	Maintenance of the terrestrial bird population in the RSA.	Viability of terrestrial bird populations.		Compare estimated doses of COPCs	
Terrestrial mammals (woodland caribou ^(a) , snowshoe hare, moose, little brown myotis ^(a) , southern red-backed vole, black bear, red fox, grey wolf)	Population	Maintenance of terrestrial mammal population in the RSA.	Viability of terrestrial mammal populations.	Radiological and toxicological doses	to growth, survival, and reproduction benchmark values (low-effect threshold doses) relevant to the assessment endpoint.	

⁽a) For species at risk, protection is at the individual level, recognizing that effects on even a few individuals represent an effect on the population. COPC = constituent of potential concern; RSA = regional study area.



6.1.4 Selection of Exposure Pathways

Exposure pathways consider the various routes by which radionuclides and/or chemicals may enter the body of the receptor, or for radionuclides, may exert effects from outside the body. Exposures to environmental media may be direct (i.e., by contact) or indirect (i.e., via contaminant transport through the food chain).

For each type of ecological receptor, Table 6-6 summarizes, the relevant exposure pathways to various environmental media including air, surface water, soil, and sediment. Direct contact or uptake exposure pathways associated with groundwater are assumed to be incomplete, as it is assumed that groundwater is inaccessible to ecological receptors, or negligible relative to other pathways. Potential effects to surface water quality due to groundwater discharges were accounted for in the IMPACT model during Project phases where groundwater discharge is identified and hydrogeological model loadings of COPCs were input to the lake nodes in IMPACT.

Airborne COPCs partition to soil and plants. For most COPCs, ingestion pathways dominate over inhalation and air immersion. The latter pathways are considered minor pathways in the EcoRA, but inhalation was included in the IMPACT model and is thus included in Table 6-6.

For fish, aquatic plants, and aquatic invertebrates, contact with water and contaminant uptake from water via bioaccumulation represents the main exposure pathway. Direct contact or uptake from sediment are also considered for benthic invertebrates and bottom-feeding fish. Individual food chain transport pathways are not calculated by the IMPACT model for aquatic organisms because exposures for aquatic receptors are determined using bioaccumulation factors (BAFs) based on surface water concentrations; these BAFs represent all operable exposure pathways. The CSA N288.6-22 recommends the use of BAFs for the estimation of COPC concentrations in plant, invertebrate, and fish tissues based on concentrations in ambient media.

For soil invertebrates and terrestrial plants, the main exposure pathway is through contact with soil and contaminant uptake from soil via bioaccumulation. Earthworms and plant roots may have the potential to be exposed to groundwater when groundwater levels are high; however, both earthworms and plants would only be exposed to groundwater occasionally as they do not reside in the saturated zone. Therefore, direct contact with groundwater (for soil invertebrates) and uptake of groundwater (for terrestrial plants) are not quantified in IMPACT.

The dominant exposure pathways for birds and mammals are expected to include uptake of contaminants via the ingestion of water, direct contact with or incidental ingestion of soil and/or sediment, and ingestion of food/prey. Direct contact with water is also considered to be a complete exposure pathway for riparian mammals and birds.



Table 6-6: Complete Exposure Pathways for All Selected Ecological Receptors to be Assessed using the IMPACT Model

Category	Ecological Receptor	Exposure Pathways	Environmental Media			
	Lichen	Direct contact	Air			
Terrestrial vegetation	Blueberry	Direct contact	In soil			
	Labrador tea	Direct contact	In soil			
Terrestrial invertebrates	Earthworm	Direct contact	In soil			
Zooplankton	Zooplankton	Direct contact	In water			
Aquatic vegetation	Macrophytes	Direct contact	In water On sediment			
q	Phytoplankton	Direct contact	In water			
	Benthic invertebrates (infaunal)	Direct contact	In water In sediment			
Benthic invertebrates	Benthic invertebrates (epifaunal)	Direct contact	In water On sediment			
Bottom-feeding fish	Lake whitefish	Direct contact	In water On sediment			
Pelagic fish	Northern pike	Direct contact	In water			
		Direct contact	On soil			
		Inhalation	Air			
	Woodland caribou	Ingestion	Water Soil Sediment Browse Lichen Macrophytes			
		Direct contact	On soil			
		Inhalation	Air			
	Snowshoe hare	Ingestion	Water Soil Browse Blueberry			
Terrestrial mammals		Direct contact	On soil			
		Inhalation	Air			
	Moose	Ingestion	Water Soil Sediment Browse Macrophytes			
		Direct contact	On soil (minimal since aerial insectivore)			
	Little brown myotis	Inhalation	Air			
		Ingestion	Water Benthic invertebrates			
	Southern red-backed vole	Direct contact	On soil			
	Southern rea-packed voie	Inhalation	Air			



Category	Ecological Receptor	Exposure Pathways	Environmental Media				
			Water Soil				
		Ingestion	Blueberry Browse				
		Direct contact					
		Direct contact	On soil				
		Inhalation	Air				
	Black bear		Water Soil				
		Ingestion	Blueberry				
			Fish				
		Direct contact	On soil				
		Inhalation	Air				
		IIIIIaiatiOII	Water				
			Soil				
Terrestrial mammals	Red fox		Vole				
l		Ingestion	Earthworm				
			Goose				
			Blueberry				
		Direct contact	On soil				
		Inhalation	Air				
			Water				
	Grey wolf		Soil				
		Ingestion	Moose				
			Caribou				
			Hare				
		Direct contact	In water				
		Direct contact	On sediment				
		Inhalation	Air				
	Muskrat		Water				
		Ingestion	Sediment				
		nigestion	Macrophyte				
			Benthic invertebrates				
			In water				
		Direct contact	On soil				
			On sediment				
Riparian Mammals		Inhalation	Air				
	Beaver		Water				
			Soil				
		Ingestion	Sediment				
			Browse Macrophytes				
			In water				
		Direct contact	On soil				
	American mink	Sirect contact	On sediment				
	American mink	Inhalation	Air				
		Ingestion	Water				



Category	Ecological Receptor	Exposure Pathways	Environmental Media			
			Soil			
			Sediment			
			Fish			
			Benthic invertebrates			
			Muskrat			
			Mallard			
		Direct contact	On soil			
		Inhalation	Air			
	Canada goose		Water			
		Ingestion	Soil			
			Browse			
		Direct contact	On soil			
		Inhalation	Air			
	Grouse		Water			
Terrestrial birds	diodse		Soil			
		Ingestion	Browse			
			Blueberry			
		Direct contact	On soil			
		Inhalation	Air			
	Rusty blackbird		Water			
	Tradity Enderterna	Ingestion	Soil			
		Ingestion	Benthic invertebrates			
			Blueberry			
		5:	In water			
		Direct contact	On sediment			
		Inhalation	Air			
	Mallard		Water			
			Sediment			
		Ingestion	Macrophytes			
iparian birds			Benthic Invertebrates			
		Direct contact	In water			
		Inhalation	Air			
	Common loon		Water			
		Ingestion	Fish			
			Benthic invertebrates			

6.1.5 Ecological Conceptual Site Model

The ecological conceptual site model illustrates how receptors are exposed to COPCs. It identifies the source of contaminants, contaminant transport mechanisms, environmental media, and the exposure pathways to be considered in the assessment for each receptor. The conceptual model for the EcoRA is illustrated in Figure 6-1.




Figure 6-1: Ecological Health Conceptual Site Model


6.2 Exposure Assessment

The exposure assessment includes identification of exposure locations and exposure factors for each ecological receptor and presentation of exposure concentrations and doses (radiological and non-radiological). Uncertainties are discussed. This subsection presents the information used in the environmental pathways model, IMPACT, at a high level. The details of the model are provided in Appendix A.

6.2.1 Exposure Locations

The conceptual model assumes that all terrestrial and aquatic receptors are present at Patterson Lake (North Arm – West Basin and South Arm), Beet Lake, and Lloyd Lake (Figure 6-2). All terrestrial and aquatic receptors were also assumed to be at Broach Lake, which was chosen as a reference location. In addition, aquatic receptors were assessed at Clearwater River upstream of Mirror River, at the edge of the RSA, as shown in Figure 6-2. Separate exposure values were estimated for each receptor in the locations where they were assumed to be present.





Note: Aquatic receptors including zooplankton, aquatic plants, benthic invertebrates, and fish are located in the lakes and are not shown as separate polygons.

Figure 6-2: Locations of Ecological Receptors Assessed in the Ecological Risk Assessment



6.2.2 Exposure Averaging

Most ecological receptors were assessed assuming 100% residency at the exposure point location. This was the case for aquatic biota, immobile terrestrial biota such as plants and soil invertebrates, and mobile terrestrial mammals and birds that have small home ranges. This assumption was also applied to migratory ecological receptors, such as waterfowl and passerine birds, that spend part of the year away from the LSA and RSA, or ecological receptors that have a small home range while nesting or rearing young. For instance, mallards are not present in the LSA and RSA for over half the year due to migration, but have a small home range during the nesting and rearing season. Therefore, it was conservatively assumed that the mallard has a 100% residency factor on one lake where the young are hatched and reared. Similarly, moose have large annual home ranges but generally remain within small seasonal home ranges of 5 km² to 10 km² during the summer and winter. For modelling purposes, these animals were associated with one location.

Animals with larger home ranges (i.e., greater than 10 km²), such as woodland caribou, black bear, and grey wolf, interact with a number of different waterbodies and land areas for feeding and water intake. For modelling purposes, these animals were associated with a central location but with residency factors applied to other adjacent locations depending on the size of their home range. In some cases, a portion of their time in the LSA or RSA may be attributed to time spent at unexposed (reference) locations. For example, due to its large home range of 1,300 km², a grey wolf located near Patterson Lake was assumed to spend 47.6% of its time at exposed locations and 52.4% of its time at unexposed locations. Further details are provided in Appendix A. Table 6-7 summarizes the residency assumptions for ecological receptors with large home ranges used for the ERA.

Table 6-7: Residency Factors for Terrestrial Ecological Receptors with Large Home Ranges

Terrestrial Ecological Receptor	Polygon	Area of Waterbody (ha)	Total Waterbody (ha)	Residency Factor
\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	Reference Lake	59		0.02
Woodland caribou at Patterson Lake	Patterson Lake North Arm – West Basin	1,256	2,910	0.43
ratterson Lake	Patterson Lake South Arm	1,595		0.55
Woodland caribou at	Reference Lake	1,119	2.150	0.52
Beet Lake	Beet Lake ^(a)	1,031 ^(a)	2,150	0.48
DI III	Reference Lake	59		0.02
Black bear at Patterson Lake	Patterson Lake North Arm – West Basin	1,256	2,910	0.43
ratterson Lake	Patterson Lake South Arm	1,595		0.55
Black bear at Beet	Reference Lake	1,119	2.150	0.52
Lake	Beet Lake ^(a)	1,031 ^(a)	2,150	0.48

Terrestrial Ecological Receptor	Polygon	Area of Waterbody (ha)	Total Waterbody (ha)	Residency Factor
	Reference Lake ^(b)	5,564 ^(b)		0.524
Grey wolf at	Patterson Lake North Arm – West Basin ^(c)	2,184 ^(c)	10.610	0.206
Patterson Lake	Patterson Lake South Arm	1,595	10,619	0.150
	Beet Lake ^(d)	1,276 ^(d)		0.120

⁽a) Woodland caribou and Black Bear at Beet Lake also use Forrest Lake - North Basin.

6.2.3 Exposure and Dose Calculations

Exposure and dose calculations for ecological receptors were completed using IMPACT version 5.6.0. IMPACT is consistent with the COPC transport equations outlined in CSA N288.1-20 and with the methods of biota dose calculation outlined in CSA N288.6-22 for both non-radiological and radiological COPCs. The equations are presented in Appendix A. Assessment of radiation exposures to ecological receptors is commonly based on estimation of the effects of the project or site. Assessments consider the radiation dose received from direct exposure to radiation as well as the dose received from ingestion of radionuclides. The radionuclide intake by ecological receptors from various pathways is converted into a dose that is presented in milligrays per day (mGy/d). The dose for each radionuclide is comprised of an internal dose component, and an external dose component, which is driven by water and sediment.

Assessment of non-radiological exposures to ecological receptors considers the dose received from ingestion of constituents of concern. This is presented as a dose in mg/kg/d for each pathway.

The inputs and assumptions used in the IMPACT model for the Project, including receptor characteristics, exposure pathways, and the derivation and identification of site-specific information used in the model are provided in Appendix A. Relevant to the exposure and dose calculations, the report:

- describes the model structure for ecological receptor assessment, specific assumptions made for the Project, and the generic equations used to calculate the transfer of constituents between environmental media and the receptors; and
- presents the development of input parameters and describes the approach used for calibration and validation based on regional monitoring data.



⁽b) Grey wolf at Reference Lake also uses Broach Lake; Lake C, Lake E, Lake G, Lake H, Unnamed Lake 1 (near Dennis Lake), and Unnamed Lake 2 (Near Naomi Lake); and Forrest Lake – South Basin.

⁽c) Grey wolf at Patterson Lake North Arm – West Basin also uses Patterson Lake North Arm – East Basin.

⁽d) Grey wolf at Beet Lake also uses Forrest Lake North Arm and Naomi Lake.

6.2.4 Exposure Factors

Exposure estimates rely on several COPC- and biota-specific exposure factors for the dose calculations. These parameters include body weights and intake rates as well as occupancy factors (OFs), DCFs, BAFs, and transfer factors (TFs).

6.2.4.1 Body Weights and Intake Rates for Ecological Receptors

Body weights and intake rates are required for the calculation of exposure to birds and mammals. Selected body weights and intake rates for bird and mammal ecological receptors are presented in Table 6-8. Body weights and intake rates were obtained, in order of preference, from CSA N288.1-20, the United States Environmental Protection Agency Wildlife Exposure Factors Handbook (USEPA 1993), the Federal Contaminated Sites Action Plan Module 3: Standardization of Wildlife Receptor Characteristics (FCSAP 2012b), and from Sample and Suter (1994), where the values were considered to be applicable. The CSA N288.6-22 recommends the use of CSA N288.1-20 and USEPA (1993) as primary literature sources for receptor characteristics, supplemented by other sources containing sufficient information to support applicability and credibility. For species not represented in the above sources, additional sources identified in Table 6-8 were consulted to identify representative body weights, and feed intake rates were calculated using allometric equations from USEPA (1993). Water intake and inhalation rates were determined using allometric equations from USEPA (1993) for birds and mammals. The incidental ingestion of soil and/or sediment was estimated from feed intake. As described by Beyer et al. (1994), incidental ingestion varied from 2% to 10.4% of dry weight food intake depending on the biota. No incidental soil or sediment ingestion was assumed for the little brown myotis, which is an aerial insectivore; nor was any sediment ingestion assumed for the Common loon, which feeds from the water column.



Table 6-8: Body Weights and Intake Rates for Bird and Mammal Ecological Receptors

Receptor	Body \	Weight		Total Fed	ed Intake		Dietary Components	Feed Typ	e Fraction	Feed Inta	ake Rate	Moisture ^(h)	Intake of Soil ^(r)	Intake of Sediment ^(r)	Intake of Soil and Sediment ^(k)	Basis of the Soil and Sediment Intake Value	Total Soil/ Sediment Intake Rate ⁽¹⁾	Water Intake Rate ^(m)	Inhalati	ion Rate
	kg	Source	kg dw/d	Source	kg fw/d	Source		fw	dw	kg dw/d	kg fw/d	unitless	%	%	%		kg dw/d	L/d	m³/d	Source
	180	(e)	3.82	(m)	12.35	n/a	Browse	0.5	0.32	1.24	6.18	0.80	5.15	1.65	6.8		0.3	10.6	34.8	(m)
Woodland caribou							Lichen	0.2	0.43	1.66	2.47	0.33				Bison				
							Macrophytes	0.3	0.24	0.927	3.71	0.75								
Snowshoe hare	1.8	(c)	0.110	(c)	0.564	n/a	Browse	0.9	0.92	0.102	0.508	0.80	3.7	0	3.7	Average of small	0.004	0.168	0.9	(c,m)
Showshoe hare							Berries (blueberry)	0.1	0.08	0.008	0.056	0.85				mammals				
	400	(b)	8.0	(b)	38.1	n/a	Browse	0.8	0.76	6.10	30.5	0.80	1.5	0.5	2.0	Moose	0.2	21.8	65.9	(m)
Moose							Macrophytes	0.2	0.24	1.90	7.62	0.75								
Little brown myotis	0.0075	(d)	0.001	n/a	0.0037	(d)	Benthic invertebrates	1	1	0.0009	0.0037	0.75	0	0	0	Assumed negligible (aerial)	0	0.00121	0.011	(m)
Southern red-backed	0.0329	(a, j)	0.00196	n/a	0.012	(a, j)	Berries (blueberry)	0.6	0.53	0.001	0.007	0.85	2.4	0	2.4	Maadayyyala	0.000047	0.00458	0.048	(a)
vole							Browse	0.4	0.47	0.001	0.005	0.80				Meadow vole				
	102.5	(o)	3.1	(b)	17.1	n/a	Berries (blueberry)	0.7	0.58	1.79	12.0	0.85	2.9	0	2.9	Average of	0.09	6.39	22.2	(m)
Black bear							Fish (northern pike)	0.3	0.42	1.28	5.13	0.75				large mammals				
	4.54	(a)	0.105	n/a	0.429	(a)	Small mammals (vole)	0.4	0.490	0.052	0.172	0.70	2.8	0	2.8	Red fox	0.00295	0.386	1.83	(m)
							Terrestrial birds (goose)	0.2	0.245	0.026	0.086	0.70								
Red fox							Terrestrial invertebrates (earthworm)	0.25	0.173	0.018	0.107	0.83								
							Berries (blueberry)	0.15	0.092	0.010	0.064	0.85								
	45	(f)	1.57	(m)	5.23	n/a	Moose	0.45	0.45	0.707	2.356	0.70	2.8	0	2.8	Red fox	0.04	3.04	11.5	(m)
Grey wolf							Caribou	0.45	0.45	0.707	2.356	0.70								
							Small mammal (hare)	0.1	0.10	0.157	0.523	0.70								
_	24	(p)	0.94	(m)	4.57	n/a	Macrophytes	0.1	0.12	0.114	0.457	0.75	1.8	0.24	2.0	Moose	0.019	1.729	6.9	(m)
Beaver							Browse	0.9	0.88	0.822	4.112	0.80								
	1.17	(a)	0.088	n/a	0.352	(a)	Macrophytes	0.8	0.8	0.070	0.282	0.75	0	3.3	3.3	Mallard	0.003	0.114	0.59	(a)
Muskrat							Benthic invertebrates	0.2	0.2	0.018	0.070	0.75								
	1.02	(a)	0.045	n/a	0.161	(a)	Fish (northern pike)	0.3	0.270	0.012	0.048	0.75	2.6	0.41	3.05		0.001	0.101	0.44	(a)
A							Benthic invertebrates	0.15	0.135	0.006	0.024	0.75				Average of mallard and				
American mink							Small mammals (muskrat)	0.45	0.486	0.022	0.072	0.70				red fox				
							Birds (mallard)	0.1	0.108	0.005	0.016	0.70								
Rusty blackbird	0.064	(g)	0.0177	n/a	0.077	(a,i)	Benthic invertebrates	0.8	0.87	0.015	0.062	0.75	10.4	0	10.4	American	0.002	0.00935	0.148	(m,n)
Nusty biackbild							Berries (blueberry)	0.2	0.13	0.002	0.015	0.85				woodcock				
Canada goose	3.70	(c)	0.0229	n/a	0.115	(a)	Browse	1	1	0.023	0.115	0.80	8.2	0	8.2	Canada goose	0.0019	0.142	1.100	(c)
Grouse	0.6	(b)	0.042	(b)	0.227	n/a	Browse	0.7	0.76	0.032	0.159	0.80	8.2	0	8.2	Canada goose	0.0034	0.0419	0.276	(m)
Grouse							Berries (blueberry)	0.3	0.24	0.010	0.068	0.85				Cariada goose				

Receptor	Body \	Weight		Total Fee	ed Intake		Dietary Components Feed Type Fraction F		Feed Int	ake Rate	Moisture ^(h)	Intake of Soil ^(r)	Intake of Sediment ^(r)	Intake of Soil and Sediment ^(k)	Basis of the Soil and Sediment Intake Value	Sediment Intake	Water Intake Rate ^(m)	Inhalati	ion Rate	
	kg	Source	kg dw/d	Source	kg fw/d	Source		fw	dw	kg dw/d	kg fw/d	unitless	%	%	%		kg dw/d	L/d	m³/d	Source
M. II.	1.13	(c)	0.06	(c)	0.240	n/a	Macrophytes	0.25	0.25	0.015	0.060	0.75	0	3.3	3.3	NA III	0.002	0.0640	0.02	(c)
Mallard							Benthic invertebrates	0.75	0.75	0.045	0.180	0.75				Mallard				
	5.3	(b)	0.159	(b)	0.636	n/a	Fish (northern pike)	0.9	0.9	0.143	0.572	0.75	0	0	0	Assumed	0	0.180	1.477	(m)
Common loon							Benthic invertebrates	0.1	0.1	0.016	0.064	0.75				negligible (sediment is washed off in water column)				

- (a) USEPA 1993; body weights, ingestion rates and inhalation rates of adults or all groups (adults and juveniles) are an average of the listed values. If only a range is given, the upper limit of the range is used. Values for the southern red-backed vole were based on the meadow vole.
- (b) FCSAP 2012b.
- (c) CSA N288.1-20.
- (d) Sample and Suter 1994.
- (e) COSEWIC Assessment and Update Status Report on the Woodland Caribou (COSEWIC 2002). Body weight calculated as a mean of the male and female upper range.
- (f) Earthroots 2020.
- (g) COSEWIC Assessment and Update Status Report on the Rusty Blackbird (COSEWIC 2006).
- (h) Sources of moisture/dry weight fraction values obtained from CSA N288.1-20 for all receptors except for earthworm (Beresford et al. 2008a), and blueberries and lichen (EIS Appendix 7A, Air Dispersion Modelling Report). The blueberry value is based on the fruit only.
- (i) The total feed intake for the American robin was used in the absence of a species-specific value.
- (j) The body weight and total feed intake for the meadow vole was used in the absence of a species-specific value.
- (k) Beyer et al. 1994.
- (I) Intake of Soil & Sediment (kg dw/d) = Total Feed Intake (kg dw/d) x Intake of Soil & Sediment (%)/100.
- (m) Calculated using allometric equations in (USEPA 1993).
- (n) Inhalation rate was multiplied by a factor of 3 for passerine birds as per (USEPA 1993).
- (o) Average of males and females from (Hinterland Who's Who 2007).
- (p) Median of adult weight range of 16 to 32 kg from (Hinterland Who's Who 2017).
- (q) For purposes of modelling the earthworm is counted to the terrestrial plants.
- (r) The % intake of soil or sediment is calculated from the combined % intake of soil and sediment, weighted to the relative proportions of terrestrial vs. aquatic dietary components for each receptor, based on the following equations.

Intake of Soil (%) = Total Intake of Soil & Sediment (%) x Feed Type Fraction Terrestrial. Intake of Sediment (%) = Total Intake of Soil & Sediment (%) x Feed Type Fraction Aquatic.

fw = fresh weight; dw = dry weight; n/a = not applicable.

6.2.4.2 Occupancy Factors, Dose Coefficients, Bioaccumulation Factors, and Transfer Factors

Short descriptions of the role of the OFs, DCFs, BAFs, and TFs are provided in Table 6-9. Additional details and the numeric factors are presented in Appendix A, as indicated in Table 6-9.

Table 6-9: Exposure Factors Used in the Rook I Project IMPACT Model

Exposure Factor	Description	Appendix A
OFs	An OF is defined as the fraction of time the receptor species spends in or on various media. The OFs are based on the experience and judgment of the risk assessor and the known behaviour of the receptor. The OFs for air, soil/sediment, soil/sediment surface, and water were used in the model.	Section 2.3.3.2, Occupancy Factor
DCFs	The DCFs represent the dose-equivalent rate per unit concentration of a radionuclide in the environment (or tissue) for a particular mode of exposure. The model used DCFs for external and internal exposures to radionuclides. Aquatic DCFs were based on values published by the ICRP for aquatic plants and northern pike (ICRP 2008), and were calculated with the ERICA Tool (Brown et al. 2008) for benthic invertebrates, zooplankton, and whitefish DCFs. Terrestrial plant and invertebrate DCFs were based on values published by the ICRP (ICRP 2008). Terrestrial animal DCFs follow the approach of ICRP (ICRP 2008).	Section 3.6.3, Dose Coefficients for Aquatic Receptors Section 3.7.3, Dose Coefficients for Terrestrial Plants and Invertebrates Section 3.7.6, Dose Coefficients for Terrestrial Animals, Birds, and Humans
TFs and BAFs	The TFs are the ratio of concentration in an animal to the animal's daily intake of a COPC. BAFs are the ratio of concentration in an organism to the concentration in an environmental medium. The TFs and BAFs are generally COPC- and biota-specific. Aquatic BAFs were generally obtained from CSA N288.1-20 and IAEA (2010), and from publicly available regional data from other uranium mine sites in northern Saskatchewan. The soil-to-plant BAFs were derived from regional data from Northern Saskatchewan. Ingestion TFs for terrestrial receptors from forage to tissue for agricultural livestock were from CSA N288.1-20. An allometric equation (transfer proportional to a -3/4 power of body weight) (CSA N288.6-22) was applied to transfer factors available for beef and poultry from CSA N288.1-20, IAEA (2010), or NCRP (1996) to estimate the transfer factors for the mammal and bird receptors, respectively. Inhalation TFs for terrestrial receptors were calculated from the ingestion transfer factor by adjusting the ingestion transfer factor by a COPC-specific inhalation/ingestion ratio (II) from CSA N288.1-20.	Section 3.6.1, Aquatic Bioaccumulation Factors Section 3.7.1, Soil-to-Plant Transfer Section 3.7.4, Ingestion Transfer Factors for Terrestrial Receptors Section 3.7.5, Inhalation Transfer Factors for Terrestrial Receptors

IAEA = International Atomic Energy Agency; BAF = bioaccumulation factor; OF = occupancy factor; TF = transfer factor; DCF = dose coefficient factor; NCRP = National Council on Radiation Protection and Measurements; ICRP = International Commission on Radiological Protection; CSA = Canadian Standards Association; COPC = constituent of potential concern.



6.2.5 Exposure Point Concentrations and Doses

This subsection presents the estimated non-radiological and radiological doses to aquatic and terrestrial ecological receptors due to releases from the Project during all phases of the Project, including the far-future projection once groundwater solutes have been released to Patterson Lake. The results are presented for both the Application Case and Upper Bound sensitivity scenario (Section 6.2.5.1, Estimated Dose to Ecological Receptors – Application Case), and the RFD Case (Section 6.2.5.2, Estimated Dose to Ecological Receptors – Reasonable Foreseeable Development Case). The estimated non-radiological and radiological concentrations in environmental media and biota tissue concentrations are shown in Appendix C.

6.2.5.1 Estimated Dose to Ecological Receptors – Application Case

6.2.5.1.1 Non-radiological Dose

Non-radiological dose was only calculated for birds and mammals, as effects to aquatic animals (fish and invertebrates) and plants and soil invertebrates are assessed based on concentrations and not doses.

The estimated non-radiological dose to the selected birds and mammals during the Project phases and the far-future projection resulting from the Application Case and reasonable upper bound sensitivity scenario are shown by COPC in Table 6-10 and Table 6-11, respectively. The doses shown represent the maximum dose by COPC over the assessment period.

For arsenic and uranium, the estimated non-radiological dose was highest during Operations, whereas for cobalt and copper, the estimated non-radiological dose was highest during the far-future projection. That is due to the additional load of cobalt and copper from groundwater flows (infiltration and seepage), primarily from the waste rock storage area and secondarily from the UGTMF in the far-future projection. The groundwater modelling assumed that the waste rock storage area and UGTMF are inexhaustible and unmitigated loading sources, causing the predicted cobalt and copper concentrations to remain elevated indefinitely in the far-future projection.



Table 6-10: Estimated Non-radiological Doses to Ecological Receptors – Operations – Application Case and Upper Bound Scenario

		Singipul Books to Ecological Receptors					ximum Dose During	Operations (mg/kg/d)				
	Biota	Location			Application Case				Į	Jpper Bound Scenario		
			Arsenic	Cobalt	Copper	Molybdenum	Uranium	Arsenic	Cobalt	Copper	Molybdenum	Uranium
		Reference (Broach Lake)	3.48 × 10 ⁻⁰³	3.94 × 10 ⁻⁰³	3.15 × 10 ⁻⁰²	1.17 × 10 ⁻⁰³	3.20 × 10 ⁻⁰³	3.48 × 10 ⁻⁰³	3.94×10^{-03}	1.94 × 10 ⁻⁰³ 3.15 × 10 ⁻⁰² 1.17 × 10 ⁻⁰³ 1.61 × 10 ⁻⁰³ 3.32 × 10 ⁻⁰² 1.88 × 10 ⁻⁰³ 1.30 × 10 ⁻⁰³ 3.20 × 10 ⁻⁰² 1.49 × 10 ⁻⁰³ 1.44 × 10 ⁻⁰³ 3.18 × 10 ⁻⁰² 1.20 × 10 ⁻⁰³ 1.77 × 10 ⁻⁰³ 2.86 × 10 ⁻⁰² 3.75 × 10 ⁻⁰⁴ 1.30 × 10 ⁻⁰³ 3.16 × 10 ⁻⁰² 4.60 × 10 ⁻⁰⁴ 1.42 × 10 ⁻⁰³ 2.94 × 10 ⁻⁰² 4.64 × 10 ⁻⁰⁴ 1.42 × 10 ⁻⁰³ 2.88 × 10 ⁻⁰² 3.92 × 10 ⁻⁰⁴ 1.42 × 10 ⁻⁰³ 2.88 × 10 ⁻⁰² 3.92 × 10 ⁻⁰⁴ 1.42 × 10 ⁻⁰³ 2.88 × 10 ⁻⁰² 3.92 × 10 ⁻⁰⁴ 1.42 × 10 ⁻⁰³ 2.88 × 10 ⁻⁰² 3.92 × 10 ⁻⁰⁴ 1.42 × 10 ⁻⁰³ 2.88 × 10 ⁻⁰² 3.92 × 10 ⁻⁰⁴ 1.10 × 10 ⁻⁰⁴ 3.31 × 10 ⁻⁰² 2.88 × 10 ⁻⁰⁴ 1.10 × 10 ⁻⁰⁴ 3.37 × 10 ⁻⁰² 3.43 × 10 ⁻⁰⁴ 1.61 × 10 ⁻⁰³ 2.28 × 10 ⁻⁰¹ 1.15 × 10 ⁻⁰³ 1.68 × 10 ⁻⁰³ 2.28 × 10 ⁻⁰¹ 1.15 × 10 ⁻⁰³ 1.89 × 10 ⁻⁰³ 2.38 × 10 ⁻⁰¹ 2.50 × 10 ⁻⁰³ 1.89 × 10 ⁻⁰³ 2.38 × 10 ⁻⁰¹ 1.28 × 10 ⁻⁰³ 1.89 × 10 ⁻⁰³ 1.61 × 10 ⁻⁰²	1.17 × 10 ⁻⁰³	3.20 × 10 ⁻⁰³
		Patterson Lake North Arm – West Basin	6.52×10^{-03}	4.59 x 10 ⁻⁰³	3.32 x 10 ⁻⁰²	1.51 x 10 ⁻⁰³	2.49 x 10 ⁻⁰²	6.61 x 10 ⁻⁰³	4.61 x 10 ⁻⁰³		1.88 x 10 ⁻⁰³	2.62 x 10 ⁻⁰²
	Beaver	Patterson Lake South Arm	3.74×10^{-03}	4.29 x 10 ⁻⁰³	3.20 x 10 ⁻⁰²	1.29 x 10 ⁻⁰³	6.39 x 10 ⁻⁰³	3.75 x 10 ⁻⁰³	4.30 x 10 ⁻⁰³		1.49 x 10 ⁻⁰³	6.73 x 10 ⁻⁰³
		Beet Lake	3.53 × 10 ⁻⁰³	4.13 x 10 ⁻⁰³	3.18 x 10 ⁻⁰²	1.23 x 10 ⁻⁰³	5.04 x 10 ⁻⁰³	3.53 x 10 ⁻⁰³	4.14 x 10 ⁻⁰³	3.18 x 10 ⁻⁰²	1.34 x 10 ⁻⁰³	5.16 x 10 ⁻⁰³
		Lloyd Lake	3.50×10^{-03}	3.97 x 10 ⁻⁰³	3.16 x 10 ⁻⁰²	1.19 x 10 ⁻⁰³	3.27 x 10 ⁻⁰³	3.50 x 10 ⁻⁰³	3.97 x 10 ⁻⁰³	3.16 x 10 ⁻⁰²	1.20 x 10 ⁻⁰³	3.28 x 10 ⁻⁰³
		Reference (Broach Lake)	2.81 × 10 ⁻⁰³	1.17 × 10 ⁻⁰³	2.86 × 10 ⁻⁰²	3.75×10^{-04}	1.02 × 10 ⁻⁰³	2.81 × 10 ⁻⁰³	1.17×10^{-03}	2.86 × 10 ⁻⁰²	3.75×10^{-04}	1.02 × 10 ⁻⁰³
	Black Bear	Patterson Lake North Arm – West Basin	5.07 × 10 ⁻⁰³	1.30 x 10 ⁻⁰³	3.15 x 10 ⁻⁰²	4.22 x 10 ⁻⁰⁴	3.90 x 10 ⁻⁰³	5.14 x 10 ⁻⁰³	1.30 x 10 ⁻⁰³	3.16 x 10 ⁻⁰²	4.60 x 10 ⁻⁰⁴	4.22 x 10 ⁻⁰³
	DIACK Dear	Beet Lake	3.12 × 10 ⁻⁰³	1.42 x 10 ⁻⁰³	2.94 x 10 ⁻⁰²	4.57 x 10 ⁻⁰⁴	1.41 x 10 ⁻⁰³	3.12 x 10 ⁻⁰³	1.42 x 10 ⁻⁰³	2.94 x 10 ⁻⁰²	4.64 x 10 ⁻⁰⁴	1.43 x 10 ⁻⁰³
		Lloyd Lake	2.83×10^{-03}	1.21 x 10 ⁻⁰³	2.88 x 10 ⁻⁰²	3.91 x 10 ⁻⁰⁴	1.04 x 10 ⁻⁰³	2.83 x 10 ⁻⁰³	1.21 x 10 ⁻⁰³	2.88 x 10 ⁻⁰²	3.92 x 10 ⁻⁰⁴	1.05 x 10 ⁻⁰³
		Reference (Broach Lake)	7.50 x 10 ⁻⁰³	6.01 x 10 ⁻⁰⁴	3.31 x 10 ⁻⁰²	2.88 x 10 ⁻⁰⁴	3.73 x 10 ⁻⁰⁴	7.50 x 10 ⁻⁰³	6.01 x 10 ⁻⁰⁴	3.31 x 10 ⁻⁰²	2.88 x 10 ⁻⁰⁴	3.73 x 10 ⁻⁰⁴
	Grey Wolf	Patterson Lake North Arm – West Basin	8.87 x 10 ⁻⁰³	6.13 x 10 ⁻⁰⁴	3.37 x 10 ⁻⁰²	3.17 x 10 ⁻⁰⁴	1.20 x 10 ⁻⁰³	8.90 x 10 ⁻⁰³	6.13 x 10 ⁻⁰⁴	3.37 x 10 ⁻⁰²	3.43 x 10 ⁻⁰⁴	1.22 x 10 ⁻⁰³
		Lloyd Lake	7.62 x 10 ⁻⁰³	6.06 x 10 ⁻⁰⁴	3.32 x 10 ⁻⁰²	2.95 x 10 ⁻⁰⁴	3.85 x 10 ⁻⁰⁴	7.62 x 10 ⁻⁰³	6.06 x 10 ⁻⁰⁴	3.32 x 10 ⁻⁰²	2.98 x 10 ⁻⁰⁴	3.85 x 10 ⁻⁰⁴
		Reference (Broach Lake)	1.15 × 10 ⁻⁰²	4.61 × 10 ⁻⁰³	2.28 × 10 ⁻⁰¹	1.15 × 10 ⁻⁰³	1.08 × 10 ⁻⁰³	1.15 × 10 ⁻⁰²	4.61×10^{-03}	2.28 × 10 ⁻⁰¹	1.15 × 10 ⁻⁰³	1.08 × 10 ⁻⁰³
		Patterson Lake North Arm – West Basin	3.46 x 10 ⁻⁰²	5.83 x 10 ⁻⁰³	2.66 x 10 ⁻⁰¹	2.80 x 10 ⁻⁰³	4.16 x 10 ⁻⁰³	3.52 x 10 ⁻⁰²	5.88 x 10 ⁻⁰³	2.67 x 10 ⁻⁰¹	6.16 x 10 ⁻⁰³	7.50 x 10 ⁻⁰³
	American Mink	Patterson Lake South Arm	1.34 x 10 ⁻⁰²	5.26 x 10 ⁻⁰³	2.47 x 10 ⁻⁰¹	1.99 x 10 ⁻⁰³	1.75 x 10 ⁻⁰³	1.35 x 10 ⁻⁰²	5.29 x 10 ⁻⁰³	2.48 x 10 ⁻⁰¹	3.77 x 10 ⁻⁰³	2.63 x 10 ⁻⁰³
		Beet Lake	1.18 x 10 ⁻⁰²	4.95 x 10 ⁻⁰³	2.38 x 10 ⁻⁰¹	1.58 x 10 ⁻⁰³	1.37 x 10 ⁻⁰³	1.18 x 10 ⁻⁰²	4.97 x 10 ⁻⁰³	2.38 x 10 ⁻⁰¹	2.50 x 10 ⁻⁰³	1.69 x 10 ⁻⁰³
		Lloyd Lake	1.15 x 10 ⁻⁰²	4.65 x 10 ⁻⁰³	2.29 x 10 ⁻⁰¹	1.19 x 10 ⁻⁰³	1.10 x 10 ⁻⁰³	1.15 x 10 ⁻⁰²	4.65 x 10 ⁻⁰³	2.29 x 10 ⁻⁰¹	1.28 x 10 ⁻⁰³	1.13 x 10 ⁻⁰³
		Reference (Broach Lake)	2.35 × 10 ⁻⁰³	2.83 × 10 ⁻⁰³	1.61 × 10 ⁻⁰²	5.80 × 10 ⁻⁰⁴	1.59 × 10 ⁻⁰³	2.35 × 10 ⁻⁰³	2.83×10^{-03}	1.61 × 10 ⁻⁰²	5.80 × 10 ⁻⁰⁴	1.59 × 10 ⁻⁰³
		Patterson Lake North Arm – West Basin	5.46 × 10 ⁻⁰³	3.46 x 10 ⁻⁰³	1.72 x 10 ⁻⁰²	8.27 x 10 ⁻⁰⁴	1.17 x 10 ⁻⁰²	5.54 × 10 ⁻⁰³	3.48 x 10 ⁻⁰³	1.73 x 10 ⁻⁰²	1.20 x 10 ⁻⁰³	1.30 x 10 ⁻⁰²
	Moose	Patterson Lake South Arm	5.43×10^{-03}	3.41 x 10 ⁻⁰³	1.72 x 10 ⁻⁰²	7.93 x 10 ⁻⁰⁴	1.14 x 10 ⁻⁰²	5.52 × 10 ⁻⁰³	3.44 x 10 ⁻⁰³	1.72 x 10 ⁻⁰²	1.12 x 10 ⁻⁰³	1.27 x 10 ⁻⁰²
		Beet Lake	2.40×10^{-03}	3.01 x 10 ⁻⁰³	1.64 x 10 ⁻⁰²	6.33 x 10 ⁻⁰⁴	2.45 x 10 ⁻⁰³	2.40×10^{-03}	3.01 x 10 ⁻⁰³	1.64 x 10 ⁻⁰²	7.33 x 10 ⁻⁰⁴	2.58 x 10 ⁻⁰³
		Lloyd Lake	2.36 × 10 ⁻⁰³	2.85 x 10 ⁻⁰³	1.62 x 10 ⁻⁰²	5.90 x 10 ⁻⁰⁴	1.63 x 10 ⁻⁰³	2.36 × 10 ⁻⁰³	2.85 x 10 ⁻⁰³	1.62 x 10 ⁻⁰²	6.01 x 10 ⁻⁰⁴	1.64 x 10 ⁻⁰³
Terrestrial Animals		Reference (Broach Lake)	2.86 × 10 ⁻⁰²	1.60 × 10 ⁻⁰²	4.08×10^{-01}	3.65×10^{-03}	5.51 × 10 ⁻⁰³	2.86 × 10 ⁻⁰²	1.60×10^{-02}	4.08 × 10 ⁻⁰¹	3.65×10^{-03}	5.51 × 10 ⁻⁰³
Ailillais		Patterson Lake North Arm –West Basin	8.77 × 10 ⁻⁰²	2.08 × 10 ⁻⁰²	4.74×10^{-01}	9.77×10^{-03}	1.82 × 10 ⁻⁰²	8.92 × 10 ⁻⁰²	2.10×10^{-02}	4.76 × 10 ⁻⁰¹	2.23×10^{-02}	4.11 × 10 ⁻⁰²
	Muskrat	Patterson Lake South Arm	3.35 × 10 ⁻⁰²	1.86 × 10 ⁻⁰²	4.42×10^{-01}	6.80×10^{-03}	8.88 × 10 ⁻⁰³	3.36 × 10 ⁻⁰²	1.87×10^{-02}	4.43 × 10 ⁻⁰¹	1.34×10^{-02}	1.49 × 10 ⁻⁰²
		Beet Lake	2.94 × 10 ⁻⁰²	1.73 × 10 ⁻⁰²	4.25 × 10 ⁻⁰¹	5.26 × 10 ⁻⁰³	6.74×10^{-03}	2.94 × 10 ⁻⁰²	1.74×10^{-02}	4.25 × 10 ⁻⁰¹	8.67×10^{-03}	8.96 × 10 ⁻⁰³
		Lloyd Lake	2.86 × 10 ⁻⁰²	1.61 × 10 ⁻⁰²	4.09×10^{-01}	3.81×10^{-03}	5.62 × 10 ⁻⁰³	2.86 × 10 ⁻⁰²	1.61×10^{-02}	4.09 × 10 ⁻⁰¹	4.16×10^{-03}	5.82 × 10 ⁻⁰³
		Reference (Broach Lake)	1.27×10^{-03}	4.90×10^{-04}	6.99×10^{-03}	3.17×10^{-04}	3.72×10^{-04}	1.27×10^{-03}	4.90×10^{-04}	6.99 × 10 ⁻⁰³	3.17×10^{-04}	3.72×10^{-04}
		Patterson Lake North Arm – West Basin	1.32 × 10 ⁻⁰³	5.12 x 10 ⁻⁰⁴	7.26 x 10 ⁻⁰³	3.79 x 10 ⁻⁰⁴	3.45 x 10 ⁻⁰³	1.32 × 10 ⁻⁰³	5.12 x 10 ⁻⁰⁴	7.26 x 10 ⁻⁰³	4.47 x 10 ⁻⁰⁴	3.49 x 10 ⁻⁰³
	Red Fox	Patterson Lake South Arm	1.28 x 10 ⁻⁰³	5.01 x 10 ⁻⁰⁴	7.04 x 10 ⁻⁰³	3.39 x 10 ⁻⁰⁴	8.14 x 10 ⁻⁰⁴	1.28 × 10 ⁻⁰³	5.02 x 10 ⁻⁰⁴	7.04 x 10 ⁻⁰³	3.74 x 10 ⁻⁰⁴	8.26 x 10 ⁻⁰⁴
		Beet Lake	1.28 x 10 ⁻⁰³	4.97 x 10 ⁻⁰⁴	7.02 x 10 ⁻⁰³	3.29 x 10 ⁻⁰⁴	6.31 x 10 ⁻⁰⁴	1.28 × 10 ⁻⁰³	4.97 x 10 ⁻⁰⁴	7.02 x 10 ⁻⁰³	3.47 x 10 ⁻⁰⁴	6.36 x 10 ⁻⁰⁴
		Lloyd Lake	1.28 × 10 ⁻⁰³	4.93 x 10 ⁻⁰⁴	7.00 x 10 ⁻⁰³	3.21 x 10 ⁻⁰⁴	3.82 x 10 ⁻⁰⁴	1.28 × 10 ⁻⁰³	4.93 x 10 ⁻⁰⁴	7.00 x 10 ⁻⁰³	3.22 x 10 ⁻⁰⁴	3.82 x 10 ⁻⁰⁴
		Reference (Broach Lake)	4.30 × 10 ⁻⁰³	3.98 × 10 ⁻⁰³	4.82 × 10 ⁻⁰²	2.01 × 10 ⁻⁰³	5.27 × 10 ⁻⁰³	4.30 × 10 ⁻⁰³	3.98 × 10 ⁻⁰³	4.82 × 10 ⁻⁰²	2.01 × 10 ⁻⁰³	5.27 × 10 ⁻⁰³
		Patterson Lake North Arm – West Basin	4.38 × 10 ⁻⁰³	4.05 x 10 ⁻⁰³	5.00 x 10 ⁻⁰²	2.31 x 10 ⁻⁰³	4.23 x 10 ⁻⁰²	4.38 × 10 ⁻⁰³	4.05 x 10 ⁻⁰³	5.00 x 10 ⁻⁰²	2.38 x 10 ⁻⁰³	4.23 x 10 ⁻⁰²
	Snowshoe Hare	Patterson Lake South Arm	4.32 × 10 ⁻⁰³	4.01 x 10 ⁻⁰³	4.86 x 10 ⁻⁰²	2.08 x 10 ⁻⁰³	1.06 x 10 ⁻⁰²	4.32 × 10 ⁻⁰³	4.02 x 10 ⁻⁰³	4.86 x 10 ⁻⁰²	2.11 x 10 ⁻⁰³	1.06 x 10 ⁻⁰²
		Beet Lake	4.31 × 10 ⁻⁰³	4.01 x 10 ⁻⁰³	4.84 x 10 ⁻⁰²	2.05 x 10 ⁻⁰³	8.39 x 10 ⁻⁰³	4.31 × 10 ⁻⁰³	4.01 x 10 ⁻⁰³	4.84 x 10 ⁻⁰²	2.07 x 10 ⁻⁰³	8.39 x 10 ⁻⁰³
		Lloyd Lake	4.31 × 10 ⁻⁰³	4.01 x 10 ⁻⁰³	4.83 x 10 ⁻⁰²	2.04 x 10 ⁻⁰³	5.39 x 10 ⁻⁰³	4.31 × 10 ⁻⁰³	4.01 x 10 ⁻⁰³	4.83 x 10 ⁻⁰²	2.04 x 10 ⁻⁰³	5.39 x 10 ⁻⁰³
		Reference (Broach Lake)	3.10 × 10 ⁻⁰³	2.94 × 10 ⁻⁰³	3.66 × 10 ⁻⁰²	1.50 × 10 ⁻⁰³	3.95 × 10 ⁻⁰³	3.10 × 10 ⁻⁰³	2.94 × 10 ⁻⁰³	3.66 × 10 ⁻⁰²	1.50 × 10 ⁻⁰³	3.95 × 10 ⁻⁰³
	6 11 6 1	Patterson Lake North Arm – West Basin	3.18 × 10 ⁻⁰³	3.00 x 10 ⁻⁰³	3.79 x 10 ⁻⁰²	1.74 x 10 ⁻⁰³	3.05 x 10 ⁻⁰²	3.18 × 10 ⁻⁰³	3.00 x 10 ⁻⁰³	3.79 x 10 ⁻⁰²	1.85 x 10 ⁻⁰³	3.06 x 10 ⁻⁰²
	Southern Red- Backed Vole	Patterson Lake South Arm	3.12 × 10 ⁻⁰³	2.97 x 10 ⁻⁰³	3.69 x 10 ⁻⁰²	1.56 x 10 ⁻⁰³	7.75 x 10 ⁻⁰³	3.12 × 10 ⁻⁰³	2.97 x 10 ⁻⁰³	3.69 x 10 ⁻⁰²	1.62 x 10 ⁻⁰³	7.76 x 10 ⁻⁰³
	Backed Voie	Beet Lake	3.11 × 10 ⁻⁰³	2.96 x 10 ⁻⁰³	3.68 x 10 ⁻⁰²	1.53 x 10 ⁻⁰³	6.18 x 10 ⁻⁰³	3.11 × 10 ⁻⁰³	2.96 x 10 ⁻⁰³	3.68 x 10 ⁻⁰²	1.56 x 10 ⁻⁰³	6.19 x 10 ⁻⁰³
		Lloyd Lake	3.11 × 10 ⁻⁰³	2.96 x 10 ⁻⁰³	3.67 x 10 ⁻⁰²	1.52 x 10 ⁻⁰³	4.03 x 10 ⁻⁰³	3.11 × 10 ⁻⁰³	2.96 x 10 ⁻⁰³	3.67 x 10 ⁻⁰²	1.52 x 10 ⁻⁰³	4.03 x 10 ⁻⁰³
		Reference (Broach Lake)	5.46 x 10 ⁻⁰³	3.94 x 10 ⁻⁰³	1.78 x 10 ⁻⁰²	1.24 x 10 ⁻⁰³	1.77 x 10 ⁻⁰³	5.46 x 10 ⁻⁰³	3.94 x 10 ⁻⁰³	1.78 x 10 ⁻⁰²	1.24 x 10 ⁻⁰³	1.77 x 10 ⁻⁰³
	Woodland Caribou	Patterson Lake North Arm – West Basin	9.67 x 10 ⁻⁰³	4.70 x 10 ⁻⁰³	1.98 x 10 ⁻⁰²	2.44 x 10 ⁻⁰³	7.09 x 10 ⁻⁰²	9.77 x 10 ⁻⁰³	4.72 x 10 ⁻⁰³	1.98 x 10 ⁻⁰²	3.11 x 10 ⁻⁰³	7.28 x 10 ⁻⁰²
		Beet Lake	5.58 x 10 ⁻⁰³	4.11 x 10 ⁻⁰³	1.81 x 10 ⁻⁰²	1.36 x 10 ⁻⁰³	7.25 x 10 ⁻⁰³	5.58 x 10 ⁻⁰³	4.12 x 10 ⁻⁰³	1.81 x 10 ⁻⁰²	1.49 x 10 ⁻⁰³	7.41 x 10 ⁻⁰³
					•		•			•		

		Maximum Dose During Operations (mg/kg/d)												
Biota	Location			Application Case				U	Jpper Bound Scenari	0				
		Arsenic	Cobalt	Copper	Molybdenum	Uranium	Arsenic	Cobalt	Copper	Molybdenum	Uranium			
	Lloyd Lake	5.60 x 10 ⁻⁰³	4.10 x 10 ⁻⁰³	1.80 x 10 ⁻⁰²	1.39 x 10 ⁻⁰³	2.21 x 10 ⁻⁰³	5.60 x 10 ⁻⁰³	4.10 x 10 ⁻⁰³	1.80 x 10 ⁻⁰²	1.41 x 10 ⁻⁰³	2.24 x 10 ⁻⁰³			
	Reference (Broach Lake)	6.44 × 10 ⁻⁰³	5.57 × 10 ⁻⁰³	5.26 × 10 ⁻⁰²	2.57 × 10 ⁻⁰³	6.34 × 10 ⁻⁰³	6.44 × 10 ⁻⁰³	5.57 × 10 ⁻⁰³	5.26 × 10 ⁻⁰²	2.57 × 10 ⁻⁰³	6.34 × 10 ⁻⁰³			
	Patterson Lake North Arm – West Basin	6.51 × 10 ⁻⁰³	5.65 x 10 ⁻⁰³	5.45 x 10 ⁻⁰²	2.88 x 10 ⁻⁰³	4.81 x 10 ⁻⁰²	6.51 × 10 ⁻⁰³	5.65 x 10 ⁻⁰³	5.45 x 10 ⁻⁰²	2.94 x 10 ⁻⁰³	4.81 x 10 ⁻⁰²			
Grouse	Patterson Lake South Arm	6.46 × 10 ⁻⁰³	5.61 x 10 ⁻⁰³	5.29 x 10 ⁻⁰²	2.64 x 10 ⁻⁰³	1.23 x 10 ⁻⁰²	6.46 × 10 ⁻⁰³	5.61 x 10 ⁻⁰³	5.29 x 10 ⁻⁰²	2.66 x 10 ⁻⁰³	1.23 x 10 ⁻⁰²			
	Beet Lake	6.45 × 10 ⁻⁰³	5.61 x 10 ⁻⁰³	5.28 x 10 ⁻⁰²	2.61 x 10 ⁻⁰³	9.86 x 10 ⁻⁰³	6.45 × 10 ⁻⁰³	5.61 x 10 ⁻⁰³	5.28 x 10 ⁻⁰²	2.62 x 10 ⁻⁰³	9.86 x 10 ⁻⁰³			
	Lloyd Lake	6.45 × 10 ⁻⁰³	5.61 x 10 ⁻⁰³	5.27 x 10 ⁻⁰²	2.60 x 10 ⁻⁰³	6.48 x 10 ⁻⁰³	6.45 × 10 ⁻⁰³	5.61 x 10 ⁻⁰³	5.27 x 10 ⁻⁰²	2.60 x 10 ⁻⁰³	6.48 x 10 ⁻⁰³			
	Reference (Broach Lake)	6.19 × 10 ⁻⁰⁴	5.56 × 10 ⁻⁰⁴	5.30 × 10 ⁻⁰³	2.58 × 10 ⁻⁰⁴	6.29 × 10 ⁻⁰⁴	6.19 × 10 ⁻⁰⁴	5.56 × 10 ⁻⁰⁴	5.30 × 10 ⁻⁰³	2.58 × 10 ⁻⁰⁴	6.29 × 10 ⁻⁰⁴			
	Patterson Lake North Arm – West Basin	6.36 × 10 ⁻⁰⁴	5.69 x 10 ⁻⁰⁴	5.49 x 10 ⁻⁰³	3.02 x 10 ⁻⁰⁴	4.86 x 10 ⁻⁰³	6.37 × 10 ⁻⁰⁴	5.70 x 10 ⁻⁰⁴	5.49 x 10 ⁻⁰³	3.32 x 10 ⁻⁰⁴	4.88 x 10 ⁻⁰³			
Canada Goose	Patterson Lake South Arm	6.22 × 10 ⁻⁰⁴	5.63 x 10 ⁻⁰⁴	5.34 x 10 ⁻⁰³	2.70 x 10 ⁻⁰⁴	1.24 x 10 ⁻⁰³	6.22 × 10 ⁻⁰⁴	5.63 x 10 ⁻⁰⁴	5.34 x 10 ⁻⁰³	2.86 x 10 ⁻⁰⁴	1.24 x 10 ⁻⁰³			
	Beet Lake	6.21 × 10 ⁻⁰⁴	5.61 x 10 ⁻⁰⁴	5.32 x 10 ⁻⁰³	2.64 x 10 ⁻⁰⁴	9.86 x 10 ⁻⁰⁴	6.21 × 10 ⁻⁰⁴	5.61 x 10 ⁻⁰⁴	5.32 x 10 ⁻⁰³	2.72 x 10 ⁻⁰⁴	9.88 x 10 ⁻⁰⁴			
	Lloyd Lake	6.21 × 10 ⁻⁰⁴	5.59 x 10 ⁻⁰⁴	5.31 x 10 ⁻⁰³	2.61 x 10 ⁻⁰⁴	6.42 x 10 ⁻⁰⁴	6.21 × 10 ⁻⁰⁴	5.59 x 10 ⁻⁰⁴	5.31 x 10 ⁻⁰³	2.62 x 10 ⁻⁰⁴	6.43 x 10 ⁻⁰⁴			
	Reference (Broach Lake)	2.77 × 10 ⁻⁰²	5.73 × 10 ⁻⁰²	3.22 × 10 ⁺⁰⁰	1.64 × 10 ⁻⁰²	4.77 × 10 ⁻⁰³	2.77 × 10 ⁻⁰²	5.73 × 10 ⁻⁰²	3.22 × 10 ⁺⁰⁰	1.64 × 10 ⁻⁰²	4.77 × 10 ⁻⁰³			
Little Brown Myotis	Patterson Lake North Arm – West Basin	8.44 × 10 ⁻⁰²	7.41 × 10 ⁻⁰²	3.74 × 10 ⁺⁰⁰	4.38 × 10 ⁻⁰²	1.57 × 10 ⁻⁰²	8.57 × 10 ⁻⁰²	7.48 × 10 ⁻⁰²	3.76 × 10 ⁺⁰⁰	1.00 × 10 ⁻⁰¹	3.52 × 10 ⁻⁰²			
	Patterson Lake South Arm	3.23 × 10 ⁻⁰²	6.62 × 10 ⁻⁰²	3.49 × 10 ⁺⁰⁰	3.05 × 10 ⁻⁰²	7.66 × 10 ⁻⁰³	3.25 × 10 ⁻⁰²	6.66 × 10 ⁻⁰²	3.50 × 10 ⁺⁰⁰	6.03 × 10 ⁻⁰²	1.28 × 10 ⁻⁰²			
	Beet Lake	2.84 × 10 ⁻⁰²	6.19 × 10 ⁻⁰²	3.36 × 10 ⁺⁰⁰	2.36 × 10 ⁻⁰²	5.83 × 10 ⁻⁰³	2.84 × 10 ⁻⁰²	6.21 × 10 ⁻⁰²	3.36 × 10 ⁺⁰⁰	3.89 × 10 ⁻⁰²	7.73 × 10 ⁻⁰³			
	Lloyd Lake	2.77 × 10 ⁻⁰²	5.77 × 10 ⁻⁰²	3.23 × 10 ⁺⁰⁰	1.71 × 10 ⁻⁰²	4.87 × 10 ⁻⁰³	2.77 × 10 ⁻⁰²	5.78 × 10 ⁻⁰²	3.23 × 10 ⁺⁰⁰	1.87 × 10 ⁻⁰²	5.04 × 10 ⁻⁰³			
	Reference (Broach Lake)	4.71 × 10 ⁻⁰³	2.18 × 10 ⁻⁰³	1.25 × 10 ⁻⁰¹	4.10 × 10 ⁻⁰⁴	3.38 × 10 ⁻⁰⁴	4.71 × 10 ⁻⁰³	2.18 × 10 ⁻⁰³	1.25 × 10 ⁻⁰¹	4.10 × 10 ⁻⁰⁴	3.38 × 10 ⁻⁰⁴			
	Patterson Lake North Arm – West Basin	1.63 × 10 ⁻⁰²	2.84 × 10 ⁻⁰³	1.46 × 10 ⁻⁰¹	1.10 × 10 ⁻⁰³	1.22 × 10 ⁻⁰³	1.66 × 10 ⁻⁰²	2.87 × 10 ⁻⁰³	1.46 × 10 ⁻⁰¹	2.50 × 10 ⁻⁰³	2.85 × 10 ⁻⁰³			
Common Loon	Patterson Lake South Arm	5.67 × 10 ⁻⁰³	2.53 × 10 ⁻⁰³	1.35 × 10 ⁻⁰¹	7.64 × 10 ⁻⁰⁴	5.72 × 10 ⁻⁰⁴	5.69 × 10 ⁻⁰³	2.54 × 10 ⁻⁰³	1.36 × 10 ⁻⁰¹	1.51 × 10 ⁻⁰³	9.85 × 10 ⁻⁰⁴			
	Beet Lake	4.87 × 10 ⁻⁰³	2.36 × 10 ⁻⁰³	1.30 × 10 ⁻⁰¹	5.91 × 10 ⁻⁰⁴	4.24 × 10 ⁻⁰⁴	4.87 × 10 ⁻⁰³	2.37 × 10 ⁻⁰³	1.30 × 10 ⁻⁰¹	9.74 × 10 ⁻⁰⁴	5.75 × 10 ⁻⁰⁴			
	Lloyd Lake	4.72 × 10 ⁻⁰³	2.20 × 10 ⁻⁰³	1.25 × 10 ⁻⁰¹	4.28 × 10 ⁻⁰⁴	3.46 × 10 ⁻⁰⁴	4.72 × 10 ⁻⁰³	2.20 × 10 ⁻⁰³	1.25 × 10 ⁻⁰¹	4.68 × 10 ⁻⁰⁴	3.60 × 10 ⁻⁰⁴			
	Reference (Broach Lake)	2.78 × 10 ⁻⁰²	2.60 × 10 ⁻⁰²	1.05 × 10 ⁺⁰⁰	6.52 × 10 ⁻⁰³	5.23 × 10 ⁻⁰³	2.78 × 10 ⁻⁰²	2.60 × 10 ⁻⁰²	1.05 × 10 ⁺⁰⁰	6.52 × 10 ⁻⁰³	5.23 × 10 ⁻⁰³			
	Patterson Lake North Arm – West Basin	8.52 × 10 ⁻⁰²	3.37 × 10 ⁻⁰²	1.23 × 10 ⁺⁰⁰	1.75 × 10 ⁻⁰²	1.73 × 10 ⁻⁰²	8.66 × 10 ⁻⁰²	3.40 × 10 ⁻⁰²	1.23 × 10 ⁺⁰⁰	3.98 × 10 ⁻⁰²	3.90 × 10 ⁻⁰²			
Mallard	Patterson Lake South Arm	3.25 × 10 ⁻⁰²	3.01 × 10 ⁻⁰²	1.14 × 10 ⁺⁰⁰	1.22 × 10 ⁻⁰²	8.42 × 10 ⁻⁰³	3.26 × 10 ⁻⁰²	3.03 × 10 ⁻⁰²	1.14 × 10 ⁺⁰⁰	2.40 × 10 ⁻⁰²	1.42 × 10 ⁻⁰²			
	Beet Lake	2.85 × 10 ⁻⁰²	2.81 × 10 ⁻⁰²	1.10 × 10 ⁺⁰⁰	9.41 × 10 ⁻⁰³	6.40 × 10 ⁻⁰³	2.85 × 10 ⁻⁰²	2.82 × 10 ⁻⁰²	1.10 × 10 ⁺⁰⁰	1.55 × 10 ⁻⁰²	8.50 × 10 ⁻⁰³			
	Lloyd Lake	2.78 × 10 ⁻⁰²	2.62 × 10 ⁻⁰²	1.06 × 10 ⁺⁰⁰	6.82 × 10 ⁻⁰³	5.33 × 10 ⁻⁰³	2.78 × 10 ⁻⁰²	2.62 × 10 ⁻⁰²	1.06 × 10 ⁺⁰⁰	7.44 × 10 ⁻⁰³	5.52 × 10 ⁻⁰³			
	Reference (Broach Lake)	7.16 × 10 ⁻⁰²	1.25 × 10 ⁻⁰¹	6.35 × 10 ⁺⁰⁰	3.66 × 10 ⁻⁰²	1.76 × 10 ⁻⁰²	7.16 × 10 ⁻⁰²	1.25 × 10 ⁻⁰¹	6.35 × 10 ⁺⁰⁰	3.66 × 10 ⁻⁰²	1.76 × 10 ⁻⁰²			
	Patterson Lake North Arm – West Basin	1.83 × 10 ⁻⁰¹	1.58 x 10 ⁻⁰¹	7.38 x 10 ⁺⁰⁰	9.06 x 10 ⁻⁰²	7.62 x 10 ⁻⁰²	1.86 × 10 ⁻⁰¹	1.60 x 10 ⁻⁰¹	7.41 x 10 ⁺⁰⁰	2.01 x 10 ⁻⁰¹	1.14 x 10 ⁻⁰¹			
Rusty Blackbird	Patterson Lake South Arm	8.08 × 10 ⁻⁰²	1.43 x 10 ⁻⁰¹	6.88 x 10 ⁺⁰⁰	6.44 x 10 ⁻⁰²	2.85 x 10 ⁻⁰²	8.11 × 10 ⁻⁰²	1.44 x 10 ⁻⁰¹	6.89 x 10 ⁺⁰⁰	1.23 x 10 ⁻⁰¹	3.87 x 10 ⁻⁰²			
	Beet Lake	7.31 × 10 ⁻⁰²	1.34 x 10 ⁻⁰¹	6.62 x 10 ⁺⁰⁰	5.08 x 10 ⁻⁰²	2.28 x 10 ⁻⁰²	7.32 × 10 ⁻⁰²	1.35 x 10 ⁻⁰¹	6.63 x 10 ⁺⁰⁰	8.08 x 10 ⁻⁰²	2.65 x 10 ⁻⁰²			
	Lloyd Lake	7.17 × 10 ⁻⁰²	1.26 x 10 ⁻⁰¹	6.38 x 10 ⁺⁰⁰	3.81 x 10 ⁻⁰²	1.79 x 10 ⁻⁰²	7.17 × 10 ⁻⁰²	1.26 x 10 ⁻⁰¹	6.38 x 10 ⁺⁰⁰	4.12 x 10 ⁻⁰²	1.82 x 10 ⁻⁰²			

Table 6-11: Estimated Non-radiological Doses to Ecological Receptors – Far-Future Projection – Application Case and Upper Bound Scenario

		Posses to Ecological Recoptors Tai Tata					aximum Dose Far-F	uture (mg/kg/d)				
	Biota	Location			Application Case				U	pper Bound Scenar	io	
			Arsenic	Cobalt	Copper	Molybdenum	Uranium	Arsenic	Cobalt	Copper	Molybdenum	Uranium
		Reference (Broach Lake)	3.48 × 10 ⁻⁰³	3.94 × 10 ⁻⁰³	3.15 × 10 ⁻⁰²	1.17 × 10 ⁻⁰³	3.20×10^{-03}	3.48 × 10 ⁻⁰³	3.94 × 10 ⁻⁰³	3.15 × 10 ⁻⁰²	1.17 × 10 ⁻⁰³	3.20 × 10 ⁻⁰³
		Patterson Lake North Arm – West Basin	3.57 × 10 ⁻⁰³	6.75 x 10 ⁻⁰³	3.76 x 10 ⁻⁰²	3.39 x 10 ⁻⁰³	7.09 x 10 ⁻⁰³	3.61 x 10 ⁻⁰³	8.24 x 10 ⁻⁰³	4.06 x 10 ⁻⁰²	9.68 x 10 ⁻⁰³	7.09 x 10 ⁻⁰³
	Beaver	Patterson Lake South Arm	3.49×10^{-03}	5.64 x 10 ⁻⁰³	3.50 x 10 ⁻⁰²	2.46 x 10 ⁻⁰³	4.10 x 10 ⁻⁰³	3.49 x 10 ⁻⁰³	6.55 x 10 ⁻⁰³	3.68 x 10 ⁻⁰²	6.11 x 10 ⁻⁰³	4.10 x 10 ⁻⁰³
		Beet Lake	3.49×10^{-03}	4.84 x 10 ⁻⁰³	3.34 x 10 ⁻⁰²	1.84 x 10 ⁻⁰³	3.57 x 10 ⁻⁰³	3.49 x 10 ⁻⁰³	5.32 x 10 ⁻⁰³	3.43 x 10 ⁻⁰²	3.75 x 10 ⁻⁰³	3.57 x 10 ⁻⁰³
		Lloyd Lake	3.48×10^{-03}	4.03 x 10 ⁻⁰³	3.17 x 10 ⁻⁰²	1.24 x 10 ⁻⁰³	3.23 x 10 ⁻⁰³	3.48 x 10 ⁻⁰³	4.08 x 10 ⁻⁰³	3.18 x 10 ⁻⁰²	1.44 x 10 ⁻⁰³	3.23 x 10 ⁻⁰³
		Reference (Broach Lake)	2.81 × 10 ⁻⁰³	1.17 × 10 ⁻⁰³	2.86 × 10 ⁻⁰²	3.75 × 10 ⁻⁰⁴	1.02×10^{-03}	2.81 × 10 ⁻⁰³	1.17 × 10 ⁻⁰³	2.86 × 10 ⁻⁰²	3.75 × 10 ⁻⁰⁴	1.02 × 10 ⁻⁰³
	Black Bear	Patterson Lake North Arm – West Basin	2.87×10^{-03}	1.63 x 10 ⁻⁰³	5.39 x 10 ⁻⁰²	6.09 x 10 ⁻⁰⁴	1.79 x 10 ⁻⁰³	2.90 x 10 ⁻⁰³	1.86 x 10 ⁻⁰³	6.66 x 10 ⁻⁰²	1.24 x 10 ⁻⁰³	1.79 x 10 ⁻⁰³
	Diack Deal	Beet Lake	3.09×10^{-03}	1.49 x 10 ⁻⁰³	3.39 x 10 ⁻⁰²	4.95 x 10 ⁻⁰⁴	1.20 x 10 ⁻⁰³	3.09 x 10 ⁻⁰³	1.53 x 10 ⁻⁰³	3.63 x 10 ⁻⁰²	6.17 x 10 ⁻⁰⁴	1.20 x 10 ⁻⁰³
		Lloyd Lake	2.82×10^{-03}	1.22 x 10 ⁻⁰³	2.97 x 10 ⁻⁰²	3.97 x 10 ⁻⁰⁴	1.03 x 10 ⁻⁰³	2.82 x 10 ⁻⁰³	1.23 x 10 ⁻⁰³	3.02 x 10 ⁻⁰²	4.23 x 10 ⁻⁰⁴	1.03 x 10 ⁻⁰³
		Reference (Broach Lake)	7.50 x 10 ⁻⁰³	6.01 x 10 ⁻⁰⁴	3.31 x 10 ⁻⁰²	2.88 x 10 ⁻⁰⁴	3.73 x 10 ⁻⁰⁴	7.50 x 10 ⁻⁰³	6.01 x 10 ⁻⁰⁴	3.31 x 10 ⁻⁰²	2.88 x 10 ⁻⁰⁴	3.73 x 10 ⁻⁰⁴
	Grey Wolf	Patterson Lake North Arm – West Basin	7.53 x 10 ⁻⁰³	6.45 x 10 ⁻⁰⁴	3.43 x 10 ⁻⁰²	4.47 x 10 ⁻⁰⁴	4.53 x 10 ⁻⁰⁴	7.55 x 10 ⁻⁰³	6.67 x 10 ⁻⁰⁴	3.49 x 10 ⁻⁰²	9.00 x 10 ⁻⁰⁴	4.53 x 10 ⁻⁰⁴
		Lloyd Lake	7.50 x 10 ⁻⁰³	6.06 x 10 ⁻⁰⁴	3.32 x 10 ⁻⁰²	3.03 x 10 ⁻⁰⁴	3.75 x 10 ⁻⁰⁴	7.50 x 10 ⁻⁰³	6.09 x 10 ⁻⁰⁴	3.33 x 10 ⁻⁰²	3.46 x 10 ⁻⁰⁴	3.75 x 10 ⁻⁰⁴
		Reference (Broach Lake)	1.15 × 10 ⁻⁰²	4.61 × 10 ⁻⁰³	2.28 × 10 ⁻⁰¹	1.15 × 10 ⁻⁰³	1.08 × 10 ⁻⁰³	1.15 × 10 ⁻⁰²	4.61 × 10 ⁻⁰³	2.28 × 10 ⁻⁰¹	1.15 × 10 ⁻⁰³	1.08 × 10 ⁻⁰³
		Patterson Lake North Arm – West Basin	1.22 × 10 ⁻⁰²	1.06 x 10 ⁻⁰²	5.90 x 10-01	2.20 x 10 ⁻⁰²	8.39 x 10 ⁻⁰³	1.25 x 10 ⁻⁰²	1.37 x 10 ⁻⁰²	7.72 x 10-01	8.14 x 10 ⁻⁰²	8.40 x 10 ⁻⁰³
	American Mink	Patterson Lake South Arm	1.16 × 10 ⁻⁰²	8.23 x 10 ⁻⁰³	4.41 x 10-01	1.33 x 10 ⁻⁰²	3.09 x 10 ⁻⁰³	1.16 x 10 ⁻⁰²	1.02 x 10 ⁻⁰²	5.47 x 10-01	4.77 x 10 ⁻⁰²	3.09 x 10 ⁻⁰³
		Beet Lake	1.15 × 10 ⁻⁰²	6.53 x 10 ⁻⁰³	3.39 x 10-01	7.46 x 10 ⁻⁰³	1.83 x 10 ⁻⁰³	1.15 x 10 ⁻⁰²	7.55 x 10 ⁻⁰³	3.95 x 10-01	2.54 x 10 ⁻⁰²	1.83 x 10 ⁻⁰³
		Lloyd Lake	1.15×10^{-02}	4.81 x 10 ⁻⁰³	2.40 x 10-01	1.79 x 10 ⁻⁰³	1.15 x 10 ⁻⁰³	1.15 x 10 ⁻⁰²	4.91 x 10 ⁻⁰³	2.45 x 10-01	3.62 x 10 ⁻⁰³	1.15 x 10 ⁻⁰³
		Reference (Broach Lake)	2.35 × 10 ⁻⁰³	2.83 × 10 ⁻⁰³	1.61 × 10 ⁻⁰²	5.80 × 10 ⁻⁰⁴	1.59×10^{-03}	2.35 × 10 ⁻⁰³	2.83 × 10 ⁻⁰³	1.61 × 10 ⁻⁰²	5.80 × 10 ⁻⁰⁴	1.59 × 10 ⁻⁰³
	Moose	Patterson Lake North Arm – West Basin	2.44×10^{-03}	5.63 x 10 ⁻⁰³	2.22 x 10 ⁻⁰²	2.78 x 10 ⁻⁰³	4.82 x 10 ⁻⁰³	2.49 x 10 ⁻⁰³	7.11 x 10 ⁻⁰³	2.52 x 10 ⁻⁰²	9.02 x 10 ⁻⁰³	4.82 x 10 ⁻⁰³
		Patterson Lake South Arm	2.43×10^{-03}	5.55 x 10 ⁻⁰³	2.20 x 10 ⁻⁰²	2.53 x 10 ⁻⁰³	4.74 x 10 ⁻⁰³	2.48 x 10 ⁻⁰³	7.01 x 10 ⁻⁰³	2.50 x 10 ⁻⁰²	8.09 x 10 ⁻⁰³	4.74 x 10 ⁻⁰³
		Beet Lake	2.36 × 10 ⁻⁰³	3.73 x 10 ⁻⁰³	1.80 x 10 ⁻⁰²	1.24 x 10 ⁻⁰³	1.91 x 10 ⁻⁰³	2.36 x 10 ⁻⁰³	4.20 x 10 ⁻⁰³	1.89 x 10 ⁻⁰²	3.13 x 10 ⁻⁰³	1.91 x 10 ⁻⁰³
Townstale		Lloyd Lake	2.35 × 10 ⁻⁰³	2.92 x 10 ⁻⁰³	1.63 x 10 ⁻⁰²	6.48 x 10 ⁻⁰⁴	1.62 x 10 ⁻⁰³	2.35 x 10 ⁻⁰³	2.97 x 10 ⁻⁰³	1.64 x 10 ⁻⁰²	8.40 x 10 ⁻⁰⁴	1.62 x 10 ⁻⁰³
Terrestrial Animals		Reference (Broach Lake)	2.86 × 10 ⁻⁰²	1.60 × 10 ⁻⁰²	4.08×10^{-01}	3.65 × 10 ⁻⁰³	5.51×10^{-03}	2.86 × 10 ⁻⁰²	1.60 × 10 ⁻⁰²	4.08 × 10 ⁻⁰¹	3.65 × 10 ⁻⁰³	5.51 × 10 ⁻⁰³
7 tillions		Patterson Lake North Arm - West Basin	3.03 × 10 ⁻⁰²	3.91 × 10 ⁻⁰²	$1.05 \times 10^{+00}$	8.14 × 10 ⁻⁰²	5.51 × 10 ⁻⁰²	3.12 × 10 ⁻⁰²	5.14 × 10 ⁻⁰²	1.38 × 10 ⁺⁰⁰	3.02 × 10 ⁻⁰¹	5.51 × 10 ⁻⁰²
	Muskrat	Patterson Lake South Arm	2.87 × 10 ⁻⁰²	3.00 × 10 ⁻⁰²	7.87 × 10 ⁻⁰¹	4.88 × 10 ⁻⁰²	1.92 × 10 ⁻⁰²	2.88 × 10 ⁻⁰²	3.75 × 10 ⁻⁰²	9.78 × 10 ⁻⁰¹	1.77 × 10 ⁻⁰¹	1.92 × 10 ⁻⁰²
		Beet Lake	2.86 × 10 ⁻⁰²	2.34 × 10 ⁻⁰²	6.06 × 10 ⁻⁰¹	2.71 × 10 ⁻⁰²	1.06 × 10 ⁻⁰²	2.86 × 10 ⁻⁰²	2.74 × 10 ⁻⁰²	7.06 × 10 ⁻⁰¹	9.39 × 10 ⁻⁰²	1.06 × 10 ⁻⁰²
		Lloyd Lake	2.86 × 10 ⁻⁰²	1.67 × 10 ⁻⁰²	4.28 × 10 ⁻⁰¹	6.04×10^{-03}	5.97×10^{-03}	2.86 × 10 ⁻⁰²	1.71 × 10 ⁻⁰²	4.38 × 10 ⁻⁰¹	1.28 × 10 ⁻⁰²	5.97 × 10 ⁻⁰³
		Reference (Broach Lake)	1.27×10^{-03}	4.90 × 10 ⁻⁰⁴	6.99 × 10 ⁻⁰³	3.17 × 10 ⁻⁰⁴	3.72×10^{-04}	1.27 × 10 ⁻⁰³	4.90 × 10 ⁻⁰⁴	6.99 × 10 ⁻⁰³	3.17 × 10 ⁻⁰⁴	3.72 × 10 ⁻⁰⁴
		Patterson Lake North Arm - West Basin	1.27×10^{-03}	5.64 x 10 ⁻⁰⁴	7.12 x 10 ⁻⁰³	6.94 x 10 ⁻⁰⁴	6.12 x 10 ⁻⁰⁴	1.27 x 10 ⁻⁰³	6.03 x 10 ⁻⁰⁴	7.18 x 10 ⁻⁰³	1.76 x 10 ⁻⁰³	6.12 x 10 ⁻⁰⁴
	Red Fox	Patterson Lake South Arm	1.27×10^{-03}	5.35 x 10 ⁻⁰⁴	7.06 x 10 ⁻⁰³	5.36 x 10 ⁻⁰⁴	4.17 x 10 ⁻⁰⁴	1.27 x 10 ⁻⁰³	5.58 x 10 ⁻⁰⁴	7.09 x 10 ⁻⁰³	1.16 x 10 ⁻⁰³	4.17 x 10 ⁻⁰⁴
		Beet Lake	1.27×10^{-03}	5.14 x 10 ⁻⁰⁴	7.03 x 10 ⁻⁰³	4.31 x 10 ⁻⁰⁴	3.94 x 10 ⁻⁰⁴	1.27 x 10 ⁻⁰³	5.26 x 10 ⁻⁰⁴	7.04 x 10 ⁻⁰³	7.54 x 10 ⁻⁰⁴	3.94 x 10 ⁻⁰⁴
		Lloyd Lake	1.27×10^{-03}	4.93 x 10 ⁻⁰⁴	6.99 x 10 ⁻⁰³	3.29 x 10 ⁻⁰⁴	3.73 x 10 ⁻⁰⁴	1.27 x 10 ⁻⁰³	4.94 x 10 ⁻⁰⁴	6.99 x 10 ⁻⁰³	3.62 x 10 ⁻⁰⁴	3.73 x 10 ⁻⁰⁴
		Reference (Broach Lake)	4.30×10^{-03}	3.98 × 10 ⁻⁰³	4.82×10^{-02}	2.01×10^{-03}	5.27×10^{-03}	4.30×10^{-03}	3.98 × 10 ⁻⁰³	4.82 × 10 ⁻⁰²	2.01 × 10 ⁻⁰³	5.27×10^{-03}
		Patterson Lake North Arm - West Basin	4.30×10^{-03}	4.07 x 10 ⁻⁰³	4.85 x 10 ⁻⁰²	2.43 × 10 ⁻⁰³	7.70 x 10 ⁻⁰³	4.30 × 10 ⁻⁰³	4.11 x 10 ⁻⁰³	4.85 x 10 ⁻⁰²	3.61 × 10 ⁻⁰³	7.70×10^{-03}
	Snowshoe Hare	Patterson Lake South Arm	4.30×10^{-03}	4.03 x 10 ⁻⁰³	4.83 x 10 ⁻⁰²	2.25×10^{-03}	5.63 x 10 ⁻⁰³	4.30×10^{-03}	4.06 x 10 ⁻⁰³	4.84 x 10 ⁻⁰²	2.94 × 10 ⁻⁰³	5.63 × 10 ⁻⁰³
		Beet Lake	4.30×10^{-03}	4.01 x 10 ⁻⁰³	4.83 x 10 ⁻⁰²	2.14×10^{-03}	5.48 x 10 ⁻⁰³	4.30×10^{-03}	4.02 x 10 ⁻⁰³	4.83 x 10 ⁻⁰²	2.50×10^{-03}	5.48 × 10 ⁻⁰³
		Lloyd Lake	4.30 × 10 ⁻⁰³	3.99 x 10 ⁻⁰³	4.82 x 10 ⁻⁰²	2.03 × 10 ⁻⁰³	5.28 x 10 ⁻⁰³	4.30 × 10 ⁻⁰³	3.99 x 10 ⁻⁰³	4.83 x 10 ⁻⁰²	2.06 × 10 ⁻⁰³	5.28 × 10 ⁻⁰³
		Reference (Broach Lake)	3.10 × 10 ⁻⁰³	2.94 × 10 ⁻⁰³	3.66 × 10 ⁻⁰²	1.50 × 10 ⁻⁰³	3.95×10^{-03}	3.10 × 10 ⁻⁰³	2.94 × 10 ⁻⁰³	3.66 × 10 ⁻⁰²	1.50 × 10 ⁻⁰³	3.95 × 10 ⁻⁰³
	Causham Balana	Patterson Lake North Arm - West Basin	3.10 × 10 ⁻⁰³	3.06 x 10 ⁻⁰³	3.69 x 10 ⁻⁰²	2.12 x 10 ⁻⁰³	5.83 x 10 ⁻⁰³	3.10 x 10 ⁻⁰³	3.13 x 10 ⁻⁰³	3.70 x 10 ⁻⁰²	3.85 x 10 ⁻⁰³	5.83 x 10 ⁻⁰³
	Southern Red-Backed Vole	Patterson Lake South Arm	3.10×10^{-03}	3.01 x 10 ⁻⁰³	3.67 x 10 ⁻⁰²	1.86 x 10 ⁻⁰³	4.23 x 10 ⁻⁰³	3.10 x 10 ⁻⁰³	3.05 x 10 ⁻⁰³	3.68 x 10 ⁻⁰²	2.87 x 10 ⁻⁰³	4.23 x 10 ⁻⁰³
	7016	Beet Lake	3.10 × 10 ⁻⁰³	2.98 x 10 ⁻⁰³	3.67 x 10 ⁻⁰²	1.69 x 10 ⁻⁰³	4.11 x 10 ⁻⁰³	3.10 x 10 ⁻⁰³	3.00 x 10 ⁻⁰³	3.67 x 10 ⁻⁰²	2.21 x 10 ⁻⁰³	4.11 x 10 ⁻⁰³
		Lloyd Lake	3.10 × 10 ⁻⁰³	2.95 x 10 ⁻⁰³	3.66 x 10 ⁻⁰²	1.52 x 10 ⁻⁰³	3.95 x 10 ⁻⁰³	3.10 x 10 ⁻⁰³	2.95 x 10 ⁻⁰³	3.66 x 10 ⁻⁰²	1.58 x 10 ⁻⁰³	3.95 x 10 ⁻⁰³
		Reference (Broach Lake)	5.46 x 10 ⁻⁰³	3.94 x 10 ⁻⁰³	1.78 x 10 ⁻⁰²	1.24 x 10 ⁻⁰³	1.77 x 10 ⁻⁰³	5.46 x 10 ⁻⁰³	3.94 x 10 ⁻⁰³	1.78 x 10 ⁻⁰²	1.24 x 10 ⁻⁰³	1.77 x 10 ⁻⁰³
	Woodland Caribou	Patterson Lake North Arm - West Basin	5.57 x 10 ⁻⁰³	6.64 x 10 ⁻⁰³	2.34 x 10 ⁻⁰²	5.54 x 10 ⁻⁰³	6.08 x 10 ⁻⁰³	5.63 x 10 ⁻⁰³	8.08 x 10 ⁻⁰³	2.63 x 10 ⁻⁰²	1.77 x 10 ⁻⁰²	6.08 x 10 ⁻⁰³
		Beet Lake	5.46 x 10 ⁻⁰³	4.49 x 10 ⁻⁰³	1.89 x 10 ⁻⁰²	2.07 x 10 ⁻⁰³	2.13 x 10 ⁻⁰³	5.46 x 10 ⁻⁰³	4.78 x 10 ⁻⁰³	1.95 x 10 ⁻⁰²	4.43 x 10 ⁻⁰³	2.13 x 10 ⁻⁰³

ECOLOGICAL RISK ASSESSMENT

					Ma	aximum Dose Far-F	uture (mg/kg/d)				
Biota	Location			Application Case				U	pper Bound Scenai	rio	
		Arsenic	Cobalt	Copper	Molybdenum	Uranium	Arsenic	Cobalt	Copper	Molybdenum	Uranium
	Lloyd Lake	5.46 x 10 ⁻⁰³	4.06 x 10 ⁻⁰³	1.80 x 10 ⁻⁰²	1.41 x 10 ⁻⁰³	1.83 x 10 ⁻⁰³	5.46 x 10 ⁻⁰³	4.12 x 10 ⁻⁰³	1.82 x 10 ⁻⁰²	1.92 x 10 ⁻⁰³	1.83 x 10 ⁻⁰³
	Reference (Broach Lake)	6.44 × 10 ⁻⁰³	5.57 × 10 ⁻⁰³	5.26 × 10 ⁻⁰²	2.57 × 10 ⁻⁰³	6.34 × 10 ⁻⁰³	6.44 × 10 ⁻⁰³	5.57 × 10 ⁻⁰³	5.26 × 10 ⁻⁰²	2.57 × 10 ⁻⁰³	6.34 × 10 ⁻⁰³
	Patterson Lake North Arm - West Basin	6.44 × 10 ⁻⁰³	5.65 x 10 ⁻⁰³	5.28 x 10 ⁻⁰²	2.88 x 10 ⁻⁰³	9.23 x 10 ⁻⁰³	6.44 x 10 ⁻⁰³	5.68 x 10 ⁻⁰³	5.28 x 10 ⁻⁰²	3.75 x 10 ⁻⁰³	9.23 x 10 ⁻⁰³
Grouse	Patterson Lake South Arm	6.44 × 10 ⁻⁰³	5.62 x 10 ⁻⁰³	5.27 x 10 ⁻⁰²	2.75 x 10 ⁻⁰³	6.76 x 10 ⁻⁰³	6.44 x 10 ⁻⁰³	5.64 x 10 ⁻⁰³	5.27 x 10 ⁻⁰²	3.26 x 10 ⁻⁰³	6.76 x 10 ⁻⁰³
	Beet Lake	6.44 × 10 ⁻⁰³	5.60 x 10 ⁻⁰³	5.26 x 10 ⁻⁰²	2.67 x 10 ⁻⁰³	6.59 x 10 ⁻⁰³	6.44 x 10 ⁻⁰³	5.61 x 10 ⁻⁰³	5.26 x 10 ⁻⁰²	2.93 x 10 ⁻⁰³	6.59 x 10 ⁻⁰³
	Lloyd Lake	6.44 × 10 ⁻⁰³	5.59 x 10 ⁻⁰³	5.26 x 10 ⁻⁰²	2.58 x 10 ⁻⁰³	6.35 x 10 ⁻⁰³	6.44 x 10 ⁻⁰³	5.59 x 10 ⁻⁰³	5.26 x 10 ⁻⁰²	2.61 x 10 ⁻⁰³	6.35 x 10 ⁻⁰³
	Reference (Broach Lake)	6.19 × 10 ⁻⁰⁴	5.56 × 10 ⁻⁰⁴	5.30 × 10 ⁻⁰³	2.58 × 10 ⁻⁰⁴	6.29 × 10 ⁻⁰⁴	6.19 × 10 ⁻⁰⁴	5.56 × 10 ⁻⁰⁴	5.30×10^{-03}	2.58×10^{-04}	6.29×10^{-04}
	Patterson Lake North Arm - West Basin	6.19 × 10 ⁻⁰⁴	5.90 x 10 ⁻⁰⁴	5.36 x 10 ⁻⁰³	4.27 x 10 ⁻⁰⁴	9.42 x 10 ⁻⁰⁴	6.20 x 10 ⁻⁰⁴	6.07 x 10 ⁻⁰⁴	5.39 x 10 ⁻⁰³	9.06 x 10 ⁻⁰⁴	9.42 x 10 ⁻⁰⁴
Canada Goose	Patterson Lake South Arm	6.19 × 10 ⁻⁰⁴	5.76 x 10 ⁻⁰⁴	5.33 x 10 ⁻⁰³	3.56 x 10 ⁻⁰⁴	6.78 x 10 ⁻⁰⁴	6.19 x 10 ⁻⁰⁴	5.87 x 10 ⁻⁰⁴	5.35 x 10 ⁻⁰³	6.34 x 10 ⁻⁰⁴	6.78 x 10 ⁻⁰⁴
	Beet Lake	6.19 × 10 ⁻⁰⁴	5.67 x 10 ⁻⁰⁴	5.32 x 10 ⁻⁰³	3.09 x 10 ⁻⁰⁴	6.56 x 10 ⁻⁰⁴	6.19 x 10 ⁻⁰⁴	5.72 x 10 ⁻⁰⁴	5.32 x 10 ⁻⁰³	4.54 x 10 ⁻⁰⁴	6.56 x 10 ⁻⁰⁴
	Lloyd Lake	6.19 × 10 ⁻⁰⁴	5.58 x 10 ⁻⁰⁴	5.30 x 10 ⁻⁰³	2.63 x 10 ⁻⁰⁴	6.30 x 10 ⁻⁰⁴	6.19 x 10 ⁻⁰⁴	5.58 x 10 ⁻⁰⁴	5.30 x 10 ⁻⁰³	2.78 x 10 ⁻⁰⁴	6.30 x 10 ⁻⁰⁴
	Reference (Broach Lake)	2.77 × 10 ⁻⁰²	5.73 × 10 ⁻⁰²	3.22 × 10 ⁺⁰⁰	1.64 × 10 ⁻⁰²	4.77 × 10 ⁻⁰³	2.77 × 10 ⁻⁰²	5.73 × 10 ⁻⁰²	3.22 × 10 ⁺⁰⁰	1.64 × 10 ⁻⁰²	4.77 × 10 ⁻⁰³
	Patterson Lake North Arm - West Basin	2.93 × 10 ⁻⁰²	1.40 × 10 ⁻⁰¹	8.33 × 10 ⁺⁰⁰	3.66 × 10 ⁻⁰¹	4.78 × 10 ⁻⁰²	3.02 × 10 ⁻⁰²	1.84 × 10 ⁻⁰¹	1.09 × 10 ⁺⁰¹	1.36 × 10 ⁺⁰⁰	4.78 × 10 ⁻⁰²
Little Brown Myotis	Patterson Lake South Arm	2.78 × 10 ⁻⁰²	1.08 × 10 ⁻⁰¹	6.22 × 10 ⁺⁰⁰	2.19 × 10 ⁻⁰¹	1.67 × 10 ⁻⁰²	2.79 × 10 ⁻⁰²	1.34 × 10 ⁻⁰¹	7.73 × 10 ⁺⁰⁰	7.95 × 10 ⁻⁰¹	1.67 × 10 ⁻⁰²
	Beet Lake	2.77 × 10 ⁻⁰²	8.39 × 10 ⁻⁰²	4.79 × 10 ⁺⁰⁰	1.22 × 10 ⁻⁰¹	9.19 × 10 ⁻⁰³	2.77 × 10 ⁻⁰²	9.81 × 10 ⁻⁰²	5.58 × 10 ⁺⁰⁰	4.22 × 10 ⁻⁰¹	9.20 × 10 ⁻⁰³
	Lloyd Lake	2.77 × 10 ⁻⁰²	6.00 × 10 ⁻⁰²	3.38 × 10 ⁺⁰⁰	2.72 × 10 ⁻⁰²	5.17 × 10 ⁻⁰³	2.77 × 10 ⁻⁰²	6.14 × 10 ⁻⁰²	3.46 × 10 ⁺⁰⁰	5.78 × 10 ⁻⁰²	5.17 × 10 ⁻⁰³
	Reference (Broach Lake)	4.71 × 10 ⁻⁰³	2.18 × 10 ⁻⁰³	1.25 × 10 ⁻⁰¹	4.10 × 10 ⁻⁰⁴	3.38 × 10 ⁻⁰⁴	4.71 × 10 ⁻⁰³	2.18 × 10 ⁻⁰³	1.25 × 10 ⁻⁰¹	4.10 × 10 ⁻⁰⁴	3.38 × 10 ⁻⁰⁴
	Patterson Lake North Arm – West Basin	5.00 × 10 ⁻⁰³	5.34 × 10 ⁻⁰³	3.22 × 10 ⁻⁰¹	9.15 × 10 ⁻⁰³	3.38 × 10 ⁻⁰³	5.15 × 10 ⁻⁰³	7.01 × 10 ⁻⁰³	4.22 × 10 ⁻⁰¹	3.40 × 10 ⁻⁰²	3.38 × 10 ⁻⁰³
Common Loon	Patterson Lake South Arm	4.74 × 10 ⁻⁰³	4.09 × 10 ⁻⁰³	2.41 × 10 ⁻⁰¹	5.48 × 10 ⁻⁰³	1.18 × 10 ⁻⁰³	4.75 × 10 ⁻⁰³	5.11 × 10 ⁻⁰³	2.99 × 10 ⁻⁰¹	1.99 × 10 ⁻⁰²	1.18 × 10 ⁻⁰³
	Beet Lake	4.72 × 10 ⁻⁰³	3.19 × 10 ⁻⁰³	1.85 × 10 ⁻⁰¹	3.05 × 10 ⁻⁰³	6.51 × 10 ⁻⁰⁴	4.72 × 10 ⁻⁰³	3.73 × 10 ⁻⁰³	2.16 × 10 ⁻⁰¹	1.05 × 10 ⁻⁰²	6.51 × 10 ⁻⁰⁴
	Lloyd Lake	4.71 × 10 ⁻⁰³	2.28 × 10 ⁻⁰³	1.31 × 10 ⁻⁰¹	6.79 × 10 ⁻⁰⁴	3.66 × 10 ⁻⁰⁴	4.71 × 10 ⁻⁰³	2.34 × 10 ⁻⁰³	1.34 × 10 ⁻⁰¹	1.44 × 10 ⁻⁰³	3.66 × 10 ⁻⁰⁴
	Reference (Broach Lake)	2.78 × 10 ⁻⁰²	2.60 × 10 ⁻⁰²	1.05 × 10 ⁺⁰⁰	6.52 × 10 ⁻⁰³	5.23 × 10 ⁻⁰³	2.78 × 10 ⁻⁰²	2.60 × 10 ⁻⁰²	1.05 × 10 ⁺⁰⁰	6.52 × 10 ⁻⁰³	5.23 × 10 ⁻⁰³
	Patterson Lake North Arm – West Basin	2.94 × 10 ⁻⁰²	6.35 × 10 ⁻⁰²	2.73 × 10 ⁺⁰⁰	1.46 × 10 ⁻⁰¹	5.23 × 10 ⁻⁰²	3.03 × 10 ⁻⁰²	8.35 × 10 ⁻⁰²	3.57 × 10 ⁺⁰⁰	5.40 × 10 ⁻⁰¹	5.23 × 10 ⁻⁰²
Mallard	Patterson Lake South Arm	2.79 × 10 ⁻⁰²	4.88 × 10 ⁻⁰²	2.04 × 10 ⁺⁰⁰	8.72 × 10 ⁻⁰²	1.82 × 10 ⁻⁰²	2.80 × 10 ⁻⁰²	6.09 × 10 ⁻⁰²	2.53 × 10 ⁺⁰⁰	3.16 × 10 ⁻⁰¹	1.83 × 10 ⁻⁰²
	Beet Lake	2.78 × 10 ⁻⁰²	3.80 × 10 ⁻⁰²	1.57 × 10 ⁺⁰⁰	4.85 × 10 ⁻⁰²	1.01 × 10 ⁻⁰²	2.78 × 10 ⁻⁰²	4.44 × 10 ⁻⁰²	1.83 × 10 ⁺⁰⁰	1.68 × 10 ⁻⁰¹	1.01 × 10 ⁻⁰²
	Lloyd Lake	2.78 × 10 ⁻⁰²	2.72 × 10 ⁻⁰²	1.11 × 10 ⁺⁰⁰	1.08 × 10 ⁻⁰²	5.66 × 10 ⁻⁰³	2.78 × 10 ⁻⁰²	2.78 × 10 ⁻⁰²	1.13 × 10 ⁺⁰⁰	2.30 × 10 ⁻⁰²	5.66 × 10 ⁻⁰³
	Reference (Broach Lake)	7.16 × 10 ⁻⁰²	1.25 × 10 ⁻⁰¹	6.35 × 10 ⁺⁰⁰	3.66 × 10 ⁻⁰²	1.76 × 10 ⁻⁰²	7.16 × 10 ⁻⁰²	1.25 × 10 ⁻⁰¹	6.35 × 10 ⁺⁰⁰	3.66 × 10 ⁻⁰²	1.76 × 10 ⁻⁰²
	Patterson Lake North Arm – West Basin	7.49 × 10 ⁻⁰²	2.88 x 10 ⁻⁰¹	1.64 x 10 ⁺⁰¹	7.22 x 10 ⁻⁰¹	1.05 x 10 ⁻⁰¹	7.67 x 10 ⁻⁰²	3.75 x 10 ⁻⁰¹	2.14 x 10 ⁺⁰¹	2.67 x 10 ⁺⁰⁰	1.06 x 10 ⁻⁰¹
Rusty Blackbird	Patterson Lake South Arm	7.19 × 10 ⁻⁰²	2.24 x 10 ⁻⁰¹	1.22 x 10 ⁺⁰¹	4.35 x 10 ⁻⁰¹	4.14 x 10 ⁻⁰²	7.20 x 10 ⁻⁰²	2.77 x 10 ⁻⁰¹	1.52 x 10 ⁺⁰¹	1.56 x 10 ⁺⁰⁰	4.14 x 10 ⁻⁰²
	Beet Lake	7.17 × 10 ⁻⁰²	1.78 x 10 ⁻⁰¹	9.43 x 10 ⁺⁰⁰	2.44 x 10 ⁻⁰¹	2.65 x 10 ⁻⁰²	7.17 x 10 ⁻⁰²	2.05 x 10 ⁻⁰¹	1.10 x 10 ⁺⁰¹	8.32 x 10 ⁻⁰¹	2.65 x 10 ⁻⁰²
	Lloyd Lake	7.16 × 10 ⁻⁰²	1.31 x 10 ⁻⁰¹	6.67 x 10 ⁺⁰⁰	5.77 x 10 ⁻⁰²	1.84 x 10 ⁻⁰²	7.16 x 10 ⁻⁰²	1.33 x 10 ⁻⁰¹	6.82 x 10 ⁺⁰⁰	1.18 x 10 ⁻⁰¹	1.84 x 10 ⁻⁰²

6.2.5.1.2 Radiological Dose

The estimated radiation doses to aquatic and terrestrial ecological receptors during the Project phases and far-future projection resulting from the Application Case and Upper Bound sensitivity scenario are shown in Table 6-12. The doses shown represent the maximum total dose from all radionuclides over the assessment period. The dose breakdown by radionuclide is shown in Appendix C.

The maximum estimated radiation dose would occur during Operations for zooplankton in the Patterson Lake North Arm – West Basin. The estimated dose to zooplankton was 0.12 milligrays per day (mGy/d) in the Application Case and 0.26 mGy/d in the reasonable upper bound sensitivity scenario. The main contributor to total dose would be from polonium-210 in water.

For terrestrial receptors, the maximum estimated radiation dose for mammals and birds occurred during Operations for the rusty blackbird eating from the Patterson Lake North Arm – West Basin. The estimated dose to the rusty blackbird was 0.10 mGy/d in the Application Case and 0.19 mGy/d in the reasonable upper bound sensitivity scenario. The main contributor to total dose would be from polonium-210 in benthic invertebrates that are eaten by the rusty blackbird. The maximum dose for terrestrial plants is to lichen (0.68 mGy/d in the Application Case and the reasonable upper bound sensitivity scenario) due to atmospheric deposition from air to the plant.

The predicted radiation dose to all ecological receptors during the far-future projection was lower than during Operations. The maximum estimated dose to zooplankton was 0.07 mGy/d in both the Application Case and reasonable upper bound sensitivity scenario, in the far-future projection. The maximum estimated dose to lichen was 0.11 mGy/d in both the Application Case and reasonable upper bound sensitivity scenario in the far-future projection.



Table 6-12: Estimated Radiation Doses to Aquatic and Terrestrial Biota – All Cases

	Biota	Location	Maximum Total Dose (mGy/d)					
			Applicat	Upper Bou	pper Bound Scenario		RFD Case	
			Operations	Far-Future	Operations	Far-Future	Operations	Far-Future
Aquatic Animals	Benthic Invertebrate	Reference (Broach Lake)	1.19×10^{-04}	1.19×10^{-04}	1.19×10^{-04}	1.19 × 10 ⁻⁰⁴	1.19 × 10 ⁻⁰⁴	1.19×10^{-04}
		Patterson Lake North Arm – West Basin	1.82×10^{-04}	1.44×10^{-04}	2.36×10^{-04}	1.44 × 10 ⁻⁰⁴	1.82 × 10 ⁻⁰⁴	1.44×10^{-04}
		Patterson Lake South Arm	1.37×10^{-04}	1.28×10^{-04}	1.41×10^{-04}	1.28 × 10 ⁻⁰⁴	1.57 × 10 ⁻⁰⁴	1.28×10^{-04}
		Beet Lake	1.26×10^{-04}	1.22×10^{-04}	1.27×10^{-04}	1.22 × 10 ⁻⁰⁴	1.33 × 10 ⁻⁰⁴	1.22×10^{-04}
		Lloyd Lake	1.19×10^{-04}	1.19×10^{-04}	1.19×10^{-04}	1.19 × 10 ⁻⁰⁴	1.20×10^{-04}	1.19×10^{-04}
	Zooplankton	Reference (Broach Lake)	6.81×10^{-02}	6.81 × 10 ⁻⁰²	6.81 × 10 ⁻⁰²	6.81 × 10 ⁻⁰²	6.81 × 10 ⁻⁰²	6.81 × 10 ⁻⁰²
		Patterson Lake North Arm – West Basin	1.17×10^{-01}	7.38×10^{-02}	2.61 × 10 ⁻⁰¹	7.38 × 10 ⁻⁰²	1.17 × 10 ⁻⁰¹	7.38×10^{-02}
		Patterson Lake South Arm	7.93×10^{-02}	6.93×10^{-02}	8.74 × 10 ⁻⁰²	6.93 × 10 ⁻⁰²	1.21 × 10 ⁻⁰¹	6.93×10^{-02}
		Beet Lake	7.31×10^{-02}	6.85×10^{-02}	7.43×10^{-02}	6.85 × 10 ⁻⁰²	8.16 × 10 ⁻⁰²	6.85 × 10 ⁻⁰²
		Lloyd Lake	6.86×10^{-02}	6.81 × 10 ⁻⁰²	6.87×10^{-02}	6.81 × 10 ⁻⁰²	6.92 × 10 ⁻⁰²	6.81 × 10 ⁻⁰²
	Northern Pike	Reference (Broach Lake)	8.33×10^{-04}	8.33 × 10 ⁻⁰⁴	8.33 × 10 ⁻⁰⁴	8.33 × 10 ⁻⁰⁴	8.33 × 10 ⁻⁰⁴	8.33 × 10 ⁻⁰⁴
		Patterson Lake North Arm – West Basin	1.54 × 10 ⁻⁰³	1.20×10^{-03}	3.05×10^{-03}	1.20 × 10 ⁻⁰³	1.54 × 10 ⁻⁰³	1.20×10^{-03}
		Patterson Lake South Arm	1.03×10^{-03}	9.36 × 10 ⁻⁰⁴	1.15×10^{-03}	9.36 × 10 ⁻⁰⁴	1.68 × 10 ⁻⁰³	9.36 × 10 ⁻⁰⁴
		Beet Lake	9.20×10^{-04}	8.72 × 10 ⁻⁰⁴	9.46 × 10 ⁻⁰⁴	8.72 × 10 ⁻⁰⁴	1.09 × 10 ⁻⁰³	8.72×10^{-04}
		Lloyd Lake	8.41 × 10 ⁻⁰⁴	8.36 × 10 ⁻⁰⁴	8.43 × 10 ⁻⁰⁴	8.36 × 10 ⁻⁰⁴	8.56 × 10 ⁻⁰⁴	8.36 × 10 ⁻⁰⁴
	Whitefish	Reference (Broach Lake)	2.58×10^{-04}	2.58×10^{-04}	2.58×10^{-04}	2.58 × 10 ⁻⁰⁴	2.58×10^{-04}	2.58×10^{-04}
		Patterson Lake North Arm – West Basin	5.57×10^{-04}	3.68×10^{-04}	6.99×10^{-04}	3.68 × 10 ⁻⁰⁴	5.57 × 10 ⁻⁰⁴	3.68×10^{-04}
		Patterson Lake South Arm	3.84×10^{-04}	2.93×10^{-04}	4.01×10^{-04}	2.93 × 10 ⁻⁰⁴	6.24 × 10 ⁻⁰⁴	2.93 × 10 ⁻⁰⁴
		Beet Lake	3.18×10^{-04}	2.72×10^{-04}	3.23×10^{-04}	2.72×10^{-04}	4.01×10^{-04}	2.72×10^{-04}
		Lloyd Lake	2.64×10^{-04}	2.59×10^{-04}	2.64×10^{-04}	2.59 × 10 ⁻⁰⁴	2.72×10^{-04}	2.59×10^{-04}
Aquatic Plants	Macrophytes	Reference (Broach Lake)	5.53×10^{-03}	5.53×10^{-03}	5.53×10^{-03}	5.53 × 10 ⁻⁰³	5.53 × 10 ⁻⁰³	5.53 × 10 ⁻⁰³
		Patterson Lake North Arm – West Basin	1.50×10^{-02}	7.25×10^{-03}	1.83×10^{-02}	7.25×10^{-03}	1.50×10^{-02}	7.25×10^{-03}
		Patterson Lake South Arm	1.02×10^{-02}	6.11×10^{-03}	1.05×10^{-02}	6.11 × 10 ⁻⁰³	1.65 × 10 ⁻⁰²	6.11 × 10 ⁻⁰³
		Beet Lake	7.88×10^{-03}	5.77×10^{-03}	7.97×10^{-03}	5.77×10^{-03}	1.01 × 10 ⁻⁰²	5.77×10^{-03}
		Lloyd Lake	5.77×10^{-03}	5.56 × 10 ⁻⁰³	5.78×10^{-03}	5.56 × 10 ⁻⁰³	5.98 × 10 ⁻⁰³	5.56×10^{-03}
	Phytoplankton	Reference (Broach Lake)	5.51×10^{-03}	5.51×10^{-03}	5.51×10^{-03}	5.51 × 10 ⁻⁰³	5.51 × 10 ⁻⁰³	5.51×10^{-03}
		Patterson Lake North Arm – West Basin	1.49×10^{-02}	7.22×10^{-03}	1.82×10^{-02}	7.22×10^{-03}	1.49 × 10 ⁻⁰²	7.22×10^{-03}
		Patterson Lake South Arm	1.01×10^{-02}	6.08×10^{-03}	1.05×10^{-02}	6.08×10^{-03}	1.65 × 10 ⁻⁰²	6.08×10^{-03}
		Beet Lake	7.84×10^{-03}	5.75×10^{-03}	7.94×10^{-03}	5.75×10^{-03}	1.00×10^{-02}	5.75×10^{-03}
		Lloyd Lake	5.75 × 10 ⁻⁰³	5.53 × 10 ⁻⁰³	5.75×10^{-03}	5.53 × 10 ⁻⁰³	5.95 × 10 ⁻⁰³	5.53 × 10 ⁻⁰³
Terrestrial Invertebrate	Terrestrial Invertebrate	Reference (Broach Lake)	5.91 × 10 ⁻⁰³	5.91 × 10 ⁻⁰³	5.91 × 10 ⁻⁰³	5.91 × 10 ⁻⁰³	5.91 × 10 ⁻⁰³	5.91 × 10 ⁻⁰³
		Patterson Lake North Arm – West Basin	2.16 x 10 ⁻⁰²	8.49 x 10 ⁻⁰³	2.16 x 10 ⁻⁰²	8.49 x 10 ⁻⁰³	2.19 x 10 ⁻⁰²	8.54 x 10 ⁻⁰³
		Patterson Lake South Arm	8.47 x 10 ⁻⁰³	6.33 x 10 ⁻⁰³	8.47 x 10 ⁻⁰³	6.33 x 10 ⁻⁰³	8.99 x 10 ⁻⁰³	6.42 x 10 ⁻⁰³
		Beet Lake	7.40 x 10 ⁻⁰³	6.16 x 10 ⁻⁰³	7.40 x 10 ⁻⁰³	6.16 x 10 ⁻⁰³	7.59 x 10 ⁻⁰³	6.19 x 10 ⁻⁰³
		Lloyd Lake	5.96 x 10 ⁻⁰³	5.92 x 10 ⁻⁰³	5.96 x 10 ⁻⁰³	5.92 x 10 ⁻⁰³	5.99 x 10 ⁻⁰³	5.93 x 10 ⁻⁰³



					Maximum Tota	I Dose (mGy/c	l)				
	Biota	Location	Applicat	tion Case	Upper Bou	nd Scenario	RFD	Case			
			Operations	Far-Future	Operations	Far-Future	Operations	Far-Future			
		Reference (Broach Lake)	8.77 x 10 ⁻⁰⁴	8.77 x 10 ⁻⁰⁴	8.77 x 10 ⁻⁰⁴	8.77 x 10 ⁻⁰⁴	8.77 x 10 ⁻⁰⁴	8.77 x 10 ⁻⁰⁴			
		Patterson Lake North Arm – West Basin	2.59 x 10 ⁻⁰³	1.17 x 10 ⁻⁰³	2.70 x 10 ⁻⁰³	1.17 x 10 ⁻⁰³	Operations 8.77 x 10 ⁻⁰⁴ 2.62 x 10 ⁻⁰³ 1.27 x 10 ⁻⁰³ 1.07 x 10 ⁻⁰³ 1.07 x 10 ⁻⁰³ 1.68 x 10 ⁻⁰³ 1.68 x 10 ⁻⁰³ 1.68 x 10 ⁻⁰³ 1.02 x 10 ⁻⁰³ 1.02 x 10 ⁻⁰³ 1.09 x 10 ⁻⁰³ 1.29 x 10 ⁻⁰³ 1.10 x 10 ⁻⁰³ 9.40 x 10 ⁻⁰⁴ 1.32 x 10 ⁻⁰³ 1.17 x 10 ⁻⁰³ 9.82 x 10 ⁻⁰⁴ 1.25 x 10 ⁻⁰³ 3.12 x 10 ⁻⁰³ 3.12 x 10 ⁻⁰³ 2.90 x 10 ⁻⁰³ 1.27 x 10 ⁻⁰³ 2.90 x 10 ⁻⁰³ 1.27 x 10 ⁻⁰³ 2.55 x 10 ⁻⁰³ 2.12 x 10 ⁻⁰³ 2.12 x 10 ⁻⁰³ 2.12 x 10 ⁻⁰³ 2.12 x 10 ⁻⁰³ 2.11 x 10 ⁻⁰³ 2.12 x 10 ⁻⁰³ 2.11 x 10 ⁻⁰⁴ 1.36 x 10 ⁻⁰⁴ 1.36 x 10 ⁻⁰⁴ 7.94 x 10 ⁻⁰⁴ 7.94 x 10 ⁻⁰⁴ 7.94 x 10 ⁻⁰⁴ 7.94 x 10 ⁻⁰⁴ 1.44 x 10 ⁻⁰³ 3.45 x 10 ⁻⁰³ 1.65 x 1	1.17 x 10 ⁻⁰³			
	Beaver	Patterson Lake South Arm	1.16 x 10 ⁻⁰³	9.27 x 10 ⁻⁰⁴	1.17 x 10 ⁻⁰³	9.27 x 10 ⁻⁰⁴	1.27 x 10 ⁻⁰³	9.36 x 10 ⁻⁰⁴			
		Beet Lake	1.04 x 10 ⁻⁰³	9.05 x 10 ⁻⁰⁴	1.04 x 10 ⁻⁰³	9.05 x 10 ⁻⁰⁴	1.07 x 10 ⁻⁰³	9.08 x 10 ⁻⁰⁴			
		Lloyd Lake	8.83 x 10 ⁻⁰⁴	8.78 x 10 ⁻⁰⁴	8.83 x 10 ⁻⁰⁴	8.78 x 10 ⁻⁰⁴	8.87 x 10 ⁻⁰⁴	8.79 x 10 ⁻⁰⁴			
		Reference (Broach Lake)	1.01 × 10 ⁻⁰³	1.01×10^{-03}	1.01×10^{-03}	1.01×10^{-03}	1.01×10^{-03}	1.01 × 10 ⁻⁰³			
	Black Bear	Patterson Lake North Arm – West Basin	1.64 x 10 ⁻⁰³	1.21 x 10 ⁻⁰³	1.70 x 10 ⁻⁰³	1.21 x 10 ⁻⁰³	1.07 x 10 ⁻⁰³ 9.08 x 1.07 x 10 ⁻⁰³ 1.01 x 10 ⁻⁰³ 1.01 x 10 ⁻⁰³ 1.01 x 10 ⁻⁰³ 1.22 x 1.31 x 10 ⁻⁰³ 1.26 x 1.02 x 10 ⁻⁰³ 1.02 x 10 ⁻⁰³ 1.02 x 10.02 x 10 ⁻⁰³ 1.09 x 10.09 x 10.	1.22 x 10 ⁻⁰³			
	ыаск веаг	Beet Lake	1.30 x 10 ⁻⁰³	1.26 x 10 ⁻⁰³	1.30 x 10 ⁻⁰³	1.26 x 10 ⁻⁰³		1.26 x 10 ⁻⁰³			
		Lloyd Lake	1.02 x 10 ⁻⁰³	1.02 x 10 ⁻⁰³	1.02 x 10 ⁻⁰³	1.02 x 10 ⁻⁰³	1.02 x 10 ⁻⁰³	1.02 x 10 ⁻⁰³			
		Reference (Broach Lake)	1.09 x 10 ⁻⁰³	1.09 x 10 ⁻⁰³	1.09 x 10 ⁻⁰³	1.09 x 10 ⁻⁰³	1.09 x 10 ⁻⁰³	x 10-04 8.77 x 10-04 x 10-03 1.17 x 10-03 x 10-03 9.36 x 10-04 x 10-04 8.79 x 10-04 x 10-03 1.01 x 10-03 x 10-03 1.22 x 10-03 x 10-03 1.26 x 10-03 x 10-03 1.02 x 10-03 x 10-03 1.09 x 10-03 x 10-03 1.09 x 10-03 x 10-03 1.09 x 10-03 x 10-03 1.09 x 10-03 x 10-03 1.01 x 10-03 x 10-03 1.01 x 10-03 x 10-04 9.40 x 10-04 x 10-03 9.51 x 10-04 x 10-04 9.40 x 10-04 x 10-03 1.25 x 10-03 x 10-03 1.25 x 10-03 x 10-03 1.56 x 10-03 x 10-03 1.26 x 10-03 x 10-03 1.26 x 10-03 x 10-03 2.00 x 10-03 x 10-03 2.00 x 10-03 x 10-03 2.01 x 10-03 x 10-03 2.01 x 10-03 x 10-03 2.02 x 10-03 x 10-03 2.01 x 10-03 x 10-03 2.02 x 10-03			
	Grey Wolf	Patterson Lake North Arm – West Basin	1.28 x 10 ⁻⁰³	1.18 x 10 ⁻⁰³	1.29 x 10 ⁻⁰³	1.18 x 10 ⁻⁰³	1.29 x 10 ⁻⁰³	1.18 x 10 ⁻⁰³			
		Lloyd Lake	1.10 x 10 ⁻⁰³	1.09 x 10 ⁻⁰³	1.10 x 10 ⁻⁰³	1.09 x 10 ⁻⁰³	1.10 x 10 ⁻⁰³	1.09 x 10 ⁻⁰³			
		Reference (Broach Lake)	9.40 x 10 ⁻⁰⁴	9.40 x 10 ⁻⁰⁴	9.40 x 10 ⁻⁰⁴	9.40 x 10 ⁻⁰⁴	9.40 x 10 ⁻⁰⁴	$\begin{array}{cccccccccccccccccccccccccccccccccccc$			
Terrestrial Animals		Patterson Lake North Arm – West Basin	1.32 x 10 ⁻⁰³	1.01 x 10 ⁻⁰³	2.51 x 10 ⁻⁰³	1.01 x 10 ⁻⁰³	1.32 x 10 ⁻⁰³	1.01 x 10 ⁻⁰³			
Animais	American Mink	Patterson Lake South Arm	9.80 x 10 ⁻⁰⁴	9.51 x 10 ⁻⁰⁴	1.06 x 10 ⁻⁰³	9.51 x 10 ⁻⁰⁴	0-03 2.62 x 10 ⁻⁰³ 1.17 x 10 ⁻⁰³ 0-04 1.27 x 10 ⁻⁰³ 9.36 x 10 ⁻⁰⁴ 0-04 1.07 x 10 ⁻⁰³ 9.08 x 10 ⁻⁰⁴ 0-04 8.87 x 10 ⁻⁰⁴ 8.79 x 10 ⁻⁰³ 0-03 1.01 x 10 ⁻⁰³ 1.01 x 10 ⁻⁰³ 0-03 1.68 x 10 ⁻⁰³ 1.22 x 10 ⁻⁰³ 0-03 1.02 x 10 ⁻⁰³ 1.02 x 10 ⁻⁰³ 0-03 1.09 x 10 ⁻⁰³ 1.09 x 10 ⁻⁰³ 0-03 1.29 x 10 ⁻⁰³ 1.09 x 10 ⁻⁰³ 0-03 1.10 x 10 ⁻⁰³ 1.09 x 10 ⁻⁰³ 0-04 9.40 x 10 ⁻⁰⁴ 9.40 x 10 ⁻⁰⁴ 0-04 9.40 x 10 ⁻⁰⁴ 9.44 x 10 ⁻⁰⁴ 0-04 9.43 x 10 ⁻⁰⁴ 9.44 x 10 ⁻⁰⁴ 0-04 9.43 x 10 ⁻⁰⁴ 9.40 x 10 ⁻⁰³ 0-03 1.25 x 10 ⁻⁰³ 1.25 x 10 ⁻⁰³ 0-03 1.25 x 10 ⁻⁰³ 1.25 x 10 ⁻⁰³ 0-03 1.29 x 10 ⁻⁰³ 1.26 x 10 ⁻⁰³ 0-03 1.29 x 10 ⁻⁰³ 1.26 x 10 ⁻⁰³ 0-03 1.29 x 10 ⁻⁰³ 1.26 x 10 ⁻⁰³ 0-03 1.29 x	9.51 x 10 ⁻⁰⁴			
		Beet Lake	9.52 x 10 ⁻⁰⁴	9.44 x 10 ⁻⁰⁴	9.63 x 10 ⁻⁰⁴	9.44 x 10 ⁻⁰⁴	9.82 x 10 ⁻⁰⁴	9.44 x 10 ⁻⁰⁴			
		Lloyd Lake	9.41 x 10 ⁻⁰⁴	9.40 x 10 ⁻⁰⁴	9.42 x 10 ⁻⁰⁴	9.40 x 10 ⁻⁰⁴	9.43 x 10 ⁻⁰⁴	8.77 × 10 ⁻⁰ · 1.17 × 10 ⁻⁰ · 9.36 × 10 ⁻⁰ · 9.08 × 10 ⁻⁰ · 1.01 × 10 ⁻⁰ · 1.22 × 10 ⁻⁰ · 1.26 × 10 ⁻⁰ · 1.09 × 10 ⁻⁰ · 1.18 × 10 ⁻⁰ · 1.09 × 10 ⁻⁰ · 1.18 × 10 ⁻⁰ · 1.01 × 10 ⁻⁰ · 9.40 × 10 ⁻⁰ · 1.17 × 10 ⁻⁰ · 9.44 × 10 ⁻⁰ · 1.25 × 10 ⁻⁰ · 1.30 × 10 ⁻⁰ · 1.26 × 10 ⁻⁰ · 2.00 × 10 ⁻⁰ · 2.18 × 10 ⁻⁰ · 2.01 × 10 ⁻⁰ · 7.28 × 10 ⁻⁰ · 7.29 × 10 ⁻⁰ · 7.29 × 10 ⁻⁰ · 1.50 × 10 ⁻⁰ · 1.50 × 10 ⁻⁰ · 1.55 × 10 ⁻⁰ · 1.50 × 10 ⁻⁰ ·			
		Reference (Broach Lake)	1.25 x 10 ⁻⁰³	1.25 x 10 ⁻⁰³	1.25 x 10 ⁻⁰³	1.25 x 10 ⁻⁰³	1.25 x 10 ⁻⁰³	1.25 x 10 ⁻⁰³			
		Patterson Lake North Arm – West Basin	3.09 x 10 ⁻⁰³	1.69 x 10 ⁻⁰³	3.33 x 10 ⁻⁰³	1.69 x 10 ⁻⁰³	9.82 x 10 ⁻⁰⁴ 9.4 9.43 x 10 ⁻⁰⁴ 9.4 1.25 x 10 ⁻⁰³ 1.2 3.3 3.12 x 10 ⁻⁰³ 1.7 3.3 2.90 x 10 ⁻⁰³ 1.5 3.4 1.48 x 10 ⁻⁰³ 1.3 1.27 x 10 ⁻⁰³ 1.2	1.70 x 10 ⁻⁰³			
	Moose	Patterson Lake South Arm	2.87 x 10 ⁻⁰³	1.55 x 10 ⁻⁰³	3.11 x 10 ⁻⁰³	1.55 x 10 ⁻⁰³		1.56 x 10 ⁻⁰³			
		Beet Lake	1.43 x 10 ⁻⁰³	1.30 x 10 ⁻⁰³	1.43 x 10 ⁻⁰³	1.30 x 10 ⁻⁰³	1.48 x 10 ⁻⁰³	10 ⁻⁰³ 1.70 x 10 ⁻⁰ 10 ⁻⁰³ 1.56 x 10 ⁻⁰ 10 ⁻⁰³ 1.30 x 10 ⁻⁰			
		Lloyd Lake	1.26 x 10 ⁻⁰³	1.26 x 10 ⁻⁰³	1.26 x 10 ⁻⁰³	1.26 x 10 ⁻⁰³	1.27 x 10 ⁻⁰³	1.26 x 10 ⁻⁰³			
		Reference (Broach Lake)	2.00 × 10 ⁻⁰³	2.00×10^{-03}	2.00×10^{-03}	2.00×10^{-03}	2.00×10^{-03}	2.00×10^{-03}			
		Patterson Lake North Arm – West Basin	2.84 × 10 ⁻⁰³	2.18×10^{-03}	5.33×10^{-03}	2.18×10^{-03}	2.84×10^{-03}	2.18×10^{-03}			
	Muskrat	Patterson Lake South Arm	2.11 × 10 ⁻⁰³	2.04×10^{-03}	2.28×10^{-03}	2.04×10^{-03}	1.07 x 10 ⁻⁰³ 9.08 x 10 8.87 x 10 ⁻⁰⁴ 8.79 x 10 1.01 x 10 ⁻⁰³ 1.01 x 10 1.68 x 10 ⁻⁰³ 1.22 x 10 1.31 x 10 ⁻⁰³ 1.26 x 10 1.02 x 10 ⁻⁰³ 1.02 x 10 1.09 x 10 ⁻⁰³ 1.09 x 10 1.29 x 10 ⁻⁰³ 1.18 x 10 1.10 x 10 ⁻⁰³ 1.09 x 10 9.40 x 10 ⁻⁰⁴ 9.40 x 10 1.32 x 10 ⁻⁰³ 1.01 x 10 1.17 x 10 ⁻⁰³ 9.51 x 10 9.82 x 10 ⁻⁰⁴ 9.44 x 10 9.43 x 10 ⁻⁰⁴ 9.40 x 10 1.25 x 10 ⁻⁰³ 1.25 x 10 3.12 x 10 ⁻⁰³ 1.56 x 10 1.27 x 10 ⁻⁰³ 1.56 x 10 1.27 x 10 ⁻⁰³ 2.00 x 10 2.90 x 10 ⁻⁰³ 1.26 x 10 2.90 x 10 ⁻⁰³ 1.26 x 10 2.12 x 10 ⁻⁰³ 2.00 x 10 2.84 x 10 ⁻⁰³ 2.00 x 10 2.12 x 10 ⁻⁰³ 2.00 x 10 2.12 x 10 ⁻⁰³ 2.00 x 10 2.12 x 10 ⁻⁰³ 1.05 x 10 2.14 x 10 ⁻⁰³ 1.05 x 10 1.36 x 10 ⁻⁰³ 1.05 x 10 1.36 x 10 ⁻⁰⁴ 7.28 x 10 1.36 x 10 ⁻⁰⁴ 7.29 x 10 1.44 x 10 ⁻⁰³ 1.44 x 10 3.45 x 10 ⁻⁰³ 1.55 x 10 1.82 x 10 ⁻⁰³ 1.50 x 10 1.82 x 10 ⁻⁰³ 1.55 x 10 1.82 x 10 ⁻⁰³ 1.55 x 10 1.85 x 10 ⁻⁰³ 1.55 x 10 1.85 x 10 ⁻⁰³ 1.55 x 10 1.85 x 10 ⁻⁰³ 1.55 x 10 1.85 x 10 ⁻⁰³ 1.55 x 10 1.85 x 10 ⁻⁰³ 1.55 x 10 1.85 x 10 ⁻⁰³ 1.55 x 10 1.85 x 10 ⁻⁰³ 1.55 x 10 1.85 x 10 ⁻⁰³ 1.55 x 10 1.85 x 10 ⁻⁰³ 1.50 x 10	2.04×10^{-03}			
		Beet Lake	2.04 × 10 ⁻⁰³	2.01×10^{-03}	2.06×10^{-03}	2.01×10^{-03}		2.01×10^{-03}			
		Lloyd Lake	2.00 × 10 ⁻⁰³	2.00×10^{-03}	2.00×10^{-03}	2.00×10^{-03}	2.01 × 10 ⁻⁰³	9.44 x 10 ⁻⁰⁴ 9.40 x 10 ⁻⁰³ 1.25 x 10 ⁻⁰³ 1.70 x 10 ⁻⁰³ 1.56 x 10 ⁻⁰³ 1.30 x 10 ⁻⁰³ 1.26 x 10 ⁻⁰³ 2.00 x 10 ⁻⁰³ 2.18 x 10 ⁻⁰³ 2.01 x 10 ⁻⁰³ 2.00 x 10 ⁻⁰³ 7.28 x 10 ⁻⁰⁴ 7.61 x 10 ⁻⁰⁴			
		Reference (Broach Lake)	7.28 × 10 ⁻⁰⁴	7.28×10^{-04}	7.28×10^{-04}	7.28×10^{-04}	7.28×10^{-04}	7.28 × 10 ⁻⁰⁴			
		Patterson Lake North Arm – West Basin	1.34 x 10 ⁻⁰³	1.04 x 10 ⁻⁰³	1.34 x 10 ⁻⁰³	1.04 x 10 ⁻⁰³	RFD Case Operations Far-8.77 x 10 ⁻⁰⁴ 8.77 8.77 x 10 ⁻⁰⁴ 8.77 2.62 x 10 ⁻⁰³ 1.17 1.27 x 10 ⁻⁰³ 9.36 1.07 x 10 ⁻⁰³ 9.08 8.87 x 10 ⁻⁰⁴ 8.79 1.01 x 10 ⁻⁰³ 1.01 1.68 x 10 ⁻⁰³ 1.22 1.31 x 10 ⁻⁰³ 1.02 1.02 x 10 ⁻⁰³ 1.09 1.09 1.09 1.29 x 10 ⁻⁰³ 1.09 1.09 1.09 1.29 x 10 ⁻⁰³ 1.01 1.17 x 10 ⁻⁰³ 9.40 1.32 x 10 ⁻⁰³ 1.01 1.17 x 10 ⁻⁰³ 9.51 9.82 x 10 ⁻⁰⁴ 9.40 1.25 x 10 ⁻⁰³ 1.25 3.12 x 10 ⁻⁰³ 1.25 3.12 x 10 ⁻⁰³ 1.26 1.25 x 10 ⁻⁰³ 1.25 3.12 x 10 ⁻⁰³ 1.26 2.90 x 10 ⁻⁰³ 1.26 2.00 x 10 ⁻⁰³ 1.26 2.00 x 10 ⁻⁰³ 1.26 2.00 x 10 ⁻⁰³ 2.00 2.84 x 10 ⁻⁰³ 1.26 2.00 x 10 ⁻⁰³ 2.00 2.12 x 10 ⁻⁰³ 2.01 2.01 2.01 2.01 <td>1.05 x 10⁻⁰³</td>	1.05 x 10 ⁻⁰³			
	Red Fox	Patterson Lake South Arm	8.28 x 10 ⁻⁰⁴	7.79 x 10 ⁻⁰⁴	8.28 x 10 ⁻⁰⁴	7.79 x 10 ⁻⁰⁴	8.49 x 10 ⁻⁰⁴	7.89 x 10 ⁻⁰⁴			
		Beet Lake	7.86 x 10 ⁻⁰⁴	7.57 x 10 ⁻⁰⁴	7.86 x 10 ⁻⁰⁴	7.57 x 10 ⁻⁰⁴	7.94 x 10 ⁻⁰⁴	9.36 x 10 ⁻⁰² 9.08 x 10 ⁻⁰³ 8.79 x 10 ⁻⁰³ 1.01 x 10 ⁻⁰³ 1.22 x 10 ⁻⁰³ 1.26 x 10 ⁻⁰³ 1.02 x 10 ⁻⁰³ 1.09 x 10 ⁻⁰³ 1.18 x 10 ⁻⁰³ 1.09 x 10 ⁻⁰³ 9.40 x 10 ⁻⁰⁴ 9.40 x 10 ⁻⁰⁴ 9.40 x 10 ⁻⁰⁴ 1.25 x 10 ⁻⁰³ 1.70 x 10 ⁻⁰³ 1.30 x 10 ⁻⁰³ 1.26 x 10 ⁻⁰³ 2.00 x 10 ⁻⁰⁴ 2.01 x 10 ⁻⁰⁴ 7.28 x 10 ⁻⁰⁶ 7.89 x 10 ⁻⁰⁶ 7.44 x 10 ⁻⁰⁶ 7.29 x 10 ⁻⁰⁶ 1.55 x 10 ⁻⁰³ 1.50 x 10 ⁻⁰³ 1.50 x 10 ⁻⁰³			
		Lloyd Lake	7.30 x 10 ⁻⁰⁴	7.29 x 10 ⁻⁰⁴	7.30 x 10 ⁻⁰⁴	7.29 x 10 ⁻⁰⁴	7.31 x 10 ⁻⁰⁴	7.29 x 10 ⁻⁰⁴			
		Reference (Broach Lake)	1.44 × 10 ⁻⁰³	1.44 × 10 ⁻⁰³	1.44 × 10 ⁻⁰³	1.44 × 10 ⁻⁰³	1.44 × 10 ⁻⁰³	8.77 x 10 ⁻⁰ · 1.17 x 10 ⁻⁰ · 9.36 x 10 ⁻⁰ · 9.08 x 10 ⁻⁰ · 1.01 x 10 ⁻⁰ · 1.22 x 10 ⁻⁰ · 1.26 x 10 ⁻⁰ · 1.09 x 10 ⁻⁰ · 1.18 x 10 ⁻⁰ · 1.18 x 10 ⁻⁰ · 1.01 x 10 ⁻⁰ · 9.40 x 10 ⁻⁰ · 9.40 x 10 ⁻⁰ · 1.25 x 10 ⁻⁰ · 1.70 x 10 ⁻⁰ · 1.26 x 10 ⁻⁰ · 2.00 x 10 ⁻⁰ · 2.18 x 10 ⁻⁰ · 2.01 x 10 ⁻⁰ · 2.01 x 10 ⁻⁰ · 7.28 x 10 ⁻⁰ · 7.29 x 10 ⁻⁰ · 7.55 x 10 ⁻⁰ · 1.55 x 10 ⁻⁰ · 1.50 x 10 ⁻⁰ · 1.50 x 10 ⁻⁰ · 1.50 x 10 ⁻⁰ · 1.50 x 10 ⁻⁰ ·			
		Patterson Lake North Arm – West Basin	3.41 x 10 ⁻⁰³	2.00 x 10 ⁻⁰³	3.41 x 10 ⁻⁰³	2.00 x 10 ⁻⁰³	3.45 x 10 ⁻⁰³	2.01 x 10 ⁻⁰³			
	Snowshoe Hare	Patterson Lake South Arm	1.76 x 10 ⁻⁰³	1.53 x 10 ⁻⁰³	1.76 x 10 ⁻⁰³	1.53 x 10 ⁻⁰³	1.82 x 10 ⁻⁰³	1.55 x 10 ⁻⁰³			
		Beet Lake	1.62 x 10 ⁻⁰³	1.49 x 10 ⁻⁰³	1.62 x 10 ⁻⁰³	1.49 x 10 ⁻⁰³	1.65 x 10 ⁻⁰³	1.50 x 10 ⁻⁰³			
		Lloyd Lake	1.44 x 10 ⁻⁰³	1.44 x 10 ⁻⁰³	1.44 x 10 ⁻⁰³	1.44 x 10 ⁻⁰³		1.44 x 10 ⁻⁰³			
		Reference (Broach Lake)	9.64 × 10 ⁻⁰⁴	9.64 × 10 ⁻⁰⁴	9.64 × 10 ⁻⁰⁴	9.64 × 10 ⁻⁰⁴		9.64 × 10 ⁻⁰⁴			



				Maximum Tota	I Dose (mGy/c	l)	
Biota	Location	Application Case			nd Scenario	RFD	Case
		Operations	Far-Future	Operations	Far-Future	Operations	Far-Future
	Patterson Lake North Arm – West Basin	1.91 x 10 ⁻⁰³	1.36 x 10 ⁻⁰³	1.91 x 10 ⁻⁰³	1.36 x 10 ⁻⁰³	1.93 x 10 ⁻⁰³	1.37 x 10 ⁻⁰³
Southern Red-Backed	Patterson Lake South Arm	1.12 x 10 ⁻⁰³	1.03 x 10 ⁻⁰³	1.12 x 10 ⁻⁰³	1.03 x 10 ⁻⁰³	1.15 x 10 ⁻⁰³	1.04 x 10 ⁻⁰³
Vole	Beet Lake	1.05 x 10 ⁻⁰³	1.00 x 10 ⁻⁰³	1.05 x 10 ⁻⁰³	1.00 x 10 ⁻⁰³	1.07 x 10 ⁻⁰³	Far-Future 1.37 x 10 ⁻⁰³
	Lloyd Lake	9.67 x 10 ⁻⁰⁴	9.65 x 10 ⁻⁰⁴	9.67 x 10 ⁻⁰⁴	9.65 x 10 ⁻⁰⁴	9.69 x 10 ⁻⁰⁴	9.66 x 10 ⁻⁰⁴
	Reference (Broach Lake)	4.28 x 10 ⁻⁰³	4.28 x 10 ⁻⁰³	4.28 x 10 ⁻⁰³	4.28 x 10 ⁻⁰³	4.28 x 10 ⁻⁰³	4.28 x 10 ⁻⁰³
Was disast Cardiana	Patterson Lake North Arm – West Basin	7.69 x 10 ⁻⁰³	4.47 x 10 ⁻⁰³	7.88 x 10 ⁻⁰³	4.47 x 10 ⁻⁰³	7.89 x 10 ⁻⁰³	4.48 x 10 ⁻⁰³
Woodland Caribou	Beet Lake	4.59 x 10 ⁻⁰³	4.31 x 10 ⁻⁰³	4.60 x 10 ⁻⁰³	4.31 x 10 ⁻⁰³	4.65 x 10 ⁻⁰³	4.32 x 10 ⁻⁰³
	Lloyd Lake	4.31 x 10 ⁻⁰³	4.29 x 10 ⁻⁰³	4.31 x 10 ⁻⁰³	4.29 x 10 ⁻⁰³	4.32 x 10 ⁻⁰³	1.37 x 10 ⁻⁰³ 1.04 x 10 ⁻⁰³ 1.01 x 10 ⁻⁰³ 9.66 x 10 ⁻⁰⁴ 4.28 x 10 ⁻⁰³ 4.48 x 10 ⁻⁰³ 4.48 x 10 ⁻⁰³ 4.29 x 10 ⁻⁰³ 2.23 x 10 ⁻⁰³ 2.26 x 10 ⁻⁰³ 2.28 x 10 ⁻⁰³ 2.28 x 10 ⁻⁰³ 2.28 x 10 ⁻⁰³ 2.28 x 10 ⁻⁰³ 2.28 x 10 ⁻⁰³ 3.22 x 10 ⁻⁰⁴ 1.19 x 10 ⁻⁰⁴ 1.19 x 10 ⁻⁰⁴ 9.04 x 10 ⁻⁰⁴ 8.71 x 10 ⁻⁰⁴ 3.23 x 10 ⁻⁰³ 3.24 x 10 ⁻⁰³ 3.24 x 10 ⁻⁰³ 3.24 x 10 ⁻⁰³ 3.24 x 10 ⁻⁰³ 3.24 x 10 ⁻⁰³ 3.24 x 10 ⁻⁰³ 3.29 x 10 ⁻⁰³ 2.98 x 10 ⁻⁰³ 2.98 x 10 ⁻⁰³ 2.98 x 10 ⁻⁰³ 2.99 x 10 ⁻⁰³ 2.99 x 10 ⁻⁰³ 2.99 x 10 ⁻⁰³ 2.99 x 10 ⁻⁰³ 2.99 x 10 ⁻⁰³ 2.99 x 10 ⁻⁰³ 2.97 x 10 ⁻⁰² 2.57 x 10 ⁻⁰² 2.57 x 10 ⁻⁰² 2.55 x 10 ⁻⁰² 7.52 x 10 ⁻⁰² 7.52 x 10 ⁻⁰²
	Reference (Broach Lake)	2.23×10^{-03}	2.23 × 10 ⁻⁰³				
	Patterson Lake North Arm – West Basin	7.15 x 10 ⁻⁰³	2.75 x 10 ⁻⁰³	7.15 x 10 ⁻⁰³	2.75 x 10 ⁻⁰³	7.24 x 10 ⁻⁰³	2.76 x 10 ⁻⁰³
Grouse	Patterson Lake South Arm	3.03 x 10 ⁻⁰³	2.31 x 10 ⁻⁰³	3.03 x 10 ⁻⁰³	2.31 x 10 ⁻⁰³	3.19 x 10 ⁻⁰³	2.33 x 10 ⁻⁰³
	Beet Lake	2.69 x 10 ⁻⁰³	2.28 x 10 ⁻⁰³	2.69 x 10 ⁻⁰³	2.28 x 10 ⁻⁰³	2.75 x 10 ⁻⁰³	2.28 x 10 ⁻⁰³
	Lloyd Lake	2.24 x 10 ⁻⁰³	2.23 x 10 ⁻⁰³	2.24 x 10 ⁻⁰³	2.23 x 10 ⁻⁰³	2.25 x 10 ⁻⁰³	2.23 x 10 ⁻⁰³
	Reference (Broach Lake)	8.70 × 10 ⁻⁰⁴	8.70 × 10 ⁻⁰⁴	8.70 × 10 ⁻⁰⁴	8.70 × 10 ⁻⁰⁴	8.70 × 10 ⁻⁰⁴	8.70 × 10 ⁻⁰⁴
	Patterson Lake North Arm – West Basin	1.95 x 10 ⁻⁰³	1.19 x 10 ⁻⁰³	1.95 x 10 ⁻⁰³	1.19 x 10 ⁻⁰³	1.97 x 10 ⁻⁰³	1.19 x 10 ⁻⁰³
Canada goose	Patterson Lake South Arm	1.05 x 10 ⁻⁰³	9.22 x 10 ⁻⁰⁴	1.05 x 10 ⁻⁰³	9.22 x 10 ⁻⁰⁴	1.08 x 10 ⁻⁰³	9.32 x 10 ⁻⁰⁴
	Beet Lake	9.73 x 10 ⁻⁰⁴	9.00 x 10 ⁻⁰⁴	9.73 x 10 ⁻⁰⁴	9.00 x 10 ⁻⁰⁴	9.86 x 10 ⁻⁰⁴	9.04 x 10 ⁻⁰⁴
	Lloyd Lake	8.73 x 10 ⁻⁰⁴	8.71 x 10 ⁻⁰⁴	8.73 x 10 ⁻⁰⁴	8.71 x 10 ⁻⁰⁴	8.75 x 10 ⁻⁰⁴	8.71 x 10 ⁻⁰⁴
	Reference (Broach Lake)	3.24 × 10 ⁻⁰³	3.23×10^{-03}	3.24×10^{-03}	3.23 × 10 ⁻⁰³	3.24×10^{-03}	3.23 × 10 ⁻⁰³
	Patterson Lake North Arm – West Basin	4.51 × 10 ⁻⁰³	3.43×10^{-03}	8.76 × 10 ⁻⁰³	3.43×10^{-03}	4.51 × 10 ⁻⁰³	3.43×10^{-03}
Little brown myotis	Patterson Lake South Arm	3.37 × 10 ⁻⁰³	3.26×10^{-03}	3.66 × 10 ⁻⁰³	3.26 × 10 ⁻⁰³	4.02×10^{-03}	3.26 × 10 ⁻⁰³
	Beet Lake	3.27 × 10 ⁻⁰³	3.24×10^{-03}	3.31 × 10 ⁻⁰³	3.24 × 10 ⁻⁰³	3.38 × 10 ⁻⁰³	3.24 × 10 ⁻⁰³
	Lloyd Lake	3.24 × 10 ⁻⁰³	3.24×10^{-03}	3.24×10^{-03}	3.24 × 10 ⁻⁰³	3.24×10^{-03}	3.24×10^{-03}
	Reference (Broach Lake)	2.98 × 10 ⁻⁰³	2.98 × 10 ⁻⁰³	2.98 × 10 ⁻⁰³	2.98 × 10 ⁻⁰³	2.98 × 10 ⁻⁰³	2.98 × 10 ⁻⁰³
	Patterson Lake North Arm – West Basin	4.14 × 10 ⁻⁰³	3.21 × 10 ⁻⁰³	8.45 × 10 ⁻⁰³	3.21 × 10 ⁻⁰³	4.14×10^{-03}	3.21 × 10 ⁻⁰³
Common loon	Patterson Lake South Arm	3.07×10^{-03}	3.02×10^{-03}	3.36×10^{-03}	3.02×10^{-03}	3.78×10^{-03}	3.02×10^{-03}
	Beet Lake	3.00×10^{-03}	2.99×10^{-03}	3.04×10^{-03}	2.99×10^{-03}	3.10×10^{-03}	2.99×10^{-03}
	Lloyd Lake	2.98 × 10 ⁻⁰³	2.98×10^{-03}	2.98×10^{-03}	2.98×10^{-03}	2.99×10^{-03}	2.98 × 10 ⁻⁰³
	Reference (Broach Lake)	2.56 × 10 ⁻⁰²	2.56 × 10 ⁻⁰²	2.56 × 10 ⁻⁰²	2.56 × 10 ⁻⁰²	2.56 × 10 ⁻⁰²	2.56 × 10 ⁻⁰²
	Patterson Lake North Arm – West Basin	3.53 × 10 ⁻⁰²	2.77 × 10 ⁻⁰²	7.10×10^{-02}	2.77 × 10 ⁻⁰²	3.53 × 10 ⁻⁰²	2.77 × 10 ⁻⁰²
Mallard	Patterson Lake South Arm	2.63×10^{-02}	2.59×10^{-02}	2.88×10^{-02}	2.59 × 10 ⁻⁰²	3.16 × 10 ⁻⁰²	2.59 × 10 ⁻⁰²
	Beet Lake	2.57 × 10 ⁻⁰²	2.57×10^{-02}	2.61 × 10 ⁻⁰²	2.57 × 10 ⁻⁰²	2.65 × 10 ⁻⁰²	2.57 × 10 ⁻⁰²
	Lloyd Lake	2.56 × 10 ⁻⁰²	2.56 × 10 ⁻⁰²	2.56 × 10 ⁻⁰²	2.56 × 10 ⁻⁰²	2.56 × 10 ⁻⁰²	2.56 × 10 ⁻⁰²
	Reference (Broach Lake)	7.07 × 10 ⁻⁰²	7.07×10^{-02}	7.07×10^{-02}	7.07×10^{-02}	7.07×10^{-02}	7.07 × 10 ⁻⁰²
	Patterson Lake North Arm – West Basin	9.85 x 10 ⁻⁰²	7.52 x 10 ⁻⁰²	1.94 x 10 ⁻⁰¹	7.52 x 10 ⁻⁰²	9.85 x 10 ⁻⁰²	7.52 x 10 ⁻⁰²
Rusty blackbird	Patterson Lake South Arm	7.29 x 10 ⁻⁰²	7.12 x 10 ⁻⁰²	7.93 x 10 ⁻⁰²	7.12 x 10 ⁻⁰²	8.69 x 10 ⁻⁰²	
	Beet Lake	7.12 x 10 ⁻⁰²	7.09 x 10 ⁻⁰²	7.21 x 10 ⁻⁰²	7.09 x 10 ⁻⁰²	7.33 x 10 ⁻⁰²	7.09 x 10 ⁻⁰²
	Lloyd Lake	7.07 x 10 ⁻⁰²	7.07 x 10 ⁻⁰²	7.08 x 10 ⁻⁰²	7.07 x 10 ⁻⁰²	7.09 x 10 ⁻⁰²	7.07 x 10 ⁻⁰²



ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT ECOLOGICAL RISK ASSESSMENT

			Maximum Total Dose (mGy/d)							
	Biota	Location	Applicat	ion Case	Upper Bou	nd Scenario	RFD Case			
			Operations	Far-Future	Operations	Far-Future	Operations	Far-Future		
	Blueberries	Patterson Lake North Arm – West Basin	1.31 x 10 ⁻⁰²	6.80 x 10 ⁻⁰³	1.31 x 10 ⁻⁰²	6.80 x 10 ⁻⁰³	1.32 x 10 ⁻⁰²	6.83 x 10 ⁻⁰³		
	blueberries	Patterson Lake South Arm	6.20 x 10 ⁻⁰³	5.17 x 10 ⁻⁰³	6.20 x 10 ⁻⁰³	5.17 x 10 ⁻⁰³	6.47 x 10 ⁻⁰³	5.23 x 10 ⁻⁰³		
	Busines	Patterson Lake North Arm – West Basin	3.18 x 10 ⁻⁰²	1.41 x 10 ⁻⁰²	3.18 x 10 ⁻⁰²	1.41 x 10 ⁻⁰²	3.22 x 10 ⁻⁰²	1.42 x 10 ⁻⁰²		
Terrestrial	Browse	Patterson Lake South Arm	1.36 x 10 ⁻⁰²	1.08 x 10 ⁻⁰²	1.36 x 10 ⁻⁰²	1.08 x 10 ⁻⁰²	1.44 x 10 ⁻⁰²	1.09 x 10 ⁻⁰²		
Plants	Labraday tas	Patterson Lake North Arm – West Basin	1.30 x 10 ⁻⁰¹	1.41 x 10 ⁻⁰²	1.30 x 10 ⁻⁰¹	1.41 x 10 ⁻⁰²	1.32 x 10 ⁻⁰¹	1.42 x 10 ⁻⁰²		
	Labrador tea	Patterson Lake South Arm	2.97 x 10 ⁻⁰²	1.08 x 10 ⁻⁰²	2.97 x 10 ⁻⁰²	1.08 x 10 ⁻⁰²	3.37 x 10 ⁻⁰²	1.09 x 10 ⁻⁰²		
	Lieban	Patterson Lake North Arm – West Basin	6.79 x 10 ⁻⁰¹	1.12 x 10 ⁻⁰¹	6.79 x 10 ⁻⁰¹	1.12 x 10 ⁻⁰¹	6.90 x 10 ⁻⁰¹	1.12 x 10 ⁻⁰¹		
	Lichen	Patterson Lake South Arm	2.05 x 10 ⁻⁰¹	1.12 x 10 ⁻⁰¹	2.05 x 10 ⁻⁰¹	1.12 x 10 ⁻⁰¹	2.23 x 10 ⁻⁰¹	1.12 x 10 ⁻⁰¹		

RFD = reasonably foreseeable development; mGy/d = milligrays per day.



6.2.5.2 Estimated Dose to Ecological Receptors – Reasonably Foreseeable Development Case

6.2.5.2.1 Non-radiological Dose

Non-radiological dose was only calculated for birds and mammals, as effects to aquatic animals (fish and invertebrates), plants, and soil invertebrates were assessed based on concentrations rather than doses.

The estimated non-radiological doses to the selected birds and mammals during the Project phases and the far-future projection resulting from the RFD Case are shown by COPC in Table 6-13. The doses shown represent the maximum dose by COPC over the assessment period.

The estimated non-radiological doses to the selected birds and mammals in the RFD Case include doses from the release of COPCs from the Fission Patterson Lake South Property. Overall, the estimated non-radiological doses from the RFD Case were similar to the results in the Application Case. Non-radiological doses for arsenic and uranium were higher in the RFD Case than in the Application Case for some terrestrial receptors. For example, during Operations for the RFD Case, the arsenic doses to the beaver, black bear, little brown myotis, common loon, mallard, American mink, muskrat, and rusty blackbird were noticeably higher in the RFD Case at Patterson Lake South Arm (Table 6-13) than in the Application Case (Table 6-10). Similarly, the uranium doses to the little brown myotis, common loon, mallard, American mink, and muskrat were noticeably higher in the RFD Case at Patterson Lake South Arm than in the Application Case. The increase in dose for arsenic and uranium to many of the riparian mammals and birds in the RFD Case is due to the expected release of effluent to Patterson Lake South Arm from the Fission Patterson Lake South Property.

The predicted dose to mammals and birds during the far-future projection in the RFD Case was similar to the that of the far-future projection in the Application Case; this is because the source term for the far-future projection is similar between the Application Case and the RFD Case.



Table 6-12: Estimated Non-radiological Doses to Ecological Receptors – Reasonably Foreseeable Development Case

			Maximum Dose RFD Case (mg/kg/d)									
	Biota	Location			Project Lifespan			\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \		Far-Future		
			Arsenic	Cobalt	Copper	Molybdenum	Uranium	Arsenic	Cobalt	Copper	Molybdenum	Uranium
		Reference (Broach Lake)	3.48×10^{-03}	3.94 × 10 ⁻⁰³	3.15 × 10 ⁻⁰²	1.17 × 10 ⁻⁰³	3.20×10^{-03}	3.48 × 10 ⁻⁰³	3.94 × 10 ⁻⁰³	3.15 × 10 ⁻⁰²	1.17 × 10 ⁻⁰³	3.20×10^{-03}
		Patterson Lake North Arm - West Basin	6.50 × 10 ⁻⁰³	4.57 × 10 ⁻⁰³	3.32 x 10 ⁻⁰²	1.51 × 10 ⁻⁰³	2.51 x 10 ⁻⁰²	3.57 × 10 ⁻⁰³	6.75 x 10 ⁻⁰³	3.76 × 10 ⁻⁰²	3.39×10^{-03}	7.10 x 10 ⁻⁰³
	Beaver	Patterson Lake South Arm	6.64 × 10 ⁻⁰³	4.60 × 10 ⁻⁰³	3.23 × 10 ⁻⁰²	1.33 × 10 ⁻⁰³	7.28 x 10 ⁻⁰³	3.49 × 10 ⁻⁰³	5.64 × 10 ⁻⁰³	3.50 × 10 ⁻⁰²	2.46 × 10 ⁻⁰³	4.13 x 10 ⁻⁰³
		Beet Lake	4.03×10^{-03}	4.24×10^{-03}	3.19 × 10 ⁻⁰²	1.26 × 10 ⁻⁰³	5.34 x 10 ⁻⁰³	3.49×10^{-03}	4.84×10^{-03}	3.33 x 10 ⁻⁰²	1.84×10^{-03}	3.58 x 10 ⁻⁰³
		Lloyd Lake	3.53×10^{-03}	3.98×10^{-03}	3.16 × 10 ⁻⁰²	1.19 × 10 ⁻⁰³	3.24×10^{-03}	3.48×10^{-03}	4.03×10^{-03}	3.17 × 10 ⁻⁰²	1.24×10^{-03}	3.22×10^{-03}
		Reference (Broach Lake)	2.81×10^{-03}	1.17×10^{-03}	2.86 × 10 ⁻⁰²	3.75×10^{-04}	1.02×10^{-03}	2.81 × 10 ⁻⁰³	1.17 × 10 ⁻⁰³	2.86 × 10 ⁻⁰²	3.75×10^{-04}	1.02×10^{-03}
	Black Bear	Patterson Lake North Arm - West Basin	9.39 × 10 ⁻⁰³	1.31 × 10 ⁻⁰³	3.19 × 10 ⁻⁰²	4.23 × 10 ⁻⁰⁴	4.15 x 10 ⁻⁰³	2.87 × 10 ⁻⁰³	1.63 × 10 ⁻⁰³	5.39 × 10 ⁻⁰²	6.09 × 10 ⁻⁰⁴	1.80 x 10 ⁻⁰³
	Diack Deal	Beet Lake	3.79 × 10 ⁻⁰³	1.43 × 10 ⁻⁰³	2.98 × 10 ⁻⁰²	4.59 × 10 ⁻⁰⁴	1.49 x 10 ⁻⁰³	3.09×10^{-03}	1.49 × 10 ⁻⁰³	3.39 × 10 ⁻⁰²	4.95 × 10 ⁻⁰⁴	1.21 x 10 ⁻⁰³
		Lloyd Lake	2.91 × 10 ⁻⁰³	1.21 × 10 ⁻⁰³	2.88 × 10 ⁻⁰²	3.91 × 10 ⁻⁰⁴	1.04×10^{-03}	2.82 × 10 ⁻⁰³	1.22 × 10 ⁻⁰³	2.97 × 10 ⁻⁰²	3.97×10^{-04}	1.03×10^{-03}
		Reference (Broach Lake)	7.50 x 10 ⁻⁰³	6.01 x 10 ⁻⁰⁴	3.31 x 10 ⁻⁰²	2.88 x 10 ⁻⁰⁴	3.73 x 10 ⁻⁰⁴	7.50 x 10 ⁻⁰³	6.01 x 10 ⁻⁰⁴	3.31 x 10 ⁻⁰²	2.88 x 10 ⁻⁰⁴	3.73 x 10 ⁻⁰⁴
	Grey Wolf	Patterson Lake North Arm - West Basin	9.02 x 10 ⁻⁰³	6.13 x 10 ⁻⁰⁴	3.37 x 10 ⁻⁰²	3.18 x 10 ⁻⁰⁴	1.22 x 10 ⁻⁰³	7.53 x 10 ⁻⁰³	6.45 x 10 ⁻⁰⁴	3.43 x 10 ⁻⁰²	4.47 x 10 ⁻⁰⁴	4.54 x 10 ⁻⁰⁴
		Lloyd Lake	7.66 x 10 ⁻⁰³	6.07 x 10 ⁻⁰⁴	3.32 x 10 ⁻⁰²	2.96 x 10 ⁻⁰⁴	3.78 x 10 ⁻⁰⁴	7.50 x 10 ⁻⁰³	6.06 x 10 ⁻⁰⁴	3.32 x 10 ⁻⁰²	3.03 x 10 ⁻⁰⁴	3.75 x 10 ⁻⁰⁴
		Reference (Broach Lake)	1.15 × 10 ⁻⁰²	4.61 × 10 ⁻⁰³	2.28 × 10 ⁻⁰¹	1.15 × 10 ⁻⁰³	1.08×10^{-03}	1.15 × 10 ⁻⁰²	4.61 × 10 ⁻⁰³	2.28 × 10 ⁻⁰¹	1.15 × 10 ⁻⁰³	1.08×10^{-03}
		Patterson Lake North Arm - West Basin	3.46 × 10 ⁻⁰²	5.83 × 10 ⁻⁰³	2.66 × 10 ⁻⁰¹	2.80 × 10 ⁻⁰³	4.17 x 10 ⁻⁰³	1.22 × 10 ⁻⁰²	1.06 × 10 ⁻⁰²	5.90 × 10 ⁻⁰¹	2.20 × 10 ⁻⁰²	8.39 x 10 ⁻⁰³
	Mink	Patterson Lake South Arm	3.14×10^{-02}	5.52 × 10 ⁻⁰³	2.56 × 10 ⁻⁰¹	2.27 × 10 ⁻⁰³	2.56 x 10 ⁻⁰³	1.16 × 10 ⁻⁰²	8.23 × 10 ⁻⁰³	4.41 × 10 ⁻⁰¹	1.33 × 10 ⁻⁰²	3.09 × 10 ⁻⁰³
		Beet Lake	1.49×10^{-02}	5.09 × 10 ⁻⁰³	2.43 × 10 ⁻⁰¹	1.73 × 10 ⁻⁰³	1.64 x 10 ⁻⁰³	1.15 × 10 ⁻⁰²	6.53 × 10 ⁻⁰³	3.39 × 10 ⁻⁰¹	7.46 × 10 ⁻⁰³	1.83 × 10 ⁻⁰³
		Lloyd Lake	1.17 × 10 ⁻⁰²	4.66 × 10 ⁻⁰³	2.30 × 10 ⁻⁰¹	1.21 × 10 ⁻⁰³	1.13 × 10 ⁻⁰³	1.15 × 10 ⁻⁰²	4.81 × 10 ⁻⁰³	2.40 × 10 ⁻⁰¹	1.79 × 10 ⁻⁰³	1.15 × 10 ⁻⁰³
		Reference (Broach Lake)	2.35×10^{-03}	2.83 × 10 ⁻⁰³	1.61 × 10 ⁻⁰²	5.80 × 10 ⁻⁰⁴	1.59 × 10 ⁻⁰³	2.35 × 10 ⁻⁰³	2.83 × 10 ⁻⁰³	1.61 × 10 ⁻⁰²	5.80 × 10 ⁻⁰⁴	1.59 × 10 ⁻⁰³
		Patterson Lake North Arm - West Basin	5.45×10^{-03}	3.45 × 10 ⁻⁰³	1.73 x 10 ⁻⁰²	8.29 x 10 ⁻⁰⁴	1.18 x 10 ⁻⁰²	2.44 × 10 ⁻⁰³	5.62 × 10 ⁻⁰³	2.22 x 10 ⁻⁰²	2.78 × 10 ⁻⁰³	4.83 x 10 ⁻⁰³
	Moose	Patterson Lake South Arm	5.42 × 10 ⁻⁰³	3.41 × 10 ⁻⁰³	1.72 × 10 ⁻⁰²	7.95 x 10 ⁻⁰⁴	1.15 x 10 ⁻⁰²	2.43 × 10 ⁻⁰³	5.55 × 10 ⁻⁰³	2.20 × 10 ⁻⁰²	2.53 × 10 ⁻⁰³	4.75 x 10 ⁻⁰³
		Beet Lake	2.90 × 10 ⁻⁰³	3.12 × 10 ⁻⁰³	1.65 × 10 ⁻⁰²	6.54 × 10 ⁻⁰⁴	2.68 x 10 ⁻⁰³	2.36 × 10 ⁻⁰³	3.73 × 10 ⁻⁰³	1.80 × 10 ⁻⁰²	1.24 × 10 ⁻⁰³	1.92 x 10 ⁻⁰³
		Lloyd Lake	2.39 × 10 ⁻⁰³	2.86 × 10 ⁻⁰³	1.62 × 10 ⁻⁰²	5.92 × 10 ⁻⁰⁴	1.62 × 10 ⁻⁰³	2.35 × 10 ⁻⁰³	2.92 × 10 ⁻⁰³	1.63 × 10 ⁻⁰²	6.48 × 10 ⁻⁰⁴	1.62 × 10 ⁻⁰³
		Reference (Broach Lake)	2.86×10^{-02}	1.60 × 10 ⁻⁰²	4.08 × 10 ⁻⁰¹	3.65 × 10 ⁻⁰³	5.51 × 10 ⁻⁰³	2.86 × 10 ⁻⁰²	1.60 × 10 ⁻⁰²	4.08 × 10 ⁻⁰¹	3.65 × 10 ⁻⁰³	5.51 × 10 ⁻⁰³
		Patterson Lake North Arm - West Basin	8.77 × 10 ⁻⁰²	2.08 × 10 ⁻⁰²	4.74 × 10 ⁻⁰¹	9.77 × 10 ⁻⁰³	1.82 × 10 ⁻⁰²	3.03 × 10 ⁻⁰²	3.91 × 10 ⁻⁰²	$1.05 \times 10^{+00}$	8.14 × 10 ⁻⁰²	5.51 × 10 ⁻⁰²
	Muskrat	Patterson Lake South Arm	6.98×10^{-02}	2.01 × 10 ⁻⁰²	4.58 × 10 ⁻⁰¹	7.84 × 10 ⁻⁰³	1.37 × 10 ⁻⁰²	2.87 × 10 ⁻⁰²	3.00 × 10 ⁻⁰²	7.87 × 10 ⁻⁰¹	4.88 × 10 ⁻⁰²	1.92 × 10 ⁻⁰²
Terrestrial		Beet Lake	3.56×10^{-02}	1.80 × 10 ⁻⁰²	4.33 × 10 ⁻⁰¹	5.82 × 10 ⁻⁰³	8.49 × 10 ⁻⁰³	2.86 × 10 ⁻⁰²	2.34 × 10 ⁻⁰²	6.06 × 10 ⁻⁰¹	2.71 × 10 ⁻⁰²	1.06 × 10 ⁻⁰²
Animals		Lloyd Lake	2.90×10^{-02}	1.62 × 10 ⁻⁰²	4.10 × 10 ⁻⁰¹	3.87 × 10 ⁻⁰³	5.78 × 10 ⁻⁰³	2.86 × 10 ⁻⁰²	1.67 × 10 ⁻⁰²	4.28 × 10 ⁻⁰¹	6.04 × 10 ⁻⁰³	5.97 × 10 ⁻⁰³
71111110115		Reference (Broach Lake)	1.27 × 10 ⁻⁰³	4.90 × 10 ⁻⁰⁴	6.99 × 10 ⁻⁰³	3.17 × 10 ⁻⁰⁴	3.72 × 10 ⁻⁰⁴	1.27 × 10 ⁻⁰³	4.90 × 10 ⁻⁰⁴	6.99 × 10 ⁻⁰³	3.17 × 10 ⁻⁰⁴	3.72 × 10 ⁻⁰⁴
		Patterson Lake North Arm - West Basin	1.31×10^{-03}	5.09 × 10 ⁻⁰⁴	7.26 x 10 ⁻⁰³	3.80 x 10 ⁻⁰⁴	3.48 x 10 ⁻⁰³	1.27 × 10 ⁻⁰³	5.64 x 10 ⁻⁰⁴	7.12 × 10 ⁻⁰³	6.94 × 10 ⁻⁰⁴	6.13 x 10 ⁻⁰⁴
	Red Fox	Patterson Lake South Arm	1.33 × 10 ⁻⁰³	5.10 x 10 ⁻⁰⁴	7.02 × 10 ⁻⁰³	3.51 × 10 ⁻⁰⁴	9.02 x 10 ⁻⁰⁴	1.27 × 10 ⁻⁰³	5.35 × 10 ⁻⁰⁴	7.06 × 10 ⁻⁰³	5.36 × 10 ⁻⁰⁴	4.20 x 10 ⁻⁰⁴
		Beet Lake	1.29 × 10 ⁻⁰³	5.00 × 10 ⁻⁰⁴	7.01 × 10 ⁻⁰³	3.34 × 10 ⁻⁰⁴	6.60 x 10 ⁻⁰⁴	1.27 × 10 ⁻⁰³	5.14 × 10 ⁻⁰⁴	7.02 × 10 ⁻⁰³	4.31 × 10 ⁻⁰⁴	3.94 x 10 ⁻⁰⁴
		Lloyd Lake	1.28 × 10 ⁻⁰³	4.93 × 10 ⁻⁰⁴	7.00 × 10 ⁻⁰³	3.21 × 10 ⁻⁰⁴	3.76 × 10 ⁻⁰⁴	1.27 × 10 ⁻⁰³	4.93 × 10 ⁻⁰⁴	6.99 × 10 ⁻⁰³	3.29 × 10 ⁻⁰⁴	3.73 × 10 ⁻⁰⁴
		Reference (Broach Lake)	4.30 × 10 ⁻⁰³	3.98 × 10 ⁻⁰³	4.82 × 10 ⁻⁰²	2.01 × 10 ⁻⁰³	5.27 × 10 ⁻⁰³	4.30 × 10 ⁻⁰³	3.98 × 10 ⁻⁰³	4.82 × 10 ⁻⁰²	2.01 × 10 ⁻⁰³	5.27 × 10 ⁻⁰³
		Patterson Lake North Arm - West Basin	4.34 × 10 ⁻⁰³	4.02 × 10 ⁻⁰³	5.00 x 10 ⁻⁰²	2.31 × 10 ⁻⁰³	4.26 x 10 ⁻⁰²	4.30 × 10 ⁻⁰³	4.06 × 10 ⁻⁰³	4.85 × 10 ⁻⁰²	2.43 × 10 ⁻⁰³	7.73 x 10 ⁻⁰³
	Snowshoe Hare	Patterson Lake South Arm	4.37 × 10 ⁻⁰³	4.02 × 10 ⁻⁰³	4.84 x 10 ⁻⁰²	2.07 × 10 ⁻⁰³	1.13 x 10 ⁻⁰²	4.30 × 10 ⁻⁰³	4.03 × 10 ⁻⁰³	4.83 × 10 ⁻⁰²	2.25 × 10 ⁻⁰³	5.67 x 10 ⁻⁰³
		Beet Lake	4.32 × 10 ⁻⁰³	4.01 × 10 ⁻⁰³	4.83 × 10 ⁻⁰²	2.05 × 10 ⁻⁰³	8.62 x 10 ⁻⁰³	4.30 × 10 ⁻⁰³	4.01 × 10 ⁻⁰³	4.83 × 10 ⁻⁰²	2.14 × 10 ⁻⁰³	5.49 x 10 ⁻⁰³
		Lloyd Lake	4.31 × 10 ⁻⁰³	4.01 x 10 ⁻⁰³	4.83 × 10 ⁻⁰²	2.04 × 10 ⁻⁰³	5.31 × 10 ⁻⁰³	4.30 × 10 ⁻⁰³	3.99 × 10 ⁻⁰³	4.82 × 10 ⁻⁰²	2.03 × 10 ⁻⁰³	5.28 × 10 ⁻⁰³
		Reference (Broach Lake)	3.10 × 10 ⁻⁰³	2.94 × 10 ⁻⁰³	3.66 × 10 ⁻⁰²	1.50 × 10 ⁻⁰³	3.95 × 10 ⁻⁰³	3.10 × 10 ⁻⁰³	2.94 × 10 ⁻⁰³	3.66 × 10 ⁻⁰²	1.50 × 10 ⁻⁰³	3.95 × 10 ⁻⁰³
	Southern Red-	Patterson Lake North Arm - West Basin	3.16 × 10 ⁻⁰³	2.98 × 10 ⁻⁰³	3.79 x 10 ⁻⁰²	1.74 × 10 ⁻⁰³	3.08 x 10 ⁻⁰²	3.10 × 10 ⁻⁰³	3.06 × 10 ⁻⁰³	3.69 × 10 ⁻⁰²	2.12 × 10 ⁻⁰³	5.84 x 10 ⁻⁰³
	Backed Vole	Patterson Lake South Arm	3.19 × 10 ⁻⁰³	2.98 × 10 ⁻⁰³	3.67 × 10 ⁻⁰²	1.57 × 10 ⁻⁰³	8.29 x 10 ⁻⁰³	3.10 × 10 ⁻⁰³	3.01 × 10 ⁻⁰³	3.67 × 10 ⁻⁰²	1.86 × 10 ⁻⁰³	4.26 x 10 ⁻⁰³
		Beet Lake	3.13 × 10 ⁻⁰³	2.97 × 10 ⁻⁰³	3.67 × 10 ⁻⁰²	1.54 × 10 ⁻⁰³	6.36 x 10 ⁻⁰³	3.10 × 10 ⁻⁰³	2.98 × 10 ⁻⁰³	3.67 × 10 ⁻⁰²	1.69 × 10 ⁻⁰³	4.12 x 10 ⁻⁰³
		Lloyd Lake	3.11 × 10 ⁻⁰³	2.96 × 10 ⁻⁰³	3.67 × 10 ⁻⁰²	1.52 × 10 ⁻⁰³	3.98 × 10 ⁻⁰³	3.10 × 10 ⁻⁰³	2.95 x 10 ⁻⁰³	3.66 × 10 ⁻⁰²	1.52×10^{-03}	3.95 × 10 ⁻⁰³
	W	Reference (Broach Lake)	5.46 x 10 ⁻⁰³	3.94 x 10 ⁻⁰³	1.78 x 10 ⁻⁰²	1.24 x 10 ⁻⁰³	1.77 x 10 ⁻⁰³	5.46 x 10 ⁻⁰³	3.94 x 10 ⁻⁰³	1.78 x 10 ⁻⁰²	1.24 x 10 ⁻⁰³	1.77 x 10 ⁻⁰³
	WoodLand	Patterson Lake North Arm - West Basin	1.16 x 10 ⁻⁰²	4.71 x 10 ⁻⁰³	1.97 x 10 ⁻⁰²	2.42 x 10 ⁻⁰³	7.32 x 10 ⁻⁰²	5.57 x 10 ⁻⁰³	6.64 x 10 ⁻⁰³	2.34 x 10 ⁻⁰²	5.54 x 10 ⁻⁰³	6.08 x 10 ⁻⁰³
	Caribou	Beet Lake	6.06 x 10 ⁻⁰³	4.18 x 10 ⁻⁰³	1.81 x 10 ⁻⁰²	1.39 x 10 ⁻⁰³	7.78 x 10 ⁻⁰³	5.46 x 10 ⁻⁰³	4.49 x 10 ⁻⁰³	1.89 x 10 ⁻⁰²	2.07 x 10 ⁻⁰³	2.13 x 10 ⁻⁰³
		Lloyd Lake	5.66 x 10 ⁻⁰³	4.11 x 10 ⁻⁰³	1.80 x 10 ⁻⁰²	1.39 x 10 ⁻⁰³	1.95 x 10 ⁻⁰³	5.46 x 10 ⁻⁰³	4.06 x 10 ⁻⁰³	1.80 x 10 ⁻⁰²	1.41 x 10 ⁻⁰³	1.83 x 10 ⁻⁰³
		Reference (Broach Lake)	6.44 × 10 ⁻⁰³	5.57 × 10 ⁻⁰³	5.26 × 10 ⁻⁰²	2.57 × 10 ⁻⁰³	6.34 × 10 ⁻⁰³	6.44 × 10 ⁻⁰³	5.57 × 10 ⁻⁰³	5.26 × 10 ⁻⁰²	2.57 × 10 ⁻⁰³	6.34 × 10 ⁻⁰³
	Graves	Patterson Lake North Arm - West Basin	6.48 × 10 ⁻⁰³	5.62 × 10 ⁻⁰³	5.45 x 10 ⁻⁰²	2.89 × 10 ⁻⁰³	4.85 x 10 ⁻⁰²	6.44×10^{-03}	5.64 × 10 ⁻⁰³	5.28 × 10 ⁻⁰²	2.88 × 10 ⁻⁰³	9.26 x 10 ⁻⁰³
	Grouse	Patterson Lake South Arm	6.49 × 10 ⁻⁰³	5.62 × 10 ⁻⁰³	5.27 × 10 ⁻⁰²	2.62 × 10 ⁻⁰³	1.31 x 10 ⁻⁰²	6.44 × 10 ⁻⁰³	5.62 × 10 ⁻⁰³	5.27 × 10 ⁻⁰²	2.75 × 10 ⁻⁰³	6.82 x 10 ⁻⁰³
		Beet Lake	6.46 × 10 ⁻⁰³	5.61 × 10 ⁻⁰³	5.27 × 10 ⁻⁰²	2.61 × 10 ⁻⁰³	1.01 x 10 ⁻⁰²	6.44 × 10 ⁻⁰³	5.60 × 10 ⁻⁰³	5.26 × 10 ⁻⁰²	2.67 × 10 ⁻⁰³	6.60 x 10 ⁻⁰³
		Lloyd Lake	6.45×10^{-03}	5.61 × 10 ⁻⁰³	5.27 × 10 ⁻⁰²	2.60 × 10 ⁻⁰³	6.39 × 10 ⁻⁰³	6.44 × 10 ⁻⁰³	5.59 x 10 ⁻⁰³	5.26 × 10 ⁻⁰²	2.58 × 10 ⁻⁰³	6.35 × 10 ⁻⁰³
	Canada Goose	Reference (Broach Lake)	6.19 × 10 ⁻⁰⁴	5.56 × 10 ⁻⁰⁴	5.30 × 10 ⁻⁰³	2.58 × 10 ⁻⁰⁴	6.29 × 10 ⁻⁰⁴	6.19 × 10 ⁻⁰⁴	5.56 × 10 ⁻⁰⁴	5.30 × 10 ⁻⁰³	2.58 × 10 ⁻⁰⁴	6.29 × 10 ⁻⁰⁴
Calla		Patterson Lake North Arm - West Basin	6.33 × 10 ⁻⁰⁴	5.66 × 10 ⁻⁰⁴	5.50 x 10 ⁻⁰³	3.03 x 10 ⁻⁰⁴	4.91 x 10 ⁻⁰³	6.19 × 10 ⁻⁰⁴	5.89 × 10 ⁻⁰⁴	5.36×10^{-03}	4.27×10^{-04}	9.45 x 10 ⁻⁰⁴

						Maximum Dose F	RFD Case (mg/kg/d)								
Biota	Location			Project Lifespan					Far-Future						
		Arsenic	Cobalt	Copper	Molybdenum	Uranium	Arsenic	Cobalt	Copper	Molybdenum	Uranium				
	Patterson Lake South Arm	6.43 × 10 ⁻⁰⁴	5.66 × 10 ⁻⁰⁴	5.32 × 10 ⁻⁰³	2.74 × 10 ⁻⁰⁴	1.33 x 10 ⁻⁰³	6.19 × 10 ⁻⁰⁴	5.76 × 10 ⁻⁰⁴	5.33 × 10 ⁻⁰³	3.56 × 10 ⁻⁰⁴	6.83 x 10 ⁻⁰⁴				
	Beet Lake	6.25 × 10 ⁻⁰⁴	5.62 × 10 ⁻⁰⁴	5.31 × 10 ⁻⁰³	2.67 × 10 ⁻⁰⁴	1.02 x 10 ⁻⁰³	6.19 × 10 ⁻⁰⁴	5.67 × 10 ⁻⁰⁴	5.32 x 10 ⁻⁰³	3.09×10^{-04}	6.58 x 10 ⁻⁰⁴				
	Lloyd Lake	6.21 × 10 ⁻⁰⁴	5.59 × 10 ⁻⁰⁴	5.31 × 10 ⁻⁰³	2.61 × 10 ⁻⁰⁴	6.34 × 10 ⁻⁰⁴	6.19 × 10 ⁻⁰⁴	5.58 x 10 ⁻⁰⁴	5.30 × 10 ⁻⁰³	2.63 × 10 ⁻⁰⁴	6.29 × 10 ⁻⁰⁴				
	Reference (Broach Lake)	2.77 × 10 ⁻⁰²	5.73 × 10 ⁻⁰²	3.22 × 10 ⁺⁰⁰	1.64 × 10 ⁻⁰²	4.77 × 10 ⁻⁰³	2.77 × 10 ⁻⁰²	5.73 × 10 ⁻⁰²	3.22 × 10 ⁺⁰⁰	1.64 × 10 ⁻⁰²	4.77×10^{-03}				
Litale Donor	Patterson Lake North Arm - West Basin	8.44×10^{-02}	7.41 × 10 ⁻⁰²	$3.74 \times 10^{+00}$	4.38 × 10 ⁻⁰²	1.57 × 10 ⁻⁰²	2.93 × 10 ⁻⁰²	1.40 × 10 ⁻⁰¹	8.33 × 10 ⁺⁰⁰	3.66 × 10 ⁻⁰¹	4.78×10^{-02}				
Little Brown	Patterson Lake South Arm	6.35×10^{-02}	6.97 × 10 ⁻⁰²	$3.61 \times 10^{+00}$	3.50 × 10 ⁻⁰²	1.11 × 10 ⁻⁰²	2.78 × 10 ⁻⁰²	1.08 × 10 ⁻⁰¹	6.22 × 10 ⁺⁰⁰	2.19 × 10 ⁻⁰¹	1.67×10^{-02}				
Myotis	Beet Lake	3.38×10^{-02}	6.39×10^{-02}	$3.42 \times 10^{+00}$	2.61 × 10 ⁻⁰²	7.10 × 10 ⁻⁰³	2.77 × 10 ⁻⁰²	8.39 × 10 ⁻⁰²	$4.79 \times 10^{+00}$	1.22 × 10 ⁻⁰¹	9.19 × 10 ⁻⁰³				
	Lloyd Lake	2.80×10^{-02}	5.79 × 10 ⁻⁰²	$3.24 \times 10^{+00}$	1.74 × 10 ⁻⁰²	4.99 × 10 ⁻⁰³	2.77 × 10 ⁻⁰²	6.00 × 10 ⁻⁰²	3.38 × 10 ⁺⁰⁰	2.72 × 10 ⁻⁰²	5.17×10^{-03}				
	Reference (Broach Lake)	4.71×10^{-03}	2.18×10^{-03}	1.25 × 10 ⁻⁰¹	4.10×10^{-04}	3.38 × 10 ⁻⁰⁴	4.71 × 10 ⁻⁰³	2.18 × 10 ⁻⁰³	1.25 × 10 ⁻⁰¹	4.10 × 10 ⁻⁰⁴	3.38 × 10 ⁻⁰⁴				
	Patterson Lake North Arm - West Basin	1.63 × 10 ⁻⁰²	2.84×10^{-03}	1.46 × 10 ⁻⁰¹	1.10 × 10 ⁻⁰³	1.22 × 10 ⁻⁰³	5.00 × 10 ⁻⁰³	5.34 × 10 ⁻⁰³	3.22 × 10 ⁻⁰¹	9.15 × 10 ⁻⁰³	3.38×10^{-03}				
Loon	Patterson Lake South Arm	2.41 × 10 ⁻⁰²	2.66 × 10 ⁻⁰³	1.40 × 10 ⁻⁰¹	8.78 × 10 ⁻⁰⁴	1.41 × 10 ⁻⁰³	4.74×10^{-03}	4.09 × 10 ⁻⁰³	2.41 × 10 ⁻⁰¹	5.48 × 10 ⁻⁰³	1.18 × 10 ⁻⁰³				
	Beet Lake	8.01 × 10 ⁻⁰³	2.43×10^{-03}	1.33 × 10 ⁻⁰¹	6.53 × 10 ⁻⁰⁴	7.06 × 10 ⁻⁰⁴	4.72 × 10 ⁻⁰³	3.19 × 10 ⁻⁰³	1.85 × 10 ⁻⁰¹	3.05 × 10 ⁻⁰³	6.51 × 10 ⁻⁰⁴				
	Lloyd Lake	4.92×10^{-03}	2.21×10^{-03}	1.25 × 10 ⁻⁰¹	4.35 × 10 ⁻⁰⁴	3.72 × 10 ⁻⁰⁴	4.71 × 10 ⁻⁰³	2.28 × 10 ⁻⁰³	1.31 × 10 ⁻⁰¹	6.79 × 10 ⁻⁰⁴	3.66×10^{-04}				
	Reference (Broach Lake)	2.78×10^{-02}	2.60 × 10 ⁻⁰²	$1.05 \times 10^{+00}$	6.52 × 10 ⁻⁰³	5.23 × 10 ⁻⁰³	2.78 × 10 ⁻⁰²	2.60 × 10 ⁻⁰²	$1.05 \times 10^{+00}$	6.52 × 10 ⁻⁰³	5.23×10^{-03}				
	Patterson Lake North Arm - West Basin	8.52×10^{-02}	3.37×10^{-02}	1.23 × 10 ⁺⁰⁰	1.75 × 10 ⁻⁰²	1.73 × 10 ⁻⁰²	2.94 × 10 ⁻⁰²	6.35 × 10 ⁻⁰²	$2.73 \times 10^{+00}$	1.46 × 10 ⁻⁰¹	5.23 × 10 ⁻⁰²				
Mallard	Patterson Lake South Arm	6.74×10^{-02}	3.22×10^{-02}	$1.18 \times 10^{+00}$	1.40 × 10 ⁻⁰²	1.29 × 10 ⁻⁰²	2.79 × 10 ⁻⁰²	4.88 × 10 ⁻⁰²	$2.04 \times 10^{+00}$	8.72 × 10 ⁻⁰²	1.82×10^{-02}				
	Beet Lake	3.45×10^{-02}	2.91 × 10 ⁻⁰²	1.12 × 10 ⁺⁰⁰	1.04 × 10 ⁻⁰²	8.04×10^{-03}	2.78 × 10 ⁻⁰²	3.80 × 10 ⁻⁰²	1.57 × 10 ⁺⁰⁰	4.85 × 10 ⁻⁰²	1.01 × 10 ⁻⁰²				
	Lloyd Lake	2.82×10^{-02}	2.63×10^{-02}	1.06 × 10 ⁺⁰⁰	6.92 × 10 ⁻⁰³	5.48 × 10 ⁻⁰³	2.78 × 10 ⁻⁰²	2.72 × 10 ⁻⁰²	1.11 × 10 ⁺⁰⁰	1.08 × 10 ⁻⁰²	5.66 × 10 ⁻⁰³				
	Reference (Broach Lake)	7.16×10^{-02}	1.25×10^{-01}	6.35 × 10 ⁺⁰⁰	3.66 × 10 ⁻⁰²	1.76 × 10 ⁻⁰²	7.16 × 10 ⁻⁰²	1.25 × 10 ⁻⁰¹	$6.35 \times 10^{+00}$	3.66 × 10 ⁻⁰²	1.76 × 10 ⁻⁰²				
	Patterson Lake North Arm - West Basin	1.83×10^{-01}	1.58×10^{-01}	$7.38 \times 10^{+00}$	9.06 × 10 ⁻⁰²	7.66 x 10 ⁻⁰²	7.49 × 10 ⁻⁰²	2.88 × 10 ⁻⁰¹	$1.64 \times 10^{+01}$	7.22 × 10 ⁻⁰¹	1.06 x 10 ⁻⁰¹				
Rusty Blackbird	Patterson Lake South Arm	1.42 × 10 ⁻⁰¹	1.50 × 10 ⁻⁰¹	$7.12 \times 10^{+00}$	7.30 × 10 ⁻⁰²	3.59 x 10 ⁻⁰²	7.19 × 10 ⁻⁰²	2.24 × 10 ⁻⁰¹	1.22 × 10 ⁺⁰¹	4.35 × 10 ⁻⁰¹	4.15 x 10 ⁻⁰²				
	Beet Lake	8.36×10^{-02}	1.38 × 10 ⁻⁰¹	$6.75 \times 10^{+00}$	5.56 × 10 ⁻⁰²	2.55 x 10 ⁻⁰²	7.17 × 10 ⁻⁰²	1.78 × 10 ⁻⁰¹	9.43 × 10 ⁺⁰⁰	2.44×10^{-01}	2.66 x 10 ⁻⁰²				
	Lloyd Lake	7.24 × 10 ⁻⁰²	1.27 × 10 ⁻⁰¹	6.39 × 10 ⁺⁰⁰	3.86 × 10 ⁻⁰²	1.80 × 10 ⁻⁰²	7.16 × 10 ⁻⁰²	1.31 × 10 ⁻⁰¹	6.67 × 10 ⁺⁰⁰	5.77 × 10 ⁻⁰²	1.84 × 10 ⁻⁰²				

RFD = reasonably foreseeable development.

6.2.5.2.2 Radiological Dose

The estimated radiation doses to aquatic and terrestrial ecological receptors during the Project phases and far-future projection resulting from the RFD Case are shown in Table 6-12. The doses shown represent the maximum total dose from all radionuclides over the assessment period. The dose breakdown by radionuclide is shown in Appendix C.

The estimated radiation dose to aquatic and terrestrial ecological receptors in the RFD Case includes a dose from the release of COPCs from the Fission Patterson Lake South Property. Thus, the radiation doses were slightly higher in the RFD Case at Patterson Lake South Arm than in the Application Case; however, the radiation dose would decrease to levels that are similar to the Application Case in the downstream environment.

In the RFD Case, the maximum estimated radiation dose of 0.12 mGy/d would occur during Operations for zooplankton in the Patterson Lake South Arm, and would decrease to 0.07 mGy/d in Lloyd Lake. The main contributor to total dose would be from polonium-210 in water. Similarly, the maximum estimated radiation doses for northern pike, lake whitefish, macrophytes, and phytoplankton were higher in the Patterson Lake South Arm in the RFD Case than in the Patterson Lake North Arm – West Basin.

For terrestrial receptors in the RFD Case, the maximum estimated radiation dose for birds and mammals still would occur during Operations for the rusty blackbird eating from the Patterson Lake North Arm – West Basin, which is 0.10 mGy/d in the Application Case and the RFD Case. The maximum dose for terrestrial plants in the RFD Case is to lichen (0.69 mGy/d) due to atmospheric deposition from air to the plant.

In the RFD Case, the predicted radiation dose to all ecological receptors during the far-future projection was lower than during Operations. The maximum estimated dose in the far-future projection to zooplankton was 0.07 mGy/d, to rusty blackbird was 0.08 mGy/d in the Patterson Lake North Arm – West Basin, and to lichen was 0.11 mGy/d in the Patterson Lake North Arm – West Basin.



6.2.6 Uncertainty in Exposure Assessment

For each model scenario and Project phase, it was conservatively assumed that ecological receptors are exposed to the maximum exposure concentrations at their location. The duration of exposure was assumed to be sufficient for each receptor to be in equilibrium with their environment. This resulted in conservatively high:

- a. direct exposure estimates for aquatic biota exposed to COPCs in water, and terrestrial plants and soil invertebrates exposed to COPCs in soil;
- b. predicted uptakes of COPCs by ecological receptors in the food chain; and
- c. estimated doses of COPCs by ecological receptors through the food chain.

The assumptions to address uncertainties in the exposure assessment are anticipated to produce conservative exposure estimates for ecological receptors. The risk that the exposure assessment underestimates potential exposure of ecological receptors to COPCs from the Project is low. That said, the following provides more detail about assessment uncertainty and how it was addressed.

6.2.6.1 Uncertainties in Uptake and Exposure Factors

Wildlife exposure factors, such as intake rates and diets, are a potential source of uncertainty. Reputable sources were used for these factors, and the factors are considered to be representative of the organisms assessed. Feed, water, and inhalation intake rates were obtained or calculated based on the following primary sources: Federal Contaminated Sites Action Plan (FCSAP 2012b), USEPA (1993) and Sample and Suter (1994). These documents have undergone several stages of review and are considered appropriate literature values for use in this assessment; therefore, the uncertainty in these values is considered acceptable.

The BAFs were used to calculate uptake into tissues (fish, invertebrates, plants). Bioaccumulation factors from regional biota and water were used for a number of aquatic BAFs in this assessment. There is inherent uncertainty in using field data to calculate BAFs from metal concentrations in tissues of aquatic biota and surface water concentrations, because the actual exposure history of the organisms is unknown. Unless it is known that a metal concentration in surface water is at a steady state for an extended period of time, the use of tissue and water concentrations sampled at the same time from the same location may not reflect the average exposure of an organism. In addition, as a result of physiological control, intracellular storage and different excretion mechanisms, biota have an ability to actively regulate the body burden of many metals and maintain homeostatic control over a range of exposures (Hamilton and Mehrle 1986; Chapman et al. 1996; Wood and Port 2000). These homeostatic controls can produce a non-linear relationship between the steady-state tissue concentration and the environmental exposure (Newman and Unger 2002). As a result, the validity of assuming a linear relationship between water and tissue concentrations is an area of uncertainty. In most cases it is difficult to assess whether non-linear relationships may exist; therefore, linear relationships are assumed by default. However, these complicating issues do not diminish the importance of BAFs in assessing the environmental hazard associated with metals.



6.2.6.2 Uncertainties in Dose Coefficients

Dose coefficients were obtained from reputable sources for reference organisms, but have not been derived specifically for all the organisms assessed. Dose coefficients for surrogate organisms were often used. They were selected with attention to similar body size and exposure habits and are believed to adequately represent the organism assessed. Dose coefficients for each receptor were not adjusted for body size and dimensions, which represents a possible source of uncertainty. For the maximum exposed receptors, the dose is primarily delivered through an alpha decay as over 95% of the dose can be attributed to the decay of polonium-210. The geometry-scaling factor of alpha particles is 1 for all organisms and geometries; as such, geometry assumptions are expected to have very little effect on the total radiation dose.

6.3 Effects Assessment

The effects assessment characterized the nature of potential effects observed or expected as a result of the Project and the Project plus the Fission Patterson Lake property. The effects assessment considers potential health risks to ecological receptors from long-term (chronic) exposures to COPCs in the environment. The potential for adverse effects during sensitive life stages for ecological receptors, including effects on growth, reproduction, and survival, were considered.

The ecological toxicity assessment considered potential adverse ecological effects from non-radiological and radiological exposures. The selected TRVs represent concentrations or weight-normalized daily doses of COPCs, or radiological dose benchmarks, below which ecologically relevant effects on growth, reproduction, or survival are not expected to occur.

6.3.1 Toxicological Benchmarks

For assessment of non-radiological COPCs, a TRV was used. A TRV is a toxicological index associating specific effects with a level of exposure to a chemical. The TRVs for aquatic biota are based on concentrations in water, while TRVs for mammals and birds are weight-normalized daily oral doses.

Arsenic, chloride, cobalt, copper, molybdenum, sulphate, and uranium were identified in Section 4.0, Model Integration and Evaluation of Sources, for further evaluation in the EcoRA for aquatic biota, and arsenic, cobalt, copper, molybdenum, and uranium were identified for further evaluation in the EcoRA for terrestrial biota.

No COPCs in air were identified for further evaluation of potential risks for ecological health; therefore, TRVs for direct contact with air were not included in the toxicity assessment. Deposition of COPCs in dust to soil was evaluated; however, no COPCs in soil were identified for further evaluation; therefore, toxicity via direct contact with COPCs in soil for plants and soil invertebrates was not included in the toxicity assessment.



6.3.1.1 Toxicity Reference Values for Aquatic Biota

Water concentration-based TRVs for aquatic biota are considered chronic effects from long-term exposures. They are concentrations below which health risks to receptors are not anticipated. The TRVs derived for aquatic biota included six categories: forage fish (lake whitefish), predator fish (northern pike), zooplankton, benthic invertebrates, phytoplankton, and aquatic plants.

The selected TRVs were 20% ECs (i.e., EC_{20} values), which are concentrations at which only 20% of the test organisms respond. The EC_{20} value was preferred because 20% is near the level at which effects become statistically discernible or measurable in both laboratory and field studies (Suter 1996; Environment Canada and Health Canada 2003), and therefore can be reliably reproduced. However, chronic EC_{20} values are not always readily available; therefore, a protocol shown in Table 6-14 was established to derive chronic EC_{20} values from available data.

Table 6-13: Procedure for Adjusting Test Endpoints to Chronic 20% Effect Concentration

Test Endpoint ^(a)	Adjustment to Chronic EC ₂₀
Chronic EC ₁₀	Multiply by 2
Chronic EC ₂₅	Multiply by 2/2.5
Chronic EC ₃₀	Multiply by 2/3
Chronic EC ₅₀	Multiply by 2/5
Chronic LC ₂₅	Multiply by 0.5
Chronic LC ₅₀	Divide by 4

(a) IC endpoints were treated as EC endpoints.

EC = effect concentration; LC = lethal concentration; IC = inhibition concentration.

Toxicity data for effect endpoints involving growth, reproduction, and survival were selected because they are considered to be relevant to the persistence of aquatic populations. Chronic toxicity data were preferred, and acute data were only considered when chronic data were not sufficient (a minimum of 2 values). If more than 20 chronic EC₂₀ values were available in each taxonomic group, a 5th percentile of the EC₂₀ values was used as a selected TRV. If there were less than 20 chronic EC₂₀ values, the lowest EC₂₀ was used as a selected TRV for the taxonomic category. Calculated values that fell below the CCME or provincial guideline were not considered appropriate as TRVs for aquatic biota and the CCME or provincial values were selected in their place. The selected TRVs for aquatic biota groups are summarized in Table 6-15. For aquatic TRVs that were based on the lowest chronic EC₂₀ value, the reference is provided in Table 6-15.

In some cases, site-specific modifying factors (ambient conditions) may influence the toxicity of a chemical. For example, these modifying factors include water hardness for copper. In these cases, the TRV must be appropriate to the ambient condition.

ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT

ECOLOGICAL RISK ASSESSMENT

The USEPA ECOTOXicology database (ECOTOX) was generally used for the selection of TRVs for aquatic organisms. There were sufficient data available from ECOTOX to derive TRVs for arsenic, copper, and molybdenum. There were limited data available in the USEPA ECOTOX database pertaining to the effects of cobalt, chloride, sulphate, and uranium exposure on aquatic biota. The TRVs for cobalt were selected from a recently published review of toxicological data by Stubblefield et al. (2020), in which a species sensitivity distribution approach was used. The TRVs for chloride were obtained from the CCME (2011a) and the TRVs for sulphate were obtained from the BC MOE (2013). The TRVs for uranium were derived from data available from toxicological reports.



Table 6-14: Selected Toxicity Reference Values for Aquatic Biota

СОРС	Biotic Group	TRV	/ Unit Rationale		Data Source
	Forage fish	0.123	mg/L	Lowest estimated chronic EC ₂₀ (survival)	ECOTOX (Birge et al. 1979)
	Predator fish	0.630	mg/L	5th percentile of estimated chronic EC ₂₀ distribution (n=50)	ЕСОТОХ
Arsenic	Zooplankton	0.340	mg/L	Lowest estimated chronic EC ₂₀ (intoxication)	ECOTOX (May-Passino and Novak 1984)
Arsenic	Benthic invertebrates	0.122	mg/L	5th percentile of estimated chronic EC ₂₀ distribution (n=27)	ЕСОТОХ
	Phytoplankton	0.0192	mg/L	Lowest estimated chronic EC ₂₀ (growth)	ECOTOX (Vocke et al. 1980)
	Aquatic plants	0.252	mg/L	Lowest estimated chronic EC ₂₀ (population)	ECOTOX (Jenner and Janssen-Mommen 1993)
	Forage fish	693	mg/L	Chronic LOEC (survival)	CCME 2011a (Birge et al. 1985 in Elphick et al. 2011a)
	Predator fish	989	mg/L	Chronic EC ₂₅ (reproduction)	CCME 2011a (Beak 1999)
Chloride	Zooplankton	421	mg/L	Chronic EC ₂₅ (reproduction)	CCME 2011a (Elphick et al. 2011a)
	Benthic invertebrates	421	mg/L	Chronic EC ₂₅ (growth)	CCME 2011a (Bartlett 2009 unpublished)
	Phytoplankton	6,066	mg/L	Chronic MATC (growth)	CCME 2011a (Kessler 1974)
	Aquatic plants	3,150	mg/L	Chronic EC ₅₀ (population)	CCME 2011a (Buckley et al. 1996)
	Forage fish	0.314	mg/L	Lowest chronic EC ₂₀ (biomass). Normalized to a hardness of 52 mg/L (lower end of the equation)	Stubblefield et al. 2020
Cobalt	Predator fish	1.791	mg/L	Lowest chronic EC ₂₀ (survival) for Rainbow Trout. Normalized to a hardness of 52 mg/L (lower end of the equation)	Stubblefield et al. 2020
	Zooplankton	0.0111	mg/L	Lowest chronic EC ₂₀ (reproduction)	Stubblefield et al. 2020
	Benthic invertebrates	0.0122	mg/L	Lowest chronic EC ₂₀ (growth). Normalized to a hardness of 52 mg/L (lower end of the equation)	Stubblefield et al. 2020



СОРС	Biotic Group	TRV	Unit	Rationale	Data Source	
	Phytoplankton	0.046	mg/L	Lowest estimated EC ₂₀ (growth)	Stubblefield et al. 2020	
	Aquatic plants	0.0096	mg/L	lowest estimated EC_{20} (growth). Normalized to a hardness of 52 mg/L (lower end of the equation)	Stubblefield et al. 2020	
	Forage fish	0.002	mg/L	5th percentile of estimated chronic EC ₂₀ distribution (n=237)	ECOTOX	
	Predator fish	0.003	mg/L	5th percentile of estimated chronic EC ₂₀ distribution (n=89)	ECOTOX	
Community (h c d)	Zooplankton	0.002	mg/L	5th percentile of estimated chronic EC ₂₀ distribution (n=117)	ECOTOX; CCME	
Copper ^(b,c,d)	Benthic invertebrates	0.002	mg/L	5th percentile of estimated chronic EC ₂₀ distribution (n=264)	ECOTOX; CCME	
	Phytoplankton ^(a)	0.0092	mg/L	5th percentile of estimated chronic EC ₂₀ distribution (n=101)	ECOTOX	
	Aquatic plants ^(a)	0.038	mg/L	5th percentile of estimated chronic EC ₂₀ distribution (n=28)	ECOTOX	
	Forage fish	31	mg/L	Saskatchewan Water Quality Guideline	WSA 2017	
	Predator fish	80	mg/L	lowest estimated EC ₂₀ value	ECOTOX; Goettl et al. (1976); McConnell (1977)	
Molybdenum	Zooplankton	31	mg/L	Saskatchewan Water Quality Guideline	WSA 2017	
	Benthic invertebrates	31	mg/L	Saskatchewan Water Quality Guideline	WSA 2017	
	Phytoplankton	31	mg/L	Saskatchewan Water Quality Guideline	WSA 2017	
	Aquatic plants	31	mg/L	Saskatchewan Water Quality Guideline	WSA 2017	
	Forage fish	2,999	mg/L	Lowest chronic EC ₂₅ (biomass)	BC MOE 2013; PESC data	
	Predator fish	502	mg/L	Lowest chronic LC ₂₅ (survival)	BC MOE 2013; PESC data	
Sulphate	Zooplankton	425	mg/L	Lowest chronic EC ₂₅ (reproduction)	BC MOE 2013; Elphick et al. 2011b	
	Benthic invertebrates	730	mg/L	Lowest chronic LC ₂₅ (survival)	BC MOE 2013; PESC data	
	Phytoplankton	2,660	mg/L	Lowest chronic EC ₂₅ (cell yield)	BC MOE 2013; Elphick et al. 2011b	



СОРС	COPC Biotic Group TRV Unit		Rationale	Data Source	
	Aquatic plants	2,310	mg/L	Lowest chronic EC ₁₀ (frond increase)	BC MOE 2013; PESC data
	Forage fish	1.50	mg/L	Lowest estimated chronic EC ₂₀ (growth)	VST 2004; Liber et al. 2007
	Predator fish	0.550	mg/L	Lowest estimated chronic EC ₂₀ (growth)	VST 2004; Liber et al. 2007
Uranium ^(e)	Zooplankton	0.060	mg/L	Lowest estimated chronic EC ₂₀ (growth)	VST 2004; Liber et al. 2007
Oranium*,	Benthic invertebrates	0.027	mg/L	Lowest estimated chronic EC ₂₀ (growth)	VST 2004; Liber et al. 2007
	Phytoplankton	0.440	mg/L	Geometric mean of 2 EC ₂₅ values	VST 2004; Liber et al. 2007
	Aquatic plants	5.50	mg/L	Geometric mean of 2 EC ₂₅ values	VST 2004; Liber et al. 2007

⁽a) Study specific hardness data were not available for the adjustment of TRVs.



⁽b) The TRV is hardness dependent and is presented as dissolved metal in soft water (hardness of 25 mg CaCO₃/L).

⁽c) Hardness dependent TRVs are presented for hardness of 25 mg CaCO₃/L and may be converted to reflect specific site hardness conditions using the equations presented.

⁽d) The TRVs presented in italics are CCME guidelines used as a default when estimated TRVs are below the recommended guideline.

⁽e) The TRVs are based on hardness of 60 mg/L, other than phytoplankton which is based on hardness of 120 mg/L.

 EC_{XX} = effect concentration for XX% response; LOEC = lowest observed effect concentration; MATC = maximum acceptable toxicant concentration; CaCO₃ = calcium carbonate; TRV = toxicity reference value; PESC = Pacific Environmental Science Centre; CCME = Canadian Council of Ministers of the Environment.

6.3.1.1.1 Arsenic

The TRVs for arsenic were estimated chronic EC_{20} values that were selected based on EC_{50} and lethal concentration (LC)₅₀ values obtained from the USEPA ECOTOX database (Table 6-15). The results suggest that phytoplankton, benthic invertebrates, and forage fish are the most sensitive to arsenic exposure while predator fish are the least sensitive.

6.3.1.1.2 Chloride

Toxicity records were taken from data selected by CCME (2011a) to derive the Canadian Water Quality Guideline value for the protection of fresh water aquatic life. The studies met the minimum primary or secondary requirements for data quality. No EC₂₀ concentrations were available from the CCME data. Low effects levels including EC₂₅, LOEC, and maximum acceptable toxicant concentration were selected preferentially, and the lowest for each aquatic group were selected (Table 6-15). The results suggest that benthic invertebrates are the most sensitive to chloride exposure while phytoplankton are the least sensitive.

6.3.1.1.3 Cobalt

The TRVs for cobalt are chronic EC_{20} values for aquatic animal groups and estimated chronic EC_{20} values for aquatic plant groups from Stubblefield et at. (2020). Stubblefield et at. (2020) conducted a series of acute and chronic toxicity tests with the primary objective to generate data needed to derive international water quality guidelines for cobalt based on USEPA and European Union requirements. Early life stage tests were conducted on three fish species, one zooplankton species, three aquatic invertebrate species, one alga, and one aquatic macrophyte. The study produced chronic EC_{20} values for the aquatic animal species and chronic EC_{10} values for aquatic plants. The TRVs for phytoplankton and aquatic plants were derived from EC_{10} values using a factor of 2 to adjust to an EC_{20} (Table 6-15). The results suggest that zooplankton, benthic invertebrates, and macrophytes are among the most sensitive to cobalt and fish are the least sensitive. The TRVs for fish, benthic invertebrates, and aquatic plants have been adjusted to the lowest hardness (i.e., 52 mg/L) based on the hardness slope from the Federal Environmental Quality Guideline (FEQG) (Environment Canada 2017).

6.3.1.1.4 Copper

The TRVs for copper for forage fish, predator fish, phytoplankton, and aquatic plants are estimated chronic EC₂₀ values. They were estimated based on EC₁₀, EC₂₅, EC₃₀, EC₅₀, LC₂₅, LC₅₀, inhibition concentration (IC₁₀), IC₂₅, and IC₅₀ values obtained from the USEPA ECOTOX database. In the case of zooplankton and benthic invertebrates, the derived TRVs were lower than the existing CCME guideline values; therefore, the CCME values were selected as the TRVs because the CCME guidelines are conservative and protective of all life forms of aquatic species in Canada. The selected TRVs are summarized in Table 6-15 for dissolved copper in soft water conditions (hardness of 25 milligrams of calcium carbonate per litre [mg CaCO₃/L]). The results suggest that zooplankton, benthic invertebrates, forage fish, and predator fish are the most sensitive to copper exposure while aquatic plants are the least sensitive.



According to the USEPA (2007a), the copper hardness-dependent criterion continuous concentration (CCC) can be calculated from the following:

$$CCC = e^{0.8545 \times ln(hardness) - 1.702} \times 0.96$$

A conversion factor of 0.96 was used to convert the total mean concentration of copper to a dissolved copper concentration. For studies on chronic toxicity of copper to aquatic biota that reported the hardness of the water, H1 (in mg/L CaCO₃), the CCC for the hardness used in the study was calculated. The CCC values for two hardness levels provide a ratio that can be used to adjust an EC₂₀ at the test hardness (H1) to an EC₂₀ at some standard hardness (H2), which is relevant to the site. This was used to determine the hardness adjusted TRV at 25 mg CaCO₃/L, H2, using the following relationship:

$$EC_{20}H2 = \frac{CCC_{H2}}{CCC_{H1}} \times EC_{20}H1$$

The EC₂₀ values derived for copper, based on a hardness of 25 mg CaCO₃/L, were compared to the CCME water quality guidelines for the same hardness condition prior to selecting the appropriate TRVs (Table 6-15).

6.3.1.1.5 Sulphate

Toxicity records were taken from data selected by the British Columbia Ministry of Environment (BC MOE 2013) to derive British Columbia's Ambient Water Quality Guideline value for sulphate for the protection of fresh water aquatic life. The BC MOE used data from the Pacific Environmental Science Centre and Dr. Chris Kennedy (Simon Fraser University) and Elphick et al. (2011b). The tests selected by BC MOE were conducted over a range of hardness levels. The BC MOE determined that, though dose-response curves of many organisms were influenced by water hardness, a consistent relationship among the species could not be established. The selected TRVs are LC₂₅ or EC₂₅ values except for the aquatic plant TRV, which is an EC₁₀ value (Table 6-15). The results suggest that aquatic animals are generally more sensitive to sulphate exposure than aquatic plants.

6.3.1.1.6 Molybdenum

For forage fish, zooplankton, benthic invertebrates, phytoplankton, and aquatic plants, the Saskatchewan Water Security Agency WQO for the protection of aquatic life (WSA 2017) was selected as the TRV for molybdenum. The WQO is based on current understanding of aquatic toxicity of molybdenum to fresh-water aquatic organisms as discussed in Section 4.2.3, Screening for Constituents of Potential Concern in Aquatic Environment. The TRV for predator fish is the lowest of three estimated chronic EC₂₀ values, derived from LC₅₀ values for rainbow trout (*Oncorhynchus mykiss*), obtained from the USEPA ECOTOX database. The selected TRV is from Goettl et al. (1976) and was republished in 1977 by McConnell, which provided detailed documentation as to the methods used to generate the data (cited in Tetra Tech 2008).



6.3.1.1.7 Uranium

Limited data were available in the USEPA ECOTOX database pertaining to the effects of uranium exposure on aquatic biota. Data were instead obtained from two reports that investigated the toxicity of uranium to aquatic biota in Northern Saskatchewan, VST (2004) and Liber et al. (2007). The TRVs for uranium are all estimated chronic EC_{20} values derived from EC_{25} , EC_{50} and EC_{25} , and EC_{25} , and EC_{25} , and EC_{25} values.

While uranium speciation and toxicity in fresh water are strongly determined by water characteristics such as hardness, pH, and temperature, the CCME (2011b) does not consider that there is sufficient information available to quantitatively evaluate the influence of these factors. Therefore, the CCME recommends a water quality guideline for uranium that is not hardness dependent. The TRVs in Table 6-15 are therefore considered appropriate for use across a range of hardness and may be conservative for hard water environments because they were derived from tests conducted under soft water conditions (water hardness of 60 mg CaCO₃/L from test studies other than phytoplankton which was based on a study using water hardness of 120 mg/L). The results suggest that zooplankton and benthic invertebrates are more sensitive to uranium exposure and phytoplankton and aquatic plants are less sensitive.

6.3.1.2 Toxicity Reference Values for Terrestrial Biota

Chronic dose-based TRVs for non-radiological COPCs were derived for birds and mammals based on endpoints (i.e., growth and reproduction) considered relevant for assessing the health of wildlife populations. The selected TRVs, shown in Table 6-16, are chronic daily intakes that are not expected to cause adverse effects to a particular ecological receptor. Where the TRV is based a single LOAEL, the specific reference is provided in Table 6-16. Lowest observed adverse effect levels were selected for each COPC. The LOAEL is the lowest exposure level at which the response of a test species in a toxicity study was statistically discernible.

The LOAELs were used in the ERA to identify a threshold of exposure below which adverse effects are not expected. Exceeding a LOAEL does not mean that effects would necessarily occur; rather, it means that effects may occur. Particularly in large populations, localized effects on a few individuals can be compensated such that there is no discernible effect on the population as a whole.

Toxicity data for bird and mammal species were preferentially selected from the USEPA ecological soil screening levels (Eco-SSL) database (USEPA 2005a). There were no data available in the USEPA Eco-SSL database pertaining to the effects of uranium exposure on birds and mammals, so TRVs were derived from data available in toxicological reports previously used in risk assessments for uranium mines in northern Saskatchewan.

Toxicity reference values were derived from the selected data for several test species of avian and mammalian wildlife. When possible, a test species was selected with a close taxonomic relationship to the ecological receptor in the risk assessment, such as within the same order, family, genus, or species. If there were several potential test species relevant to an ecological receptor, consideration was given to similar diet and body size. A sensitive test species of the

same class was selected to represent an ecological receptor when no data were available for species with a closer taxonomic relationship.

Table 6-15: Selected Toxicity Reference Values for Terrestrial Biota

Ecological Receptor	Arsenic (mg/kg/d)	Cobalt (mg/kg/d)	Copper (mg/kg/d)	Molybdenum (mg/kg/d)	Uranium (mg/kg/d)
Mammals					
Black bear	3.1	13	12	2.6	5.6
Beaver	14	13	119	3.8	5.6
Woodland caribou	14	13	1.5	4.1	5.6
Red fox	3.1	13	12	2.6	5.6
Snowshoe hare	3.0	13	46	30	5.6
Little brown myotis	3.1	13	12	2.6	5.6
American mink	3.1	13	12	2.6	5.6
Moose	14	13	1.5	4.1	5.6
Muskrat	14	13	119	3.8	5.6
Southern red-backed vole	21	28	296	2.6	5.6
Grey wolf	3.1	13	12	2.6	5.6
Birds	·				
Grouse	3.6	14	35	39	16
Common loon	3.6	14	27	21	16
Mallard	5.1	14	75	21	16
Canada goose	3.6	14	27	21	16
Rusty blackbird	3.6	14	27	21	16

6.3.1.2.1 Arsenic

A summary of the TRVs selected for mammalian and avian species for arsenic is shown in Table 6-17.

Mammalian Toxicity Reference Values

Data for growth and reproduction for mammalian species were obtained from the Eco-SSL document for arsenic exposure (USEPA 2005b). The data were based on a total of 14 LOAEL values from studies with dogs, goats, guinea pigs, mice, pigs, rabbits, and rats. The geometric means of the LOAELs within species ranged from 0.84 mg/kg/d for a guinea pig to 20.7 mg/kg/d for a mouse. Each of the species mean values of LOAEL can be considered as a TRV for arsenic for other mammals. In the event that a species has no closely related test species, the second lowest LOAEL value of 3.0 mg/kg/d for rabbit and dog can be used as a conservative default for the arsenic TRV. Although this LOAEL is not the minimum of the species LOAELs, it was selected over the minimum LOAEL of 0.84 mg/kg/d for guinea pig because the latter value was essentially at the same level as the minimum NOAEL from the same dataset. Of the total 14 LOAEL values used to derive the species LOAELs only two are below 3.0 mg/kg/d. As such, the LOAEL of 3.0 mg/kg/d was selected as the default LOAEL as it is more representative of the LOAEL data overall.



Avian Toxicity Reference Values

Data for growth and reproduction for avian species were obtained from the Eco-SSL document for arsenic exposure (USEPA 2005b). The document was based on studies with chickens and ducks. After review of the data, two LOAEL values were retained for ducks and one was retained for chicken. The selected avian TRVs are 3.6 mg/kg/d for chickens based on a single LOAEL value, and 5.1 mg/kg/d for duck based on the geometric mean of two LOAEL values.

Table 6-16: Summary of Selected Toxicity Reference Values for Mammal and Bird Test Species - Arsenic

Test Species TRV (mg/kg/d)		Rationale				
Mammal						
Dog (Canis familiaris)	3.1	Geometric mean of two LOAELs for the species				
Goat (Capra hircus)	14	Geometric mean of two LOAELs for the species				
Guinea pig (Cavia porcellus)	0.84	Single LOAEL for the species (Hunder et al. 1999)				
Mouse (Mus musculus)	21	Geometric mean of three LOAELs for the species				
Pig (Sus scrofa)	9.4	Single LOAEL for the species (Morrison and Chavez 1983)				
Rabbit (Oryctolagus cuniculus)	3.0	Single LOAEL for the species (Nemec et al. 1998)				
Rat (Rattus norvegicus)	14	Geometric mean of four LOAELs for the species				
Bird						
Mallard duck (Anas platyrhynchos)	5.1	Geometric mean of two LOAELs for the species				
Chicken (Gallus sp.)	3.6	Single LOAEL for the species (Howell and Hill 1978)				

LOAEL = lowest observed adverse effect level; TRV = toxicity reference value.

6.3.1.2.2 Cobalt

A summary of the TRVs selected for mammalian and avian species for cobalt is shown in Table 6-18.

Mammalian Toxicity Reference Values

Two mammal species were represented in the Eco-SSL database with growth or reproduction endpoints and with LOAEL values reported (USEPA 2005c). The species geometric mean values of LOAELs ranged from 13 mg/kg/d for rat to 28 mg/kg/d for mouse, from 6 and 7 LOAEL values respectively.

Avian Toxicity Reference Values

Data for growth and reproduction for avian species were obtained from the Eco-SSL database for cobalt exposure (USEPA 2005a). The document was based on studies with chickens and ducks. Eight LOAEL values were retained for chicken. The data for ducks were not retained because the LOAEL value was above the mortality endpoint. The selected avian TRV is 14 mg/kg/d for chickens based on the geometric mean of eight LOAEL values.



Table 6-17: Summary of Selected Toxicity Reference Values for Mammal and Bird Test Species - Cobalt

Test Species	TRV (mg/kg/d)	Rationale
Mammal		
Mouse (Mus musculus)	28	Geometric mean of six LOAELs for the species
Rat (Rattus norvegicus)	13	Geometric mean of seven LOAELs for the species
Bird	·	
Chicken (Gallus sp.)	14	Geometric mean of eight LOAELs for the species

LOAEL = lowest observed adverse effect level; TRV = toxicity reference value.

6.3.1.2.3 Copper

A summary of the TRVs selected for mammalian and avian species for copper is shown in Table 6-19.

Mammalian Toxicity Reference Values

The mammalian data for growth and reproduction were obtained from the data presented in the Eco-SSL document for copper exposure (USEPA 2007b). The data were based on studies with goats, minks, mice, pigs, rabbits, rats, and sheep. The TRVs for goat, rabbit, and sheep are each based on a single LOAEL value. The TRVs for mink, mouse, pig, and rat are geometric means of the LOAEL data for each test species. For rats, the geometric mean of the NOAEL values was larger than the geometric mean of the LOAEL values due to the larger dataset for the LOAEL values. For this species, the LOAEL was therefore derived only from the studies that had both LOAEL and NOAEL values. The geometric mean of the LOAELs ranged from 1.47 mg/kg/d for a goat to 296 mg/kg/d for a mouse. In the event that a species had no closely related test species, the LOAEL of 8.8 mg/kg/d for pig was used as the default mammalian TRV. It is not the lowest LOAEL; however, it can be used as the default mammalian TRV as it is the lowest species LOAEL above the lowest NOAEL from the same dataset.

Avian Toxicity Reference Values

The avian data for growth and reproduction were obtained from the data presented in the Eco-SSL document for copper exposure (USEPA 2007b). The data were based on studies with chickens, ducks, and turkeys. The geometric means of the LOAELs within species were selected to serve as the TRVs. The geometric means of the LOAELs for chickens, ducks, and turkeys were 34.9 mg/kg/d, 75.2 mg/kg/d, and 27 mg/kg/d respectively, based on 78, 3, and 9 LOAEL values, respectively. These values were used as TRVs for other similar species in the EcoRA.



Table 6-18: Summary of Selected Toxicity Reference Values for Mammal and Bird Test Species - Copper

Test Species	TRV (mg/kg/d)	Rationale
Mammal		
Goat (Capra hircus)	1.5	Single LOAEL for the species (Solaiman et al. 2001)
Mink (Neovison vison)	12	Geometric mean of two LOAELs for the species
Mouse (Mus musculus)	296	Geometric mean of five LOAELs for the species
Pig (Sus scrofa)	8.8	Geometric mean of four LOAELs for the species
Rabbit (Oryctolagus cuniculus)	46	Single LOAEL for the species (Grobner et al. 1986)
Rat (Rattus norvegicus)	119	Geometric mean of five LOAELs for the species
Sheep (Ovis aires)	3.0	Single LOAEL for the species (Ortolani et al. 2003)
Bird		
Chicken (Gallus sp.)	35	Geometric mean of 78 LOAELs for the species
Duck (Anas platyrhynchos)	75	Geometric mean of three LOAELs for the species
Turkey (Meleagris gallopavo)	27	Geometric mean of nine LOAELs for the species

LOAEL = lowest observed adverse effect level; TRV = toxicity reference value.

6.3.1.2.4 Molybdenum

A summary of the TRVs selected for mammalian and avian species for molybdenum is shown in Table 6-20.

Mammalian Toxicity Reference Values

The selected TRVs for mammals are from studies with reported LOAEL values for growth and reproduction endpoints. Relevant LOAEL values for molybdenum were obtained from literature for four mammal species: rabbit (*Oryctolagus cunculus*), mouse (*Mus musculus*), cow (*Bos taurus*), and rat (*Rattus norvegicus*). The TRVs for rabbit, mouse, and cow are based on one LOAEL value and the TRV for rat is the geometric mean of two LOAEL values. The TRVs range from 2.6 mg/kg/d for a mouse to 30 mg/kg/d for a rabbit. In the event that a species had no closely related test species, the LOAEL of 2.6 mg/kg/d for mouse was used as the default mammalian TRV.

Avian Toxicity Reference Values

The selected TRVs for birds are from studies with reported LOAEL values for growth and reproduction endpoints. Relevant LOAEL values for molybdenum were obtained from literature for two avian species: chicken (*Gallus* sp.) and turkey (*Meleagris gallopavo*). The TRV for chicken is the geometric mean of three LOAEL values and the TRV for turkey is based on one LOAEL value from Underwood (1971). The TRVs range from 21 mg/kg/d for a turkey to 39 mg/kg/d for a chicken. In the event that a species had no closely related test species, the LOAEL of 21 mg/kg/d for turkey was used as the default avian TRV.



Table 6-19: Summary of Selected Toxicity Reference Values for Mammal and Bird Test Species – Molybdenum

Test Species	TRV (mg/kg/d)	Rationale
Mammal		
Rabbit (Oryctolagus cunculus)	30	Single LOAEL for the species (Arrington and Davis 1953)
Mouse (Mus musculus)	2.6	Single LOAEL for the species (Schroeder and Mitchener 1971)
Cow (Bos taurus)	4.1	Single LOAEL for the species (Thomas and Moss 1951)
Rat (Rattus norvegicus)	3.8	Geometric mean of two LOAELs for the species
Bird		
Chicken (Gallus domesticus)	39	Geometric mean of three LOAELs for the species
Turkey (Melagris gallopavo)	21	Single LOAEL for the species (Underwood 1971)

LOAEL = lowest observed adverse effect level; TRV = toxicity reference value.

6.3.1.2.5 Uranium

A summary of the TRVs selected for mammalian and avian species for uranium is shown in Table 6-21.

Mammalian Toxicity Reference Values

There is no Eco-SSL document for uranium. Previous risk assessments have used TRVs for mammalian exposure to uranium derived from the data of Sample et al. (1996). The Sample et al. (1996) data for mammalian species were based on a study by Paternain et al. (1989) related to reproduction in mice. Sample et al. derived a LOAEL of 6.13 mg/kg/d from the study. The TRV quoted by Sample et al. (1996) contains a small unit conversion error. Instead of 6.13 mg/kg/d as reported, the value should be 5.6 mg/kg/d. The difference arises from Sample's use of uranyl acetate molecular weight rather that uranyl acetate dehydrate molecular weight in converting the molecular dose to uranium dose. The correct value (5.6 mg/kg/d) can be found in the ATSDR toxicity profile for uranium. It represents the oral dose (to parents) at which viability of F1 offspring was reduced. Since the LOAEL value from Paternain et al. (1989) is the only mammalian data available, this LOAEL value of 5.6 mg/kg/d was selected for evaluating uranium toxicity to mammalian species.

Avian Toxicity Reference Values

No data were available in the Eco-SSL database related to uranium exposure in avian species. Previous risk assessments used TRVs for avian exposure to uranium derived from the data of Sample et al. (1996). The Sample et al. data were based on a study by Haseltine and Sileo (1983) related to mortality, body weight, liver, and kidney effects in ducks; Sample et al. derived a NOAEL of 16 mg/kg/d, but no LOAEL from the study. There are no other avian data available for uranium. The study used powdered metallic uranium. Uranium in this form would likely be oxidized to ionic form in the gut since uranium is strongly reducing in aqueous systems (Durante and Pugliese 2002). Uranium in the environment similarly exists in an oxidized ionic form. Solubility differences



among ionic forms in the gut can be bounded. The ICRP (1994) has determined that some oxidized species in the gut may be an order of magnitude less soluble than the most soluble species. Any reduced solubility in the gut would be offset by the fact that a NOAEL value is used in the absence of a LOAEL value. Since the NOAEL value from Haseltine and Sileo (1983) is the only avian data available, this NOAEL value of 16 mg/kg/d was selected as the TRV for evaluating uranium toxicity to avian species.

Table 6-20: Summary of Selected Toxicity Reference Values for Mammal and Bird Test Species - Uranium

Test Species	TRV (mg/kg/d)	Rationale
Mammal		
Mouse (Mus musculus)	5.6	Paternain et al. 1989, from Sample et al. 1996
Bird		
Black duck (Anas rubripes)	16	Haseltine and Sileo 1983, from Sample et al. 1996

TRV = toxicity reference value.

6.3.2 Radiation Benchmarks

Radiation dose benchmarks of 0.4 mGy/h (9.6 mGy/d) and 0.1 mGy/h (2.4 mGy/d; UNSCEAR [2008] were selected for the assessment of effects on aquatic biota and terrestrial biota, respectively, as recommended in CSA N288.6-22), and are summarized in Table 6-22. These are total dose benchmarks; therefore, the doses to biota due to each radionuclide of concern are summed to compare against these benchmarks. The aquatic biota considered by the United Nations Scientific Committee on Effects of Atomic Radiation are organisms such as fish and benthic invertebrates that reside in water. Birds and mammals with riparian habits are considered to be terrestrial biota. Dose calculations in this ERA follow the same convention.

Exceedance of either of these benchmarks is considered to indicate the potential for adverse effects to occur, and the need for more detailed assessment.

Table 6-21: Radiological Dose Benchmarks for Aquatic and Terrestrial Ecological Receptors

Ecological Receptor	Dose Rate Limit (mGy/d)
Aquatic Organism	
Aquatic plants (algae and macrophytes)	9.6
Benthic invertebrates	9.6
Fish	9.6
Terrestrial Organism	
Terrestrial animals	2.4
Terrestrial plants	2.4

mGy/d = milligrays per day.



6.3.3 Uncertainty in the Effects Assessment

Uncertainties associated with the estimation of ecotoxicological effect levels for COPCs are inherent in the ERA process. Many uncertainties are associated with the use of literature-based TRVs. These uncertainties may include: extrapolation of results from laboratory tests to the field, differences in sensitivity between the test organism and resident organisms, laboratory conditions that are not representative of field conditions, and the form of the COPC used in toxicity testing, which may not be representative of the form that would be found at the Project.

The use of TRVs from laboratory studies tends to be conservative because these studies are typically chemical-specific and use highly bioavailable forms of the COPC studied. In field situations, the chemical form of the same COPC may be less bioavailable, and toxicity-modifying factors may be present that were not acting in laboratory tests.

The EC₂₀ values were used for aquatic biota to reduce uncertainty by representing a standard threshold level of low magnitude effects. Depending on the size of the available dataset, selection of the 5th percentile or lowest EC₂₀ value as the TRV for an aquatic biota group was intended to reduce the likelihood that risks would be underestimated.

There is inherent uncertainty associated with the use of LELs such as LOAEL values as TRVs for birds and mammals as these values are not precisely related to a particular magnitude of effect. However, LOAEL values have widespread use in the risk assessment community and the science is not currently available to change this approach to TRVs. Defaulting to the most conservative TRV for ecological receptors that are not closely related taxonomically to the test species was meant to reduce the likelihood that risks would be underestimated.

6.4 Risk Characterization

Risk assessment is the process of estimating the likelihood of undesirable effects on ecological health resulting from exposure to chemical or radiological contaminants. Three components must be present for risks to ecological health to exist:

- The COPC must be present at concentrations sufficient to cause a possible adverse effect.
- A receptor must be present.
- There must be a complete exposure pathway by which the receptor can come into contact with the COPC.

6.4.1 Risk Estimation

Risk characterization is the process in the EcoRA that integrated the results from the exposure and effects assessments to estimate the risk of adverse effects on ecological receptors. The risk characterization also evaluated the uncertainties associated with the overall conclusion of risk.



ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT ECOLOGICAL RISK ASSESSMENT

The hazard quotient (HQ) is a simple approach that provides a quantitative estimate of overall risk. The HQ is the ratio between the exposure estimate and a TRV:

$HQ = \underline{Exposure (Dose) Estimate}$ TRV

If the HQ is less than or equal to 1, this suggests low risk to the ecological receptor because exposure estimates are below levels known to cause adverse effects. If the HQ is greater than 1, adverse effects may be possible, and further investigation of the assumptions of the exposure and effects assessments could be considered to reduce the conservatism inherent in the EcoRA. Risk assessment is often an iterative process of refining information, and the identified risk in the first iteration is often an artefact of conservative assumptions. If further evaluation under more realistic assumptions confirms the risk, this information can be used to inform mitigation to avoid, eliminate, or reduce the source of the risk.

6.4.1.1 Estimated Risk to Ecological Receptors – Application Case

This subsection presents the estimated risks to aquatic and terrestrial ecological receptors due to releases from the Project during all phases, including the far-future projection once groundwater solutes have been released to Patterson Lake. The results are presented for both the Application Case and the Upper Bound sensitivity scenario.

6.4.1.1.1 Non-radiological Risk

The estimated HQs for all aquatic and terrestrial ecological receptors at all assessed locations during Operations and the far-future projection are shown in Table 6-23 and Table 6-24. The HQs represent the maximum HQ over the Project phase for the COPCs of interest, which is a conservative representation of risk.

No significant adverse effect on either aquatic or terrestrial populations or communities as a result of releases from the Project would be likely during Operations, for both the Application Case and reasonable Upper Bound sensitivity scenario. All estimated HQs for all COPCs (i.e., chloride, sulphate, arsenic, cobalt, copper, molybdenum, and uranium) for all ecological receptors remain below the HQ benchmark of 1.

For assessment of risk to benthic invertebrates, risk was calculated based on toxicity benchmarks as water concentrations. However, considering that benthic invertebrates also reside in sediment, a comparison of predicted sediment concentrations against sediment toxicity benchmarks was warranted. This only applied to arsenic and molybdenum in sediment, as no other COPC exceeded sediment screening values (Table 4-3). Arsenic in sediment at Patterson Lake North Arm – West Basin was predicted to exceed the REF screening value from Burnett-Seidel and Liber (2013), but was below the NE2 value for the Application Case and reasonable upper bound sensitivity scenario. Arsenic was below the sediment screening values at all other downstream locations. Concentrations below the NE2 value for arsenic indicate that benthic invertebrate community metrics (abundance, richness, and evenness) downstream of discharges are not expected to differ significantly (i.e., by less than 20%) from those observed at natural background conditions.



ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT ECOLOGICAL RISK ASSESSMENT

Since there were no HQ exceedances for any of the aquatic or terrestrial receptors during Operations, individual species at risk would also be considered protected.

For the far-future projection, once groundwater solutes reach Patterson Lake North Arm – West Basin HQs would remain below the benchmark of 1 for all COPCs with the exception of copper for both the Application Case and reasonable upper bound sensitivity scenario. Exceedances of the HQ benchmark value of 1 for the far-future projection were limited spatially to Patterson Lake and were limited in magnitude to a maximum of 1.4 times the benchmark for the reasonable upper bound sensitivity scenario. Exceedances of the HQ benchmark were predicted for some groups of aquatic animals but not for aquatic plants:

- 1) The estimated HQ for copper exceeded 1 in Patterson Lake North Arm West Basin for lake whitefish (1.1), aquatic invertebrates (1.1), and zooplankton (1.1) in the far-future projection for the Application Case. All other waterbodies in the downstream environment have HQs for copper below 1.
- 2) The estimated HQ for copper exceeded 1 in Patterson Lake North Arm West Basin for lake whitefish (1.4), aquatic invertebrates (1.4), and zooplankton (1.4) in the far-future projection for the reasonable upper bound sensitivity scenario. In Patterson Lake South Arm the HQ is at 1 for lake whitefish, aquatic invertebrates, and zooplankton. All other waterbodies in the downstream environment had HQs for copper below 1.
- The estimated HQs for all COPCs for all birds and mammals at all locations remained below 1 for both the Application Case and reasonable upper bound sensitivity scenario.
- 4) Molybdenum in sediment at Patterson Lake North Arm West Basin was predicted to exceed its REF screening value from Burnett-Seidel and Liber (2013) but was below its NE2 value for the far-future projection reasonable upper bound sensitivity scenario. Molybdenum was below the sediment screening values at all other downstream locations. Concentrations below the NE2 value for molybdenum indicate that benthic invertebrate community metrics (abundance, richness, and evenness) downstream of discharges are not expected to differ significantly (i.e., by less than 20%) from those observed at natural background conditions.

An HQ that is slightly above a value of one is not indicative of impending risk to aquatic animal communities at Patterson Lake North Arm – West Basin. The fresh water copper TRVs used for lake whitefish, benthic invertebrates, and zooplankton were all 0.002 mg/L (Section 6.3.1, Toxicological Benchmarks). These TRV correspond to the CCME's default water quality guideline for copper.

The CCME water quality guideline value is a hardness-based equation, but the default water quality guideline of 0.002 mg/L applies to soft water conditions, below a hardness value of 82 mg CaCO₃/L. In the far-future projection, water hardness in Patterson Lake North Arm – West Basin is expected to be near baseline levels, at or below 25 mg CaCO₃/L; therefore, the default CCME water quality guideline is applicable. It is a conservative benchmark for copper below which adverse



ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT ECOLOGICAL RISK ASSESSMENT

population level effects for most aquatic biota are not expected to occur. The most sensitive species include fresh water snails (*Lymnaea stagnalis* and *Pyrgulopsis robusta*), bivalves (*Villosa iris*), and water flea (*Daphnia magna*; ECCC 2021). The most sensitive endpoints for chronic exposure to copper include growth for benthic invertebrates, reproduction for zooplankton, and growth and reproduction for fish (ECCC 2021). The CCME water quality guideline is based on no or low effect levels for the most sensitive species; thus, the marginal TRV exceedances predicted by HQs just above 1 indicate a possible slight reduction in growth or reproduction for exposed individuals of sensitive species in Patterson Lake North Arm – West Basin. Additionally, since the HQ exceedance is localized to Patterson Lake North Arm – West Basin, population level effects are not expected.

For benthic invertebrates that are in contact with lake sediments, the far-future projection for copper in sediment in Patterson Lake North Arm – West Basin (6.51 mg/kg dw for Application Case and 8.52 mg/kg dw for Upper Bound) was well below the sediment quality LEL guideline for copper (22 mg/kg dw; Thompson et al. 2005), indicating that benthic invertebrates are unlikely to be adversely affected by copper at the predicted exposure level.

Additional evaluation of this copper risk was completed in the aquatic health assessment (EIS Appendix 11A, Aquatic Health Assessment of the Potential for Adverse Effects of Predicted Far-Future Copper Concentrations in Patterson Lake), with the objective of informing an adaptive management plan for assessing and reducing uncertainty related to waste rock storage area seepage and increasing mitigations if necessary.



Table 6-22: Non-radiological Risk to Ecological Receptors – Application Case and Upper Bound Scenario – Operations

		Thore to Eoological Recopt	1.						Maximum HC	Operations						
	Biota	Location				Application Cas	e		· · · · · · · · · · · · · · · · · · ·	Operations		Upp	er Bound Scenar	io		
			Chloride	Sulphate	Arsenic	Cobalt	Copper	Molybdenum	Uranium	Chloride	Sulphate	Arsenic	Cobalt	Copper	Molybdenum	Uranium
		Reference (Broach Lake)	1.00 × 10 ⁻⁰³	1.45 × 10 ⁻⁰³	8.50 × 10 ⁻⁰⁴	4.75 x 10 ⁻⁰²	4.18 × 10 ⁻⁰¹	6.61 × 10 ⁻⁰⁶	3.57 × 10 ⁻⁰³	1.00 × 10 ⁻⁰³	1.45 × 10 ⁻⁰³	8.50 × 10 ⁻⁰⁴	4.75 x 10 ⁻⁰²	4.18 × 10 ⁻⁰¹	6.61 × 10 ⁻⁰⁶	3.57 × 10 ⁻⁰³
	Benthic	Patterson Lake North Arm – West Basin	1.03 × 10 ⁻⁰²	2.93 × 10 ⁻⁰¹	2.59 × 10 ⁻⁰³	6.15 x 10 ⁻⁰²	4.86 × 10 ⁻⁰¹	1.77 × 10 ⁻⁰⁵	1.17 × 10 ⁻⁰²	2.38 × 10 ⁻⁰¹	2.96 × 10 ⁻⁰¹	2.64 × 10 ⁻⁰³	6.20 x 10 ⁻⁰²	4.88 × 10 ⁻⁰¹	4.03 × 10 ⁻⁰⁵	2.63 × 10 ⁻⁰²
	Invertebrate	Patterson Lake South Arm	7.52 × 10 ⁻⁰³	1.99 × 10 ⁻⁰¹	9.94 × 10 ⁻⁰⁴	5.49 x 10 ⁻⁰²	4.53 × 10 ⁻⁰¹	1.23 × 10 ⁻⁰⁵	5.73 × 10 ⁻⁰³	1.70 × 10 ⁻⁰¹	2.01 × 10 ⁻⁰¹	9.98 × 10 ⁻⁰⁴	5.52 x 10 ⁻⁰²	4.54 × 10 ⁻⁰¹	2.43 × 10 ⁻⁰⁵	9.60 × 10 ⁻⁰³
		Beet Lake	4.69 × 10 ⁻⁰³	1.13 × 10 ⁻⁰¹	8.74 × 10 ⁻⁰⁴	5.14 x 10 ⁻⁰²	4.36 × 10 ⁻⁰¹	9.53 × 10 ⁻⁰⁶	4.36 × 10 ⁻⁰³	9.63 × 10 ⁻⁰²	1.14 × 10 ⁻⁰¹	8.74 × 10 ⁻⁰⁴	5.15 x 10 ⁻⁰²	4.37 × 10 ⁻⁰¹	1.57 × 10 ⁻⁰⁵	5.78 × 10 ⁻⁰³
		Lloyd Lake	1.39 × 10 ⁻⁰³	1.33 × 10 ⁻⁰²	8.51 × 10 ⁻⁰⁴	4.79 x 10 ⁻⁰²	4.20 × 10 ⁻⁰¹	6.91 × 10 ⁻⁰⁶	3.64×10^{-03}	1.10 × 10 ⁻⁰²	1.34 × 10 ⁻⁰²	8.52 × 10 ⁻⁰⁴	4.79 x 10 ⁻⁰²	4.20×10^{-01}	7.54 × 10 ⁻⁰⁶	3.77×10^{-03}
		Reference (Broach Lake)	1.00 × 10 ⁻⁰³	2.50×10^{-03}	3.64 × 10 ⁻⁰⁴	5.25 x 10 ⁻⁰²	4.21 × 10 ⁻⁰¹	6.66 × 10 ⁻⁰⁶	1.67×10^{-03}	1.00 × 10 ⁻⁰³	2.50×10^{-03}	3.64 × 10 ⁻⁰⁴	5.25 x 10 ⁻⁰²	4.21×10^{-01}	6.66 × 10 ⁻⁰⁶	1.67×10^{-03}
		Patterson Lake North Arm – West Basin	1.03 × 10 ⁻⁰²	5.04 × 10 ⁻⁰¹	1.28 × 10 ⁻⁰³	6.94 x 10 ⁻⁰²	4.97 × 10 ⁻⁰¹	1.93 × 10 ⁻⁰⁵	6.39 × 10 ⁻⁰³	2.38 × 10 ⁻⁰¹	5.09 × 10 ⁻⁰¹	1.31 × 10 ⁻⁰³	7.01 x 10 ⁻⁰²	5.00 × 10 ⁻⁰¹	4.48 × 10 ⁻⁰⁵	1.54 × 10 ⁻⁰²
	Zooplankton	Patterson Lake South Arm	7.52 × 10 ⁻⁰³	3.42 × 10 ⁻⁰¹	4.40 × 10 ⁻⁰⁴	6.17 x 10 ⁻⁰²	4.61 × 10 ⁻⁰¹	1.31 × 10 ⁻⁰⁵	2.94×10^{-03}	1.70 × 10 ⁻⁰¹	3.45 × 10 ⁻⁰¹	4.42 × 10 ⁻⁰⁴	6.21 x 10 ⁻⁰²	4.62 × 10 ⁻⁰¹	2.63 × 10 ⁻⁰⁵	5.16 × 10 ⁻⁰³
		Beet Lake	4.69 × 10 ⁻⁰³	1.94×10^{-01}	3.76 × 10 ⁻⁰⁴	5.73 x 10 ⁻⁰²	4.42 × 10 ⁻⁰¹	9.97 × 10 ⁻⁰⁶	2.14×10^{-03}	9.63 × 10 ⁻⁰²	1.96 × 10 ⁻⁰¹	3.77×10^{-04}	5.75 x 10 ⁻⁰²	4.42×10^{-01}	1.68 × 10 ⁻⁰⁵	2.95×10^{-03}
Aquatic		Lloyd Lake	1.39 × 10 ⁻⁰³	2.29×10^{-02}	3.65 × 10 ⁻⁰⁴	5.30 x 10 ⁻⁰²	4.23×10^{-01}	7.00×10^{-06}	1.71×10^{-03}	1.10 × 10 ⁻⁰²	2.30×10^{-02}	3.65 × 10 ⁻⁰⁴	5.30 x 10 ⁻⁰²	4.24×10^{-01}	7.70×10^{-06}	1.79×10^{-03}
Animals		Reference (Broach Lake)	4.27 × 10 ⁻⁰⁴	2.11×10^{-03}	1.96 × 10 ⁻⁰⁴	3.26 x 10 ⁻⁰⁴	2.81 × 10 ⁻⁰¹	2.58 × 10 ⁻⁰⁶	1.82×10^{-04}	4.27×10^{-04}	2.11 × 10 ⁻⁰³	1.96 × 10 ⁻⁰⁴	3.26 x 10 ⁻⁰⁴	2.81 × 10 ⁻⁰¹	2.58 × 10 ⁻⁰⁶	1.82×10^{-04}
	Nambana Bila	Patterson Lake North Arm – West Basin	4.39 × 10 ⁻⁰³	4.27 × 10 ⁻⁰¹	6.92 × 10 ⁻⁰⁴	4.30 x 10 ⁻⁰⁴	3.32 × 10 ⁻⁰¹	7.48 × 10 ⁻⁰⁶	6.97 × 10 ⁻⁰⁴	1.01 × 10 ⁻⁰¹	4.31 × 10 ⁻⁰¹	7.09 × 10 ⁻⁰⁴	4.34 x 10 ⁻⁰⁴	3.33 × 10 ⁻⁰¹	1.74 × 10 ⁻⁰⁵	1.68 × 10 ⁻⁰³
	Northern Pike	Patterson Lake South Arm	3.20 × 10 ⁻⁰³	2.90×10^{-01}	2.37×10^{-04}	3.82 x 10 ⁻⁰⁴	3.08×10^{-01}	5.08 × 10 ⁻⁰⁶	3.21×10^{-04}	7.24 × 10 ⁻⁰²	2.92 × 10 ⁻⁰¹	2.38×10^{-04}	3.85 x 10 ⁻⁰⁴	3.08×10^{-01}	1.02 × 10 ⁻⁰⁵	5.63×10^{-04}
		Beet Lake	2.00 × 10 ⁻⁰³	1.64 × 10 ⁻⁰¹	2.03 × 10 ⁻⁰⁴	3.55 x 10 ⁻⁰⁴	2.95 × 10 ⁻⁰¹	3.86 × 10 ⁻⁰⁶	2.33×10^{-04}	4.10 × 10 ⁻⁰²	1.66 × 10 ⁻⁰¹	2.03 × 10 ⁻⁰⁴	3.56 x 10 ⁻⁰⁴	2.95 × 10 ⁻⁰¹	6.51 × 10 ⁻⁰⁶	3.22×10^{-04}
		Lloyd Lake	5.92 × 10 ⁻⁰⁴	1.93 × 10 ⁻⁰²	1.97 × 10 ⁻⁰⁴	3.29 x 10 ⁻⁰⁴	2.82 × 10 ⁻⁰¹	2.71 × 10 ⁻⁰⁶	1.87×10^{-04}	4.70×10^{-03}	1.95 × 10 ⁻⁰²	1.97 × 10 ⁻⁰⁴	3.29 x 10 ⁻⁰⁴	2.82 × 10 ⁻⁰¹	2.98 × 10 ⁻⁰⁶	1.95 × 10 ⁻⁰⁴
		Reference (Broach Lake)	6.09 × 10 ⁻⁰⁴	3.54×10^{-04}	9.65 × 10 ⁻⁰⁴	1.85 x 10 ⁻⁰³	4.21 × 10 ⁻⁰¹	6.64 x 10 ⁻⁰⁶	6.62×10^{-05}	6.09 × 10 ⁻⁰⁴	3.54 × 10 ⁻⁰⁴	9.65 × 10 ⁻⁰⁴	1.85 x 10 ⁻⁰³	4.21 × 10 ⁻⁰¹	6.64 x 10 ⁻⁰⁶	6.62×10^{-05}
	Whitefish	Patterson Lake North Arm – West Basin	6.27 × 10 ⁻⁰³	7.14 × 10 ⁻⁰²	3.30 × 10 ⁻⁰³	2.44 x 10 ⁻⁰³	4.95 × 10 ⁻⁰¹	1.89 x 10 ⁻⁰⁵	2.44 × 10 ⁻⁰⁴	1.45 × 10 ⁻⁰¹	7.21 × 10 ⁻⁰²	3.38 × 10 ⁻⁰³	2.46 x 10 ⁻⁰³	4.97 × 10 ⁻⁰¹	4.37 x 10 ⁻⁰⁵	5.81 × 10 ⁻⁰⁴
	willtelisii	Patterson Lake South Arm	4.57 × 10 ⁻⁰³	4.85 × 10 ⁻⁰²	1.16 × 10 ⁻⁰³	2.17 x 10 ⁻⁰³	4.59 × 10 ⁻⁰¹	1.29 x 10 ⁻⁰⁵	1.14 × 10 ⁻⁰⁴	1.03 × 10 ⁻⁰¹	4.89 × 10 ⁻⁰²	1.16 × 10 ⁻⁰³	2.18 x 10 ⁻⁰³	4.60×10^{-01}	2.58 x 10 ⁻⁰⁵	1.98×10^{-04}
		Beet Lake	2.85 × 10 ⁻⁰³	2.75 × 10 ⁻⁰²	9.97 × 10 ⁻⁰⁴	2.02 x 10 ⁻⁰³	4.40×10^{-01}	9.86 x 10 ⁻⁰⁶	8.38 × 10 ⁻⁰⁵	5.85 × 10 ⁻⁰²	2.78 × 10 ⁻⁰²	9.97 × 10 ⁻⁰⁴	2.02 x 10 ⁻⁰³	4.41×10^{-01}	1.65 x 10 ⁻⁰⁵	1.15 × 10 ⁻⁰⁴
		Lloyd Lake	8.45 × 10 ⁻⁰⁴	3.24×10^{-03}	9.67 × 10 ⁻⁰⁴	1.87 x 10 ⁻⁰³	4.23 × 10 ⁻⁰¹	6.98 x 10 ⁻⁰⁶	6.78×10^{-05}	6.71 × 10 ⁻⁰³	3.27×10^{-03}	9.67 × 10 ⁻⁰⁴	1.87 x 10 ⁻⁰³	4.23×10^{-01}	7.66 x 10 ⁻⁰⁶	7.07×10^{-05}
		Reference (Broach Lake)	1.34 × 10 ⁻⁰⁴	4.59 × 10 ⁻⁰⁴	4.91 × 10 ⁻⁰⁴	6.07 x 10 ⁻⁰²	2.22 × 10 ⁻⁰²	6.66 × 10 ⁻⁰⁶	1.82×10^{-05}	1.34 × 10 ⁻⁰⁴	4.59 × 10 ⁻⁰⁴	4.91 × 10 ⁻⁰⁴	6.07 x 10 ⁻⁰²	2.22 × 10 ⁻⁰²	6.66 × 10 ⁻⁰⁶	1.82×10^{-05}
	Macrophytes	Patterson Lake North Arm – West Basin	1.38 × 10 ⁻⁰³	9.26 × 10 ⁻⁰²	1.73 × 10 ⁻⁰³	8.03 x 10 ⁻⁰²	2.62 × 10 ⁻⁰²	1.93 × 10 ⁻⁰⁵	6.97 × 10 ⁻⁰⁵	3.18 × 10 ⁻⁰²	9.34 × 10 ⁻⁰²	1.77 × 10 ⁻⁰³	8.11 x 10 ⁻⁰²	2.63 × 10 ⁻⁰²	4.48 × 10 ⁻⁰⁵	1.68 × 10 ⁻⁰⁴
	Macrophytes	Patterson Lake South Arm	1.01 × 10 ⁻⁰³	6.28 × 10 ⁻⁰²	5.93 × 10 ⁻⁰⁴	7.13 x 10 ⁻⁰²	2.43 × 10 ⁻⁰²	1.31 × 10 ⁻⁰⁵	3.21 × 10 ⁻⁰⁵	2.27 × 10 ⁻⁰²	6.34 × 10 ⁻⁰²	5.96 × 10 ⁻⁰⁴	7.17 x 10 ⁻⁰²	2.43×10^{-02}	2.63 × 10 ⁻⁰⁵	5.63×10^{-05}
		Beet Lake	6.27 × 10 ⁻⁰⁴	3.56 × 10 ⁻⁰²	5.08 × 10 ⁻⁰⁴	6.62 x 10 ⁻⁰²	2.33 × 10 ⁻⁰²	9.97 × 10 ⁻⁰⁶	2.33 × 10 ⁻⁰⁵	1.29 × 10 ⁻⁰²	3.60 × 10 ⁻⁰²	5.08 × 10 ⁻⁰⁴	6.65 x 10 ⁻⁰²	2.33 × 10 ⁻⁰²	1.68 × 10 ⁻⁰⁵	3.22×10^{-05}
Aquatic		Lloyd Lake	1.86 × 10 ⁻⁰⁴	4.20×10^{-03}	4.92 × 10 ⁻⁰⁴	6.13 x 10 ⁻⁰²	2.23 × 10 ⁻⁰²	7.00 × 10 ⁻⁰⁶	1.87 × 10 ⁻⁰⁵	1.48 × 10 ⁻⁰³	4.23×10^{-03}	4.92 × 10 ⁻⁰⁴	6.13 x 10 ⁻⁰²	2.23 × 10 ⁻⁰²	7.70×10^{-06}	1.95 × 10 ⁻⁰⁵
Plants		Reference (Broach Lake)	6.96 × 10 ⁻⁰⁵	3.99×10^{-04}	6.44 × 10 ⁻⁰³	1.27 x 10 ⁻⁰²	9.16 × 10 ⁻⁰²	6.66 × 10 ⁻⁰⁶	2.28 × 10 ⁻⁰⁴	6.96 × 10 ⁻⁰⁵	3.99×10^{-04}	6.44 × 10 ⁻⁰³	1.27 x 10 ⁻⁰²	9.16 × 10 ⁻⁰²	6.66 × 10 ⁻⁰⁶	2.28×10^{-04}
	Phytoplankton	Patterson Lake North Arm – West Basin	7.16 × 10 ⁻⁰⁴	8.05 × 10 ⁻⁰²	2.27 × 10 ⁻⁰²	1.68 x 10 ⁻⁰²	1.08 × 10 ⁻⁰¹	1.93 × 10 ⁻⁰⁵	8.72 × 10 ⁻⁰⁴	1.65 × 10 ⁻⁰²	8.13 × 10 ⁻⁰²	2.33 × 10 ⁻⁰²	1.69 x 10 ⁻⁰²	1.09 × 10 ⁻⁰¹	4.48 × 10 ⁻⁰⁵	2.10 × 10 ⁻⁰³
	Thytoplankton	Patterson Lake South Arm	5.22 × 10 ⁻⁰⁴	5.47 × 10 ⁻⁰²	7.79 × 10 ⁻⁰³	1.49 x 10 ⁻⁰²	1.00 × 10 ⁻⁰¹	1.31 × 10 ⁻⁰⁵	4.01×10^{-04}	1.18 × 10 ⁻⁰²	5.52 × 10 ⁻⁰²	7.83×10^{-03}	1.50 x 10 ⁻⁰²	1.01×10^{-01}	2.63 × 10 ⁻⁰⁵	7.03×10^{-04}
		Beet Lake	3.26 × 10 ⁻⁰⁴	3.10 × 10 ⁻⁰²	6.66 × 10 ⁻⁰³	1.38 x 10 ⁻⁰²	9.61 × 10 ⁻⁰²	9.97 × 10 ⁻⁰⁶	2.92 × 10 ⁻⁰⁴	6.68 × 10 ⁻⁰³	3.13 × 10 ⁻⁰²	6.67 × 10 ⁻⁰³	1.39 x 10 ⁻⁰²	9.62 × 10 ⁻⁰²	1.68 × 10 ⁻⁰⁵	4.03 × 10 ⁻⁰⁴
		Lloyd Lake	1.78 × 10 ⁻⁰⁴	1.34 × 10 ⁻⁰²	6.46 × 10 ⁻⁰³	1.28 x 10 ⁻⁰²	9.21 × 10 ⁻⁰²	7.00 × 10 ⁻⁰⁶	2.34 × 10 ⁻⁰⁴	7.66 × 10 ⁻⁰⁴	1.35 × 10 ⁻⁰²	6.51 × 10 ⁻⁰³	1.28 x 10 ⁻⁰²	9.21 × 10 ⁻⁰²	7.70 × 10 ⁻⁰⁶	2.44 × 10 ⁻⁰⁴
		Reference (Broach Lake)	n/a	n/a	2.45 × 10 ⁻⁰⁴	2.94 × 10 ⁻⁰⁴	2.65 × 10 ⁻⁰⁴	3.09 × 10 ⁻⁰⁴	5.71 × 10 ⁻⁰⁴	n/a	n/a	2.45 × 10 ⁻⁰⁴	2.94 × 10 ⁻⁰⁴	2.65 × 10 ⁻⁰⁴	3.09 × 10 ⁻⁰⁴	5.71 × 10 ⁻⁰⁴
	Beaver	Patterson Lake North Arm – West Basin	n/a	n/a	4.59 × 10 ⁻⁰⁴	3.43 × 10 ⁻⁰⁴	2.79 × 10 ⁻⁰⁴	3.96 × 10 ⁻⁰⁴	4.45 × 10 ⁻⁰³	n/a	n/a	4.65 × 10 ⁻⁰⁴	3.45 × 10 ⁻⁰⁴	2.79 × 10 ⁻⁰⁴	4.94 × 10 ⁻⁰⁴	4.68 × 10 ⁻⁰³
		Patterson Lake South Arm	n/a	n/a	2.64 × 10 ⁻⁰⁴	3.20 × 10 ⁻⁰⁴	2.69 × 10 ⁻⁰⁴	3.40 × 10 ⁻⁰⁴	1.14 × 10 ⁻⁰³	n/a	n/a	2.64 × 10 ⁻⁰⁴	3.21 × 10 ⁻⁰⁴	2.69 × 10 ⁻⁰⁴	3.92 × 10 ⁻⁰⁴	1.20 × 10 ⁻⁰³
		Beet Lake	n/a	n/a	2.49×10^{-04}	3.08 × 10 ⁻⁰⁴	2.67 × 10 ⁻⁰⁴	3.25×10^{-04}	8.99 × 10 ⁻⁰⁴	n/a	n/a	2.49 × 10 ⁻⁰⁴	3.09×10^{-04}	2.67 × 10 ⁻⁰⁴	3.51 × 10 ⁻⁰⁴	9.22 × 10 ⁻⁰⁴
		Lloyd Lake	n/a	n/a	2.46×10^{-04} 9.22×10^{-04}	2.97 × 10 ⁻⁰⁴	2.65 × 10 ⁻⁰⁴	3.14 × 10 ⁻⁰⁴	5.84×10^{-04} 1.82×10^{-04}	n/a	n/a	2.46 × 10 ⁻⁰⁴	2.97 × 10 ⁻⁰⁴	2.65 × 10 ⁻⁰⁴	3.16 × 10 ⁻⁰⁴	5.86 × 10 ⁻⁰⁴
		Reference (Broach Lake)	n/a	n/a	9.22 × 10 °	8.71 × 10 ⁻⁰⁵	2.49 × 10 ⁻⁰³	1.44 × 10 ⁻⁰⁴	1.82 × 10 °	n/a	n/a	9.22 × 10 ⁻⁰⁴	8.71 × 10 ⁻⁰⁵	2.49 × 10 ⁻⁰³	1.44 × 10 ⁻⁰⁴	1.82 × 10 ⁻⁰⁴
	rial Black Bear	Patterson Lake North Arm – West Basin	n/a	n/a	1.66 × 10 ⁻⁰³	9.70 × 10 ⁻⁰⁵	2.74 × 10 ⁻⁰³	1.62 × 10 ⁻⁰⁴	6.96 × 10 ⁻⁰⁴	n/a	n/a	1.69 × 10 ⁻⁰³	9.73 × 10 ⁻⁰⁵	2.75 × 10 ⁻⁰³	1.77 × 10 ⁻⁰⁴	7.54 × 10 ⁻⁰⁴
Animals		Beet Lake	n/a	n/a	1.02 × 10 ⁻⁰³	1.06 × 10 ⁻⁰⁴	2.56 × 10 ⁻⁰³	1.76 × 10 ⁻⁰⁴	2.51 × 10 ⁻⁰⁴	n/a	n/a	1.02 × 10 ⁻⁰³	1.06 × 10 ⁻⁰⁴	2.56 × 10 ⁻⁰³	1.78 × 10 ⁻⁰⁴	2.55 × 10 ⁻⁰⁴
		Lloyd Lake	n/a	n/a	9.27 × 10 ⁻⁰⁴	9.04 × 10 ⁻⁰⁵	2.50 × 10 ⁻⁰³	1.50 × 10 ⁻⁰⁴	1.86 × 10 ⁻⁰⁴	n/a	n/a	9.27 × 10 ⁻⁰⁴	9.04 × 10 ⁻⁰⁵	2.50 × 10 ⁻⁰³	1.51 × 10 ⁻⁰⁴	1.87 × 10 ⁻⁰⁴
		Reference (Broach Lake)	n/a	n/a	2.46 x 10 ⁻⁰³	4.49 x 10 ⁻⁰⁵	2.88 x 10 ⁻⁰³	1.11 x 10 ⁻⁰⁴	6.66 x 10 ⁻⁰⁵	n/a	n/a	2.46 x 10 ⁻⁰³	4.49 x 10 ⁻⁰⁵	2.88 x 10 ⁻⁰³	1.11 x 10 ⁻⁰⁴	6.66 x 10 ⁻⁰⁵
	Grey Wolf	Patterson Lake North Arm – West Basin	n/a	n/a	2.91 x 10 ⁻⁰³	4.58 x 10 ⁻⁰⁵	2.93 x 10 ⁻⁰³	1.22 x 10 ⁻⁰⁴	2.14 x 10 ⁻⁰⁴	n/a	n/a	2.92 x 10 ⁻⁰³	4.58 x 10 ⁻⁰⁵	2.93 x 10 ⁻⁰³	1.32 x 10 ⁻⁰⁴	2.18 x 10 ⁻⁰⁴
		Lloyd Lake	n/a	n/a	2.50 x 10 ⁻⁰³	4.53 x 10 ⁻⁰⁵	2.89 x 10 ⁻⁰³	1.14 x 10 ⁻⁰⁴	6.87 x 10 ⁻⁰⁵	n/a	n/a	2.50 x 10 ⁻⁰³	4.53 x 10 ⁻⁰⁵	2.89 x 10 ⁻⁰³	1.14 x 10 ⁻⁰⁴	6.88 x 10 ⁻⁰⁵
	American Mink	Reference (Broach Lake) Patterson Lake North Arm	n/a	n/a	3.77 × 10 ⁻⁰³	3.45 × 10 ⁻⁰⁴	1.99 × 10 ⁻⁰²	4.40×10^{-04} 1.08×10^{-03}	1.93 × 10 ⁻⁰⁴ 7.43 x 10 ⁻⁰⁴	n/a	n/a	3.77 × 10 ⁻⁰³	3.45 × 10 ⁻⁰⁴	1.99 × 10 ⁻⁰²	4.40 × 10 ⁻⁰⁴	1.93 × 10 ⁻⁰⁴
		– West Basin	n/a	n/a	1.13 x 10 ⁻⁰²	4.36 x 10 ⁻⁰⁴	2.31 x 10 ⁻⁰²	1.08 X 10 °	7.43 X 10 °	n/a	n/a	1.15 x 10 ⁻⁰²	4.39 x 10 ⁻⁰⁴	2.32 x 10 ⁻⁰²	2.37 x 10 ⁻⁰³	1.34 x 10 ⁻⁰³

								Maximum HQ	Operations						
Biota	Location				Application Case						Upp	er Bound Scenar	io		•
		Chloride	Sulphate	Arsenic	Cobalt	Copper	Molybdenum	Uranium	Chloride	Sulphate	Arsenic	Cobalt	Copper	Molybdenum	Urani
	Patterson Lake South Arm	n/a	n/a	4.39 x 10 ⁻⁰³	3.93 x 10 ⁻⁰⁴	2.15 x 10 ⁻⁰²	7.67 x 10 ⁻⁰⁴	3.12 x 10 ⁻⁰⁴	n/a	n/a	4.41 x 10 ⁻⁰³	3.95 x 10 ⁻⁰⁴	2.16 x 10 ⁻⁰²	1.45 x 10 ⁻⁰³	4.70 x
	Beet Lake	n/a	n/a	3.87 x 10 ⁻⁰³	3.70 x 10 ⁻⁰⁴	2.07 x 10 ⁻⁰²	6.08 x 10 ⁻⁰⁴	2.44 x 10 ⁻⁰⁴	n/a	n/a	3.87 x 10 ⁻⁰³	3.71 x 10 ⁻⁰⁴	2.07 x 10 ⁻⁰²	9.60 x 10 ⁻⁰⁴	3.02 x
	Lloyd Lake	n/a	n/a	3.78 x 10 ⁻⁰³	3.47 x 10 ⁻⁰⁴	2.00 x 10 ⁻⁰²	4.58 x 10 ⁻⁰⁴	1.97 x 10 ⁻⁰⁴	n/a	n/a	3.78 x 10 ⁻⁰³	3.47 x 10 ⁻⁰⁴	2.00 x 10 ⁻⁰²	4.94 x 10 ⁻⁰⁴	2.02 x
	Reference (Broach Lake)	n/a	n/a	1.63 × 10 ⁻⁰⁴	2.11 × 10 ⁻⁰⁴	1.07 × 10 ⁻⁰²	1.41 × 10 ⁻⁰⁴	2.84×10^{-04}	n/a	n/a	1.63 × 10 ⁻⁰⁴	2.11 × 10 ⁻⁰⁴	1.07 × 10 ⁻⁰²	1.41 × 10 ⁻⁰⁴	2.84 ×
	Patterson Lake North Arm – West Basin	n/a	n/a	3.79 × 10 ⁻⁰⁴	2.58 × 10 ⁻⁰⁴	1.15×10^{-02}	2.02 × 10 ⁻⁰⁴	2.08 × 10 ⁻⁰³	n/a	n/a	3.85×10^{-04}	2.60 × 10 ⁻⁰⁴	1.15 × 10 ⁻⁰²	2.92 × 10 ⁻⁰⁴	2.32 ×
Moose	Patterson Lake South Arm	n/a	n/a	3.77 × 10 ⁻⁰⁴	2.55 × 10 ⁻⁰⁴	1.15 × 10 ⁻⁰²	1.94 × 10 ⁻⁰⁴	2.03 × 10 ⁻⁰³	n/a	n/a	3.83 × 10 ⁻⁰⁴	2.57× 10 ⁻⁰⁴	1.15 × 10 ⁻⁰²	2.73 × 10 ⁻⁰⁴	2.27 ×
	Beet Lake	n/a	n/a	1.67 × 10 ⁻⁰⁴	2.25 × 10 ⁻⁰⁴	1.09 × 10 ⁻⁰²	1.54 × 10 ⁻⁰⁴	4.37 × 10 ⁻⁰⁴	n/a	n/a	1.67 × 10 ⁻⁰⁴	2.25 × 10 ⁻⁰⁴	1.09 × 10 ⁻⁰²	1.79 × 10 ⁻⁰⁴	4.60 >
	Lloyd Lake	n/a	n/a	1.64 × 10 ⁻⁰⁴	2.13×10^{-04}	1.08 × 10 ⁻⁰²	1.44 × 10 ⁻⁰⁴	2.91 × 10 ⁻⁰⁴	n/a	n/a	1.64 × 10 ⁻⁰⁴	2.13 × 10 ⁻⁰⁴	1.08 × 10 ⁻⁰²	1.46 × 10 ⁻⁰⁴	2.93
	Reference (Broach Lake)	n/a	n/a	2.01 × 10 ⁻⁰³	1.19 × 10 ⁻⁰³	3.42×10^{-03}	9.60 × 10 ⁻⁰⁴	9.83 × 10 ⁻⁰⁴	n/a	n/a	2.01 × 10 ⁻⁰³	1.19 × 10 ⁻⁰³	3.42×10^{-03}	9.60 × 10 ⁻⁰⁴	9.83
	Patterson Lake North Arm – West Basin	n/a	n/a	6.18 × 10 ⁻⁰³	1.56 × 10 ⁻⁰³	3.98 × 10 ⁻⁰³	2.57 × 10 ⁻⁰³	3.25 × 10 ⁻⁰³	n/a	n/a	6.28 × 10 ⁻⁰³	1.57 × 10 ⁻⁰³	4.00 × 10 ⁻⁰³	5.87 × 10 ⁻⁰³	7.34
Muskrat	Patterson Lake South Arm	n/a	n/a	2.36 × 10 ⁻⁰³	1.39×10^{-03}	3.71×10^{-03}	1.79 × 10 ⁻⁰³	1.59 × 10 ⁻⁰³	n/a	n/a	2.36 × 10 ⁻⁰³	1.40 × 10 ⁻⁰³	3.72×10^{-03}	3.53×10^{-03}	2.67
	Beet Lake	n/a	n/a	2.07 × 10 ⁻⁰³	1.29×10^{-03}	3.57 × 10 ⁻⁰³	1.39 × 10 ⁻⁰³	1.20×10^{-03}	n/a	n/a	2.07×10^{-03}	1.30 × 10 ⁻⁰³	3.58 × 10 ⁻⁰³	2.28×10^{-03}	1.60
	Lloyd Lake	n/a	n/a	2.02 × 10 ⁻⁰³	1.20×10^{-03}	3.44 × 10 ⁻⁰³	1.00 × 10 ⁻⁰³	1.00 × 10 ⁻⁰³	n/a	n/a	2.02 × 10 ⁻⁰³	1.20 × 10 ⁻⁰³	3.44×10^{-03}	1.10×10^{-03}	1.04
	Reference (Broach Lake)	n/a	n/a	4.18 × 10 ⁻⁰⁴	3.66×10^{-05}	6.08 × 10 ⁻⁰⁴	1.22 × 10 ⁻⁰⁴	6.64×10^{-05}	n/a	n/a	4.18 × 10 ⁻⁰⁴	3.66 × 10 ⁻⁰⁵	6.08 × 10 ⁻⁰⁴	1.22×10^{-04}	6.64
	Patterson Lake North Arm – West Basin	n/a	n/a	4.32 × 10 ⁻⁰⁴	3.82 × 10 ⁻⁰⁵	6.31 × 10 ⁻⁰⁴	1.46 × 10 ⁻⁰⁴	6.16 × 10 ⁻⁰⁴	n/a	n/a	4.32 × 10 ⁻⁰⁴	3.83 × 10 ⁻⁰⁵	6.31 × 10 ⁻⁰⁴	1.72 × 10 ⁻⁰⁴	6.24
Red Fox	Patterson Lake South Arm	n/a	n/a	4.20 x 10 ⁻⁰⁴	3.74 x 10 ⁻⁰⁵	6.12 x 10 ⁻⁰⁴	1.30 x 10 ⁻⁰⁴	1.45 x 10 ⁻⁰⁴	n/a	n/a	4.20 × 10 ⁻⁰⁴	3.75 x 10 ⁻⁰⁵	6.12 x 10 ⁻⁰⁴	1.44 x 10 ⁻⁰⁴	1.47
	Beet Lake	n/a	n/a	4.19 x 10 ⁻⁰⁴	3.71 x 10 ⁻⁰⁵	6.11 x 10 ⁻⁰⁴	1.26 x 10 ⁻⁰⁴	1.13 x 10 ⁻⁰⁴	n/a	n/a	4.19 x 10 ⁻⁰⁴	3.71 x 10 ⁻⁰⁵	6.11 x 10 ⁻⁰⁴	1.33 x 10 ⁻⁰⁴	1.13
	Lloyd Lake	n/a	n/a	4.19 × 10 ⁻⁰⁴	3.68×10^{-05}	6.09 × 10 ⁻⁰⁴	1.23 × 10 ⁻⁰⁴	6.81×10^{-05}	n/a	n/a	4.19 × 10 ⁻⁰⁴	3.68×10^{-05}	6.09 × 10 ⁻⁰⁴	1.24 × 10 ⁻⁰⁴	6.82
	Reference (Broach Lake)	n/a	n/a	1.43×10^{-03}	2.97×10^{-04}	1.06 × 10 ⁻⁰³	6.71 × 10 ⁻⁰⁵	9.42 × 10 ⁻⁰⁴	n/a	n/a	1.43×10^{-03}	2.97 × 10 ⁻⁰⁴	1.06×10^{-03}	6.71 × 10 ⁻⁰⁵	9.42
	Patterson Lake North Arm – West Basin	n/a	n/a	1.46 × 10 ⁻⁰³	3.02 × 10 ⁻⁰⁴	1.09 × 10 ⁻⁰³	7.69 × 10 ⁻⁰⁵	7.54 × 10 ⁻⁰³	n/a	n/a	1.46 × 10 ⁻⁰³	3.03 × 10 ⁻⁰⁴	1.09 × 10 ⁻⁰³	7.94 × 10 ⁻⁰⁵	7.55
Snowshoe Hare	Patterson Lake South Arm	n/a	n/a	1.44 × 10 ⁻⁰³	3.00 × 10 ⁻⁰⁴	1.06 × 10 ⁻⁰³	6.92 × 10 ⁻⁰⁵	1.89 × 10 ⁻⁰³	n/a	n/a	1.44 × 10 ⁻⁰³	3.00 × 10 ⁻⁰⁴	1.06 × 10 ⁻⁰³	7.05 × 10 ⁻⁰⁵	1.89
	Beet Lake	n/a	n/a	1.44 × 10 ⁻⁰³	2.99 × 10 ⁻⁰⁴	1.06 × 10 ⁻⁰³	6.82 × 10 ⁻⁰⁵	1.50 × 10 ⁻⁰³	n/a	n/a	1.44 × 10 ⁻⁰³	3.00 × 10 ⁻⁰⁴	1.06 × 10 ⁻⁰³	6.88 × 10 ⁻⁰⁵	1.50
	Lloyd Lake	n/a	n/a	1.44 × 10 ⁻⁰³	2.99 × 10 ⁻⁰⁴	1.06 × 10 ⁻⁰³	6.79 × 10 ⁻⁰⁵	9.63 × 10 ⁻⁰⁴	n/a	n/a	1.44 × 10 ⁻⁰³	2.99 × 10 ⁻⁰⁴	1.06 × 10 ⁻⁰³	6.79×10^{-05}	9.61
	Reference (Broach Lake)	n/a	n/a	1.50 × 10 ⁻⁰⁴	1.05×10^{-04}	1.24 × 10 ⁻⁰⁴	5.79 × 10 ⁻⁰⁴	7.05×10^{-04}	n/a	n/a	1.50 × 10 ⁻⁰⁴	1.05×10^{-04}	1.24 × 10 ⁻⁰⁴	5.79×10^{-04}	7.05
Southern Red-	Patterson Lake North Arm – West Basin	n/a	n/a	1.53 × 10 ⁻⁰⁴	1.07 × 10 ⁻⁰⁴	1.28 × 10 ⁻⁰⁴	6.68 × 10 ⁻⁰⁴	5.45 × 10 ⁻⁰³	n/a	n/a	1.54 × 10 ⁻⁰⁴	1.07 × 10 ⁻⁰⁴	1.28 × 10 ⁻⁰⁴	7.10 × 10 ⁻⁰⁴	5.46
Backed Vole	Patterson Lake South Arm	n/a	n/a	1.51 × 10 ⁻⁰⁴	1.06 × 10 ⁻⁰⁴	1.24 × 10 ⁻⁰⁴	6.01 × 10 ⁻⁰⁴	1.38 × 10 ⁻⁰³	n/a	n/a	1.51 × 10 ⁻⁰⁴	1.06 × 10 ⁻⁰⁴	1.24 × 10 ⁻⁰⁴	6.23 × 10 ⁻⁰⁴	1.39
	Beet Lake	n/a	n/a	1.50 × 10 ⁻⁰⁴	1.06 × 10 ⁻⁰⁴	1.24 × 10 ⁻⁰⁴	5.90 × 10 ⁻⁰⁴	1.10 × 10 ⁻⁰³	n/a	n/a	1.50 × 10 ⁻⁰⁴	1.06 × 10 ⁻⁰⁴	1.24 × 10 ⁻⁰⁴	6.01 × 10 ⁻⁰⁴	1.11
	Lloyd Lake	n/a	n/a	1.50 × 10 ⁻⁰⁴	1.06×10^{-04}	1.24 × 10 ⁻⁰⁴	5.85 × 10 ⁻⁰⁴	7.20×10^{-04}	n/a	n/a	1.50×10^{-04}	1.06 × 10 ⁻⁰⁴	1.24 × 10 ⁻⁰⁴	5.86 × 10 ⁻⁰⁴	7.20
	Reference (Broach Lake)	n/a	n/a	3.79 x 10 ⁻⁰⁴	2.94 x 10 ⁻⁰⁴	1.19 x 10 ⁻⁰²	3.02 x 10 ⁻⁰⁴	3.16 x 10 ⁻⁰⁴	n/a	n/a	3.79 x 10 ⁻⁰⁴	2.94 x 10 ⁻⁰⁴	1.19 x 10 ⁻⁰²	3.02 x 10 ⁻⁰⁴	3.16
Woodland	Patterson Lake North Arm – West Basin	n/a	n/a	6.71 x 10 ⁻⁰⁴	3.51 x 10 ⁻⁰⁴	1.32 x 10 ⁻⁰²	5.95 x 10 ⁻⁰⁴	1.27 x 10 ⁻⁰²	n/a	n/a	6.79 x 10 ⁻⁰⁴	3.52 x 10 ⁻⁰⁴	1.32 x 10 ⁻⁰²	7.58 x 10 ⁻⁰⁴	1.30
Caribou	Beet Lake	n/a	n/a	3.87 x 10 ⁻⁰⁴	3.07 x 10 ⁻⁰⁴	1.20 x 10 ⁻⁰²	3.33 x 10 ⁻⁰⁴	1.30 x 10 ⁻⁰³	n/a	n/a	3.87 x 10 ⁻⁰⁴	3.07 x 10 ⁻⁰⁴	1.20 x 10 ⁻⁰²	3.62 x 10 ⁻⁰⁴	1.32
	Lloyd Lake	n/a	n/a	3.89 x 10 ⁻⁰⁴	3.06 x 10 ⁻⁰⁴	1.20 x 10 ⁻⁰²	3.39 x 10 ⁻⁰⁴	3.95 x 10 ⁻⁰⁴	n/a	n/a	3.89 x 10 ⁻⁰⁴	3.06 x 10 ⁻⁰⁴	1.20 x 10 ⁻⁰²	3.45 x 10 ⁻⁰⁴	4.00
	Reference (Broach Lake)	n/a	n/a	1.79 × 10 ⁻⁰³	3.95 × 10 ⁻⁰⁴	1.51 × 10 ⁻⁰³	6.67 × 10 ⁻⁰⁵	3.96×10^{-04}	n/a	n/a	1.79×10^{-03}	3.95×10^{-04}	1.51 × 10 ⁻⁰³	6.67×10^{-05}	3.96
	Patterson Lake North Arm – West Basin	n/a	n/a	1.81 × 10 ⁻⁰³	4.00 × 10 ⁻⁰⁴	1.56 × 10 ⁻⁰³	7.46 × 10 ⁻⁰⁵	3.00 × 10 ⁻⁰³	n/a	n/a	1.81 × 10 ⁻⁰³	4.00 × 10 ⁻⁰⁴	1.56 × 10 ⁻⁰³	7.61 × 10 ⁻⁰⁵	3.01
Grouse	Patterson Lake South Arm	n/a	n/a	1.79 × 10 ⁻⁰³	3.97 × 10 ⁻⁰⁴	1.52 × 10 ⁻⁰³	6.83 × 10 ⁻⁰⁵	7.69 × 10 ⁻⁰⁴	n/a	n/a	1.79 × 10 ⁻⁰³	3.97 × 10 ⁻⁰⁴	1.52 × 10 ⁻⁰³	6.90 × 10 ⁻⁰⁵	7.70
	Beet Lake	n/a	n/a	1.79 × 10 ⁻⁰³	3.97×10^{-04}	1.52 × 10	6.75 × 10 ⁻⁰⁵	6.16×10^{-04}	n/a	n/a	1.79×10^{-03}	3.97×10^{-04}	1.52 × 10	6.79 × 10 ⁻⁰⁵	6.16
	Lloyd Lake	n/a	n/a	1.79×10^{-03}	3.97×10^{-04}	1.51 × 10 ⁻⁰³	6.73×10^{-05}	4.05×10^{-04}	n/a	n/a	1.79×10^{-03}	3.97×10^{-04}	1.51 × 10 ⁻⁰³	6.74×10^{-05}	4.05
	Reference (Broach Lake)	n/a	n/a	1.72 × 10 ⁻⁰⁴	3.93×10^{-05}	1.96 × 10 ⁻⁰⁴	1.24×10^{-05}	3.93×10^{-05}	n/a	n/a	1.73 × 10 1.72 × 10 ⁻⁰⁴	3.93×10^{-05}	1.96 × 10 ⁻⁰⁴	1.24×10^{-05}	3.93
	Patterson Lake North Arm – West Basin	n/a	n/a	1.77 × 10 ⁻⁰⁴	4.03 × 10 ⁻⁰⁵	2.03 × 10 ⁻⁰⁴	1.45 × 10 ⁻⁰⁵	3.04 × 10 ⁻⁰⁴	n/a	n/a	1.77 × 10 ⁻⁰⁴	4.03 × 10 ⁻⁰⁵	2.03 × 10 ⁻⁰⁴	1.60 × 10 ⁻⁰⁵	3.05
Canada Goose	Patterson Lake South Arm	n/a	n/a	1.73 × 10 ⁻⁰⁴	3.98 × 10 ⁻⁰⁵	1.98 × 10 ⁻⁰⁴	1.30 × 10 ⁻⁰⁵	7.72 × 10 ⁻⁰⁵	n/a	n/a	1.73 × 10 ⁻⁰⁴	3.98 × 10 ⁻⁰⁵	1.97 × 10 ⁻⁰⁴	1.37 × 10 ⁻⁰⁵	7.75
	Beet Lake	n/a	n/a	1.73 × 10 ⁻⁰⁴	3.97×10^{-05}	1.98 × 10 1.97 × 10 ⁻⁰⁴	1.27×10^{-05}	6.16×10^{-05}	n/a	n/a	1.73×10^{-04}	3.97×10^{-05}	1.97 × 10 ⁻⁰⁴	1.31 × 10 ⁻⁰⁵	6.17
	Lloyd Lake	n/a	n/a	1.73 × 10 1.72 × 10 ⁻⁰⁴	3.96×10^{-05}	1.97 × 10 ⁻⁰⁴	1.27×10^{-05}	4.02×10^{-05}	n/a	n/a	1.73×10^{-04}	3.96×10^{-05}	1.97 × 10 ⁻⁰⁴	1.26 × 10 ⁻⁰⁵	4.02
	Reference (Broach Lake)	n/a	n/a	9.07×10^{-03}	4.28×10^{-03}	2.80×10^{-01}	6.31 × 10 ⁻⁰³	8.52 × 10 ⁻⁰⁴	n/a	n/a	9.07×10^{-03}	4.28×10^{-03}	2.80×10^{-01}	6.31 × 10 ⁻⁰³	8.52
Little Brown	Patterson Lake North Arm – West Basin	n/a	n/a	2.77 × 10 ⁻⁰²	5.53 × 10 ⁻⁰³	3.26 × 10 ⁻⁰¹	1.69 × 10 ⁻⁰²	2.80 × 10 ⁻⁰³	n/a	n/a	2.81 × 10 ⁻⁰²	5.59 × 10 ⁻⁰³	3.27 × 10 ⁻⁰¹	3.84 × 10 ⁻⁰²	6.28
Myotis	Patterson Lake South Arm	n/a	n/a	1.06 × 10 ⁻⁰²	4.95 × 10 ⁻⁰³	3.03 × 10 ⁻⁰¹	1.17 × 10 ⁻⁰²	1.37 × 10 ⁻⁰³	n/a	n/a	1.06 × 10 ⁻⁰²	4.97 × 10 ⁻⁰³	3.04 × 10 ⁻⁰¹	2.32 × 10 ⁻⁰²	2.29
	Beet Lake	n/a	n/a	9.32 × 10 ⁻⁰³	4.62×10^{-03}	2.92 × 10 ⁻⁰¹	9.10 × 10 ⁻⁰³	1.04×10^{-03}	n/a	n/a	9.32 × 10 ⁻⁰³	4.64×10^{-03}	2.92 × 10 ⁻⁰¹	1.50 × 10 ⁻⁰²	1.38

ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT

ECOLOGICAL RISK ASSESSMENT

								Maximum HC	Operations						
Biota	Location				Application Cas	е					Upp	er Bound Scenar	io		
		Chloride	Sulphate	Arsenic	Cobalt	Copper	Molybdenum	Uranium	Chloride	Sulphate	Arsenic	Cobalt	Copper	Molybdenum	Uranium
	Lloyd Lake	n/a	n/a	9.08×10^{-03}	4.31×10^{-03}	2.81 × 10 ⁻⁰¹	6.59 × 10 ⁻⁰³	8.70×10^{-04}	n/a	n/a	9.08 × 10 ⁻⁰³	4.31×10^{-03}	2.81 × 10 ⁻⁰¹	7.20×10^{-03}	9.00 × 10 ⁻⁰⁴
	Reference (Broach Lake)	n/a	n/a	1.31×10^{-03}	1.54×10^{-04}	4.62×10^{-03}	1.97×10^{-05}	2.11 × 10 ⁻⁰⁵	n/a	n/a	1.31 × 10 ⁻⁰³	1.54×10^{-04}	4.62×10^{-03}	1.97 × 10 ⁻⁰⁵	2.11 × 10 ⁻⁰⁵
	Patterson Lake North Arm – West Basin	n/a	n/a	4.52 × 10 ⁻⁰³	2.01 × 10 ⁻⁰⁴	5.39 × 10 ⁻⁰³	5.27 × 10 ⁻⁰⁵	7.64 × 10 ⁻⁰⁵	n/a	n/a	4.62 × 10 ⁻⁰³	2.03 × 10 ⁻⁰⁴	5.42 × 10 ⁻⁰³	1.20 × 10 ⁻⁰⁴	1.78 × 10 ⁻⁰⁴
Common Loon	Patterson Lake South Arm	n/a	n/a	1.58 × 10 ⁻⁰³	1.79 × 10 ⁻⁰⁴	5.01 × 10 ⁻⁰³	3.67 × 10 ⁻⁰⁵	3.57 × 10 ⁻⁰⁵	n/a	n/a	1.58 × 10 ⁻⁰³	1.80 × 10 ⁻⁰⁴	5.02 × 10 ⁻⁰³	7.24 × 10 ⁻⁰⁵	6.16 × 10 ⁻⁰⁵
	Beet Lake	n/a	n/a	1.35 × 10 ⁻⁰³	1.67 × 10 ⁻⁰⁴	4.82×10^{-03}	2.84 × 10 ⁻⁰⁵	2.65 × 10 ⁻⁰⁵	n/a	n/a	1.35 × 10 ⁻⁰³	1.68 × 10 ⁻⁰⁴	4.82 × 10 ⁻⁰³	4.68 × 10 ⁻⁰⁵	3.60 × 10 ⁻⁰⁵
	Lloyd Lake	n/a	n/a	1.31 × 10 ⁻⁰³	1.56 × 10 ⁻⁰⁴	4.64 × 10 ⁻⁰³	2.06 × 10 ⁻⁰⁵	2.16 × 10 ⁻⁰⁵	n/a	n/a	1.31 × 10 ⁻⁰³	1.56 × 10 ⁻⁰⁴	4.64 × 10 ⁻⁰³	2.25 × 10 ⁻⁰⁵	2.25 × 10 ⁻⁰⁵
	Reference (Broach Lake)	n/a	n/a	5.44 × 10 ⁻⁰³	1.85 × 10 ⁻⁰³	1.40 × 10 ⁻⁰²	3.13 × 10 ⁻⁰⁴	3.27×10^{-04}	n/a	n/a	5.44 × 10 ⁻⁰³	1.85 × 10 ⁻⁰³	1.40 × 10 ⁻⁰²	3.13 × 10 ⁻⁰⁴	3.27 × 10 ⁻⁰⁴
	Patterson Lake North Arm – West Basin	n/a	n/a	1.67 × 10 ⁻⁰²	2.41 × 10 ⁻⁰³	1.63 × 10 ⁻⁰²	8.38 × 10 ⁻⁰⁴	1.08 × 10 ⁻⁰³	n/a	n/a	1.70 × 10 ⁻⁰²	2.43 × 10 ⁻⁰³	1.64 × 10 ⁻⁰²	1.91 × 10 ⁻⁰³	2.44 × 10 ⁻⁰³
Mallard	Patterson Lake South Arm	n/a	n/a	6.37 × 10 ⁻⁰³	2.15 × 10 ⁻⁰³	1.52 × 10 ⁻⁰²	5.84 × 10 ⁻⁰⁴	5.26 × 10 ⁻⁰⁴	n/a	n/a	6.40 × 10 ⁻⁰³	2.16 × 10 ⁻⁰³	1.52 × 10 ⁻⁰²	1.15 × 10 ⁻⁰³	8.85 × 10 ⁻⁰⁴
	Beet Lake	n/a	n/a	5.59 × 10 ⁻⁰³	2.01 × 10 ⁻⁰³	1.46 × 10 ⁻⁰²	4.52 × 10 ⁻⁰⁴	4.00×10^{-04}	n/a	n/a	5.60 × 10 ⁻⁰³	2.01 × 10 ⁻⁰³	1.46 × 10 ⁻⁰²	7.44 × 10 ⁻⁰⁴	5.31 × 10 ⁻⁰⁴
	Lloyd Lake	n/a	n/a	5.45 × 10 ⁻⁰³	1.87 × 10 ⁻⁰³	1.41 × 10 ⁻⁰²	3.27 × 10 ⁻⁰⁴	3.33×10^{-04}	n/a	n/a	5.45 × 10 ⁻⁰³	1.87×10^{-03}	1.41 × 10 ⁻⁰²	3.57 × 10 ⁻⁰⁴	3.45 × 10 ⁻⁰⁴
Dusty Plackbird	Reference (Broach Lake)	n/a	n/a	1.99 × 10 ⁻⁰²	8.87 × 10 ⁻⁰³	2.35 × 10 ⁻⁰¹	1.76 × 10 ⁻⁰³	1.10 × 10 ⁻⁰³	n/a	n/a	1.99 × 10 ⁻⁰²	8.87 × 10 ⁻⁰³	2.35 × 10 ⁻⁰¹	1.76 × 10 ⁻⁰³	1.10 × 10 ⁻⁰³
	Patterson Lake North Arm – West Basin	n/a	n/a	5.08 × 10 ⁻⁰²	1.12 × 10 ⁻⁰²	2.73 × 10 ⁻⁰¹	4.35 × 10 ⁻⁰³	4.76 × 10 ⁻⁰³	n/a	n/a	5.16 × 10 ⁻⁰²	1.13 × 10 ⁻⁰²	2.74 × 10 ⁻⁰¹	9.63 × 10 ⁻⁰³	7.15 × 10 ⁻⁰³
Rusty Blackbird	Patterson Lake South Arm	n/a	n/a	2.24 × 10 ⁻⁰²	1.01 × 10 ⁻⁰²	2.55 × 10 ⁻⁰¹	3.09 × 10 ⁻⁰³	1.78 × 10 ⁻⁰³	n/a	n/a	2.25 × 10 ⁻⁰²	1.02 × 10 ⁻⁰²	2.55 × 10 ⁻⁰¹	5.89 × 10 ⁻⁰³	2.24 × 10 ⁻⁰³
	Beet Lake	n/a	n/a	2.03 × 10 ⁻⁰²	9.52 × 10 ⁻⁰³	2.45 × 10 ⁻⁰¹	2.44 × 10 ⁻⁰³	1.42 × 10 ⁻⁰³	n/a	n/a	2.03 × 10 ⁻⁰²	9.54 × 10 ⁻⁰³	2.46 × 10 ⁻⁰¹	3.88 × 10 ⁻⁰³	1.66 × 10 ⁻⁰³
_	Lloyd Lake	n/a	n/a	1.99 × 10 ⁻⁰²	8.93 × 10 ⁻⁰³	2.36 × 10 ⁻⁰¹	1.83 × 10 ⁻⁰³	1.12 × 10 ⁻⁰³	n/a	n/a	1.99 × 10 ⁻⁰²	8.94 × 10 ⁻⁰³	2.36 × 10 ⁻⁰¹	1.98 × 10 ⁻⁰³	1.14 × 10 ⁻⁰³

n/a = not applicable; HQ = hazard quotient.

Table 6-23: Non-radiological Risk to Ecological Receptors – Application Case and Upper Bound Scenario – Far-Future Projection

									Maximum H	O Far-Future						
	Biota	Location				Application Cas	e					Ur	per Bound Sce	nario		
	Diota	200011011	Chloride	Sulphate	Arsenic	Cobalt	Copper	Molybdenum	Uranium	Chloride	Sulphate	Arsenic	Cobalt	Copper	Molybdenum	Uranium
		Reference (Broach Lake)	1.00 × 10 ⁻⁰³	1.45 × 10 ⁻⁰³	8.50 × 10 ⁻⁰⁴	4.75 x 10 ⁻⁰²	4.18 × 10 ⁻⁰¹	6.61 × 10 ⁻⁰⁶	3.57 × 10 ⁻⁰³	1.00×10^{-03}	1.45 × 10 ⁻⁰³	8.50 × 10 ⁻⁰⁴	4.75 x 10 ⁻⁰²	4.18 × 10 ⁻⁰¹	6.61 × 10 ⁻⁰⁶	3.57 × 10 ⁻⁰³
		Patterson Lake North Arm –	1.99 × 10 ⁻⁰³	2.95 × 10 ⁻⁰³	9.01 × 10 ⁻⁰⁴	1.16 x 10 ⁻⁰¹	1.08 × 10 ⁺⁰⁰	1.47 × 10 ⁻⁰⁴	3.57 × 10 ⁻⁰²	2.48 × 10 ⁻⁰³	3.82 × 10 ⁻⁰³	9.29 × 10 ⁻⁰⁴	1.53 x 10 ⁻⁰¹	1.42 × 10 ⁺⁰⁰	5.47 × 10 ⁻⁰⁴	3.58 × 10 ⁻⁰²
	Benthic Invertebrate	West Basin Patterson Lake South Arm	1.73 × 10 ⁻⁰³	2.55 × 10 ⁻⁰³	8.54 × 10 ⁻⁰⁴	8.92 x 10 ⁻⁰²	8.08 × 10 ⁻⁰¹	8.83 × 10 ⁻⁰⁵	1.25 × 10 ⁻⁰²	2.08 × 10 ⁻⁰³	3.19 × 10 ⁻⁰³	8.57 × 10 ⁻⁰⁴	1.11 x 10 ⁻⁰¹	1.00 × 10 ⁺⁰⁰	3.20 × 10 ⁻⁰⁴	1.25 × 10 ⁻⁰²
		Beet Lake	1.41 × 10 ⁻⁰³	2.07 × 10 ⁻⁰³	8.51 × 10 ⁻⁰⁴	6.96 x 10 ⁻⁰²	6.22 × 10 ⁻⁰¹	4.92 × 10 ⁻⁰⁵	6.88 × 10 ⁻⁰³	1.61 × 10 ⁻⁰³	2.43×10^{-03}	8.51 × 10 ⁻⁰⁴	8.13 x 10 ⁻⁰²	7.25 × 10 ⁻⁰¹	1.70 × 10 ⁻⁰⁴	6.88 × 10 ⁻⁰³
		Lloyd Lake	1.04 × 10 ⁻⁰³	1.52 × 10 ⁻⁰³	8.50 × 10 ⁻⁰⁴	4.98 x 10 ⁻⁰²	4.39 × 10 ⁻⁰¹	1.09 × 10 ⁻⁰⁵	3.87×10^{-03}	1.07 × 10 ⁻⁰³	1.56 × 10 ⁻⁰³	8.50 × 10 ⁻⁰⁴	5.10 x 10 ⁻⁰²	4.50 × 10 ⁻⁰¹	2.33 × 10 ⁻⁰⁵	3.87×10^{-03}
		Reference (Broach Lake)	1.00 × 10 ⁻⁰³	2.50 × 10 ⁻⁰³	3.64 × 10 ⁻⁰⁴	5.25 x 10 ⁻⁰²	4.21 × 10 ⁻⁰¹	6.66 × 10 ⁻⁰⁶	1.67 × 10 ⁻⁰³	1.00 × 10 ⁻⁰³	2.50 × 10 ⁻⁰³	3.64 × 10 ⁻⁰⁴	5.25 x 10 ⁻⁰²	4.21 × 10 ⁻⁰¹	6.66 × 10 ⁻⁰⁶	1.67 × 10 ⁻⁰³
	Zooplankton	Patterson Lake North Arm — West Basin	1.99 × 10 ⁻⁰³	5.07 × 10 ⁻⁰³	3.86 × 10 ⁻⁰⁴	1.29 x 10 ⁻⁰¹	1.09 × 10 ⁺⁰⁰	1.49 × 10 ⁻⁰⁴	1.67 × 10 ⁻⁰²	2.48 × 10 ⁻⁰³	6.57 × 10 ⁻⁰³	3.98 × 10 ⁻⁰⁴	1.69 x 10 ⁻⁰¹	1.43 × 10 ⁺⁰⁰	5.51 × 10 ⁻⁰⁴	1.67 × 10 ⁻⁰²
	Zoopialiktoli	Patterson Lake South Arm	1.73 × 10 ⁻⁰³	4.38×10^{-03}	3.66 × 10 ⁻⁰⁴	9.87 x 10 ⁻⁰²	8.14 × 10 ⁻⁰¹	8.90 × 10 ⁻⁰⁵	5.84 × 10 ⁻⁰³	2.08 × 10 ⁻⁰³	5.48×10^{-03}	3.67 × 10 ⁻⁰⁴	1.23 x 10 ⁻⁰¹	1.01 × 10 ⁺⁰⁰	3.23 × 10 ⁻⁰⁴	5.84×10^{-03}
		Beet Lake	1.41 × 10 ⁻⁰³	3.56 × 10 ⁻⁰³	3.64 × 10 ⁻⁰⁴	7.70 x 10 ⁻⁰²	6.27 × 10 ⁻⁰¹	4.95 × 10 ⁻⁰⁵	3.22 × 10 ⁻⁰³	1.61 × 10 ⁻⁰³	4.18 × 10 ⁻⁰³	3.64 × 10 ⁻⁰⁴	8.99 x 10 ⁻⁰²	7.30 × 10 ⁻⁰¹	1.71 × 10 ⁻⁰⁴	3.22 × 10 ⁻⁰³
Amustis		Lloyd Lake	1.04 × 10 ⁻⁰³	2.61 × 10 ⁻⁰³	3.64 × 10 ⁻⁰⁴	5.50 x 10 ⁻⁰²	4.42 × 10 ⁻⁰¹	1.10 × 10 ⁻⁰⁵	1.81 × 10 ⁻⁰³	1.07 × 10 ⁻⁰³	2.67 × 10 ⁻⁰³	3.64 × 10 ⁻⁰⁴	5.64 x 10 ⁻⁰²	4.53 × 10 ⁻⁰¹	2.34 × 10 ⁻⁰⁵	1.81 × 10 ⁻⁰³
Aquatic		Reference (Broach Lake)	4.27×10^{-04}	2.11 × 10 ⁻⁰³	1.96 × 10 ⁻⁰⁴	3.26 x 10 ⁻⁰⁴	2.81 × 10 ⁻⁰¹	2.58 × 10 ⁻⁰⁶	1.82 × 10 ⁻⁰⁴	4.27×10^{-04}	2.11 × 10 ⁻⁰³	1.96 × 10 ⁻⁰⁴	3.26 x 10 ⁻⁰⁴	2.81 × 10 ⁻⁰¹	2.58 × 10 ⁻⁰⁶	1.82 × 10 ⁻⁰⁴
Animals	Northern Pike	Patterson Lake North Arm — West Basin	8.49 × 10 ⁻⁰⁴	4.30 × 10 ⁻⁰³	2.08 × 10 ⁻⁰⁴	7.97 x 10 ⁻⁰⁴	7.27 × 10 ⁻⁰¹	5.75 × 10 ⁻⁰⁵	1.83 × 10 ⁻⁰³	1.05 × 10 ⁻⁰³	5.56 × 10 ⁻⁰³	2.15 × 10 ⁻⁰⁴	1.05 x 10 ⁻⁰³	9.51 × 10 ⁻⁰¹	2.14 × 10 ⁻⁰⁴	1.83 × 10 ⁻⁰³
		Patterson Lake South Arm	7.35×10^{-04}	3.71×10^{-03}	1.97 × 10 ⁻⁰⁴	6.12 x 10 ⁻⁰⁴	5.43×10^{-01}	3.45×10^{-05}	6.37×10^{-04}	8.86×10^{-04}	4.64×10^{-03}	1.98×10^{-04}	7.64 x 10 ⁻⁰⁴	6.74×10^{-01}	1.25 × 10 ⁻⁰⁴	6.37×10^{-04}
		Beet Lake	6.01 × 10 ⁻⁰⁴	3.01 × 10 ⁻⁰³	1.97 × 10 ⁻⁰⁴	4.77 x 10 ⁻⁰⁴	4.18 × 10 ⁻⁰¹	1.92 × 10 ⁻⁰⁵	3.51 × 10 ⁻⁰⁴	6.85 × 10 ⁻⁰⁴	3.53 × 10 ⁻⁰³	1.97 × 10 ⁻⁰⁴	5.57 x 10 ⁻⁰⁴	4.87 × 10 ⁻⁰¹	6.64 × 10 ⁻⁰⁵	3.52 × 10 ⁻⁰⁴
		Lloyd Lake	4.45 × 10 ⁻⁰⁴	2.21 × 10 ⁻⁰³	1.96 × 10 ⁻⁰⁴	3.41 x 10 ⁻⁰⁴	2.95 × 10 ⁻⁰¹	4.27 × 10 ⁻⁰⁶	1.98 × 10 ⁻⁰⁴	4.54 × 10 ⁻⁰⁴	2.26 × 10 ⁻⁰³	1.96 × 10 ⁻⁰⁴	3.49 x 10 ⁻⁰⁴	3.02 × 10 ⁻⁰¹	9.08 × 10 ⁻⁰⁶	1.98 × 10 ⁻⁰⁴
		Reference (Broach Lake)	6.09 × 10 ⁻⁰⁴	3.54×10^{-04}	9.65 × 10 ⁻⁰⁴	1.85 x 10 ⁻⁰³	4.21 × 10 ⁻⁰¹	6.64 x 10 ⁻⁰⁶	6.62 × 10 ⁻⁰⁵	6.09 × 10 ⁻⁰⁴	3.54 × 10 ⁻⁰⁴	9.65 × 10 ⁻⁰⁴	1.85 x 10 ⁻⁰³	4.21 × 10 ⁻⁰¹	6.64 x 10 ⁻⁰⁶	6.62 × 10 ⁻⁰⁵
	Whitefish	Patterson Lake North Arm — West Basin	1.21 × 10 ⁻⁰³	7.19 × 10 ⁻⁰⁴	1.02 × 10 ⁻⁰³	4.54 x 10 ⁻⁰³	1.09 × 10 ⁺⁰⁰	1.48 x 10 ⁻⁰⁴	6.63 × 10 ⁻⁰⁴	1.50 × 10 ⁻⁰³	9.31 × 10 ⁻⁰⁴	1.05 × 10 ⁻⁰³	5.97 x 10 ⁻⁰³	1.42 × 10 ⁺⁰⁰	5.50 x 10 ⁻⁰⁴	6.63 × 10 ⁻⁰⁴
	Willensii	Patterson Lake South Arm	1.05 × 10 ⁻⁰³	6.21 × 10 ⁻⁰⁴	9.70 × 10 ⁻⁰⁴	3.48 x 10 ⁻⁰³	8.12 × 10 ⁻⁰¹	8.88 x 10 ⁻⁰⁵	2.31 × 10 ⁻⁰⁴	1.26 × 10 ⁻⁰³	7.76 × 10 ⁻⁰⁴	9.73 × 10 ⁻⁰⁴	4.35 x 10 ⁻⁰³	1.01 × 10 ⁺⁰⁰	3.22 x 10 ⁻⁰⁴	2.31 × 10 ⁻⁰⁴
		Beet Lake	8.57 × 10 ⁻⁰⁴	5.04 × 10 ⁻⁰⁴	9.66 × 10 ⁻⁰⁴	2.72 x 10 ⁻⁰³	6.25 × 10 ⁻⁰¹	4.94 x 10 ⁻⁰⁵	1.28 × 10 ⁻⁰⁴	9.78 × 10 ⁻⁰⁴	5.92 × 10 ⁻⁰⁴	9.66 × 10 ⁻⁰⁴	3.17 x 10 ⁻⁰³	7.28 × 10 ⁻⁰¹	1.71 x 10 ⁻⁰⁴	1.28 × 10 ⁻⁰⁴
		Lloyd Lake	6.35 × 10 ⁻⁰⁴	3.70×10^{-04}	9.65 × 10 ⁻⁰⁴	1.94 x 10 ⁻⁰³	4.42×10^{-01}	1.10 x 10 ⁻⁰⁵	7.18×10^{-05}	6.47×10^{-04}	3.79×10^{-04}	9.65 × 10 ⁻⁰⁴	1.99 x 10 ⁻⁰³	4.52 × 10 ⁻⁰¹	2.34 x 10 ⁻⁰⁵	7.18 × 10 ⁻⁰⁵
	Macrophytes	Reference (Broach Lake)	1.34×10^{-04}	4.59×10^{-04}	4.91 × 10 ⁻⁰⁴	6.07 x 10 ⁻⁰²	2.22 × 10 ⁻⁰²	6.66 × 10 ⁻⁰⁶	1.82 × 10 ⁻⁰⁵	1.34×10^{-04}	4.59×10^{-04}	4.91 × 10 ⁻⁰⁴	6.07 x 10 ⁻⁰²	2.22 × 10 ⁻⁰²	6.66 × 10 ⁻⁰⁶	1.82 × 10 ⁻⁰⁵
		Patterson Lake North Arm — West Basin	2.66 × 10 ⁻⁰⁴	9.32 × 10 ⁻⁰⁴	5.21 × 10 ⁻⁰⁴	1.49 x 10 ⁻⁰¹	5.74 × 10 ⁻⁰²	1.49 × 10 ⁻⁰⁴	1.83 × 10 ⁻⁰⁴	3.31 × 10 ⁻⁰⁴	1.21 × 10 ⁻⁰³	5.37 × 10 ⁻⁰⁴	1.95 x 10 ⁻⁰¹	7.51 × 10 ⁻⁰²	5.51 × 10 ⁻⁰⁴	1.83 × 10 ⁻⁰⁴
	Macrophytes	Patterson Lake South Arm	2.31 × 10 ⁻⁰⁴	8.05 × 10 ⁻⁰⁴	4.93 × 10 ⁻⁰⁴	1.14 x 10 ⁻⁰¹	4.28 × 10 ⁻⁰²	8.90 × 10 ⁻⁰⁵	6.37 × 10 ⁻⁰⁵	2.78 × 10 ⁻⁰⁴	1.01 × 10 ⁻⁰³	4.95 × 10 ⁻⁰⁴	1.43 x 10 ⁻⁰¹	5.32 × 10 ⁻⁰²	3.23 × 10 ⁻⁰⁴	6.37 × 10 ⁻⁰⁵
		Beet Lake	1.89 × 10 ⁻⁰⁴	6.54×10^{-04}	4.91 × 10 ⁻⁰⁴	8.90 x 10 ⁻⁰²	3.30 × 10 ⁻⁰²	4.95×10^{-05}	3.51 × 10 ⁻⁰⁵	2.15×10^{-04}	7.67×10^{-04}	4.92 × 10 ⁻⁰⁴	1.04 x 10 ⁻⁰¹	3.84 × 10 ⁻⁰²	1.71×10^{-04}	3.52×10^{-05}
Agustic		Lloyd Lake	1.40×10^{-04}	4.79×10^{-04}	4.91 × 10 ⁻⁰⁴	6.36 x 10 ⁻⁰²	2.33 × 10 ⁻⁰²	1.10 × 10 ⁻⁰⁵	1.98 × 10 ⁻⁰⁵	1.42×10^{-04}	4.91 × 10 ⁻⁰⁴	4.91 × 10 ⁻⁰⁴	6.52 x 10 ⁻⁰²	2.38 × 10 ⁻⁰²	2.34×10^{-05}	1.98 × 10 ⁻⁰⁵
Aquatic																+
Plants		Reference (Broach Lake) Patterson Lake North Arm — West Basin	6.96×10^{-05} 1.38×10^{-04}	3.99 × 10 ⁻⁰⁴ 8.11 × 10 ⁻⁰⁴	6.44×10^{-03} 6.83×10^{-03}	1.27 x 10 ⁻⁰² 3.10 x 10 ⁻⁰²	9.16×10^{-02} 2.37×10^{-01}	6.66×10^{-06} 1.49×10^{-04}	2.28 × 10 ⁻⁰⁴ 2.28 × 10 ⁻⁰³	6.96×10^{-05} 1.72×10^{-04}	3.99 × 10 ⁻⁰⁴ 1.05 × 10 ⁻⁰³	6.44×10^{-03} 7.04×10^{-03}	1.27 x 10 ⁻⁰² 4.08 x 10 ⁻⁰²	9.16×10^{-02} 3.10×10^{-01}	6.66×10^{-06} 5.51×10^{-04}	2.28 × 10 ⁻⁰⁴ 2.28 × 10 ⁻⁰³
	Phytoplankton		1.20 × 10 ⁻⁰⁴	7.00 × 10 ⁻⁰⁴	6.48 × 10 ⁻⁰³	2.38 x 10 ⁻⁰²	1.77 × 10 ⁻⁰¹	8.90 × 10 ⁻⁰⁵	7.96 × 10 ⁻⁰⁴	1.44 × 10 ⁻⁰⁴	8.75 × 10 ⁻⁰⁴	6.49 × 10 ⁻⁰³	2.97 x 10 ⁻⁰²	2.20 × 10 ⁻⁰¹	3.23 × 10 ⁻⁰⁴	7.97 × 10 ⁻⁰⁴
		Patterson Lake South Arm														
		Beet Lake	9.79 × 10 ⁻⁰⁵	5.69 × 10 ⁻⁰⁴	6.45 × 10 ⁻⁰³	1.86 x 10 ⁻⁰²	1.36 × 10 ⁻⁰¹	4.95 × 10 ⁻⁰⁵	4.39 × 10 ⁻⁰⁴	1.12 × 10 ⁻⁰⁴	6.67 × 10 ⁻⁰⁴	6.45 × 10 ⁻⁰³	2.17 x 10 ⁻⁰²	1.59 × 10 ⁻⁰¹	1.71 × 10 ⁻⁰⁴	4.39 × 10 ⁻⁰⁴
		Lloyd Lake	8.15 × 10 ⁻⁰⁵		6.44 × 10 ⁻⁰³	1.33 x 10 ⁻⁰²	9.62 × 10 ⁻⁰²	1.10 × 10 ⁻⁰⁵	2.47 × 10 ⁻⁰⁴		5.12 × 10 ⁻⁰⁴	6.44 × 10 ⁻⁰³				2.47 × 10 ⁻⁰⁴
		Reference (Broach Lake) Patterson Lake North Arm —	n/a n/a	n/a n/a	2.45 × 10 ⁻⁰⁴ 2.51 × 10 ⁻⁰⁴	2.94 × 10 ⁻⁰⁴ 5.04 × 10 ⁻⁰⁴	2.65 × 10 ⁻⁰⁴ 3.16 x 10 ⁻⁰⁴	3.09 × 10 ⁻⁰⁴ 8.92 × 10 ⁻⁰⁴	5.71 × 10 ⁻⁰⁴ 1.27 × 10 ⁻⁰³	n/a n/a	n/a n/a	2.45×10^{-04} 2.54×10^{-04}	2.94 × 10 ⁻⁰⁴ 6.15 x 10 ⁻⁰⁴	2.65 × 10 ⁻⁰⁴ 3.41 x 10 ⁻⁰⁴	3.09×10^{-04} 2.55×10^{-03}	5.71 × 10 ⁻⁰⁴ 1.27 × 10 ⁻⁰³
	Beaver	West Basin					2.									
		Patterson Lake South Arm	n/a	n/a	2.46 × 10 ⁻⁰⁴	4.21 x 10 ⁻⁰⁴	2.94 x 10 ⁻⁰⁴	6.47 × 10 ⁻⁰⁴	7.32 x 10 ⁻⁰⁴	n/a	n/a	2.46 × 10 ⁻⁰⁴	4.89 x 10 ⁻⁰⁴	3.09 x 10 ⁻⁰⁴	1.61 × 10 ⁻⁰³	7.32 x 10 ⁻⁰⁴
		Beet Lake	n/a	n/a	2.45 × 10 ⁻⁰⁴	3.62 x 10 ⁻⁰⁴	2.80 x 10 ⁻⁰⁴	4.85 × 10 ⁻⁰⁴	6.38 x 10 ⁻⁰⁴	n/a	n/a	2.45 × 10 ⁻⁰⁴	3.97 x 10 ⁻⁰⁴	2.88 x 10 ⁻⁰⁴	9.86 × 10 ⁻⁰⁴	6.38 x 10 ⁻⁰⁴
		Lloyd Lake	n/a	n/a	2.45 × 10 ⁻⁰⁴	3.01 x 10 ⁻⁰⁴	2.66 x 10 ⁻⁰⁴	3.27 × 10 ⁻⁰⁴	5.76 x 10 ⁻⁰⁴	n/a	n/a	2.45×10^{-04}	3.05 x 10 ⁻⁰⁴	2.67 x 10 ⁻⁰⁴	3.78 × 10 ⁻⁰⁴	5.76 x 10 ⁻⁰⁴
		Reference (Broach Lake)	n/a	n/a	9.22 × 10 ⁻⁰⁴	8.71×10^{-05}	2.49×10^{-03}	1.44 × 10 ⁻⁰⁴	1.82×10^{-04}	n/a	n/a	9.22×10^{-04}	8.71×10^{-05}	2.49×10^{-03}	1.44 × 10 ⁻⁰⁴	1.82×10^{-04}
Terrestrial Animals	Black Bear	Patterson Lake North Arm — West Basin	n/a	n/a	9.42 × 10 ⁻⁰⁴	1.22 x 10 ⁻⁰⁴	4.69 x 10 ⁻⁰³	2.34 × 10 ⁻⁰⁴	3.20 x 10 ⁻⁰⁴	n/a	n/a	9.52 × 10 ⁻⁰⁴	1.39 x 10 ⁻⁰⁴	5.79 x 10 ⁻⁰³	4.77 × 10 ⁻⁰⁴	3.20 x 10 ⁻⁰⁴
	Didek Deal	Beet Lake	n/a	n/a	1.01 × 10 ⁻⁰³	1.11 x 10 ⁻⁰⁴	2.94 x 10 ⁻⁰³	1.91 × 10 ⁻⁰⁴	2.15 x 10 ⁻⁰⁴	n/a	n/a	1.01 × 10 ⁻⁰³	1.15 x 10 ⁻⁰⁴	3.16 x 10 ⁻⁰³	2.37 × 10 ⁻⁰⁴	2.15 x 10 ⁻⁰⁴
		Lloyd Lake	n/a	n/a	9.25 × 10 ⁻⁰⁴	9.13 x 10 ⁻⁰⁵	2.58 x 10 ⁻⁰³	1.53 × 10 ⁻⁰⁴	1.85 x 10 ⁻⁰⁴	n/a	n/a	9.25×10^{-04}	9.21 x 10 ⁻⁰⁵	2.63 x 10 ⁻⁰³	1.63 × 10 ⁻⁰⁴	1.85 x 10 ⁻⁰⁴
		Reference (Broach Lake)	n/a	n/a	2.46 x 10 ⁻⁰³	4.49 x 10 ⁻⁰⁵	2.88 x 10 ⁻⁰³	1.11 x 10 ⁻⁰⁴	6.66 x 10 ⁻⁰⁵	n/a	n/a	2.46 x 10 ⁻⁰³	4.49 x 10 ⁻⁰⁵	2.88 x 10 ⁻⁰³	1.11 x 10 ⁻⁰⁴	6.66 x 10 ⁻⁰⁵
	Grey Wolf	Patterson Lake North Arm — West Basin	n/a	n/a	2.47 x 10 ⁻⁰³	4.81 x 10 ⁻⁰⁵	2.98 x 10 ⁻⁰³	1.72 x 10 ⁻⁰⁴	8.09 x 10 ⁻⁰⁵	n/a	n/a	2.48 x 10 ⁻⁰³	4.98 x 10 ⁻⁰⁵	3.03 x 10 ⁻⁰³	3.46 x 10 ⁻⁰⁴	8.09 x 10 ⁻⁰⁵
		Lloyd Lake	n/a	n/a	2.46 x 10 ⁻⁰³	4.53 x 10 ⁻⁰⁵	2.89 x 10 ⁻⁰³	1.17 x 10 ⁻⁰⁴	6.69 x 10 ⁻⁰⁵	n/a	n/a	2.46 x 10 ⁻⁰³	4.55 x 10 ⁻⁰⁵	2.89 x 10 ⁻⁰³	1.33 x 10 ⁻⁰⁴	6.69 x 10 ⁻⁰⁵
		•														

								Maximum HC	Q Far-Future						
Biota	Location				Application Cas		1					per Bound Sce	nario		
	Reference (Broach Lake)	Chloride n/a	Sulphate n/a	Arsenic 3.77 × 10 ⁻⁰³	Cobalt 3.45 × 10 ⁻⁰⁴	Copper 1.99 × 10 ⁻⁰²	Molybdenum 4.40 × 10 ⁻⁰⁴	Uranium 1.93 × 10 ⁻⁰⁴	Chloride n/a	Sulphate n/a	Arsenic 3.77 × 10 ⁻⁰³	Cobalt 3.45 × 10 ⁻⁰⁴	Copper 1.99 × 10 ⁻⁰²	Molybdenum 4.40 × 10 ⁻⁰⁴	Uranium 1.93 × 10 ⁻⁰⁴
	Patterson Lake North Arm — West Basin	n/a	n/a	3.98 × 10 ⁻⁰³	7.90 x 10 ⁻⁰⁴	5.13 x 10 ⁻⁰²	8.48 × 10 ⁻⁰³	1.50 x 10 ⁻⁰³	n/a	n/a	4.10 × 10 ⁻⁰³	1.03 x 10 ⁻⁰³	6.71 x 10 ⁻⁰²	3.13 × 10 ⁻⁰²	1.50 x 10 ⁻⁰³
American Mink	Patterson Lake South Arm	n/a	n/a	3.79 × 10 ⁻⁰³	6.15 x 10 ⁻⁰⁴	3.83 x 10 ⁻⁰²	5.11 × 10 ⁻⁰³	5.52 x 10 ⁻⁰⁴	n/a	n/a	3.80 × 10 ⁻⁰³	7.58 x 10 ⁻⁰⁴	4.76 x 10 ⁻⁰²	1.84 × 10 ⁻⁰²	5.52 x 10 ⁻⁰⁴
	Beet Lake	n/a	n/a	3.77×10^{-03}	4.87 x 10 ⁻⁰⁴	2.95 x 10 ⁻⁰²	2.87 × 10 ⁻⁰³	3.27 x 10 ⁻⁰⁴	n/a	n/a	3.78 × 10 ⁻⁰³	5.64 x 10 ⁻⁰⁴	3.44 x 10 ⁻⁰²	9.77 × 10 ⁻⁰³	3.27 x 10 ⁻⁰⁴
	Lloyd Lake	n/a	n/a	3.77×10^{-03}	3.59 x 10 ⁻⁰⁴	2.08 x 10 ⁻⁰²	6.88 × 10 ⁻⁰⁴	2.05 x 10 ⁻⁰⁴	n/a	n/a	3.77×10^{-03}	3.67 x 10 ⁻⁰⁴	2.13 x 10 ⁻⁰²	1.39 × 10 ⁻⁰³	2.05 x 10 ⁻⁰⁴
	Reference (Broach Lake)	n/a	n/a	1.63 × 10 ⁻⁰⁴	2.11 × 10 ⁻⁰⁴	1.07 × 10 ⁻⁰²	1.41 × 10 ⁻⁰⁴	2.84 × 10 ⁻⁰⁴	n/a	n/a	1.63 × 10 ⁻⁰⁴	2.11 × 10 ⁻⁰⁴	1.07 × 10 ⁻⁰²	1.41 × 10 ⁻⁰⁴	2.84 × 10 ⁻⁰⁴
Mana	Patterson Lake North Arm — West Basin	n/a	n/a	1.69 × 10 ⁻⁰⁴	4.20 x 10 ⁻⁰⁴	1.48 x 10 ⁻⁰²	6.77 × 10 ⁻⁰⁴	8.61 × 10 ⁻⁰⁴	n/a	n/a	1.73 × 10 ⁻⁰⁴	5.31 x 10 ⁻⁰⁴	1.68 x 10 ⁻⁰²	2.20 × 10 ⁻⁰³	8.62 x 10 ⁻⁰⁴
Moose	Patterson Lake South Arm	n/a	n/a	1.69 × 10 ⁻⁰⁴	4.14 x 10 ⁻⁰⁴	1.47 x 10 ⁻⁰²	6.16 × 10 ⁻⁰⁴	8.46 × 10 ⁻⁰⁴	n/a	n/a	1.72 × 10 ⁻⁰⁴	5.24 x 10 ⁻⁰⁴	1.67 x 10 ⁻⁰²	1.97 × 10 ⁻⁰³	8.47 x 10 ⁻⁰⁴
	Beet Lake	n/a	n/a	1.64 × 10 ⁻⁰⁴	2.78 x 10 ⁻⁰⁴	1.20 x 10 ⁻⁰²	3.03 × 10 ⁻⁰⁴	3.41 × 10 ⁻⁰⁴	n/a	n/a	1.64 × 10 ⁻⁰⁴	3.14 x 10 ⁻⁰⁴	1.26 x 10 ⁻⁰²	7.63 × 10 ⁻⁰⁴	3.42 x 10 ⁻⁰⁴
	Lloyd Lake	n/a	n/a	1.63 × 10 ⁻⁰⁴	2.18 x 10 ⁻⁰⁴	1.09 x 10 ⁻⁰²	1.58 × 10 ⁻⁰⁴	2.89 × 10 ⁻⁰⁴	n/a	n/a	1.63 × 10 ⁻⁰⁴	2.22 x 10 ⁻⁰⁴	1.09 x 10 ⁻⁰²	2.05 × 10 ⁻⁰⁴	2.89 x 10 ⁻⁰⁴
	Reference (Broach Lake)	n/a	n/a	2.01 × 10 ⁻⁰³	1.19 × 10 ⁻⁰³	3.42 × 10 ⁻⁰³	9.60 × 10 ⁻⁰⁴	9.83 × 10 ⁻⁰⁴	n/a	n/a	2.01 × 10 ⁻⁰³	1.19 × 10 ⁻⁰³	3.42 × 10 ⁻⁰³	9.60 × 10 ⁻⁰⁴	9.83 × 10 ⁻⁰⁴
Muskrat	Patterson Lake North Arm — West Basin	n/a	n/a	2.13 × 10 ⁻⁰³	2.92 × 10 ⁻⁰³	8.86 × 10 ⁻⁰³	2.14 × 10 ⁻⁰²	9.84 × 10 ⁻⁰³	n/a	n/a	2.20 × 10 ⁻⁰³	3.84 × 10 ⁻⁰³	1.16 × 10 ⁻⁰²	7.95 × 10 ⁻⁰²	9.85 × 10 ⁻⁰³
	Patterson Lake South Arm	n/a	n/a	2.02 × 10 ⁻⁰³	2.24×10^{-03}	6.61 × 10 ⁻⁰³	1.28 × 10 ⁻⁰²	3.43×10^{-03}	n/a	n/a	2.03 × 10 ⁻⁰³	2.80 × 10 ⁻⁰³	8.22 × 10 ⁻⁰³	4.65 × 10 ⁻⁰²	3.44 × 10 ⁻⁰³
	Beet Lake	n/a	n/a	2.01 × 10 ⁻⁰³	1.75×10^{-03}	5.09 × 10 ⁻⁰³	7.14 × 10 ⁻⁰³	1.89 × 10 ⁻⁰³	n/a	n/a	2.01 × 10 ⁻⁰³	2.04 × 10 ⁻⁰³	5.93 × 10 ⁻⁰³	2.47 × 10 ⁻⁰²	1.90 × 10 ⁻⁰³
	Lloyd Lake	n/a	n/a	2.01 × 10 ⁻⁰³	1.25 × 10 ⁻⁰³	3.59 × 10 ⁻⁰³	1.59 × 10 ⁻⁰³	1.07 × 10 ⁻⁰³	n/a	n/a	2.01 × 10 ⁻⁰³	1.28 × 10 ⁻⁰³	3.68 × 10 ⁻⁰³	3.38 × 10 ⁻⁰³	1.07 × 10 ⁻⁰³
	Reference (Broach Lake)	n/a	n/a	4.18 × 10 ⁻⁰⁴	3.66×10^{-05}	6.08×10^{-04}	1.22 × 10 ⁻⁰⁴	6.64×10^{-05}	n/a	n/a	4.18×10^{-04}	3.66×10^{-05}	6.08 × 10 ⁻⁰⁴	1.22 × 10 ⁻⁰⁴	6.64×10^{-05}
Red Fox	Patterson Lake North Arm — West Basin	n/a	n/a	4.18 × 10 ⁻⁰⁴	4.21 x 10 ⁻⁰⁵	6.19 x 10 ⁻⁰⁴	2.67 × 10 ⁻⁰⁴	1.09 x 10 ⁻⁰⁴	n/a	n/a	4.18 × 10 ⁻⁰⁴	4.50 x 10 ⁻⁰⁵	6.24 x 10 ⁻⁰⁴	6.78 × 10 ⁻⁰⁴	1.09 x 10 ⁻⁰⁴
	Patterson Lake South Arm	n/a	n/a	4.18 × 10 ⁻⁰⁴	3.99 x 10 ⁻⁰⁵	6.14 x 10 ⁻⁰⁴	2.06 × 10 ⁻⁰⁴	7.44 x 10 ⁻⁰⁵	n/a	n/a	4.18 × 10 ⁻⁰⁴	4.17 x 10 ⁻⁰⁵	6.17 x 10 ⁻⁰⁴	4.45 × 10 ⁻⁰⁴	7.44 x 10 ⁻⁰⁵
	Beet Lake	n/a	n/a	4.18 × 10 ⁻⁰⁴	3.84 x 10 ⁻⁰⁵	6.11 x 10 ⁻⁰⁴	1.66 × 10 ⁻⁰⁴	7.03 x 10 ⁻⁰⁵	n/a	n/a	4.18 × 10 ⁻⁰⁴	3.93 x 10 ⁻⁰⁵	6.12 x 10 ⁻⁰⁴	2.90 × 10 ⁻⁰⁴	7.03 x 10 ⁻⁰⁵
	Lloyd Lake	n/a	n/a	4.18 × 10 ⁻⁰⁴	3.68 x 10 ⁻⁰⁵	6.08 x 10 ⁻⁰⁴	1.27 × 10 ⁻⁰⁴	6.66 x 10 ⁻⁰⁵	n/a	n/a	4.18 × 10 ⁻⁰⁴	3.69 x 10 ⁻⁰⁵	6.08 x 10 ⁻⁰⁴	1.39 × 10 ⁻⁰⁴	6.66 x 10 ⁻⁰⁵
	Reference (Broach Lake)	n/a	n/a	1.43 × 10 ⁻⁰³	2.97×10^{-04}	1.06 × 10 ⁻⁰³	6.71 × 10 ⁻⁰⁵	9.42 × 10 ⁻⁰⁴	n/a	n/a	1.43 × 10 ⁻⁰³	2.97×10^{-04}	1.06 × 10 ⁻⁰³	6.71 × 10 ⁻⁰⁵	9.42 × 10 ⁻⁰⁴
Snowshoe Hare	Patterson Lake North Arm — West Basin	n/a	n/a	1.43 × 10 ⁻⁰³	3.04 x 10 ⁻⁰⁴	1.06 x 10 ⁻⁰³	8.10 × 10 ⁻⁰⁵	1.38 x 10 ⁻⁰³	n/a	n/a	1.43 × 10 ⁻⁰³	3.07 x 10 ⁻⁰⁴	1.06 x 10 ⁻⁰³	1.20 × 10 ⁻⁰⁴	1.38 x 10 ⁻⁰³
	Patterson Lake South Arm	n/a	n/a	1.43 × 10 ⁻⁰³	3.01 x 10 ⁻⁰⁴	1.06 x 10 ⁻⁰³	7.52 × 10 ⁻⁰⁵	1.01 x 10 ⁻⁰³	n/a	n/a	1.43 × 10 ⁻⁰³	3.03 x 10 ⁻⁰⁴	1.06 x 10 ⁻⁰³	9.80 × 10 ⁻⁰⁵	1.01 x 10 ⁻⁰³
	Beet Lake	n/a	n/a	1.43 × 10 ⁻⁰³	2.99 x 10 ⁻⁰⁴	1.06 x 10 ⁻⁰³	7.13 × 10 ⁻⁰⁵	9.78 x 10 ⁻⁰⁴	n/a	n/a	1.43 × 10 ⁻⁰³	3.00 x 10 ⁻⁰⁴	1.06 x 10 ⁻⁰³	8.32 × 10 ⁻⁰⁵	9.78 x 10 ⁻⁰⁴
	Lloyd Lake	n/a	n/a	1.43 × 10 ⁻⁰³	2.98 x 10 ⁻⁰⁴	1.06 x 10 ⁻⁰³	6.76 × 10 ⁻⁰⁵	9.43 x 10 ⁻⁰⁴	n/a	n/a	1.43 × 10 ⁻⁰³	2.98 x 10 ⁻⁰⁴	1.06 x 10 ⁻⁰³	6.88 × 10 ⁻⁰⁵	9.43 x 10 ⁻⁰⁴
	Reference (Broach Lake)	n/a	n/a	1.50 × 10 ⁻⁰⁴	1.05 × 10 ⁻⁰⁴	1.24 × 10 ⁻⁰⁴	5.79 × 10 ⁻⁰⁴	7.05 × 10 ⁻⁰⁴	n/a	n/a	1.50 × 10 ⁻⁰⁴	1.05 × 10 ⁻⁰⁴	1.24 × 10 ⁻⁰⁴	5.79 × 10 ⁻⁰⁴	7.05 × 10 ⁻⁰⁴
Southern Red-	Patterson Lake North Arm — West Basin	n/a	n/a	1.50 × 10 ⁻⁰⁴	1.10 x 10 ⁻⁰⁴	1.25 x 10 ⁻⁰⁴	8.14 × 10 ⁻⁰⁴	1.04 x 10 ⁻⁰³	n/a	n/a	1.50 × 10 ⁻⁰⁴	1.12 x 10 ⁻⁰⁴	1.25 x 10 ⁻⁰⁴	1.48 × 10 ⁻⁰³	1.04 x 10 ⁻⁰³
Backed Vole	Patterson Lake South Arm	n/a	n/a	1.50 × 10 ⁻⁰⁴	1.08 x 10 ⁻⁰⁴	1.24 x 10 ⁻⁰⁴	7.15 × 10 ⁻⁰⁴	7.56 x 10 ⁻⁰⁴	n/a	n/a	1.50×10^{-04}	1.09 x 10 ⁻⁰⁴	1.24 x 10 ⁻⁰⁴	1.10 × 10 ⁻⁰³	7.56 x 10 ⁻⁰⁴
	Beet Lake	n/a	n/a	1.50 × 10 ⁻⁰⁴	1.07 x 10 ⁻⁰⁴	1.24 x 10 ⁻⁰⁴	6.50 × 10 ⁻⁰⁴	7.33 x 10 ⁻⁰⁴	n/a	n/a		1.07 x 10 ⁻⁰⁴	1.24 x 10 ⁻⁰⁴	8.52 × 10 ⁻⁰⁴	7.33 x 10 ⁻⁰⁴
	Lloyd Lake	n/a	n/a	1.50 × 10 ⁻⁰⁴	1.05 x 10 ⁻⁰⁴	1.24 x 10 ⁻⁰⁴	5.86 × 10 ⁻⁰⁴	7.06 x 10 ⁻⁰⁴	n/a	n/a	1.50×10^{-04}	1.05 x 10 ⁻⁰⁴	1.24 x 10 ⁻⁰⁴	6.06 × 10 ⁻⁰⁴	7.06 x 10 ⁻⁰⁴
	Reference (Broach Lake)	n/a	n/a	3.79 x 10 ⁻⁰⁴	2.94 x 10 ⁻⁰⁴	1.19 x 10 ⁻⁰²	3.02 x 10 ⁻⁰⁴	3.16 x 10 ⁻⁰⁴	n/a	n/a	3.79 x 10 ⁻⁰⁴	2.94 x 10 ⁻⁰⁴	1.19 x 10 ⁻⁰²	3.02 x 10 ⁻⁰⁴	3.16 x 10 ⁻⁰⁴
Woodland Caribou	Patterson Lake North Arm — West Basin	n/a	n/a	3.87 x 10 ⁻⁰⁴	4.96 x 10 ⁻⁰⁴	1.56 x 10 ⁻⁰²	1.35 x 10 ⁻⁰³	1.09 x 10 ⁻⁰³	n/a	n/a	3.91 x 10 ⁻⁰⁴	6.04 x 10 ⁻⁰⁴	1.75 x 10 ⁻⁰²	4.33 x 10 ⁻⁰³	1.09 x 10 ⁻⁰³
	Beet Lake	n/a	n/a	3.79 x 10 ⁻⁰⁴	3.35 x 10 ⁻⁰⁴	1.26 x 10 ⁻⁰²	5.05 x 10 ⁻⁰⁴	3.80 x 10 ⁻⁰⁴	n/a	n/a	3.79 x 10 ⁻⁰⁴	3.57 x 10 ⁻⁰⁴	1.30 x 10 ⁻⁰²	1.08 x 10 ⁻⁰³	3.80 x 10 ⁻⁰⁴
	Lloyd Lake	n/a	n/a	3.79 x 10 ⁻⁰⁴	3.03 x 10 ⁻⁰⁴	1.20 x 10 ⁻⁰²	3.45 x 10 ⁻⁰⁴	3.28 x 10 ⁻⁰⁴	n/a	n/a	3.79 x 10 ⁻⁰⁴	3.08 x 10 ⁻⁰⁴	1.21 x 10 ⁻⁰²	4.67 x 10 ⁻⁰⁴	3.28 x 10 ⁻⁰⁴
	Reference (Broach Lake)	n/a	n/a	1.79 × 10 ⁻⁰³	3.95 × 10 ⁻⁰⁴	1.51 × 10 ⁻⁰³	6.67 × 10 ⁻⁰⁵	3.96 × 10 ⁻⁰⁴	n/a	n/a	1.79 × 10 ⁻⁰³	3.95 × 10 ⁻⁰⁴	1.51 × 10 ⁻⁰³	6.67 × 10 ⁻⁰⁵	3.96 × 10 ⁻⁰⁴
Grouse	Patterson Lake North Arm — West Basin	n/a	n/a	1.79 × 10 ⁻⁰³	4.00 x 10 ⁻⁰⁴	1.51 x 10 ⁻⁰³	7.47 × 10 ⁻⁰⁵	5.77 x 10 ⁻⁰⁴	n/a	n/a	1.79 × 10 ⁻⁰³	4.02 x 10 ⁻⁰⁴	1.51 x 10 ⁻⁰³	9.73 × 10 ⁻⁰⁵	5.77 x 10 ⁻⁰⁴
	Patterson Lake South Arm	n/a	n/a	1.79 × 10 ⁻⁰³	3.98 x 10 ⁻⁰⁴	1.51 x 10 ⁻⁰³	7.13 × 10 ⁻⁰⁵	4.23 x 10 ⁻⁰⁴	n/a	n/a	1.79 × 10 ⁻⁰³	3.99 x 10 ⁻⁰⁴	1.51 x 10 ⁻⁰³	8.44 × 10 ⁻⁰⁵	4.23 x 10 ⁻⁰⁴
	Beet Lake	n/a	n/a	1.79 × 10 ⁻⁰³	3.96 x 10 ⁻⁰⁴	1.51 x 10 ⁻⁰³	6.91 × 10 ⁻⁰⁵	4.12 x 10 ⁻⁰⁴	n/a	n/a	1.79 × 10 ⁻⁰³	3.97 x 10 ⁻⁰⁴	1.51 x 10 ⁻⁰³	7.59 × 10 ⁻⁰⁵	4.12 x 10 ⁻⁰⁴
	Lloyd Lake	n/a	n/a	1.79 × 10 ⁻⁰³	3.95 x 10 ⁻⁰⁴	1.51 x 10 ⁻⁰³	6.69 × 10 ⁻⁰⁵	3.97 x 10 ⁻⁰⁴	n/a	n/a	1.79 × 10 ⁻⁰³	3.95 x 10 ⁻⁰⁴	1.51 x 10 ⁻⁰³	6.76 × 10 ⁻⁰⁵	3.97 x 10 ⁻⁰⁴
	Reference (Broach Lake)	n/a	n/a	1.72 × 10 ⁻⁰⁴	3.93 × 10 ⁻⁰⁵	1.96 × 10 ⁻⁰⁴	1.24 × 10 ⁻⁰⁵	3.93 × 10 ⁻⁰⁵	n/a	n/a	1.72 × 10 ⁻⁰⁴	3.93×10^{-05}	1.96 × 10 ⁻⁰⁴	1.24 × 10 ⁻⁰⁵	3.93 × 10 ⁻⁰⁵
Canada Goose	Patterson Lake North Arm — West Basin	n/a	n/a	1.72 × 10 ⁻⁰⁴	4.17 x 10 ⁻⁰⁵	1.99 x 10 ⁻⁰⁴	2.05 × 10 ⁻⁰⁵	5.89 x 10 ⁻⁰⁵	n/a	n/a	1.72 × 10 ⁻⁰⁴	4.30 x 10 ⁻⁰⁵	2.00 x 10 ⁻⁰⁴	4.35 × 10 ⁻⁰⁵	5.89 x 10 ⁻⁰⁵
	Patterson Lake South Arm	n/a	n/a	1.72 × 10 ⁻⁰⁴	4.08 x 10 ⁻⁰⁵	1.97 x 10 ⁻⁰⁴	1.71 × 10 ⁻⁰⁵	4.24 x 10 ⁻⁰⁵	n/a	n/a	1.72 × 10 ⁻⁰⁴	4.15 x 10 ⁻⁰⁵	1.98 x 10 ⁻⁰⁴	3.04 × 10 ⁻⁰⁵	4.24 x 10 ⁻⁰⁵
	Beet Lake	n/a	n/a	1.72×10^{-04}	4.01 x 10 ⁻⁰⁵	1.97 x 10 ⁻⁰⁴	1.48×10^{-05}	4.10 x 10 ⁻⁰⁵	n/a	n/a	1.72×10^{-04}	4.05 x 10 ⁻⁰⁵	1.97 x 10 ⁻⁰⁴	2.18×10^{-05}	4.10 x 10 ⁻⁰⁵

ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT

ECOLOGICAL RISK ASSESSMENT

								Maximum HO) Far-Future						
Biota	Location				Application Cas	e					Uŗ	per Bound Sce	nario		
		Chloride	Sulphate	Arsenic	Cobalt	Copper	Molybdenum	Uranium	Chloride	Sulphate	Arsenic	Cobalt	Copper	Molybdenum	Uranium
	Lloyd Lake	n/a	n/a	1.72 × 10 ⁻⁰⁴	3.95 x 10 ⁻⁰⁵	1.96 x 10 ⁻⁰⁴	1.26 × 10 ⁻⁰⁵	3.94 x 10 ⁻⁰⁵	n/a	n/a	1.72 × 10 ⁻⁰⁴	3.95 x 10 ⁻⁰⁵	1.96 x 10 ⁻⁰⁴	1.33 × 10 ⁻⁰⁵	3.94 x 10 ⁻⁰⁵
	Reference (Broach Lake)	n/a	n/a	9.07×10^{-03}	4.28×10^{-03}	2.80×10^{-01}	6.31×10^{-03}	8.52 × 10 ⁻⁰⁴	n/a	n/a	9.07×10^{-03}	4.28×10^{-03}	2.80×10^{-01}	6.31 × 10 ⁻⁰³	8.52 × 10 ⁻⁰⁴
Little Brown	Patterson Lake North Arm — West Basin	n/a	n/a	9.61 × 10 ⁻⁰³	1.05 × 10 ⁻⁰²	7.24 × 10 ⁻⁰¹	1.41 × 10 ⁻⁰¹	8.53 × 10 ⁻⁰³	n/a	n/a	9.91 × 10 ⁻⁰³	1.38 × 10 ⁻⁰²	9.48 × 10 ⁻⁰¹	5.23 × 10 ⁻⁰¹	8.54 × 10 ⁻⁰³
Myotis	Patterson Lake South Arm	n/a	n/a	9.11 × 10 ⁻⁰³	8.03 × 10 ⁻⁰³	5.41 × 10 ⁻⁰¹	8.43 × 10 ⁻⁰²	2.98 × 10 ⁻⁰³	n/a	n/a	9.13 × 10 ⁻⁰³	1.00 × 10 ⁻⁰²	6.72 × 10 ⁻⁰¹	3.06 × 10 ⁻⁰¹	2.98 × 10 ⁻⁰³
	Beet Lake	n/a	n/a	9.07 × 10 ⁻⁰³	6.26 × 10 ⁻⁰³	4.16 × 10 ⁻⁰¹	4.69 × 10 ⁻⁰²	1.64 × 10 ⁻⁰³	n/a	n/a	9.08 × 10 ⁻⁰³	7.32 × 10 ⁻⁰³	4.85 × 10 ⁻⁰¹	1.62 × 10 ⁻⁰¹	1.64 × 10 ⁻⁰³
	Lloyd Lake	n/a	n/a	9.07 × 10 ⁻⁰³	4.48×10^{-03}	2.94 × 10 ⁻⁰¹	1.04 × 10 ⁻⁰²	9.24 × 10 ⁻⁰⁴	n/a	n/a	9.07 × 10 ⁻⁰³	4.59 × 10 ⁻⁰³	3.01 × 10 ⁻⁰¹	2.22 × 10 ⁻⁰²	9.24 × 10 ⁻⁰⁴
	Reference (Broach Lake)	n/a	n/a	1.31 × 10 ⁻⁰³	1.54 × 10 ⁻⁰⁴	4.62 × 10 ⁻⁰³	1.97 × 10 ⁻⁰⁵	2.11 × 10 ⁻⁰⁵	n/a	n/a	1.31 × 10 ⁻⁰³	1.54 × 10 ⁻⁰⁴	4.62 × 10 ⁻⁰³	1.97 × 10 ⁻⁰⁵	2.11 × 10 ⁻⁰⁵
Common Loon	Patterson Lake North Arm — West Basin	n/a	n/a	1.39 × 10 ⁻⁰³	3.78 × 10 ⁻⁰⁴	1.19 × 10 ⁻⁰²	4.39 × 10 ⁻⁰⁴	2.11 × 10 ⁻⁰⁴	n/a	n/a	1.43 × 10 ⁻⁰³	4.96 × 10 ⁻⁰⁴	1.56 × 10 ⁻⁰²	1.63 × 10 ⁻⁰³	2.11 × 10 ⁻⁰⁴
	Patterson Lake South Arm	n/a	n/a	1.32 × 10 ⁻⁰³	2.90 × 10 ⁻⁰⁴	8.92 × 10 ⁻⁰³	2.63 × 10 ⁻⁰⁴	7.37 × 10 ⁻⁰⁵	n/a	n/a	1.32 × 10 ⁻⁰³	3.62 × 10 ⁻⁰⁴	1.11 × 10 ⁻⁰²	9.54 × 10 ⁻⁰⁴	7.38 × 10 ⁻⁰⁵
	Beet Lake	n/a	n/a	1.31 × 10 ⁻⁰³	2.26 × 10 ⁻⁰⁴	6.86 × 10 ⁻⁰³	1.46 × 10 ⁻⁰⁴	4.07 × 10 ⁻⁰⁵	n/a	n/a	1.31 × 10 ⁻⁰³	2.64 × 10 ⁻⁰⁴	7.99 × 10 ⁻⁰³	5.06 × 10 ⁻⁰⁴	4.07 × 10 ⁻⁰⁵
	Lloyd Lake	n/a	n/a	1.31 × 10 ⁻⁰³	1.62 × 10 ⁻⁰⁴	4.85 × 10 ⁻⁰³	3.26×10^{-05}	2.29 × 10 ⁻⁰⁵	n/a	n/a	1.31 × 10 ⁻⁰³	1.65 × 10 ⁻⁰⁴	4.96 × 10 ⁻⁰³	6.93 × 10 ⁻⁰⁵	2.29×10^{-05}
	Reference (Broach Lake)	n/a	n/a	5.44×10^{-03}	1.85×10^{-03}	1.40×10^{-02}	3.13×10^{-04}	3.27×10^{-04}	n/a	n/a	5.44×10^{-03}	1.85 × 10 ⁻⁰³	1.40×10^{-02}	3.13×10^{-04}	3.27×10^{-04}
Mallard	Patterson Lake North Arm — West Basin	n/a	n/a	5.77 × 10 ⁻⁰³	4.54 × 10 ⁻⁰³	3.63 × 10 ⁻⁰²	6.99 × 10 ⁻⁰³	3.27 × 10 ⁻⁰³	n/a	n/a	5.95 × 10 ⁻⁰³	5.97 × 10 ⁻⁰³	4.74 × 10 ⁻⁰²	2.59 × 10 ⁻⁰²	3.27 × 10 ⁻⁰³
ivialiaiu	Patterson Lake South Arm	n/a	n/a	5.47×10^{-03}	3.48×10^{-03}	2.71 × 10 ⁻⁰²	4.19 × 10 ⁻⁰³	1.14 × 10 ⁻⁰³	n/a	n/a	5.48 × 10 ⁻⁰³	4.35 × 10 ⁻⁰³	3.36 × 10 ⁻⁰²	1.52 × 10 ⁻⁰²	1.14 × 10 ⁻⁰³
	Beet Lake	n/a	n/a	5.45 × 10 ⁻⁰³	2.72 × 10 ⁻⁰³	2.08 × 10 ⁻⁰²	2.33 × 10 ⁻⁰³	6.29 × 10 ⁻⁰⁴	n/a	n/a	5.45 × 10 ⁻⁰³	3.17 × 10 ⁻⁰³	2.43 × 10 ⁻⁰²	8.06 × 10 ⁻⁰³	6.29 × 10 ⁻⁰⁴
	Lloyd Lake	n/a	n/a	5.44 × 10 ⁻⁰³	1.94 × 10 ⁻⁰³	1.47 × 10 ⁻⁰²	5.19 × 10 ⁻⁰⁴	3.54 × 10 ⁻⁰⁴	n/a	n/a	5.44 × 10 ⁻⁰³	1.99 × 10 ⁻⁰³	1.51 × 10 ⁻⁰²	1.10 × 10 ⁻⁰³	3.54 × 10 ⁻⁰⁴
	Reference (Broach Lake)	n/a	n/a	1.99 × 10 ⁻⁰²	8.87 × 10 ⁻⁰³	2.35 × 10 ⁻⁰¹	1.76 × 10 ⁻⁰³	1.10 × 10 ⁻⁰³	n/a	n/a	1.99 × 10 ⁻⁰²	8.87 × 10 ⁻⁰³	2.35 × 10 ⁻⁰¹	1.76 × 10 ⁻⁰³	1.10 × 10 ⁻⁰³
Decate: Dischlaring	Patterson Lake North Arm — West Basin	n/a	n/a	2.08 × 10 ⁻⁰²	2.04 x 10 ⁻⁰²	6.07 x 10 ⁻⁰¹	3.47 × 10 ⁻⁰²	6.59 x 10 ⁻⁰³	n/a	n/a	2.13 × 10 ⁻⁰²	2.65 x 10 ⁻⁰²	7.94 x 10 ⁻⁰¹	1.28 × 10 ⁻⁰¹	6.60 x 10 ⁻⁰³
Rusty Blackbird	Patterson Lake South Arm	n/a	n/a	2.00 × 10 ⁻⁰²	1.59 x 10 ⁻⁰²	4.53 x 10 ⁻⁰¹	2.09 × 10 ⁻⁰²	2.59 x 10 ⁻⁰³	n/a	n/a	2.00 × 10 ⁻⁰²	1.96 x 10 ⁻⁰²	5.63 x 10 ⁻⁰¹	7.51 × 10 ⁻⁰²	2.59 x 10 ⁻⁰³
	Beet Lake	n/a	n/a	1.99 × 10 ⁻⁰²	1.26 x 10 ⁻⁰²	3.49 x 10 ⁻⁰¹	1.17 × 10 ⁻⁰²	1.66 x 10 ⁻⁰³	n/a	n/a	1.99 × 10 ⁻⁰²	1.45 x 10 ⁻⁰²	4.07 x 10 ⁻⁰¹	4.00 × 10 ⁻⁰²	1.66 x 10 ⁻⁰³
	Lloyd Lake	n/a	n/a	1.99 × 10 ⁻⁰²	9.25 x 10 ⁻⁰³	2.47 x 10 ⁻⁰¹	2.77 × 10 ⁻⁰³	1.15 x 10 ⁻⁰³	n/a	n/a	1.99 × 10 ⁻⁰²	9.45 x 10 ⁻⁰³	2.53 x 10 ⁻⁰¹	5.65 × 10 ⁻⁰³	1.15 x 10 ⁻⁰³

Note: **Bold** indicates the HQ is greater than 1.

n/a = not applicable; HQ = hazard quotient.

6.4.1.1.2 Radiological Risk

An HQ is not typically calculated for radiological risk; however, a comparison against the ecological dose benchmarks is discussed.

There were no predicted exceedances of the 9.6 mGy/d radiation dose benchmark for aquatic biota in all basins of Patterson Lake and all downstream waterbodies for the Application Case and reasonable upper bound sensitivity scenario during both the Project phases and far-future projection. Additionally, there were no predicted exceedances of the 2.4 mGy/d radiation dose benchmark for terrestrial and riparian biota at or near the Project site, LSA, or RSA for the Application Case and reasonable upper bound sensitivity scenario during both the Project phases and far-future projection.

6.4.1.2 Estimated Risk to Ecological Receptors – Reasonably Foreseeable Development Case

6.4.1.2.1 Non-radiological Risk

With the addition of the Fission Patterson Lake South Property, there were no predicted exceedances of the HQ of 1 for any of the ecological receptors in Operations (Table 6-25).

For assessment of risk to benthic invertebrates, risk was calculated based on toxicity benchmarks as water concentrations. However, considering that benthic invertebrates also reside in sediment, a comparison of predicted sediment concentrations against sediment toxicity benchmarks was warranted for arsenic based on the sediment screening conducted in Table 4-3. Arsenic concentrations in sediment at Patterson Lake North Arm – West Basin and at Patterson Lake South Arm were predicted to exceed the REF screening value from Burnett-Seidel and Liber (2013) but were below the NE2 value for the RFD Case at all waterbodies.

The results for the far-future projection in the RFD Case were the same as for the far-future projection in the Application Case, as discussed in Section 6.4.1.1.1, Non-radiological Risk.

6.4.1.2.2 Radiological Risk

A HQ is not typically calculated for radiological risk; however, a comparison against the ecological dose benchmarks is discussed.

With the addition of the Fission Patterson Lake South Property, there were no predicted exceedances of the 9.6 mGy/d radiation dose benchmark for aquatic biota, in all basins of Patterson Lake and all downstream waterbodies. Additionally, there were no exceedances of the 2.4 mGy/d radiation dose benchmark for terrestrial and riparian biota at or near the Project site, LSA, or RSA for the RFD Case.



Table 6-24: Non-radiological Risk to Ecological Receptors – Reasonably Foreseeable Development Case – Operations and Far-Future Projection

	Biota	Location				Operation	,			n HQ RFD Case			Far-Futur	Δ		
	DiOla	Location	Chloride	Sulphate	Arsenic	Cobalt	Copper	Molybdenum	Uranium	Chloride	Sulphate	Arsenic	Cobalt	Copper	Molybdenum	Uranium
		Reference (Broach Lake)	1.00 × 10 ⁻⁰³	1.45 × 10 ⁻⁰³		4.75 x 10 ⁻⁰²	4.18 × 10 ⁻⁰¹	6.61 × 10 ⁻⁰⁶	3.57 × 10 ⁻⁰³	1.00 × 10 ⁻⁰³		8.50 × 10 ⁻⁰⁴	4.75 x 10 ⁻⁰²	4.18 × 10 ⁻⁰¹	6.61 × 10 ⁻⁰⁶	3.57 × 10
	Benthic Invertebrate	Patterson Lake North Arm — West Basin	1.03 × 10 ⁻⁰²	2.93 × 10 ⁻⁰¹	2.59 × 10 ⁻⁰³	6.15 x 10 ⁻⁰²	4.86 × 10 ⁻⁰¹	1.77 × 10 ⁻⁰⁵	1.17 × 10 ⁻⁰²	1.99 × 10 ⁻⁰³	2.95 × 10 ⁻⁰³	9.01 × 10 ⁻⁰⁴	1.16 x 10 ⁻⁰¹	1.08 × 10 ⁺⁰⁰	1.47 × 10 ⁻⁰⁴	3.57 × 10
	bentinc invertebrate	Patterson Lake South Arm	1.01 × 10 ⁻⁰²	2.92 × 10 ⁻⁰¹	1.95 × 10 ⁻⁰³	5.78 x 10 ⁻⁰²	4.69 × 10 ⁻⁰¹	1.41 × 10 ⁻⁰⁵	8.26 × 10 ⁻⁰³	1.73 × 10 ⁻⁰³	2.55 × 10 ⁻⁰³	8.54 × 10 ⁻⁰⁴	8.92 x 10 ⁻⁰²	8.08 × 10 ⁻⁰¹	8.83 × 10 ⁻⁰⁵	1.25 × 10
		Beet Lake	5.63 × 10 ⁻⁰³	1.49 × 10 ⁻⁰¹	1.04 × 10 ⁻⁰³	5.30 x 10 ⁻⁰²	4.45 × 10 ⁻⁰¹	1.05 × 10 ⁻⁰⁵	5.30 × 10 ⁻⁰³	1.41 × 10 ⁻⁰³	2.07×10^{-03}	8.51 × 10 ⁻⁰⁴	6.96 x 10 ⁻⁰²	6.22 × 10 ⁻⁰¹	4.92 × 10 ⁻⁰⁵	6.88 × 10
		Lloyd Lake	1.49 × 10 ⁻⁰³	1.71 × 10 ⁻⁰²	8.61 × 10 ⁻⁰⁴	4.81 x 10 ⁻⁰²	4.21 × 10 ⁻⁰¹	7.01 × 10 ⁻⁰⁶	3.73 × 10 ⁻⁰³	1.04 × 10 ⁻⁰³	1.52 × 10 ⁻⁰³	8.50 × 10 ⁻⁰⁴	4.98 x 10 ⁻⁰²	4.39 × 10 ⁻⁰¹	1.09 × 10 ⁻⁰⁵	3.87 × 10
		Reference (Broach Lake)	1.00×10^{-03}	2.50×10^{-03}	3.64×10^{-04}	5.25 x 10 ⁻⁰²	4.21 × 10 ⁻⁰¹	6.66×10^{-06}	1.67 × 10 ⁻⁰³	1.00×10^{-03}	2.50×10^{-03}	3.64×10^{-04}	5.25 x 10 ⁻⁰²	4.21 × 10 ⁻⁰¹	6.66 × 10 ⁻⁰⁶	1.67 × 10
	Zooplankton	Patterson Lake North Arm — West Basin	1.03 × 10 ⁻⁰²	5.04 × 10 ⁻⁰¹	1.28 × 10 ⁻⁰³	6.94 x 10 ⁻⁰²	4.97 × 10 ⁻⁰¹	1.93 × 10 ⁻⁰⁵	6.39 × 10 ⁻⁰³	1.99 × 10 ⁻⁰³	5.07 × 10 ⁻⁰³	3.86 × 10 ⁻⁰⁴	1.29 x 10 ⁻⁰¹	1.09 × 10 ⁺⁰⁰	1.49 × 10 ⁻⁰⁴	1.67 × 10
	Zoopialiktoli	Patterson Lake South Arm	1.01×10^{-02}	5.04 × 10 ⁻⁰¹	2.04×10^{-03}	7.04 x 10 ⁻⁰²	5.04 × 10 ⁻⁰¹	1.84×10^{-05}	8.94 × 10 ⁻⁰³	1.73 × 10 ⁻⁰³	4.38×10^{-03}	3.66×10^{-04}	9.87 x 10 ⁻⁰²	8.14 × 10 ⁻⁰¹	8.90 × 10 ⁻⁰⁵	5.84 × 10
		Beet Lake	5.63×10^{-03}	2.56 × 10 ⁻⁰¹	6.50×10^{-04}	6.04 x 10 ⁻⁰²	4.57×10^{-01}	1.19 × 10 ⁻⁰⁵	4.20×10^{-03}	1.41 × 10 ⁻⁰³	3.56×10^{-03}	3.64×10^{-04}	7.70 x 10 ⁻⁰²	6.27×10^{-01}	4.95 × 10 ⁻⁰⁵	3.22 × 10
Aquatic		Lloyd Lake	1.49×10^{-03}	2.94 × 10 ⁻⁰²	3.81×10^{-04}	5.33 x 10 ⁻⁰²	4.25 × 10 ⁻⁰¹	7.18×10^{-06}	1.91 × 10 ⁻⁰³	1.04×10^{-03}	2.61×10^{-03}	3.64×10^{-04}	5.50 x 10 ⁻⁰²	4.42×10^{-01}	1.10 × 10 ⁻⁰⁵	1.81 × 1
Animals		Reference (Broach Lake)	4.27×10^{-04}	2.11 × 10 ⁻⁰³	1.96 × 10 ⁻⁰⁴	3.26 x 10 ⁻⁰⁴	2.81 × 10 ⁻⁰¹	2.58 × 10 ⁻⁰⁶	1.82×10^{-04}	4.27×10^{-04}	2.11 × 10 ⁻⁰³	1.96 × 10 ⁻⁰⁴	3.26 x 10 ⁻⁰⁴	2.81 × 10 ⁻⁰¹	2.58 × 10 ⁻⁰⁶	1.82 × 10
	Northern Pike	Patterson Lake North Arm — West Basin	4.39 × 10 ⁻⁰³	4.27 × 10 ⁻⁰¹	6.92 × 10 ⁻⁰⁴	4.30 x 10 ⁻⁰⁴	3.32 × 10 ⁻⁰¹	7.48 × 10 ⁻⁰⁶	6.97 × 10 ⁻⁰⁴	8.49 × 10 ⁻⁰⁴	4.30 × 10 ⁻⁰³	2.08 × 10 ⁻⁰⁴	7.97 x 10 ⁻⁰⁴	7.27 × 10 ⁻⁰¹	5.75 × 10 ⁻⁰⁵	1.83 × 1
	Northern Fike	Patterson Lake South Arm	4.32 × 10 ⁻⁰³	4.27 × 10 ⁻⁰¹	1.10 × 10 ⁻⁰³	4.36 x 10 ⁻⁰⁴	3.36 × 10 ⁻⁰¹	7.15 × 10 ⁻⁰⁶	9.76 × 10 ⁻⁰⁴	7.35 × 10 ⁻⁰⁴	3.71 × 10 ⁻⁰³	1.97 × 10 ⁻⁰⁴	6.12 x 10 ⁻⁰⁴	5.43 × 10 ⁻⁰¹	3.45 × 10 ⁻⁰⁵	6.37 × 1
		Beet Lake	2.40×10^{-03}	2.17 × 10 ⁻⁰¹	3.51 × 10 ⁻⁰⁴	3.75 x 10 ⁻⁰⁴	3.05 × 10 ⁻⁰¹	4.62 × 10 ⁻⁰⁶	4.58 × 10 ⁻⁰⁴	6.01 × 10 ⁻⁰⁴	3.01 × 10 ⁻⁰³	1.97 × 10 ⁻⁰⁴	4.77 x 10 ⁻⁰⁴	4.18 × 10 ⁻⁰¹	1.92 × 10 ⁻⁰⁵	3.51 × 1
		Lloyd Lake	6.35 × 10 ⁻⁰⁴	2.49 × 10 ⁻⁰²	2.06×0^{-04}	3.30 x 10 ⁻⁰⁴	2.83 × 10 ⁻⁰¹	2.78 × 10 ⁻⁰⁶	2.08 × 10 ⁻⁰⁴	4.45 × 10 ⁻⁰⁴	2.21 × 10 ⁻⁰³	1.96 × 10 ⁻⁰⁴	3.41 x 10 ⁻⁰⁴	2.95 × 10 ⁻⁰¹	4.27 × 10 ⁻⁰⁶	1.98 × 1
		Reference (Broach Lake)	6.09 × 10 ⁻⁰⁴	3.54 × 10 ⁻⁰⁴	9.65 × 10 ⁻⁰⁴	1.85 x 10 ⁻⁰³	4.21 × 10 ⁻⁰¹	6.64 x 10 ⁻⁰⁶	6.62 × 10 ⁻⁰⁵	6.09 × 10 ⁻⁰⁴	3.54 × 10 ⁻⁰⁴	9.65 × 10 ⁻⁰⁴	1.85 x 10 ⁻⁰³	4.21 × 10 ⁻⁰¹	6.64 x 10 ⁻⁰⁶	6.62 × 1
	Whitefish	Patterson Lake North Arm — West Basin	6.27 × 10 ⁻⁰³	7.14 × 10 ⁻⁰²	3.30 × 10 ⁻⁰³	2.44 x 10 ⁻⁰³	4.95 × 10 ⁻⁰¹	1.89 x 10 ⁻⁰⁵	2.44 × 10 ⁻⁰⁴	1.21 × 10 ⁻⁰³	7.19 × 10 ⁻⁰⁴	1.02 × 10 ⁻⁰³	4.54 x 10 ⁻⁰³	1.09 × 10 ⁺⁰⁰	1.48 x 10 ⁻⁰⁴	6.63 × 1
	willensii	Patterson Lake South Arm	6.15 × 10 ⁻⁰³	7.14 × 10 ⁻⁰²	4.71 × 10 ⁻⁰³	2.43 x 10 ⁻⁰³	4.95 × 10 ⁻⁰¹	1.74 x 10 ⁻⁰⁵	3.05 × 10 ⁻⁰⁴	1.05 × 10 ⁻⁰³	6.21 × 10 ⁻⁰⁴	9.70 × 10 ⁻⁰⁴	3.48 x 10 ⁻⁰³	8.12 × 10 ⁻⁰¹	8.88 x 10 ⁻⁰⁵	2.31 × 1
		Beet Lake	3.42×10^{-03}	3.63 × 10 ⁻⁰²	1.60 × 10 ⁻⁰³	2.12 x 10 ⁻⁰³	4.54 × 10 ⁻⁰¹	1.16 x 10 ⁻⁰⁵	1.50 × 10 ⁻⁰⁴	8.57 × 10 ⁻⁰⁴	5.04 × 10 ⁻⁰⁴	9.66 × 10 ⁻⁰⁴	2.72 x 10 ⁻⁰³	6.25 × 10 ⁻⁰¹	4.94 x 10 ⁻⁰⁵	1.28 × 1
		Lloyd Lake	9.07×10^{-04}	4.17×10^{-03}	1.00×10^{-03}	1.88 x 10 ⁻⁰³	4.24×10^{-01}	7.14 x 10 ⁻⁰⁶	7.40×10^{-05}	6.35 × 10 ⁻⁰⁴	3.70×10^{-04}	9.65 × 10 ⁻⁰⁴	1.94 x 10 ⁻⁰³	4.42×10^{-01}	1.10 x 10 ⁻⁰⁵	7.18 × 1
		Reference (Broach Lake)	1.34×10^{-04}	4.59 × 10 ⁻⁰⁴	4.91 × 10 ⁻⁰⁴	6.07 x 10 ⁻⁰²	2.22 × 10 ⁻⁰²	6.66 × 10 ⁻⁰⁶	1.82 × 10 ⁻⁰⁵	1.34 × 10 ⁻⁰⁴	4.59×10^{-04}	4.91 × 10 ⁻⁰⁴	6.07 x 10 ⁻⁰²	2.22 × 10 ⁻⁰²	6.66 × 10 ⁻⁰⁶	1.82 × 1
	Macrophytes	Patterson Lake North Arm — West Basin	1.38 × 10 ⁻⁰³	9.26 × 10 ⁻⁰²	1.73 × 10 ⁻⁰³	8.03 x 10 ⁻⁰²	2.62 × 10 ⁻⁰²	1.93 × 10 ⁻⁰⁵	6.97 × 10 ⁻⁰⁵	2.66 × 10 ⁻⁰⁴	9.32 × 10 ⁻⁰⁴	5.21 × 10 ⁻⁰⁴	1.49 x 10 ⁻⁰¹	5.74 × 10 ⁻⁰²	1.49 × 10 ⁻⁰⁴	1.83 × 1
	macrophytes	Patterson Lake South Arm	1.35 × 10 ⁻⁰³	9.26 × 10 ⁻⁰²	2.75×10^{-03}	8.14 x 10 ⁻⁰²	2.65 × 10 ⁻⁰²	1.84 × 10 ⁻⁰⁵	9.76 × 10 ⁻⁰⁵	2.31 × 10 ⁻⁰⁴	8.05 × 10 ⁻⁰⁴	4.93×10^{-04}	1.14 x 10 ⁻⁰¹	4.28 × 10 ⁻⁰²	8.90 × 10 ⁻⁰⁵	6.37 × 1
		Beet Lake	7.53×10^{-04}	4.71 × 10 ⁻⁰²	8.76 × 10 ⁻⁰⁴	6.99 x 10 ⁻⁰²	2.41 × 10 ⁻⁰²	1.19 × 10 ⁻⁰⁵	4.58 × 10 ⁻⁰⁵	1.89 × 10 ⁻⁰⁴	6.54 × 10 ⁻⁰⁴	4.91 × 10 ⁻⁰⁴	8.90 x 10 ⁻⁰²	3.30 × 10 ⁻⁰²	4.95 × 10 ⁻⁰⁵	3.51 × 1
Aquatic		Lloyd Lake	2.00×10^{-04}	5.40×10^{-03}	5.15 × 10 ⁻⁰⁴	6.17 x 10 ⁻⁰²	2.24 × 10 ⁻⁰²	7.18 × 10 ⁻⁰⁶	2.08 × 10 ⁻⁰⁵	1.40×10^{-04}	4.79×10^{-04}	4.91 × 10 ⁻⁰⁴	6.36 x 10 ⁻⁰²	2.33 × 10 ⁻⁰²	1.10 × 10 ⁻⁰⁵	1.98 × 1
Plants		Reference (Broach Lake)	6.96 × 10 ⁻⁰⁵	3.99×10^{-04}	6.44×10^{-03}	1.27 x 10 ⁻⁰²	9.16 × 10 ⁻⁰²	6.66 × 10 ⁻⁰⁶	2.28 × 10 ⁻⁰⁴	6.96 × 10 ⁻⁰⁵	3.99 × 10 ⁻⁰⁴	6.44 × 10 ⁻⁰³	1.27 x 10 ⁻⁰²	9.16 × 10 ⁻⁰²	6.66 × 10 ⁻⁰⁶	2.28 × 1
	Phytoplankton	Patterson Lake North Arm — West Basin	7.16 × 10 ⁻⁰⁴	8.05 × 10 ⁻⁰²	2.27 × 10 ⁻⁰²	1.68 x 10 ⁻⁰²	1.08 × 10 ⁻⁰¹	1.93 × 10 ⁻⁰⁵	8.72 × 10 ⁻⁰⁴	1.38 × 10 ⁻⁰⁴	8.11 × 10 ⁻⁰⁴	6.83 × 10 ⁻⁰³	3.10 x 10 ⁻⁰²	2.37 × 10 ⁻⁰¹	1.49 × 10 ⁻⁰⁴	2.28 × 1
	, .op	Patterson Lake South Arm	7.04×10^{-04}	8.06 × 10 ⁻⁰²		1.70 x 10 ⁻⁰²	1.10 × 10 ⁻⁰¹	1.84 × 10 ⁻⁰⁵	1.22 × 10 ⁻⁰³	1.20 × 10 ⁻⁰⁴	7.00×10^{-04}	6.48×10^{-03}	2.38 x 10 ⁻⁰²	1.77 × 10 ⁻⁰¹	8.90 × 10 ⁻⁰⁵	7.96 × 1
		Beet Lake				1.46 x 10 ⁻⁰²		1.19 × 10 ⁻⁰⁵			5.69 × 10 ⁻⁰⁴	6.45×10^{-03}	1.86 x 10 ⁻⁰²	1.36 × 10 ⁻⁰¹	4.95 × 10 ⁻⁰⁵	4.39 × 1
		Lloyd Lake	2.06 × 10 ⁻⁰⁴	1.76 × 10 ⁻⁰²		1.29 x 10 ⁻⁰²	9.24 × 10 ⁻⁰²	7.18 × 10 ⁻⁰⁶	2.60 × 10 ⁻⁰⁴	8.15 × 10 ⁻⁰⁵	4.70×10^{-04}	6.44×10^{-03}	1.33 x 10 ⁻⁰²	9.62 × 10 ⁻⁰²	1.10 × 10 ⁻⁰⁵	2.47 × 1
		Reference (Broach Lake)	n/a	n/a	2.45×10^{-04}	2.94 × 10 ⁻⁰⁴	2.65 × 10 ⁻⁰⁴	3.09 × 10 ⁻⁰⁴	5.71 × 10 ⁻⁰⁴	n/a	n/a	2.45×10^{-04}	2.94 × 10 ⁻⁰⁴	2.65 × 10 ⁻⁰⁴	3.09 × 10 ⁻⁰⁴	5.71 × 1
	Beaver	Patterson Lake North Arm — West Basin	n/a	n/a	4.58 × 10 ⁻⁰⁴	3.42 x 10 ⁻⁰⁴	2.79 x 10 ⁻⁰⁴	3.97 x 10 ⁻⁰⁴	4.49 x 10 ⁻⁰³	n/a	n/a	2.51 × 10 ⁻⁰⁴	5.04 x 10 ⁻⁰⁴	3.16 x 10 ⁻⁰⁴	8.92 × 10 ⁻⁰⁴	1.27 x 1
	Beaver	Patterson Lake South Arm	n/a	n/a	4.68×10^{-04}	3.43 x 10 ⁻⁰⁴	2.71 x 10 ⁻⁰⁴	3.50 x 10 ⁻⁰⁴	1.30 x 10 ⁻⁰³	n/a	n/a	2.46×10^{-04}	4.21 x 10 ⁻⁰⁴	2.94 x 10 ⁻⁰⁴	6.47 × 10 ⁻⁰⁴	7.37 x 10
		Beet Lake	n/a	n/a	2.84 × 10 ⁻⁰⁴	3.17 x 10 ⁻⁰⁴	2.68 x 10 ⁻⁰⁴	3.31 x 10 ⁻⁰⁴	9.53 x 10 ⁻⁰⁴	n/a	n/a	2.45×10^{-04}	3.62 x 10 ⁻⁰⁴	2.80 x 10 ⁻⁰⁴	4.85 × 10 ⁻⁰⁴	6.39 x 10
		Lloyd Lake	n/a	n/a	2.48 × 10 ⁻⁰⁴	2.97 x 10 ⁻⁰⁴	2.65 x 10 ⁻⁰⁴	3.14 x 10 ⁻⁰⁴	5.79 x 10 ⁻⁰⁴	n/a	n/a	2.45×10^{-04}	3.01 x 10 ⁻⁰⁴	2.66 x 10 ⁻⁰⁴	3.27 × 10 ⁻⁰⁴	5.76 x 1
		Reference (Broach Lake)	n/a	n/a	9.22 × 10 ⁻⁰⁴	8.71 × 10 ⁻⁰⁵	2.49×10^{-03}	1.44 × 10 ⁻⁰⁴	1.82 × 10 ⁻⁰⁴	n/a	n/a	9.22 × 10 ⁻⁰⁴	8.71 × 10 ⁻⁰⁵	2.49 × 10 ⁻⁰³	1.44 × 10 ⁻⁰⁴	1.82 × 1
errestrial Animals	Black Bear	Patterson Lake North Arm — West Basin	n/a	n/a	3.08 × 10 ⁻⁰³	9.77 x 10 ⁻⁰⁵	2.77 x 10 ⁻⁰³	1.63 x 10 ⁻⁰⁴	7.41 x 10 ⁻⁰⁴	n/a	n/a	9.42 × 10 ⁻⁰⁴	1.22 × 10 ⁻⁰⁴	4.69 × 10 ⁻⁰³	2.34 × 10 ⁻⁰⁴	3.21 x 10
		Beet Lake	n/a	n/a	1.24 × 10 ⁻⁰³	1.07 x 10 ⁻⁰⁴	2.59 x 10 ⁻⁰³	1.77 x 10 ⁻⁰⁴	2.65 x 10 ⁻⁰⁴	n/a	n/a	1.01 × 10 ⁻⁰³	1.11 × 10 ⁻⁰⁴	2.94 × 10 ⁻⁰³	1.91 × 10 ⁻⁰⁴	2.15 x 1
		Lloyd Lake	n/a	n/a	9.55 × 10 ⁻⁰⁴	9.06 x 10 ⁻⁰⁵	2.51 x 10 ⁻⁰³	1.50 x 10 ⁻⁰⁴	1.86 x 10 ⁻⁰⁴	n/a	n/a	9.25 × 10 ⁻⁰⁴	9.13 × 10 ⁻⁰⁵	2.58 × 10 ⁻⁰³	1.53 × 10 ⁻⁰⁴	1.84 x 10
		Reference (Broach Lake)	n/a	n/a	2.46 x 10 ⁻⁰³	4.49 x 10 ⁻⁰⁵	2.88 x 10 ⁻⁰³	1.11 x 10 ⁻⁰⁴	6.66 x 10 ⁻⁰⁵	n/a	n/a	2.46 x 10 ⁻⁰³	4.49 x 10 ⁻⁰⁵	2.88 x 10 ⁻⁰³	1.11 x 10 ⁻⁰⁴	6.66 x 1
	Grey Wolf	Patterson Lake North Arm — West Basin	n/a	n/a	2.96 x 10 ⁻⁰³	4.58 x 10 ⁻⁰⁵	2.93 x 10 ⁻⁰³	1.22 x 10 ⁻⁰⁴	2.18 x 10 ⁻⁰⁴	n/a	n/a	2.47 x 10 ⁻⁰³	4.81 x 10 ⁻⁰⁵	2.98 x 10 ⁻⁰³	1.72 x 10 ⁻⁰⁴	8.10 x 10
		Lloyd Lake	n/a	n/a	2.51 x 10 ⁻⁰³	4.53 x 10 ⁻⁰⁵	2.89 x 10 ⁻⁰³	1.14 x 10 ⁻⁰⁴	6.75 x 10 ⁻⁰⁵	n/a	n/a	2.46 x 10 ⁻⁰³	4.53 x 10 ⁻⁰⁵	2.89 x 10 ⁻⁰³	1.17 x 10 ⁻⁰⁴	6.69 x 1

								Maximun	n HQ RFD Case	e					
Biota	Location				Operations	S						Far-Futur	e		
		Chloride	Sulphate	Arsenic	Cobalt	Copper	Molybdenum	Uranium	Chloride	Sulphate	Arsenic	Cobalt	Copper	Molybdenum	Uranium
	Reference (Broach Lake)	n/a	n/a	3.77×10^{-03}	3.45×10^{-04}	1.99 × 10 ⁻⁰²	4.40 × 10 ⁻⁰⁴	1.93×10^{-04}	n/a	n/a	3.77 × 10 ⁻⁰³	3.45 × 10 ⁻⁰⁴	1.99 × 10 ⁻⁰²	4.40×10^{-04}	1.93 × 10 ⁻⁰⁴
	Patterson Lake North Arm — West Basin	n/a	n/a	1.13 x 10 ⁻⁰²	4.36 x 10 ⁻⁰⁴	2.31 x 10 ⁻⁰²	1.08 x 10 ⁻⁰³	7.45 x 10 ⁻⁰⁴	n/a	n/a	3.98 × 10 ⁻⁰³	7.89 x 10 ⁻⁰⁴	5.13 x 10 ⁻⁰²	8.48 × 10 ⁻⁰³	1.50 x 10 ⁻⁰³
American Mink	Patterson Lake South Arm	n/a	n/a	1.03 × 10 ⁻⁰²	4.12 x 10 ⁻⁰⁴	2.23 x 10 ⁻⁰²	8.72 x 10 ⁻⁰⁴	4.56 x 10 ⁻⁰⁴	n/a	n/a	3.79 × 10 ⁻⁰³	6.15 x 10 ⁻⁰⁴	3.83 x 10 ⁻⁰²	5.11 × 10 ⁻⁰³	5.52 x 10 ⁻⁰⁴
	Beet Lake	n/a	n/a	4.87×10^{-03}	3.80 x 10 ⁻⁰⁴	2.11 x 10 ⁻⁰²	6.64 x 10 ⁻⁰⁴	2.94 x 10 ⁻⁰⁴	n/a	n/a	3.77×10^{-03}	4.87 x 10 ⁻⁰⁴	2.95 x 10 ⁻⁰²	2.87 × 10 ⁻⁰³	3.27 x 10 ⁻⁰⁴
	Lloyd Lake	n/a	n/a	3.84 × 10 ⁻⁰³	3.48 x 10 ⁻⁰⁴	2.00 x 10 ⁻⁰²	4.64 x 10 ⁻⁰⁴	2.01 x 10 ⁻⁰⁴	n/a	n/a	3.77 × 10 ⁻⁰³	3.59 x 10 ⁻⁰⁴	2.08 x 10 ⁻⁰²	6.88 × 10 ⁻⁰⁴	2.05 x 10 ⁻⁰⁴
	Reference (Broach Lake)	n/a	n/a	1.63 × 10 ⁻⁰⁴	2.11 × 10 ⁻⁰⁴	1.07 × 10 ⁻⁰²	1.41 × 10 ⁻⁰⁴	2.84 × 10 ⁻⁰⁴	n/a	n/a	1.63 × 10 ⁻⁰⁴	2.11 × 10 ⁻⁰⁴	1.07 × 10 ⁻⁰²	1.41 × 10 ⁻⁰⁴	2.84 × 10 ⁻⁰⁴
Moose	Patterson Lake North Arm — West Basin	n/a	n/a	3.78 × 10 ⁻⁰⁴	2.58 x 10 ⁻⁰⁴	1.15 x 10 ⁻⁰²	2.02 x 10 ⁻⁰⁴	2.10 x 10 ⁻⁰³	n/a	n/a	1.69 × 10 ⁻⁰⁴	4.20 x 10 ⁻⁰⁴	1.48 x 10 ⁻⁰²	6.77 × 10 ⁻⁰⁴	8.62 x 10 ⁻⁰⁴
Wioose	Patterson Lake South Arm	n/a	n/a	3.77 × 10 ⁻⁰⁴	2.55 x 10 ⁻⁰⁴	1.15 x 10 ⁻⁰²	1.94 x 10 ⁻⁰⁴	2.05 x 10 ⁻⁰³	n/a	n/a	1.69 × 10 ⁻⁰⁴	4.14 x 10 ⁻⁰⁴	1.47 x 10 ⁻⁰²	6.16 × 10 ⁻⁰⁴	8.47 x 10 ⁻⁰⁴
	Beet Lake	n/a	n/a	2.02×10^{-04}	2.33 x 10 ⁻⁰⁴	1.10 x 10 ⁻⁰²	1.59 x 10 ⁻⁰⁴	4.79 x 10 ⁻⁰⁴	n/a	n/a	1.64 × 10 ⁻⁰⁴	2.78 x 10 ⁻⁰⁴	1.20 x 10 ⁻⁰²	3.03×10^{-04}	3.42 x 10 ⁻⁰⁴
	Lloyd Lake	n/a	n/a	1.66 × 10 ⁻⁰⁴	2.14 x 10 ⁻⁰⁴	1.08 x 10 ⁻⁰²	1.44 x 10 ⁻⁰⁴	2.90 x 10 ⁻⁰⁴	n/a	n/a	1.63 × 10 ⁻⁰⁴	2.18 x 10 ⁻⁰⁴	1.09 x 10 ⁻⁰²	1.58 × 10 ⁻⁰⁴	2.89 x 10 ⁻⁰⁴
	Reference (Broach Lake)	n/a	n/a	2.01 × 10 ⁻⁰³	1.19 × 10 ⁻⁰³	3.42 × 10 ⁻⁰³	9.60 × 10 ⁻⁰⁴	9.83×10^{-04}	n/a	n/a	2.01 × 10 ⁻⁰³	1.19 × 10 ⁻⁰³	3.42×10^{-03}	9.60 × 10 ⁻⁰⁴	9.83×10^{-04}
Muskrat	Patterson Lake North Arm — West Basin	n/a	n/a	6.18 × 10 ⁻⁰³	1.56 × 10 ⁻⁰³	3.98 × 10 ⁻⁰³	2.57 × 10 ⁻⁰³	3.25×10^{-03}	n/a	n/a	2.13 × 10 ⁻⁰³	2.92 × 10 ⁻⁰³	8.86 × 10 ⁻⁰³	2.14 × 10 ⁻⁰²	9.84 × 10 ⁻⁰³
Wuskiat	Patterson Lake South Arm	n/a	n/a	4.91 × 10 ⁻⁰³	1.50 × 10 ⁻⁰³	3.85 × 10 ⁻⁰³	2.06 × 10 ⁻⁰³	2.44 × 10 ⁻⁰³	n/a	n/a	2.02 × 10 ⁻⁰³	2.24 × 10 ⁻⁰³	6.61 × 10 ⁻⁰³	1.28 × 10 ⁻⁰²	3.43×10^{-03}
	Beet Lake	n/a	n/a	2.51 × 10 ⁻⁰³	1.35 × 10 ⁻⁰³	3.64 × 10 ⁻⁰³	1.53 × 10 ⁻⁰³	1.52 × 10 ⁻⁰³	n/a	n/a	2.01×10^{-03}	1.75×10^{-03}	5.09×10^{-03}	7.14×10^{-03}	1.89 × 10 ⁻⁰³
	Lloyd Lake	n/a	n/a	2.04×10^{-03}	1.21×10^{-03}	3.45×10^{-03}	1.02×10^{-03}	1.03×10^{-03}	n/a	n/a	2.01×10^{-03}	1.25×10^{-03}	3.59×10^{-03}	1.59 × 10 ⁻⁰³	1.07×10^{-03}
	Reference (Broach Lake)	n/a	n/a	4.18 × 10 ⁻⁰⁴	3.66 × 10 ⁻⁰⁵	6.08 × 10 ⁻⁰⁴	1.22 × 10 ⁻⁰⁴	6.64×10^{-05}	n/a	n/a	4.18 × 10 ⁻⁰⁴	3.66 × 10 ⁻⁰⁵	6.08 × 10 ⁻⁰⁴	1.22 × 10 ⁻⁰⁴	6.64×10^{-05}
Red Fox	Patterson Lake North Arm — West Basin	n/a	n/a	4.29 × 10 ⁻⁰⁴	3.80 x 10 ⁻⁰⁵	6.32 x 10 ⁻⁰⁴	1.46 x 10 ⁻⁰⁴	6.21 x 10 ⁻⁰⁴	n/a	n/a	4.18 × 10 ⁻⁰⁴	4.21 x 10 ⁻⁰⁵	6.19 x 10 ⁻⁰⁴	2.67 × 10 ⁻⁰⁴	1.10 x 10 ⁻⁰⁴
	Patterson Lake South Arm	n/a	n/a	4.36 × 10 ⁻⁰⁴	3.81 x 10 ⁻⁰⁵	6.10 x 10 ⁻⁰⁴	1.35 x 10 ⁻⁰⁴	1.61 x 10 ⁻⁰⁴	n/a	n/a	4.18 × 10 ⁻⁰⁴	3.99 x 10 ⁻⁰⁵	6.14 x 10 ⁻⁰⁴	2.06 × 10 ⁻⁰⁴	7.49 x 10 ⁻⁰⁵
	Beet Lake	n/a	n/a	4.22 × 10 ⁻⁰⁴	3.74 x 10 ⁻⁰⁵	6.09 x 10 ⁻⁰⁴	1.28 x 10 ⁻⁰⁴	1.18 x 10 ⁻⁰⁴	n/a	n/a	4.18 × 10 ⁻⁰⁴	3.84 x 10 ⁻⁰⁵	6.11 x 10 ⁻⁰⁴	1.66 × 10 ⁻⁰⁴	7.04 x 10 ⁻⁰⁵
	Lloyd Lake	n/a	n/a	4.19 × 10 ⁻⁰⁴	3.68 x 10 ⁻⁰⁵	6.09 x 10 ⁻⁰⁴	1.24 x 10 ⁻⁰⁴	6.72 x 10 ⁻⁰⁵	n/a	n/a	4.18×10^{-04}	3.68 x 10 ⁻⁰⁵	6.08 x 10 ⁻⁰⁴	1.27 × 10 ⁻⁰⁴	6.66 x 10 ⁻⁰⁵
	Reference (Broach Lake)	n/a	n/a	1.43 × 10 ⁻⁰³	2.97 × 10 ⁻⁰⁴	1.06 × 10 ⁻⁰³	6.71 × 10 ⁻⁰⁵	9.42 × 10 ⁻⁰⁴	n/a	n/a	1.43 × 10 ⁻⁰³	2.97×10^{-04}	1.06 × 10 ⁻⁰³	6.71 × 10 ⁻⁰⁵	9.42 × 10 ⁻⁰⁴
Snowshoe Hare	Patterson Lake North Arm — West Basin	n/a	n/a	1.45 × 10 ⁻⁰³	3.00 x 10 ⁻⁰⁴	1.09 x 10 ⁻⁰³	7.71 x 10 ⁻⁰⁵	7.61 x 10 ⁻⁰³	n/a	n/a	1.43 × 10 ⁻⁰³	3.04 x 10 ⁻⁰⁴	1.06 x 10 ⁻⁰³	8.10 × 10 ⁻⁰⁵	1.38 x 10 ⁻⁰³
Showshoe Hare	Patterson Lake South Arm	n/a	n/a	1.46×10^{-03}	3.01 x 10 ⁻⁰⁴	1.06 x 10 ⁻⁰³	6.90 x 10 ⁻⁰⁵	2.01 x 10 ⁻⁰³	n/a	n/a	1.43×10^{-03}	3.01 x 10 ⁻⁰⁴	1.06 x 10 ⁻⁰³	7.52 × 10 ⁻⁰⁵	1.01 x 10 ⁻⁰³
	Beet Lake	n/a	n/a	1.44×10^{-03}	3.00 x 10 ⁻⁰⁴	1.06 x 10 ⁻⁰³	6.84 x 10 ⁻⁰⁵	1.54 x 10 ⁻⁰³	n/a	n/a	1.43 × 10 ⁻⁰³	2.99 x 10 ⁻⁰⁴	1.06 x 10 ⁻⁰³	7.13 × 10 ⁻⁰⁵	9.81 x 10 ⁻⁰⁴
	Lloyd Lake	n/a	n/a	1.44 × 10 ⁻⁰³	2.99 x 10 ⁻⁰⁴	1.06 x 10 ⁻⁰³	6.79 x 10 ⁻⁰⁵	9.49 x 10 ⁻⁰⁴	n/a	n/a	1.43 × 10 ⁻⁰³	2.98 x 10 ⁻⁰⁴	1.06 x 10 ⁻⁰³	6.76 × 10 ⁻⁰⁵	9.42 x 10 ⁻⁰⁴
	Reference (Broach Lake)	n/a	n/a	1.50 × 10 ⁻⁰⁴	1.05 × 10 ⁻⁰⁴	1.24 × 10 ⁻⁰⁴	5.79 × 10 ⁻⁰⁴	7.05×10^{-04}	n/a	n/a	1.50 × 10 ⁻⁰⁴	1.05 × 10 ⁻⁰⁴	1.24 × 10 ⁻⁰⁴	5.79 × 10 ⁻⁰⁴	7.05×10^{-04}
Southern Red-Backed	Patterson Lake North Arm — West Basin	n/a	n/a	1.52 × 10 ⁻⁰⁴	1.07 x 10 ⁻⁰⁴	1.28 x 10 ⁻⁰⁴	6.69 x 10 ⁻⁰⁴	5.49 x 10 ⁻⁰³	n/a	n/a	1.50 × 10 ⁻⁰⁴	1.09 x 10 ⁻⁰⁴	1.25 x 10 ⁻⁰⁴	8.14 × 10 ⁻⁰⁴	1.04 x 10 ⁻⁰³
Vole	Patterson Lake South Arm	n/a	n/a	1.54 × 10 ⁻⁰⁴	1.07 x 10 ⁻⁰⁴	1.24 x 10 ⁻⁰⁴	6.04 x 10 ⁻⁰⁴	1.48 x 10 ⁻⁰³	n/a	n/a	1.50 × 10 ⁻⁰⁴	1.08 x 10 ⁻⁰⁴	1.24 x 10 ⁻⁰⁴	7.15 × 10 ⁻⁰⁴	7.62 x 10 ⁻⁰⁴
	Beet Lake	n/a	n/a	1.51 × 10 ⁻⁰⁴		1.24 x 10 ⁻⁰⁴	5.93 x 10 ⁻⁰⁴	1.14 x 10 ⁻⁰³	n/a	n/a	1.50 × 10 ⁻⁰⁴	1.07 x 10 ⁻⁰⁴	1.24 x 10 ⁻⁰⁴	6.50 × 10 ⁻⁰⁴	7.35 x 10 ⁻⁰⁴
	Lloyd Lake	n/a	n/a	1.50 × 10 ⁻⁰⁴	1.06 x 10 ⁻⁰⁴	1.24 x 10 ⁻⁰⁴	5.85 x 10 ⁻⁰⁴	7.10 x 10 ⁻⁰⁴	n/a	n/a	1.50 × 10 ⁻⁰⁴	1.05 x 10 ⁻⁰⁴	1.24 x 10 ⁻⁰⁴	5.86 × 10 ⁻⁰⁴	7.05 x 10 ⁻⁰⁴
	Reference (Broach Lake)	n/a	n/a	3.79 x 10 ⁻⁰⁴	2.94 x 10 ⁻⁰⁴	1.19 x 10 ⁻⁰²	3.02 x 10 ⁻⁰⁴	3.16 x 10 ⁻⁰⁴	n/a	n/a	3.79 x 10 ⁻⁰⁴	2.94 x 10 ⁻⁰⁴	1.19 x 10 ⁻⁰²	3.02 × 10 ⁻⁰⁴	3.16 x 10 ⁻⁰⁴
Woodland Caribou	Patterson Lake North Arm — West Basin	n/a	n/a	8.03 x 10 ⁻⁰⁴	3.52 x 10 ⁻⁰⁴	1.31 x 10 ⁻⁰²	5.90 x 10 ⁻⁰⁴	1.31 x 10 ⁻⁰²	n/a	n/a	3.87 x 10 ⁻⁰⁴	4.96 x 10 ⁻⁰⁴	1.56 x 10 ⁻⁰²	1.35 × 10 ⁻⁰³	1.09 x 10 ⁻⁰³
	Beet Lake	n/a	n/a	4.21 x 10 ⁻⁰⁴	3.12 x 10 ⁻⁰⁴	1.20 x 10 ⁻⁰²	3.38 x 10 ⁻⁰⁴	1.39 x 10 ⁻⁰³	n/a	n/a	3.79 x 10 ⁻⁰⁴	3.35 x 10 ⁻⁰⁴	1.26 x 10 ⁻⁰²	5.05 × 10 ⁻⁰⁴	3.80 x 10 ⁻⁰⁴
	Lloyd Lake	n/a	n/a	3.93 x 10 ⁻⁰⁴	3.07 x 10 ⁻⁰⁴	1.20 x 10 ⁻⁰²	3.40 x 10 ⁻⁰⁴	3.49 x 10 ⁻⁰⁴	n/a	n/a	3.79 x 10 ⁻⁰⁴	3.03 x 10 ⁻⁰⁴	1.20 x 10 ⁻⁰²	3.45 × 10 ⁻⁰⁴	3.27 x 10 ⁻⁰⁴
	Reference (Broach Lake)	n/a	n/a	1.79 × 10 ⁻⁰³	3.95 × 10 ⁻⁰⁴	1.51 × 10 ⁻⁰³	6.67 × 10 ⁻⁰⁵	3.96×10^{-04}	n/a	n/a	1.79 × 10 ⁻⁰³	3.95 × 10 ⁻⁰⁴	1.51 × 10 ⁻⁰³	6.67 × 10 ⁻⁰⁵	3.96×10^{-04}
Grouse	Patterson Lake North Arm — West Basin	n/a	n/a	1.80 × 10 ⁻⁰³	3.98 x 10 ⁻⁰⁴	1.56 x 10 ⁻⁰³	7.48 x 10 ⁻⁰⁵	3.03 x 10 ⁻⁰³	n/a	n/a	1.79 × 10 ⁻⁰³	3.99 x 10 ⁻⁰⁴	1.51 x 10 ⁻⁰³	7.47 × 10 ⁻⁰⁵	5.79 x 10 ⁻⁰⁴
-0.0030	Patterson Lake South Arm	n/a	n/a	1.80 × 10 ⁻⁰³	3.98 x 10 ⁻⁰⁴	1.51 x 10 ⁻⁰³	6.80 x 10 ⁻⁰⁵	8.19 x 10 ⁻⁰⁴	n/a	n/a	1.79 × 10 ⁻⁰³	3.98 x 10 ⁻⁰⁴	1.51 x 10 ⁻⁰³	7.13 × 10 ⁻⁰⁵	4.26 x 10 ⁻⁰⁴
	Beet Lake	n/a	n/a	1.79 × 10 ⁻⁰³	3.97 x 10 ⁻⁰⁴	1.51 x 10 ⁻⁰³	6.76 x 10 ⁻⁰⁵	6.32 x 10 ⁻⁰⁴	n/a	n/a	1.79 × 10 ⁻⁰³	3.96 x 10 ⁻⁰⁴	1.51 x 10 ⁻⁰³	6.91 × 10 ⁻⁰⁵	4.13 x 10 ⁻⁰⁴
	Lloyd Lake	n/a	n/a	1.79 × 10 ⁻⁰³	3.97 x 10 ⁻⁰⁴	1.51 x 10 ⁻⁰³	6.73 x 10 ⁻⁰⁵	3.99 x 10 ⁻⁰⁴	n/a	n/a	1.79 × 10 ⁻⁰³	3.95 x 10 ⁻⁰⁴	1.51 x 10 ⁻⁰³	6.69 × 10 ⁻⁰⁵	3.97 x 10 ⁻⁰⁴
	Reference (Broach Lake)	n/a	n/a	1.72 × 10 ⁻⁰⁴	3.93×10^{-05}	1.96 × 10 ⁻⁰⁴	1.24 × 10 ⁻⁰⁵	3.93×10^{-05}	n/a	n/a	1.72 × 10 ⁻⁰⁴	3.93×10^{-05}	1.96 × 10 ⁻⁰⁴	1.24 × 10 ⁻⁰⁵	3.93×10^{-05}
Canada Goose	Patterson Lake North Arm — West Basin	n/a	n/a	1.76 × 10 ⁻⁰⁴	4.01 x 10 ⁻⁰⁵	2.04 x 10 ⁻⁰⁴	1.45 x 10 ⁻⁰⁵	3.07 x 10 ⁻⁰⁴	n/a	n/a	1.72 × 10 ⁻⁰⁴	4.17 x 10 ⁻⁰⁵	1.99 x 10 ⁻⁰⁴	2.05 × 10 ⁻⁰⁵	5.90 x 10 ⁻⁰⁵
	Patterson Lake South Arm	n/a	n/a	1.79 × 10 ⁻⁰⁴	4.01 x 10 ⁻⁰⁵	1.97 x 10 ⁻⁰⁴	1.32 x 10 ⁻⁰⁵	8.30 x 10 ⁻⁰⁵	n/a	n/a	1.72 × 10 ⁻⁰⁴	4.08 x 10 ⁻⁰⁵	1.97 x 10 ⁻⁰⁴	1.71 × 10 ⁻⁰⁵	4.27 x 10 ⁻⁰⁵
	Beet Lake	n/a	n/a	1.74×10^{-04}	3.98 x 10 ⁻⁰⁵	1.97 x 10 ⁻⁰⁴	1.28 x 10 ⁻⁰⁵	6.35 x 10 ⁻⁰⁵	n/a	n/a	1.72×10^{-04}	4.01 x 10 ⁻⁰⁵	1.97 x 10 ⁻⁰⁴	1.48×10^{-05}	4.11 x 10 ⁻⁰⁵

MAINOMMENTAL VISIV ASSESS	IVIEINT FOR THE ROOK I PROJECT
	ECOLOGICAL RISK ASSESSMENT

								Maximum	HQ RFD Case	е					
Biota	Location		Operations					Far-Future							
		Chloride	Sulphate	Arsenic	Cobalt	Copper	Molybdenum	Uranium	Chloride	Sulphate	Arsenic	Cobalt	Copper	Molybdenum	Uranium
	Lloyd Lake	n/a	n/a	1.73×10^{-04}	3.96 x 10 ⁻⁰⁵	1.97 x 10 ⁻⁰⁴	1.25 x 10 ⁻⁰⁵	3.96 x 10 ⁻⁰⁵	n/a	n/a	1.72×10^{-04}	3.95 x 10 ⁻⁰⁵	1.96 x 10 ⁻⁰⁴	1.26 × 10 ⁻⁰⁵	3.93 x 10 ⁻⁰⁵
	Reference (Broach Lake)	n/a	n/a	9.07×10^{-03}	4.28×10^{-03}	2.80×10^{-01}	6.31 × 10 ⁻⁰³	8.52 × 10 ⁻⁰⁴	n/a	n/a	9.07×10^{-03}	4.28×10^{-03}	2.80×10^{-01}	6.31 × 10 ⁻⁰³	8.52×10^{-0}
Little Brown Myotis	Patterson Lake North Arm — West Basin	n/a	n/a	2.77 × 10 ⁻⁰²	5.53 × 10 ⁻⁰³	3.26 × 10 ⁻⁰¹	1.69 × 10 ⁻⁰²	2.80 × 10 ⁻⁰³	n/a	n/a	9.61 × 10 ⁻⁰³	1.05 × 10 ⁻⁰²	7.24 × 10 ⁻⁰¹	1.41 × 10 ⁻⁰¹	8.53 × 10 ⁻⁰
Little Brown Myotis	Patterson Lake South Arm	n/a	n/a	2.08 × 10 ⁻⁰²	5.21 × 10 ⁻⁰³	3.14×10^{-01}	1.35 × 10 ⁻⁰²	1.98 × 10 ⁻⁰³	n/a	n/a	9.11 × 10 ⁻⁰³	8.03 × 10 ⁻⁰³	5.41 × 10 ⁻⁰¹	8.43 × 10 ⁻⁰²	2.98 × 10 ⁻⁰
	Beet Lake	n/a	n/a	1.11 × 10 ⁻⁰²	4.77×10^{-03}	2.98 × 10 ⁻⁰¹	1.00 × 10 ⁻⁰²	1.27 × 10 ⁻⁰³	n/a	n/a	9.07 × 10 ⁻⁰³	6.26 × 10 ⁻⁰³	4.16 × 10 ⁻⁰¹	4.69 × 10 ⁻⁰²	1.64 × 10 ⁻⁰
	Lloyd Lake	n/a	n/a	9.19 × 10 ⁻⁰³	4.33 × 10 ⁻⁰³	2.82 × 10 ⁻⁰¹	6.69 × 10 ⁻⁰³	8.90 × 10 ⁻⁰⁴	n/a	n/a	9.07 × 10 ⁻⁰³	4.48 × 10 ⁻⁰³	2.94 × 10 ⁻⁰¹	1.04 × 10 ⁻⁰²	9.24 × 10 ⁻⁰
	Reference (Broach Lake)	n/a	n/a	1.31 × 10 ⁻⁰³	1.54 × 10 ⁻⁰⁴	4.62 × 10 ⁻⁰³	1.97 × 10 ⁻⁰⁵	2.11 × 10 ⁻⁰⁵	n/a	n/a	1.31 × 10 ⁻⁰³	1.54 × 10 ⁻⁰⁴	4.62 × 10 ⁻⁰³	1.97 × 10 ⁻⁰⁵	2.11 × 10 ⁻⁰
Common Loon	Patterson Lake North Arm — West Basin	n/a	n/a	4.52 × 10 ⁻⁰³	2.01 × 10 ⁻⁰⁴	5.39 × 10 ⁻⁰³	5.27 × 10 ⁻⁰⁵	7.64 × 10 ⁻⁰⁵	n/a	n/a	1.39 × 10 ⁻⁰³	3.78 × 10 ⁻⁰⁴	1.19 × 10 ⁻⁰²	4.39 × 10 ⁻⁰⁴	2.11 × 10
Common Loon	Patterson Lake South Arm	n/a	n/a	6.69 × 10 ⁻⁰³	1.88 × 10 ⁻⁰⁴	5.18 × 10 ⁻⁰³	4.22 × 10 ⁻⁰⁵	8.81 × 10 ⁻⁰⁵	n/a	n/a	1.32 × 10 ⁻⁰³	2.90 × 10 ⁻⁰⁴	8.92 × 10 ⁻⁰³	2.63 × 10 ⁻⁰⁴	7.37 × 10 ⁻¹
	Beet Lake	n/a	n/a	2.23 × 10 ⁻⁰³	1.72 × 10 ⁻⁰⁴	4.91 × 10 ⁻⁰³	3.13 × 10 ⁻⁰⁵	4.41 × 10 ⁻⁰⁵	n/a	n/a	1.31 × 10 ⁻⁰³	2.26 × 10 ⁻⁰⁴	6.86 × 10 ⁻⁰³	1.46 × 10 ⁻⁰⁴	4.07 × 10
	Lloyd Lake	n/a	n/a	1.37 × 10 ⁻⁰³	1.56 × 10 ⁻⁰⁴	4.65 × 10 ⁻⁰³	2.09 × 10 ⁻⁰⁵	2.32 × 10 ⁻⁰⁵	n/a	n/a	1.31 × 10 ⁻⁰³	1.62 × 10 ⁻⁰⁴	4.85 × 10 ⁻⁰³	3.26 × 10 ⁻⁰⁵	2.29 × 10
	Reference (Broach Lake)	n/a	n/a	5.44 × 10 ⁻⁰³	1.85 × 10 ⁻⁰³	1.40 × 10 ⁻⁰²	3.13 × 10 ⁻⁰⁴	3.27 × 10 ⁻⁰⁴	n/a	n/a	5.44 × 10 ⁻⁰³	1.85 × 10 ⁻⁰³	1.40 × 10 ⁻⁰²	3.13 × 10 ⁻⁰⁴	3.27 × 10
Mallard	Patterson Lake North Arm — West Basin	n/a	n/a	1.67 × 10 ⁻⁰²	2.41 × 10 ⁻⁰³	1.63 × 10 ⁻⁰²	8.38 × 10 ⁻⁰⁴	1.08 × 10 ⁻⁰³	n/a	n/a	5.77 × 10 ⁻⁰³	4.54 × 10 ⁻⁰³	3.63 × 10 ⁻⁰²	6.99 × 10 ⁻⁰³	3.27 × 10
Ivialiaiu	Patterson Lake South Arm	n/a	n/a	1.32 × 10 ⁻⁰²	2.30×10^{-03}	1.57 × 10 ⁻⁰²	6.70 × 10 ⁻⁰⁴	8.07 × 10 ⁻⁰⁴	n/a	n/a	5.47 × 10 ⁻⁰³	3.48×10^{-03}	2.71 × 10 ⁻⁰²	4.19 × 10 ⁻⁰³	1.14 × 10
	Beet Lake	n/a	n/a	6.77 × 10 ⁻⁰³	2.08 × 10 ⁻⁰³	1.49 × 10 ⁻⁰²	4.99 × 10 ⁻⁰⁴	5.02 × 10 ⁻⁰⁴	n/a	n/a	5.45 × 10 ⁻⁰³	2.72 × 10 ⁻⁰³	2.08 × 10 ⁻⁰²	2.33 × 10 ⁻⁰³	6.29 × 10
	Lloyd Lake	n/a	n/a	5.52 × 10 ⁻⁰³	1.88 × 10 ⁻⁰³	1.41 × 10 ⁻⁰²	3.32 × 10 ⁻⁰⁴	3.43×10^{-04}	n/a	n/a	5.44 × 10 ⁻⁰³	1.94 × 10 ⁻⁰³	1.47 × 10 ⁻⁰²	5.19 × 10 ⁻⁰⁴	3.54 × 10
	Reference (Broach Lake)	n/a	n/a	1.99 × 10 ⁻⁰²	8.87×10^{-03}	2.35 × 10 ⁻⁰¹	1.76 × 10 ⁻⁰³	1.10 × 10 ⁻⁰³	n/a	n/a	1.99 × 10 ⁻⁰²	8.87 × 10 ⁻⁰³	2.35 × 10 ⁻⁰¹	1.76 × 10 ⁻⁰³	1.10 × 10
Rusty Blackbird	Patterson Lake North Arm — West Basin	n/a	n/a	5.08 × 10 ⁻⁰²	1.12 x 10 ⁻⁰²	2.73 x 10 ⁻⁰¹	4.35 x 10 ⁻⁰³	4.79 x 10 ⁻⁰³	n/a	n/a	2.08 × 10 ⁻⁰²	2.04 x 10 ⁻⁰²	6.07 x 10 ⁻⁰¹	3.47 × 10 ⁻⁰²	6.59 x 10 ⁻
Rasty Diackbira	Patterson Lake South Arm	n/a	n/a	3.94 × 10 ⁻⁰²	1.06 x 10 ⁻⁰²	2.64 x 10 ⁻⁰¹	3.51 x 10 ⁻⁰³	2.24 x 10 ⁻⁰³	n/a	n/a	2.00 × 10 ⁻⁰²	1.59 x 10 ⁻⁰²	4.53 x 10 ⁻⁰¹	2.09 × 10 ⁻⁰²	2.59 x 10 ⁻¹
	Beet Lake	n/a	n/a	2.32 × 10 ⁻⁰²	9.78 x 10 ⁻⁰³	2.50 x 10 ⁻⁰¹	2.67 x 10 ⁻⁰³	1.59 x 10 ⁻⁰³	n/a	n/a	1.99 × 10 ⁻⁰²	1.26 x 10 ⁻⁰²	3.49 x 10 ⁻⁰¹	1.17 × 10 ⁻⁰²	1.66 x 10 ⁻
	Lloyd Lake	n/a	n/a	2.01 × 10 ⁻⁰²	8.96 x 10 ⁻⁰³	2.37 x 10 ⁻⁰¹	1.85 x 10 ⁻⁰³	1.13 x 10 ⁻⁰³	n/a	n/a	1.99 × 10 ⁻⁰²	9.25 x 10 ⁻⁰³	2.47 x 10 ⁻⁰¹	2.77 × 10 ⁻⁰³	1.15 x 10 ⁻⁰

Note: **Bold** indicates the HQ is greater than 1.

n/a = not applicable; HQ = hazard quotient; RFD = reasonably foreseeable development.

6.4.2 Uncertainty in the Risk Characterization

Since the risk characterization is dependent on the problem formulation and the exposure and effects assessments, any uncertainty identified in these assessments propagates uncertainty into the risk estimates. In general, the uncertainties are expected to cause an overestimation, not an underestimation of risk due to the conservative approaches employed in the ERA, including the use of:

- maximum predicted concentrations for COPCs in media for each exposure scenario;
- exposure of mobile ecological receptors to COPCs in the environment for chronic periods of time and during sensitive life stages; and
- effect levels based on low-effect threshold concentrations and doses.

The assumptions to address uncertainties in the ERA are anticipated to produce overly conservative estimates of risk, as discussed below.

For the calculation of risk to environmental receptors, there are uncertainties associated with the use of literature-based TRVs. These uncertainties may include: extrapolation of results from laboratory tests to the field, differences in sensitivity between the test organism and resident organisms, laboratory conditions that are not representative of field conditions, and the form of the COPC used in toxicity testing which may not be representative of the form found at the site.

The use of TRVs from laboratory studies tends to be conservative because these studies are typically chemical-specific and use highly bioavailable forms of the COPC. In field situations, the chemical form of the COPC may be less bioavailable, and toxicity-modifying factors may be present that were not acting in laboratory tests.

There is inherent uncertainty associated with the use of LELs such as LOAEL values as TRVs as these values are not precisely related to biologically relevant thresholds and do not provide information about the actual magnitude of effects in the reported studies. However, LOAEL values have widespread use in the risk assessment community and the science is not currently available to change this approach to TRVs.

Taken together, these approaches are anticipated to produce a risk characterization has not underestimated risk; the resulting HQs are either overestimates or realistic estimates of risk, both of which are considered acceptable.



7.0 QUALITY ASSURANCE AND SENSITIVITY ANALYSIS

7.1 Quality Assurance and Quality Control

Throughout the planning and preparation of the ERA, all staff worked under Ecometrix ISO 9001:2015 certified Quality Management System. All work was internally reviewed and verified. Reviews included verification of input data in the IMPACT files against the source documents and verification of selected results with independent calculation spreadsheets, as well as review of report content. Comments have been addressed as appropriate by report revisions. The review process has been documented through a paper trail of review comments and responses. Examples of the independent calculation spreadsheets are provided in Appendix C.

The software used for the ERA was IMPACT, the non-derived release limits Version 5.6.0, which aligns with the guidance for modelling radionuclides in the environment that is referred to in the Canadian Standards Association Group (CSA) standard N288.1-20 (CSA 2020), CSA N288.6-22 (CSA 2022), and supporting documentation. When utilizing IMPACT for this Project, all inputs to IMPACT were checked, along with an overall verification of IMPACT scenario files. Checks were performed on data and calculations to verify that transcription errors and formula errors, if any, were caught and addressed. Checks of the model structure, algorithms and functions have been made repeatedly throughout the model development history as it has been used in several related applications that underwent multiple layers of review.

The ERICA Tool, version 1.3.1, was used as a source of biota dose coefficients. Its parameters, including dose coefficients, have been subject to validation through numerous intercomparison exercises, as described by Brown et al., (2003, 2008, 2016) and have generally compared well with other sources. The intercomparisons of dose coefficients are described by Vives I Batlle et al. (2007, 2011). The external dose predictions for small mammals have been validated against dosimetric measurements (Beresford et al. 2008b). The code and database are updated from time to time, as described in its documented version history.

Quality assurance activities for other models used as inputs into the ERA are discussed in EIS Appendix 7A (Air Dispersion Modelling Report), TSD XIV (Groundwater Flow and Solute Transport Modelling Reporting) EIS Appendix 9A (Hydrological Modelling Summary Report) and EIS Appendix 10A (Surface Water Quality Modelling Report). As a quality control step, results from the other models were compared to IMPACT results wherever there were similar predictions in terms of nodes and parameters to confirm they were similar.

7.2 Sensitivity Analysis

A sensitivity analysis of key model parameters was undertaken to understand the degree to which the results or conclusions of the risk assessment would vary if these parameters differed from what was assumed.



7.2.1 Effects of Annual Weather Patterns

Dose and risk presented in Section 5.0 and Section 6.0 were calculated using average annual flows in all waterbodies. The effects of flow variation over each year were investigated separately in this sensitivity analysis. Average monthly flows at the model nodes are shown in Appendix A, Table 3-2). The change in flow is discussed using copper and polonium-210 as examples for a non-radiological and radiological COPC that either have the highest concentrations in waterbodies or have the highest contribution to dose of the most exposed human and ecological receptors. Average monthly and average annual outflows of all modelled waterbodies are provided in Table 7-1 below.

Figure 7-1 shows concentrations of copper and polonium-210 in water in Patterson Lake North Arm – West Basin during Operations. Concentrations were compared between a simulation using monthly flows and a simulation using annual average flows. The variation of water concentrations during Operations is mainly attributed to the changing patterns of liquid emissions as opposed to monthly flows. As treated effluent flow rates are reduced towards the end of Operations, the difference between the average and monthly flows becomes more noticeable (after year 30 of the total simulation time). Using monthly flows, the maximum deviation from the annual average for copper is -0.2% and +0.3%. For polonium-210 the maximum deviation from the concentrations calculated using annual average flows is -5.8% and +5.1%. The maximum deviation between annual average flows and monthly flows occurs while treated effluent is not discharged.

Similar deviations can be found when calculating dose and risk for humans. When considering monthly average flows, the total radiation dose to the subsistence harvester (one-year-old; i.e., the receptor with the highest exposure) increases by a maximum of 1% (Figure 7-2). This occurs after 42 years, when the total dose using constant flows is calculated to be 0.831 mSv/yr and the total dose using monthly flows is calculated to be 0.838 mSv/yr. Similar trends can be extrapolated for other human receptors and COPCs.

For ecological receptors, the increase in dose, when considering monthly flow patterns compared to annual averages, is a maximum of 0.5% for the total dose (i.e., sum of all pathways and radionuclides) during Operations. Risk (i.e., HQ) from copper from all pathways was estimated to be at a maximum of 0.2% greater when considering monthly flow patterns.

In summary, the doses and risks to human and ecological receptors have the potential to fluctuate around the average value presented in the report when monthly flow patterns are considered. In general, the maximum deviation of dose or risk between the monthly flows and annual average flows is under 1%. As the incremental dose to humans over the duration of a year is well below the public dose limit, this simplification of the calculation in the model does not affect conclusions for human receptors.



Table 7-1: Average Monthly Outflows and Average Annual Outflows of the Modelled Waterbodies

	Average Monthly Flows (m³/s)							Annual Average					
Lake Name	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Flows (m ³ /s)
Broach Lake	0.1	0.1	0.1	0.3	0.5	0.4	0.3	0.2	0.2	0.2	0.2	0.2	0.3
Clearwater River above Patterson Lake	0.2	0.2	0.3	0.8	1.2	0.9	0.7	0.6	0.6	0.6	0.5	0.3	0.6
Patterson Lake North Arm – East Basin	0.4	0.4	0.5	0.9	1.3	1.1	0.9	0.8	0.8	0.8	0.7	0.5	0.8
Patterson Lake North Arm – West Basin	0.7	0.7	0.7	1.0	1.4	1.3	1.2	1.1	1.0	1.0	0.9	0.7	1.0
Patterson Lake South Arm	1.1	1.0	1.1	1.2	1.7	1.7	1.6	1.5	1.5	1.4	1.2	1.1	1.4
Forrest Lake – North Basin	1.9	1.7	1.7	1.8	2.7	2.8	2.6	2.4	2.3	2.3	2.3	2.1	2.2
Beet Lake	2.0	1.8	1.8	2.1	3.3	3.0	2.7	2.6	2.5	2.5	2.4	2.2	2.4
Naomi Lake	2.4	2.2	2.1	2.6	4.6	4.0	3.5	3.3	3.2	3.2	3.0	2.7	3.1
Clearwater River above Mirror River confluence	3.5	3.5	3.5	6.0	10.0	8.0	6.6	6.0	6.0	6.0	5.4	4.0	5.7
Clearwater River above Lloyd Lake	11.7	11.6	11.7	20.0	33.1	26.5	22.1	20.0	19.9	20.0	17.8	13.2	19.0
Lloyd Lake	17.6	16.1	15.5	19.4	34.8	33.2	28.4	24.2	22.6	23.5	21.3	19.5	23.0

 m^3/s = cubic metre per second.






Figure 7-1: Concentrations of Copper and Polonium-210 in Patterson Lake North Arm – West Basin during Operations
Cu = copper; Po-210 = polonium-210; Bq/L = becquerels per litre.




Figure 7-2: Total Dose to the Subsistence Harvester (One-Year-Old) at Patterson Lake South Arm during Operations mSv/yr = millisieverts per year.

7.2.2 Deposition of Constituents of Potential Concern on Soil and Plants

IMPACT follows the CSA N288.1-20 model, which uses a number of input parameters that are specified in N288.1. The CSA N288.1-20 air dispersion model can be used in IMPACT to estimate average concentrations in air. In this assessment, modelled air concentrations were used from another model (AERMOD) and only the wet and dry deposition on plants and soil was calculated in IMPACT. This deposition was added to a soil baseline estimated from sampling results of soil and plants.

A number of assumptions (Table 7-2) based on default parameters from CSA N288.1-20 were made in modelling deposition of airborne contaminants in this assessment, which can be a source of uncertainty. Dry deposition velocity, washout ratio to soil and plants, and the fraction of time precipitation falls when the wind is from a certain sector are not site-specific, but the default values recommended by CSA N288.1-20 were used. For example, in the air model (EIS Appendix 7A), the dry deposition rates are location-specific. The dry deposition rates were calculated from the air concentrations for a given constituent and the deposition velocity, at a reference height. The wet deposition rates were calculated from the air concentrations for a given constituent and the washout ratio. The default dry deposition velocity of 0.0014 m/s was used for all metals (CSA 2020). Since concentrations decrease with distance from the emission sources, deposition rates also decrease with increasing distance from the emission sources, even with the assumption of a constant deposition velocity.

Table 7-2: Parameters Used in Modelling Deposition of Airborne Constituents of Potential Concern on Soil and Plants

Model Parameter	Unit	Value used in IMPACT (a)	Value used in AERMOD
Dry deposition velocity	m/s	0.0014	Greater than 0.002
Wet deposition washout ratio to soil	unitless	5.5E+6	Varied hourly with meteorological conditions
Wet deposition washout ratio to plants	unitless	6.3E+5	Same as washout ratio to soil
Precipitation rate	mm/yr	800	Varied hourly; annual precipitation was 573.6 mm/yr averaged over the five modelling years (2012 – 2016)
Fraction of time precipitation falls when the wind is from a certain sector	unitless	0.36	Not applicable; model uses hourly wind data and precipitation rate

(a) Values used in IMPACT model are the default parameters specified in CSA N288.1-20, assuming sandy soil. Values presented are for all elements other than noble gases, iodine, and N-13.

m/s = metres per second; mm/yr = millimetres per year.



7.2.3 Traditional Food Assumptions for Subsistence Harvesters

In the HHRA, the subsistence harvester was assumed to collect 50% of the Traditional Foods in the diet from locations potentially impacted by the Project, and the other 50% of the Traditional Foods in the diet from a reference location unaffected by the Project. This assumption was considered representative of how a subsistence harvester would collect Traditional Foods, by moving around and collecting from more than one location.

To evaluate the potential changes to the predicted cancer risk to the subsistence harvester, a sensitivity analysis was conducted assuming the subsistence harvester obtained 100% of the Traditional Foods in the overall diet from the area of the Project.

Additionally, as indicated in Section 5.2.3.2, Bioavailability, bioaccessibility factors (Table 7-3) for moose meat and organs were applied to the exposure doses, and exposure doses from fish were also adjusted to account for only the inorganic form of arsenic. In this sensitivity analysis these adjustments were removed as well.

Based on the assumption that the subsistence harvester would obtain Traditional Foods entirely from the area of the Project, the arsenic ILCR would exceed the negligible cancer risk level of 1 in 100,000 for the subsistence harvester harvesting Traditional Foods from Patterson Lake South Arm and would be at the negligible cancer risk level for the subsistence harvester at Beet Lake during Operations for all cases (Table 7-4). The arsenic ILCR would be below the negligible cancer risk level of 1 in 100,000 for the subsistence harvester at Lloyd Lake. During the far-future projection, the arsenic ILCR would generally be at or below the negligible cancer risk level at all locations.

This sensitivity scenario has been presented as a worst-case scenario, that would be considered unlikely, since a subsistence harvester is likely to obtain Traditional Foods from multiple locations.



Table 7-3: Comparison of Arsenic Bioavailability Assumptions and the Local Intake Fractions of Traditional Food for Adult Subsistence Harvester

Food Time	Arsenic Bioaccessibility Factors (unitless)						
Food Type	Assessed Scenario	Sensitivity Scenario					
Moose organs	0.19	1					
Moose meat	0.59	1					
Fish	0.1	1					
	Local Intake Fractions of Traditional Food (%)						

	Local Intake Fractions of Traditional Food (%)							
Food Type	Assesso	ed Scenario	Sensitivit	y Scenario				
rood type	Reference Location	Exposure Location	Reference Location	Exposure Location				
Northern pike	25.00	25.00	0	50.00				
Lake whitefish	25.00	25.00	0	50.00				
Store foods	41.59	41.59	0	83.18				
Moose	1.69	1.69	0	3.38				
Moose organs	0.75	0.75	0	1.50				
Beaver	4.68	4.68	0	9.36				
Mallard	0.28	0.28	0	0.56				
Grouse	1.01	1.01	0	2.02				
Labrador tea	9.88	9.88	0	19.76				
Fruits and berries	40.12	40.12	0	80.24				

% = percent.

Table 7-4: Estimated Incremental Lifetime Cancer Risk from Arsenic to Subsistence Harvester – Sensitivity Scenario

Receptor	Ri	Case Cancer sk 00,000)	Cancer F	nd Scenario Risk (per 000)	RFD Case Cancer Risk (per 100,000)		
	Operations	Far-Future	Operations	Far-Future	Operations	Far-Future	
Subsistence harvester - Patterson Lake South Arm (composite)	28	0.7	29	1	123	0.7	
Subsistence harvester - Beet Lake (composite)	1	0.03	1	0.04	1	0.03	
Subsistence harvester - Lloyd Lake (composite)	0.4	0.002	0.4	0.003	0.4	0.002	

Note: **Bold** indicates exceedance of the negligible cancer risk level of 1 in 100,000.

7.2.4 Climate Change

As a scenario within the RFD Case, potential effects from climate change, including how natural factors may be altered resulting from climate change, were considered. Indigenous Groups indicated concerns about cumulative effects from human development, policies, and climate change (TSD II: BNDN; TSD III: BRDN; TSD IV: MN-S; TSD V.1: CRDN).

Potential climate change influences on surface water quality in a scenario presented within the RFD Case were considered as a sensitivity scenario for the surface water quality model (EIS Section 10, Surface Water Quality and Sediment Quality). The sensitivity scenario utilized the same inputs as the RFD case, except that the inputs used from hydrology were reflective of potential changes associated with climate change. The results from the RFD Case plus climate change sensitivity scenario are discussed in EIS Section 10.5.2.1.6, Climate Change Sensitivity Scenario.

Overall, the predicted COPC concentrations within the LSA would be similar between the RFD Case and the RFD Case plus climate change sensitivity scenario. During both the Project lifespan and the far-future projection, climate change effects on surface water within the LSA would be negligible to minor relative to projections from the RFD Case. As such, the effects of climate change on the results of the ERA would also be negligible to minor, and no quantitative assessment was warranted.



8.0 CONCLUSIONS AND MONITORING

The selection of human and ecological receptors for inclusion in the ERA was informed by Indigenous Knowledge primarily through JWG meetings and IKTLU Studies, as well as other Local Knowledge, information from baseline studies, as well as professional judgement. The assumptions made for the Traditional Foods diet (i.e., amounts consumed and food types) was an iterative process that was informed by JWG meetings, IKTLU Studies, as well as discussions with representatives from Saskatchewan Ministry of Environment, Saskatchewan Health Authority and the CNSC.

The ERA focused on COPCs that exceeded screening values in air and water based on predicted atmospheric releases and aqueous releases (treated effluent, treated sewage, site runoff, and groundwater solute releases) from the Rook I Project. Based on the screening of atmospheric releases, no COPCs in air were advanced for further quantitative assessment in the ERA.

Based on the screening of aqueous releases, arsenic, cobalt, copper, molybdenum, uranium, sulphate, and chloride were advanced for further quantitative assessment in the ERA. Selenium, a constituent of interest for uranium mines and mills was assessed in terms of predicted fish tissue concentrations, since selenium toxicity in the aquatic environment is primarily from bioaccumulation of selenium in the aquatic food chain and not only from exposure to selenium in water. Predicted fish tissue concentrations in waterbodies immediately downstream of the source including Patterson Lake North Arm – West Basin were below the recommended USEPA fish tissue criterion; therefore, selenium was not considered further for quantitative assessment in the ERA.

Radionuclides, including the uranium-238 series and radon were included as COPCs because these constituents are of public interest.

8.1 Human Health Risk Assessment

The HHRA estimated dose and risk to the following receptors: camp worker, subsistence harvester, seasonal resident, and future permanent resident in the far-future projection. The Application Case, reasonable upper bound sensitivity scenario, and RFD Case were assessed for all phases of the Project and the far-future projection.

8.1.1 Non-radiological Human Health Risk Assessment

For assessment of non-carcinogens, risk was estimated based on an incremental HQ. Incremental HQs were compared to a benchmark HQ value of 0.2 because exposures to background sources were not included. This approach is consistent with Health Canada's guidance on human health preliminary quantitative risk assessment (Health Canada 2021a).

No significant adverse effect on any human receptors, as a result of releases from the Project, was likely during Operations for the Application Case, reasonable upper bound sensitivity scenario, and RFD Case. All estimated incremental HQs for all non-carcinogenic COPCs (cobalt, copper,



molybdenum, and uranium) remained below the acceptable risk level of 0.2 per pathway for the one-year-old and adult for all human receptors.

For assessment of risk for carcinogens (arsenic), the ILCR was estimated and compared against the cancer risk level of 1 in 100,000 recommended by Health Canada (2021a). Incremental cancer risk was predicted to exceed the negligible cancer risk level of 1 in 100,000 for the subsistence harvester at Patterson Lake South Arm just outside the Project footprint but is not expected to exceed the negligible cancer risk within the RSA farther from the Project. These findings are based on the conservative assumption of high consumption of Traditional Foods including fish and terrestrial animals in the Project footprint and LSA.

8.1.2 Radiological Human Health Risk Assessment

The incremental radiation dose to all human receptors during all Project phases and the far-future projection are below the regulatory public dose limit of 1 mSv/yr for the Application Case, the reasonable upper bound sensitivity scenario, and the RFD Case.

If a dose constraint of 0.3 mSv/yr is applied, the dose to the subsistence harvester (one-year-old) would be less than the dose constraint for the Application Case, the reasonable upper bound sensitivity scenario and RFD Case, and well below the regulatory public dose limit.

In the far-future projection, a future permanent resident living at the location of the camp could receive a dose up to 0.07 mSv/yr, well below the regulatory public dose limit and the dose constraint.

Overall, since the radiation dose estimates would be below the public dose limit, no discernable health effects are anticipated due to exposure of these receptors to radioactive releases from the Project.

8.2 Ecological Risk Assessment

The EcoRA estimated dose and risk to representative aquatic and terrestrial receptors. The Application Case, sensitive upper bound sensitivity scenario, and RFD Case were assessed for all phases of the Project and the far-future projection.

8.2.1 Non-radiological Ecological Risk Assessment

The potential for ecological effects was assessed by comparing exposure levels to toxicological benchmarks and characterized quantitatively in terms of HQs. An HQ greater than 1 indicates adverse effects may be possible for a given ecological receptor and further investigation would be warranted.

No significant adverse effect on either aquatic or terrestrial populations or communities, as a result of releases from the Project, would be likely during Operations, for the Application Case, reasonable upper bound sensitivity scenario, and RFD Case. All estimated HQs for all COPCs



ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT CONCLUSIONS AND MONITORING

(chloride, sulphate, arsenic, cobalt, copper, molybdenum, and uranium) for all ecological receptors would remain below the HQ benchmark of 1.

For the far-future projection, once groundwater solutes reach Patterson Lake North Arm – West Basin, HQs would remain below the benchmark of 1 for all COPCs with the exception of copper for the Application Case, reasonable upper bound sensitivity scenario, and RFD Case. Exceedances of the HQ benchmark value of 1 for the far-future projection would be limited spatially to the Patterson Lake and are limited in magnitude to just above the benchmark of 1. Exceedances of the HQ benchmark were predicted for lake whitefish, aquatic invertebrates, and zooplankton.

The TRV for lake whitefish, benthic invertebrates and zooplankton corresponds to the CCME's default water quality guideline for copper. The CCME water quality guideline is a conservative benchmark for copper below which adverse effects for most aquatic biota are not expected to occur. The most sensitive species include fresh water snails (*Lymnaea stagnalis* and *Pyrgulopsis robusta*), bivalves (*Villosa iris*), and water flea (*Daphnia magna*; ECCC 2021). The most sensitive endpoints for chronic exposure to copper include growth for benthic invertebrates, reproduction for zooplankton, and growth and reproduction for fish (ECCC 2021). The CCME water quality guideline is based on no or low effect levels for the most sensitive species; thus, the marginal TRV exceedances predicted by HQs just above 1 indicate a possible slight reduction in growth or reproduction for exposed individuals of sensitive species in Patterson Lake North Arm – West Basin. However, since the HQ exceedance is localized to Patterson Lake North Arm – West Basin, population level effects are not expected.

As noted in Section 6.4.1, Risk Estimation, the evaluation of risk related to copper exposure is an iterative process. Additional evaluation of this risk was completed in the aquatic health assessment (EIS Appendix 11A), with the objective of informing an adaptive management plan for assessing and reducing uncertainty related to waste rock storage area seepage and increasing mitigations if necessary.

Species at risk were assessed using surrogate species. The rusty blackbird and little brown myotis were identified as surrogate species at risk. Since no HQs above 1 were identified for these ecological receptors, individual species at risk would also be considered protected.

8.2.2 Radiological Ecological Risk Assessment

Radiation dose benchmarks of 9.6 mGy/d and 2.4 mGy/d (UNSCEAR 2008) were selected for the assessment of effects on aquatic biota and terrestrial biota, respectively, as recommended in CSA N288.6-22.

There were no predicted exceedances of the 9.6 mGy/d radiation dose benchmark for aquatic biota and 2.4 mGy/d radiation dose benchmark for terrestrial and riparian biota for the Application Case, the reasonable upper bound sensitivity scenario, and the RFD Case, during both the Project phases and the far-future projection.



Since there are no predicted exceedances of the respective dose benchmarks for any of the aquatic or terrestrial receptors, individual species at risk would also be considered protected.

8.3 Monitoring and Follow-up

The HHRA and EcoRA were developed based on the best available information for the Rook I Project, including baseline monitoring data, assumptions on estimates of source terms, Traditional Food diet (consumption rates and food types), among others.

Monitoring would focus on collecting data to verify ERA model predictions as well as provide data to improve model predictions as the Project begins. Recommended monitoring would support NexGen's adaptive management framework with the goal of reducing uncertainty over time through an iterative process:

- **Air quality:** There were no exceedances of annual screening values for any constituents, indicating that unacceptable chronic effects from direct exposure to air are not expected. Therefore, no air COPCs were identified in the ERA. However, short-term exceedances, based on maximum predicted concentrations for the 24-hour averaging period may occur within the Project footprint at the camp and at the fence line for nitrogen dioxide and particulate matter, including uranium in TSP and PM₁₀. The predicted exceedances are infrequent, short-term, and limited to within the Project footprint and at the associated boundary. Unacceptable levels of risk are not expected from infrequent, direct short-term exposures to these constituents in air. However, these constituents would be monitored as part of the Effluent and Emissions Plan.
- Traditional foods study: The assumptions for the Traditional Food diet were initially developed from the FNFNES undertaken in Saskatchewan in 2015 (Chan et al. 2018; Chan et al. 2019) in combination with professional judgement. Assumptions were subsequently modified based on discussions during JWG meetings, information available from IKTLU studies, and discussions with representatives from Saskatchewan Ministry of Environment, Saskatchewan Health Authority, and the CNSC. NexGen would be working with local Indigenous Groups in an effort to complete a targeted Traditional Foods study to help validate or modify the dietary assumptions made in the HHRA.
- Environmental monitoring: NexGen is implementing an Environmental Monitoring Plan consistent with requirements and guidance in CSA N288.4-19: Environmental monitoring programs at nuclear facilities and uranium mines and mills (CSA 2019). Monitoring would focus on providing data to verify the predictions made by the ERA, refine the models used in the ERA, and reduce the uncertainty in the predictions made by the ERA. The Environmental Monitoring Plan would include collection of surface water, sediment, and soil samples as well as fish tissue samples, benthic invertebrate tissue samples, and country foods such as blueberries. Monitoring locations would be focused in the area of Patterson Lake, but also extend out to the LSA and RSA to confirm predictions of limited spatial extent of effects. Monitoring constituents would include those identified as COPCs in the ERA, including metals and uranium-238 series radionuclides. However, monitoring could

ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT CONCLUSIONS AND MONITORING

extend to include other constituents for other purposes, such as meeting regulatory requirements for monitoring or constituents of public interest from other uranium mines and process plants.

- Worker Monitoring: To keep exposures to ionizing radiation hazards as low as reasonably achievable during all phases of the Project, exposures to gamma radiation, long-lived radioactive dust, radon progeny, and radon gas would be routinely monitored for workers designated as nuclear energy workers. Personal dosimetry equipment would be provided for all workers who require it, and dose records would be maintained for each nuclear energy worker at the Project site. Effective (whole body) and equivalent (organ-specific) doses would be measured and recorded as applicable. Doses would be routinely tracked and compared to internal and external limits. The processes for classifying nuclear energy workers and for managing worker dosimetry would be included in the Radiation Protection Program. Chemical, physical, or biological health and safety hazards encountered by workers during all phases of the Project would be monitored in accordance with established sample collection and analysis methods to quantify exposure and risk to workers and confirm the effectiveness of applicable controls. Results from personal occupational exposure and workplace monitoring would be collected, maintained, stored, communicated, and used to identify improvement opportunities, as required. The process for identifying health and safety hazards and monitoring occupational exposures would be outlined in the Health and Safety Program.
- **Far-Future projection effects:** Localized effects to some individual ecological receptors were predicted due to copper released from groundwater into Patterson Lake North Arm West Basin in the far-future projection. As it is difficult to verify effects predicted far into the future, adaptive management would be implemented throughout Operations to reduce the potential loads from surface and underground features into the groundwater in the far-future projection.



9.0 REFERENCES

9.1 Acts and Regulations

- The Clean Air Regulations. RRS c C-12.1, Reg 1. Repealed by c E-10.22, Reg 2 effective 1 June 2015. Available at https://www.canlii.org/en/sk/laws/regu/rrs-c-c-12.1-reg-1.html.
- Canadian Environmental Protection Act, 1999. SC 1999, c 33. Current to 7 October 2021. Available at https://laws-lois.justice.gc.ca/eng/acts/c-15.31/.
- Radiation Protection Regulations. SOR/2000-203 under the *Nuclear Safety and Control Act*. Last amended 22 September 2017. Available at https://laws-lois.justice.gc.ca/eng/regulations/SOR-2000-203/index.html.
- Nuclear Safety and Control Act. SC 1997, c 9. Last amended 1 January 2017. Available at https://laws-lois.justice.gc.ca/eng/acts/n-28.3/.
- Species at Risk Act. SC. 2002, c 29. Last amended 6 October 2020. Available at: https://laws-lois.justice.gc.ca/eng/acts/s-15.3/.
- The Wild Species at Risk Regulations. RRS c W-13.11 Reg 1 under *The Wildlife Act, 1997*. Effective 20 January 1999. Available at https://www.canlii.org/en/sk/laws/regu/rrs-c-w-13.11-reg-1.html.
- The Wildlife Act, 1998. SS 1998, c W-13.12. Effective 6 March 2000. Available at https://www.canlii.org/en/sk/laws/stat/ss-1998-c-w-13.12/latest/ss-1998-c-w-13.12.html.

9.2 Literature Cited

- Alberta. 2021. Alberta Ambient Air Quality Objectives. Ambient Air Quality Objectives and supporting documents. Accessed October 2021. Available at https://www.alberta.ca/ambient-air-quality-objectives.aspx#jumplinks-5.
- ATSDR (Agency for Toxic Substances and Disease Registry). 2004. Toxicological Profile for Cobalt. Available at http://www.atsdr.cdc.gov/toxprofiles/tp33.pdf.
- ATSDR. 2007. Toxicological Profile for Arsenic. Available at https://www.atsdr.cdc.gov/ToxProfiles/tp2.pdf.
- Arrington, L.R. and Davis, G.K. 1953. Molybdenum toxicity in the rabbit. J. Nutr. 51:295-304.
- Bartlett, A. (University of Waterloo). 2009. An assessment of the chronic toxicity of sodium chloride to Hyalella azteca. Unpublished data presented at the 1st International Conference on Urban Drainage and Road Salt Management in Cold Climates: Advances In Best Practices. May.



- BC MOE (British Columbia Ministry of Environment). 2013. Ambient Water Quality Guideline for Sulphate. Technical Appendix. Update. April. Water Protection & Sustainability Branch.
- BC MECCS (British Columbia Ministry of Environment and Climate Change Strategy). 2020. B.C. Source Drinking Water Quality Guidelines: Guideline Summary.
- BC MECCS. 2021. British Columbia Approved Water Quality Guidelines: Aquatic Life, Wildlife & Agriculture. Available at https://www2.gov.bc.ca/assets/gov/environment/air-land-water/waterquality/water-quality-guidelines/approved-wqgs/wqg summary aquaticlife wildlife agri.pdf.
- Beanlands G, Duinker P. 1983. An ecological framework for environmental impact assessment in Canada. Halifax Nova Scotia: Institute for Resources and Environmental Studies, Dalhousie University 132 p.
- Beak (Beak International Inc.). 1999. Ecotoxicology test results. Environment Canada, National Water Research Institute. Unpublished report for M.S. Evans.
- Beresford, N.A., Barnett, C.L., Howard, B.J., Scott, W.A., Brown, J.E., Copplestone, D. 2008a. Derivation of transfer parameters for use within the ERICA Tool and the default concentration ratios for terrestrial biota. Journal of Environmental Radioactivity 1393–1407.
- Beresford, N.A., Gaschak, S., Barnett, C.L., Howard, B.J., Chizhevsky, I., Stromman, G., Oughton, D.H., Wright, S.M., Maksimenko, A., Copplestone, D., 2008b. Estimating the exposure of small mammals at three sites within the Chernobyl exclusion zone a test application of the ERICA Tool. Journal of Environmental Radioactivity 1496–1502.
- Beyer, W.N., Connor, E., Gerould, S. 1994. Estimates of Soil Ingestion by Wildlife. J. Wildl. Manage. 58.
- Birge, W.J., Hudson, J.E., Black, J.A., and Wasterman, A.G. 1979. Embryo-larval Bioassays on Inorganic Coal Elements and In Situ Biomonitoring of Coal-waste Effluents. In: D.E. Samuel, J.R. Stauffer, C.H. Hocutt, and W.T. Mason Jr. (eds.), Surface Mining and Fish/wildlife Needs in the Eastern United States. Fish and Wildlife Service, Office of Biological Services. FWS/OBS-78/81. U.S. Government Printing Office, Washington, D.C. pp. 97-104.
- Birge, W.J., Black, J.A., Westerman, A.G., Short, T.M., Taylor, S.B., Bruser, D.M., Wallingford, E.D. (Kentucky Natural Resources and Environmental Protection Cabinet), 1985.

 Recommendations on numerical values for regulating iron and chloride concentrations for the purpose of protecting warm water species of aquatic life in the Commonwealth of Kentucky. Memorandum of Agreement No. 5429.
- BNDN-JWG (Birch Narrows Dene Nation-Joint Working Group). 2019a. Meeting Minutes. Meeting #2. 4 December 2019.



- BNDN-JWG. 2019b. Meeting Minutes. Meeting #1. 25 October 2019.
- BNDN-JWG. 2020. Meeting Minutes. Meeting #7. 27 January 2020.
- BRDN-JWG (Buffalo River Dene Nation-Joint Working Group). 2019a. Meeting Minutes. Meeting #2. 5 December 2019.
- BRDN-JWG. 2019b. Meeting Minutes. Meeting #1. 1 November 2019.
- Brown, J., Strand, P., Hosseini, A., Borretzen, P. 2003. Handbook for Assessment of the Exposure of Biota to Ionising Radiation from Radionuclides in the Environment.
- Brown, J.E., Alfonso, B., Avila, R., Beresford, N.A., Copplestone, D., Prohl, G., Ulanovsky, A. 2008. The ERICA Tool. Journal of Environmental Radioactivity 99, 1371–1383. September.
- Brown, J., Alfonso, B., Avila, R., Beresford, N.A., Copplestone, N.A., Hosseini, A. 2016. A new version of the ERICA tool to facilitate impact assessments of radioactivity on wild plants and animals. Journal of Environmental Radioactivity 153 141–148.
- Buckley, J.A., Rustagi, K.P., Laughlin, J.D. 1996. Response of Lemna minor to sodium chloride and a statistical analysis of continuous measurements for EC50 and 95% confidence limits calculation. Bulletin of Environmental Contamination and Toxicology 57, 1003–1008.
- Burnett-Seidel, C., Liber, K., 2013. Derivation of no-effect and reference-level sediment quality values for application at Saskatchewan uranium operations. Environ. Monit. Assess. 185, 9481–9494.
- CanNorth (Canada North Environmental Services). 2018. Human Health Risk Assessment for the Eastern Athabasca Basin. Final Report. October.
- CanNorth. 2000. Hatchet Lake Dietary Survey. Final Report.
- CCME (Canadian Council of Ministers of the Environment). 1995. Protocol for the Derivation of Canadian Sediment Quality Guidelines for the Protection of Aquatic Life. CCME EPC-98E. Prepared by Environment Canada, Guidelines Division, Technical Secretariat of the CCME Task Group on Water Quality Guidelines, Ottawa.
- CCME. 1999a. Canadian Sediment Quality Guidelines for the Protection of Freshwater Aquatic Life. Updated September 2007; Accessed July 2021. Available at https://ccme.ca/en/current-activities/canadian-environmental-quality-guidelines.
- CCME. 1999b. Canadian Environmental Quality Guidelines. Available at https://ccme.ca/en/current-activities/canadian-environmental-quality-quidelines.
- CCME. 2007. A protocol for the derivation of water quality guidelines for the protection of aquatic life 2007.



- CCME. 2011a. Canadian Water Quality Guidelines for the Protection of Aquatic Life. Chloride Ion. Scientific Criteria Document Report No. PN 1460.
- CCME. 2011b. Canadian Water Quality Guidelines for the Protection of Aquatic Life. Uranium.
- CCME. 2021a. Canadian Environmental Quality Guidelines (CEQGs) provide science-based goals for the quality of aquatic and terrestrial ecosystems. Available at https://ccme.ca/en/summary-table.
- CCME. 2021b. Canadian Ambient Air Quality Standards. CAAQS. Available at https://ccme.ca/en/air-quality-report#slide-7.
- Chan, L., Receveur, O., Batal, M., Sadik, T., Schwartz, H., Ing, A., Fediuk, K., Tikhonov, C. (First Nations Food, Nutrition & Environmental Study (FNFNES). University of Ottawa.). 2019. Erratum to the First Nations Food, Nutrition and Environment Study (FNFNES): Results from Saskatchewan 2015. Ottawa: University of Ottawa, 2018.
- Chan, L., Receveur, O., Sadik, T., Schwartz, H., Ing, A., Fediuk, K., Tikhonov, C. (University of Ottawa). 2018. First Nations Food, Nutrition and Environment Study (FNFNES): Results from Saskatchewan (2015).
- Chapman, P.M., Allen, H.E., Z'Graggen, M.N. 1996. Evaluation of bioaccumulation factors in regulating metals. Environ. Sci. Technol. 30, 448A-452A.
- Chen, C.J., Chuang, Y.C., Lin, T.M., Wu, H.Y. 1985. Malignant neoplasms among residents of a blackfoot disease-endemic area in Taiwan: High-arsenic artesian well water and cancers. Cancer Research 45, 5895–5899.
- Canadian Nuclear Safety Commission (CNSC). 2020. REGDOC-2.9.1, Environmental Principles, Assessments and Protection Measures, Version 1.2. September 2020. ISBN 978-0-660-06255-6. Available at http://nuclearsafety.gc.ca/eng/pdfs/REGDOCS/REGDOC-2-9-1-Environmental-Principles-Assessments-and-Protection-Measures-eng.pdf.
- CNSC. 2021. Generic Guidelines for the Preparation of an Environmental Impact Statement pursuant to the *Canadian Environmental Assessment Act, 2012*. Available at https://nuclearsafety.gc.ca/eng/resources/environmental-protection/ceaa-2012-generic-eis-guidelines.cfm.
- COSEWIC (Committee on the Status of Endangered Wildlife in Canada.). 2002. COSEWIC assessment and update status report on the woodland caribou Rangifer tarandus caribou in Canada.
- COSEWIC. 2006. Rusty blackbird (Euphagus carolinus): COSEWIC assessment and status report 2017.
- COSEWIC. 2007. COSEWIC Assessment and Update Status Report on the Common Nighthawk.



- COSEWIC. 2011. COSEWIC Assessment and Update Status Report on the Barn Swallow.
- CRDN (Clearwater River Dene Nation). 2019a. Clearwater River Dene Nation Comments on Project Description NexGen Energy Ltd.'s Rook I Project. 31 May 2019.
- CRDN. 2019b. Comments in Respect to License Renewal for Orano Canada Inc. Cluff Lake Project. Canadian Nuclear Safety Commission. April 18, 2019. CMD 19-H3.12. File 6.01.07.
- CRDN-JWG (Clearwater River Dene Nation-Joint Working Group). 2020. Meeting Minutes. Meeting #3. 10 March 2020.
- CSA (Canadian Standards Association Group). 2019. CSA N288.4-19: Environmental monitoring programs at nuclear facilities and uranium mines and mills.
- CSA. 2020. CSA N288.1-20: Guidelines for calculating derived release limits for radioactive material in airborne or liquid effluents for normal operation of nuclear facilities.
- CSA. 2022. CSA N288.6-22: Environmental Risk Assessments at Nuclear Facilities and Uranium Mines and Mills.
- Davis, J.E., Fields, J.P. 1958. Experimental production of polycythemia in humans by administration of cobalt chloride. Proc Soc Exp Biol Med 99, 493–495.
- Durante M, Pugliese M. 2002. Estimates of radiological risk from depleted uranium weapons in war scenarios. Health Physics 82(1): 14-20.
- Earthroots. 2020. Wolves in Ontario. Accessed May 2021. Available at http://wolvesontario.org/wolves-ontario/.
- ECCC (Environment and Climate Change Canada). 2020. Federal Environmental Quality Guidelines, Strontium. July.
- ECCC. 2021. Canadian Environmental Protection Act, 1999 Federal Environmental Quality Guidelines. Copper. April 2021.
- ECCC. 2022. Federal Environmental Quality Guidelines Selenium. August.
- Elphick J.R.F., K.D. Bergh and H.C. Bailey. 2011a. Chronic toxicity of chloride to freshwater species: effects of hardness and implications for water quality guidelines. Environmental Toxicology and Chemistry. 30, 239-246.
- Elphick, J.R., Davies, M., Gilron, G., Canaria, E.C. 2011b. An aquatic toxicological evaluation of sulfate: The case for considering hardness as a modifying factor in setting water quality guidelines. Environmental Toxicology and Chemistry 30, 247–253.



- ENV (Saskatchewan Ministry of the Environment). 2021. Saskatchewan Environmental Quality Guidelines. Available at https://envrbrportal.crm.saskatchewan.ca/segg-search/.
- ENV (Saskatchewan Ministry of the Environment), 2012. Saskatchewan Air Quality Modelling Guideline.
- Environment Canada and Health Canada. 1993. Priority Substances List Assessment Report for Arsenic and its Compounds.
- Environment Canada and Health Canada. 2003. Priority Substances List Assessment Report: Releases of Radionuclides from Nuclear Facilities (Impact on Non- Human Biota). PSL2.
- Environment Canada. 2016. Federal Environmental Quality Guidelines, Vanadium. May.
- Environment Canada, 2017. Federal Environmental Quality Guidelines, Cobalt. May.
- FCSAP (Federal Contaminated Sites Action Plan). 2012a. Ecological Risk Assessment Guidance. March.
- FCSAP. 2012b. Ecological Risk Assessment Guidance, Module 3: Standardization of Wildlife Receptor Characteristics. Prepared for: Environment Canada (Government of Canada), Vancouver, B.C. March.
- Fission (Fission Uranium Corp.). 2019. Technical Report on the Pre-Feasibility Study of the Patterson Lake South Property using Underground Mining Methods, Northern Saskatchewan, Canada. NI 43-101 Report. Prepared by Roscoe Postle Associates Inc. for Fission Uranium Corp. November 2019. Available at https://www.miningnewsfeed.com/reports/PattersonLakeSouth UG PFS 09192019.pg
- Fission. 2021. Fission Project Description. Prepared by Clifton Engineering Group Inc. and Canada North Environmental Services Limited Partnership for Fission Energy Corp. November 22, 2021.
- Government of Saskatchewan. n.d. Clearwater River Provincial Parker. Accessed February 2022. Available at https://www.tourismsaskatchewan.com/provincialpark/1419/clearwater-river-provincial-park#sort=relevancy.
- Government of Saskatchewan. 2015. Table 20: Saskatchewan Ambient Air Quality Standards. Accessed August 2021. Available at https://envrbrportal.crm.saskatchewan.ca/Pages/SEQS/Table20-SEQS-SAAQS.pdf.
- Gilman, A.P., Villeneuve, D.C., Secours, V.E., Yagminas, A.P., Tracy, B.L., Quinn, J.M., Valli, V.E., Wiles, R.J., Moss, M.A. 1998. Uranyl nitrate: 28-day and 91-day toxicity studies in the Sprague-Dawley rat. Toxicological Science 41, 117–128.



- Goettl, J.P.Jr., P.H. Davies and J.R. Sinley. 1976. Water Pollution Studies. In: D.B.Cope (Ed.), Colorado Fish.Res.Rev.1972-1975, DOW-R-R-F72-75, Colorado Div.of Wildl., Boulder, CO: 68-75.
- Grobner, M. A., Cheeke, P.R., and Patton, N.M. 1986. Effect of Dietary Copper and Oxytetracycline on Growth and Mortality of Weanling Rabbits. Journal of Applied Rabbit Research. 9(2): 46-53.
- Hamilton, S.J., Mehrle, P.M. 1986. Metallothionein in fish: review of its importance in assessing stress from metal contaminants. Transactions of the American Fisheries Society 115, 596–609.
- Haseltine, S.D., Sileo, L. 1983. Response of American Black Ducks to Dietary Uranium: A Proposed Substitute for Lead Shot. Journal of Wildlife Management 47, 1124–1129.
- Health Canada, 1998. National Ambient Air Quality Objectives for Particulate Matter. Executive Summary. Part 1: Science Assessment Document.
- Health Canada, 2000. Food and Nutrition. Dietary Intakes of Contaminants and other Chemicals for Different Age-sex Groups of Canadians (1993 to 1999).
- Health Canada. 2006. Guidelines for Canadian Drinking Water Quality: Guideline Technical Document—Arsenic. Healthy Environments and Consumer Safety Branch.
- Health Canada. 2009. Government of Canada Radon Guideline. Last updated November 2009. Available at https://www.canada.ca/en/health-canada/services/environmental-workplace-health/radiation/radon/government-canada-radon-guideline.htm.
- Health Canada. 2010a. Federal Contaminated Site Risk Assessment in Canada, Part I: Guidance on Human Health Preliminary Quantitative Risk Assessment (PQRA), Version 2.0. Contaminated Sites Program. September.
- Health Canada. 2010b. Federal Contaminated Site Risk Assessment in Canada. Part VI: Guidance on Human Health Detailed Quantitative Radiological Risk Assessment (DQRARAD). September.
- Health Canada. 2010c. Part II: Health Canada Toxicological Reference Values (TRVs) and Chemical-Specific Factors, Version 2.0. September.
- Health Canada. 2011a. Food and Nutrition. Dietary Intakes of Contaminants and other Chemicals for Different Age-sex Groups of Canadians. Health Canada. Canadian Total Diet Study. Available at https://www.canada.ca/en/health-canada/services/food-nutrition/food-nutrition-surveillance/canadian-total-diet-study/dietary-intakes-contaminants-other-chemicals-different-sex-groups-canadians.html.



- Health Canada. 2011b. Canadian Guidelines for the Management of Naturally Occurring Radioactive Materials (NORM). Prepared by the Canadian NORM Working Group of the Federal Provincial Territorial Radiation Protection Committee.
- Health Canada. 2013. Interim Guidance on Human Health Risk Assessment for Short-Term Exposure to Carcinogens at Contaminated Sites. Federal Contaminated Sites Risk Assessment in Canada.
- Health Canada. 2014. Radon Reduction Guide for Canadians. Report No. 140040.
- Health Canada. 2016a. Guidance for evaluating Human Health Impacts in Environmental Assessment: Air Quality. Healthy Environments and Consumer Safety Branch, Health Canada, Ottawa, Ontario.
- Health Canada. 2016b. Human Health Risk Assessment for Ambient Nitrogen Dioxide. Healthy Environments and Consumer Safety Branch.
- Health Canada. 2016c. Human Health Risk Assessment for Coarse Particulate Matter. Healthy Environments and Consumer Safety Branch.
- Health Canada. 2018. Guidance for Evaluating Human Health Impacts in Environmental Assessments: Country Foods. Report No. H129- 54/5- 2018E-PDF.
- Health Canada. 2019a. Guidelines for Canadian Drinking Water Quality: Guideline Technical Document—Copper.
- Health Canada. 2019b. Guidelines for Canadian Drinking Water Quality: Guideline Technical Document—Uranium.
- Health Canada. 2020. Guidelines for Canadian Drinking Water Quality Summary Table. September 2020. Available at https://www.canada.ca/content/dam/hc-sc/ewh-semt/alt_formats/pdf/pubs/water-eau/sum_guide-res_recom/summary-table-EN-2020-02-11.pdf.
- Health Canada. 2021a. Federal Contaminated Site Risk Assessment in Canada: Guidance on Human Health Preliminary Quantitative Risk Assessment (PQRA). Version 3.0.
- Health Canada. 2021b. Federal Contaminated Site Risk Assessment in Canada: Toxicological Reference Values (TRVs) Version 3.0.
- Hinterland Who's Who. 2007. Profile on the Black Bear [WWW Document]. URL http://www.hww.ca/en/wildlife/mammals/black-bear.html.
- Hinterland Who's Who. 2017. Profile on the Beaver [WWW Document]. URL http://www.hww.ca/en/wildlife/mammals/beaver.html.



- Howell, G.O. and Hill, C.H., 1978. Biological Interaction of Selenium with Other Trace Elements in Chicks. Environ. Health Perspect. 25: 147-50.
- Hunder, G., Schaper J., Ademuyiwa, O., and Elsenhans, B., 1999. Species Differences in Arsenic-mediated Renal Copper Accumulation: A Comparison between Rats, Mice and Guinea Pigs. Hum. Exp. Toxicol. 18(11): 699-705.
- IAEA (International Atomic Energy Agency). 2010. Handbook of Parameter Values for the Prediction of Radionuclide Transfer in Terrestrial and Freshwater Environments., Technical Report Series No. 472.
- IAEA. 2014. Radiation Protection and Safety of Radiation Sources: International Basic Safety Standards. General Safety Requirements Part 3. No GSR Part 3.
- ICRP (International Commission on Radiological Protection). 1994. Dose Coefficients for Intakes of Radionuclides by Workers., Publication 68. Annals of the ICRP 24(4).
- ICRP. 2003. Basic Anatomical and Physiological Data for Use in Radiological Protection., ICRP Publication 89.
- ICRP. 2007. The 2007 Recommendations of the International Commission on Radiological Protection. ICRP Publication 103. Ann. ICRP 37 (2-4).
- ICRP. 2008. Environmental Protection the Concept and Use of Reference Animals and Plants., ICRP Publication 108.
- IOM. 2001. Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc. A Report of the Panel on Micronutrients, Subcommittees on Upper Reference Levels of Nutrients and of the Interpretation and Uses of Dietary Intakes, and the Standing Committee on the Scientific Evaluation of Dietary Reference Intakes. Food and Nutrition Board of the Institute of Medicine of the National Academies. National Academy Press, Washington, DC.
- InterGroup (Intergroup Consultants). 2018a. NexGen-Rook 1 Traditional Land Use Desktop Study. Draft. December.
- InterGroup. 2018b. Desktop Review of Other Land and Resources Use in the Vicinity of the Rook I Project. Draft. November.
- Jenner, H.A. and Janssen-Mommen, J.P.M. 1993. Duckweed Lemna minor as a tool for testing toxicity of coal residues and polluted sediments. Arch. Environ. Contam. Toxicol. 25(1):3-11.
- Kessler, E. 1974. Physiological and biochemical contributions to the taxonomy of the genus Chlorella. IX: Salt tolerance as a taxonomic character. Archives of Microbiology 100, 51–56.



- Laird, B.D., Chan, H.M. 2013. Bioacessibility of metals in fish, shellfish, wild game, and seaweed harvested in British Columbia, Canada. Food and Chemical Toxicology 58:381-387.
- Liber, K.S., de Rosemond, S., Budnick, K. (Toxicology Centre, University of Saskatchewan, for AREVA Resources Inc. and Cameco Corporation). 2007. Uranium Toxicity to Regionally-Representative Algae and Invertebrate Species.
- May-Passino, D.R.M. and A.J. Novak. 1984. Toxicity of arsenate and DDT to the Cladoceran Bosmina longirostris. Bull.Environ.Contam.Toxicol. 33(3):325-329.
- MECP (Ontario Ministry of the Environment, Conservation and Parks). 2020. Technical Assessment and Standards Development Branch, 2020. Ambient Air Quality Criteria.
- McConnell, R.P. 1977. Toxicity of molybdenum to rainbow trout under laboratory conditions. In: Chappell, W.R. and K.K. Peterson (Eds.). Molybdenum in the environment. Volume 2. The geochemistry, cycling, and industrial uses of molybdenum. Marcel Dekker, New York.
- McLoughlin, P.D., Stewart, K., Superbie, C., Perry, T., Tomchuk, P., Greuel, R., Singh, K., Truchon-Savard, A., Henkelman, J., Johnstone, J.F. (Department of Biology, University of Saskatchewan). 2016. Population dynamics and critical habitat of woodland caribou in the Saskatchewan Boreal Shield. Interim Project Report, 2013-2016.
- MN-S (Métis Nation-Saskatchewan). 2019. Comments concerning the NexGen Energy Ltd. Rook I Project Description. Submission to the Canadian Nuclear Safety Commission and Saskatchewan Ministry of the Environment re: NexGen Rook 1 Project. April 2019.
- MN-S-JWG (Métis Nation Saskatchewan-Joint Working Group). 2019a. Meeting Minutes. Meeting #1. 29 October 2019.
- MN-S-JWG. 2019b. Meeting Minutes. Meeting #2. 10 December 2019.
- MN-S-JWG. 2020. Meeting Minutes. Meeting #4. 27 February 2020.
- MOE (Ontario Ministry of the Environment). 1994. Water Management: Policies, guidelines, provincial water quality objectives.
- MOE. 2011. Technical Assessment and Standards Development Branch. Ontario Air Standards for Uranium and Uranium Compounds.
- Morales, K.H., Ryan, L., Kuo, T.L., Wu, M.M., Chen, C.J. 2000. Risk of internal cancers from arsenic in drinking water. Environmental Health Perspectives 108, 655–661.
- Morrison, L.L. and Chavez, E.R., 1983. Selenium-arsenic Interaction in the Weanling Pig. Canadian Journal of Animal Science 63(1): 239-246.



- NCRP (National Council on Radiation Protection and Measurements). 1996. Screening Models for Releases of Radionuclides to Atmosphere, Surface Water, and Ground Vol. I and Vol. II. Report No. 123.
- Nemec, M.D., Holson, J.F., Farr, C.H., and Hood, R.D., 1998. Developmental Toxicity Assessment of Arsenic Acid in Mice and Rabbits. Reprod. Toxicol. 12(6): 647-658.
- Newman, M.C., Unger, M.A. 2002. Fundamentals of Ecotoxicology, 2nd ed. CRC/Lewis Press, Boca Raton, FL.
- NexGen (NexGen Energy Ltd.). 2019. Community Information Session. Round One June 2019. Submitted to Saskatchewan Ministry of Environment Environmental Assessment Branch. Canadian Nuclear Safety Commission (CNSC). July 2019.
- Novak, A., Walters, B.S., and Passino, D.R.M., 1980. Toxicity of Contaminants to Invertebrate Food Organisms. Prog.Fish.Res.1980, Great Lakes Fish.Lab., U.S.Fish Wildl.Serv., Ann Arbor, MI:2.
- Olivares, M., Pizarro, F., Speisky, H., Lönnerdal, B., Uauy, R. 1998. Copper in Infant Nutrition: Safety of World Health Organization Provisional Guideline Value for Copper Content of Drinking Water. Journal of Pediatric Gastroenterology and Nutrition 26, 251–257.
- Ortolani, E.L., Machado, C.H., and Sucupira, M.C.A. 2003. Assessment of Some Clinical and Laboratory Variables for Early Diagnosis of Cumulative Copper Poisoning in Sheep. Veterinary and Human Toxicology [Vet. Hum. Toxicol.]. Vol. 45, No. 6, P. 289. Dec.
- Paternain, J.L., Domingo, J.L., Ortega, A., Llobet, J.M. 1989. The Effects of Uranium on Reproduction, Gestation, and Postnatal Survival in Mice. Ecotoxicol. Env. Saf. 17, 291–296.
- Persaud D, Jaagumagi R, Hayton A. 1993. Guidelines for the protection and management of aquatic sediment quality in Ontario. Water Resources Branch, Ontario Ministry of the Environment, Toronto, Canada.
- Richardson, G. (O'Connor Associates Environmental Inc). 1997. Compendium of Canadian Human Exposure Factors for Risk Assessment.
- Sample, B.E., Opresko, D.M., Suter, G.W.I. (United States Department of Energy). 1996. Toxicological Benchmarks for Wildlife: 1996 Revision. June.
- Sample, B.E., Suter, G.W.I. (Oak Ridges National Laboratory). 1994. Estimating Exposure of Terrestrial Wildlife to Contaminants. Report No. ES/ER/TM-125. September.
- Schroeder, H. A. and Mitchener, M. 1971. Toxic effects of trace elements on the reproduction of mice and rats. Arch. Environ. Health. 23: 102-106.



- Solaiman, S.G., Maloney, M.A., Qureshi, M.A., Davis, G., and D'Andrea, G. 2001. Effects of High Copper Supplements on Performance, Health, Plasma Copper and Enzymes in Goats. Small Ruminant Research. 41(2):127-139.
- Statistics Canada. 2004. Canadian Community Health Survey, Cycle 2.2, Nutrition.
- Stubblefield, W.A., Van Genderen, E., Cardwell, A.S., Heijerick, D.G., Janssen, C.R., De Schamphelaere, A.C. 2020. Acute and Chronic Toxicity of Cobalt to Freshwater Organisms: Using a Species Sensitivity Distribution Approach to Establish International Water Quality Standards. Environmental Toxicology and Chemistry 39, 799–811.
- Suter, G.W.I. (Oak Ridge National Laboratory). 1996. Risk Characterization for Ecological Risk Assessment of Contaminated Sites.
- Tetra Tech Inc. 2008. Aquatic Life Water Quality Criteria for Molybdenum, July 9, 2008. Prepared for Nevada Division of Environmental Protection. Bureau of Water Quality Planning.
- TCEQ (Texas Commission on Environmental Quality). 2016. Effects Screening Levels. Available at. https://www.tceq.texas.gov/toxicology/esl/ESLMain.html.
- Thomas, J W. and Moss, S. 1951. The effect of orally administered molybdenum on growth, spermatogenesis and testes histology of young dairy bulls. J. Dairy Sci. 34:929-934.
- Thompson, P.A., Kurias, J., Mihok, S. 2005. Derivation and use of sediment quality guidelines for ecological risk assessment of metals and radionuclides released to the environment from uranium mining and milling activities in Canada. Environ. Monit. Assess. 110, 71–85.
- Underwood, E.J. 1971. Trace elements in human and animal nutrition. Pages 116 to 140 in Molybdenum. Academic Press, New York.
- UNSCEAR (United Nations Scientific Committee on the Effects of Atomic Radiation). 2000. Volume 1, Annex B: Exposures from Natural Radiation Sources and Effects of Ionizing Radiation, Report to the General Assembly with Scientific Annexes.
- UNSCEAR. 2008. Sources and Effects of Ionizing Radiation. Annex E. Effects of Ionizing Radiation on Non-human Biota., Report to the General Assembly with Scientific Annexes.
- USEPA (United States Environmental Protection Agency). 1985. Proposed Standard for Radon-222 Emissions to Air from Underground Uranium Mines. Report No. EPA 520/1-85-010.
- USEPA. 1986. Standard for Radon-222 Emissions from Licensed Uranium Mill Tailings. Report No. EPA 520/1-86-009.
- USEPA. 1993. Wildlife Exposure Factors Handbook. Report No. EPA/600/R-93/187.



- USEPA. 2005a. Guidance for Developing Ecological Soil Screening Levels. OSWER Directive 9285.7-55. Revised. February.
- USEPA. 2005b. Ecological Soil Screening Levels for Arsenic Interim Final. March.
- USEPA. 2005c. Ecological Soil Screening Levels for Cobalt– Interim Fina. Interim Final. March.
- USEPA. 2007a. Aquatic Life Ambient Freshwater Quality Criteria Copper. February.
- USEPA. 2007b. Ecological Soil Screening Levels for Copper. Interim Final. February.
- USEPA. 2021. 2021 Revision* to: Aquatic Life Ambient Water Quality Criterion for Selenium Freshwater 2016. Report No. EPA 822-R-21-006. August.
- Vives I Batlle, J., Balonov, M., Beaugelin-Seiller, K., Beresford, N.A., Brown, J., Cheng, J.-J., Copplestone, D., Doi, M., Filistovic, V., Golikov, V., Horyna, J., Hosseini, A., Howard, B.J., Jones, S.R., Kamboj, S., Kryshev, A., Nedvecaite, T., Olyslaegers, G., Prohl, G., Ulanovsky, A., Vives Lynch, S., Yankovic, T., Yu, C. 2007. Inter-comparison of absorbed dose rates for non-human biota 349–373.
- Vives i Batlle, J., Beaugelin-Seiller, K., Beresford, N.A., Copplestone, D., Horyna, J., Hosseini, A., Hosseini, A., Johansen, M., Kamboj, S., Keum, D.-K., Kurosawa, N., Newsome, L., Olyslaegers, G., Vandenhove, H., Ryufuku, S., Vives Lynch, S., Wood, M.D., Yu, C. 2011. Radiation Environ. Biophys. 50:231-251.
- Vocke, R.W., Sears, K.L., O'Toole, J.J., and Wildman, R.B. 1980. Growth responses of selected freshwater algae to trace elements and scrubber ash slurry generated by coal-fired power plants. Water. Res. 14: 141-150.
- VST (Vizon SciTech Inc., for the Canadian Nuclear Safety Commission). 2004. Final Report on the Toxicity Investigation of Uranium to Aquatic Organisms.
- WHO (World Health Organization), 2017. Guidelines for drinking-water quality.
- Wood, C.M., Port, P. 2000. Intracellular pH regulation and buffer capacity in CO2/HCO3-buffered media in cultured epithelial cells from rainbow trout gills. J. Comp. Physiol. B. 170, 175–184.
- WSA (Saskatchewan Water Security Agency). 2017. Saskatchewan Water Quality Objective for the Protection of Aquatic Life Molybdenum. Fact Sheet. Report No. WSA 514.
- Wu, M.M., Kuo, T.L., Hwang, Y.H., Chen, C.J. 1989. Dose–response relation between arsenic concentration in well water and mortality from cancers and vascular diseases. American Journal of Epidemiology 130, 1123–1132.



YNLRO (Ya'thi Néné Lands and Resources Office). 2019. Comments concerning the NexGen Energy Ltd. Rook I Project Description. Written Submission from the Ya'thi Néné Lands and Resources Office In the Matter of; NexGen Energy Ltd. Rook I Project Description. May 2019.

Appendix A IMPACT Model Report





ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT - APPENDIX A - IMPACT MODEL

MODELLING REPORT

Langping Wu, Ph.D. Environmental Modeller Technical Lead

Rina Parker, M.A.Sc., P.Eng.

Rima Packer

Environmental Risk Assessment Specialist

Project Manager

Don Hart, Ph.D.

Senior Ecotoxicologist

Project Principal and Reviewer

TABLE OF CONTENTS

TAB	LE OF	CONTENTS	I
LIST	OF TA	ABLES	III
LIST	OF FIG	GURES	IV
ABB	REVIA	TIONS AND UNITS OF MEASURE	v
1.0	INTR	RODUCTION	1.1
	1.1	Overview of the IMPACT Model	1.1
	1.2	Objective of this Document	1.2
	1.3	Structure of this Document	1.3
2.0	THE	IMPACT MODEL	2.1
	2.1	The IMPACT Model Structure	2.1
	2.2	Water Polygons	2.2
	2.2.1	Ecological Receptors Residing in Water Polygons	2.3
		Aquatic Pathways	
		Radiological Dose to Aquatic Ecological Receptors	
		·	
	2.3	Land Polygons Terrestrial Ecological Receptors	
		Home Ranges and Residency Factors	
		Exposure Assumptions for Terrestrial Ecological Receptors	
		Terrestrial Pathways	
		Radiological Dose to Terrestrial Ecological Receptors	
	2.3.6	Non-radiological Risk to Terrestrial Ecological Receptors	2.25
		Exposure of Human Receptors	
		Exposure Assumptions for Human Receptors	
		Exposure Duration and Frequency Radiological Dose	
		Non-radiological Dose	
3.0	DEVI	3.1	
	3.1	Ambient Hydrology	3.1
		Lake Morphometry	3.1
	3.1.2	Surface Water Flows	3.2
	3.2	Water Quality	3.6

i

ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT - APPENDIX A - IMPACT MODEL

Table of Contents

	3.2.1	Background Water Quality	3.6
	3.3	Sediment Quality	3.16
	3.3.1	Background Sediment Quality	
	3.3.2	Water to Sediment Partitioning Coefficients	
	3.4	Air Quality	3.23
	3.5	Soil Quality	3.23
	3.5.1	Background Soil Quality	
	3.5.2	Soil Quality during Operations	
	3.6	Transfer of Constituents to Aquatic Receptors	3.24
	3.6.1	Aquatic Bioaccumulation Factors	3.25
	3.6.2	Model Validation	3.25
	3.6.3	Dose Coefficients for Aquatic Receptors	3.32
	3.7	Transfer of Constituents to Terrestrial Receptors and Humans	3.33
	3.7.1	Soil-to-Plant Transfer	
	3.7.2	Model Validation	3.34
	3.7.3	Dose Coefficients for Terrestrial Plants and Invertebrates	3.37
	3.7.4	Ingestion Transfer Factors for Terrestrial Receptors	3.38
		Inhalation Transfer Factors for Terrestrial Receptors	
	3.7.6	·	
<i>1</i> 0	DECE	:DENCES	<i>1</i> 1

LIST OF TABLES

Table 2-1: Water-Sediment Transport Modelling Parameters	2.8
Table 2-2: Estimated Home Ranges of Selected Terrestrial Ecological Receptors	2.14
Table 2-3: Residency Factors for Terrestrial Ecological Receptors	2.16
Table 2-4: Bird and Mammal Body Weights and Intake Rates	2.19
Table 2-5: Receptor Occupancy Factors	2.21
Table 2-6: Modelling Parameters for the Soil Model	2.22
Table 2-7: Summary of Receptor Characteristics for the Human Health Risk Assessment	2.26
Table 2-8: Annual Food Intakes (kg/yr) for Food Items of the Human Receptors' Diets	2.30
Table 2-9: Surrogate Ecological Receptors Used in the Traditional Foods Diet	2.31
Table 2-10: Local Intake Fractions for Human Diet	2.32
Table 2-11: Concentrations of Constituents of Potential Concern in Store Foods	2.33
Table 2-12: Summary of Residency Assumptions for Human Health Receptor Groups	2.34
Table 3-1: Lake Morphometry for all Modelled Lakes in the IMPACT Model	3.1
Table 3-2: Average Monthly Outflows at the Model Nodes	3.4
Table 3-3: Mean Monthly Local Inflows for All Waterbodies Modelled in IMPACT	3.5
Table 3-4: Calibrated Local Inflow Concentrations for Each Water Polygon Modelled in IMF	PACT
	3.7
Table 3-5: Baseline Water and Sediment Concentrations Used in the IMPACT Model	3.16
Table 3-6: Distribution Coefficients Used in the IMPACT Model	3.22
Table 3-7: Soil Baseline Concentrations	3.23
Table 3-8: Aquatic Bioaccumulation Factors (L/kg fw)	3.25
Table 3-9: External Dose Coefficients for Aquatic Receptors	3.32
Table 3-10: Internal Dose Coefficients for Aquatic Receptors	3.33
Table 3-11: Bioaccumulation Factors for Terrestrial Plants and Invertebrates	
Table 3-12: External Dose Coefficients for Terrestrial Plants and Invertebrates	3.37
Table 3-13: Internal Dose Coefficients for Terrestrial Plants and Invertebrates	3.37
Table 3-14: Ingestion Transfer Factors for Mammals (d/kg fw)	3.39
Table 3-15: Ingestion Transfer Factors for Birds (d/kg fw)	
Table 3-16: Inhalation Transfer Factors for Mammals and Birds (d/kg fw)	
Table 3-17: Internal Dose coefficients for Terrestrial Animals and Birds	3.42
Table 3-18: External Dose Coefficients for Terrestrial Animals and Birds	
Table 3-19: Human Dose Coefficients for Internal and External Exposure	3.44



LIST OF FIGURES

Figure 1-1: Rook I Project Site and Study Areas	1.3
Figure 2-1: Representation of Transfer between Aquatic Media in the IMPACT Model	2.5
Figure 2-2: Human Receptor Locations for the Rook I Model	2.27
Figure 3-1: Waterbodies and Flow Direction as Implemented in the IMPACT Model	3.3
Figure 3-2: Modelled and Measured Baseline Concentrations in Water	3.15
Figure 3-3: Modelled and Measured Baseline Concentrations in Sediment	3.21
Figure 3-4: Modelled and Measured Baseline Concentrations (fresh weight) of Radionu	clides in
Macrophytes	3.28
Figure 3-5: Modelled and Measured Baseline Concentrations (fresh weight) of Radionu	clides in
Northern Pike	3.29
Figure 3-6: Modelled and Measured Baseline Concentrations (fresh weight) of Radionu	clides in
Lake Whitefish	3.31
Figure 3-7: Modelled and Measured Concentrations (dry weight) of Radionuclides in Bl	ueberries
	3.36



ABBREVIATIONS AND UNITS OF MEASURE

LIST OF ACRONYMS

Acronym	Definition
BAF	bioaccumulation factor
COPC	constituent of potential concern
COSEWIC	Committee on the Status of Endangered Wildlife in Canada
CSA	Canadian Standards Association
DCF	dose coefficient
dw	dry weight
fw	fresh weight
HQ	hazard quotient
IMPACT	Integrated Model for the Probabilistic Assessment of Contaminant Transport
TF	transfer factor
USEPA	United States Environmental Protection Agency

LIST OF UNITS

Unit	Definition
%	percent
μGy/h	micrograys per hour
Bq/kg	becquerels per kilogram
Bq/L	becquerels per litre
Bq/m ²	becquerels per square metre
Bq/m ³	becquerels per cubic metre
cm/h	centimetres per hour
d	day
h	hour
ha	hectare
kg	kilogram
kg/L	kilograms per litre
kg/m²/s	kilograms per square metre per second
kg/yr	kilograms per year
km	kilometre
km ²	square kilometre
L	litre
L/d	litres per day



ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT - APPENDIX A - IMPACT MODEL

Abbreviations and Units of Measure

Unit	Definition
L/kg	litres per kilogram
L/m ² /s	litres per square metre per second
L/m ³	litres per cubic metre
L/s	litres per second
m	metre
m/s	metres per second
m ²	square metre
m ³	cubic metre
m³/d	cubic metres per day
m³/kg	cubic metres per kilogram
m³/yr	cubic metres per year
mg	milligram
mg/kg	milligrams per kilogram
mg/kg/d	milligrams per kilogram per day
mg/L	milligrams per litre
mg/m³	milligrams per cubic metre
mg/s	milligrams per second
mSv/yr	millisieverts per year
S	second
Sv/Bq	sieverts per becquerel
Sv/yr	sieverts per year
yr	year



1.0 Introduction

1.1 Overview of the IMPACT Model

The environmental transport and pathways model, IMPACT (Integrated Model for the Probabilistic Assessment of Contaminant Transport), is used to evaluate the transport and effects of constituents of potential concern (COPCs) on the local environment and receptors, including humans and non-human biota. The model represents a convenient platform and powerful tool to complete systematic evaluations of the risks to ecological and human receptors associated with releases of constituents to water and air from proposed or existing anthropogenic (i.e., human) activities. The scope of this appendix is to provide further detail on the modelling approach followed to calculate dose and risk to ecological and human receptors, as reported in the main body of Technical Support Document (TSD) XXI, Environmental Risk Assessment.

IMPACT is a modelling tool, the current version of which was created, and is maintained and supported, by Ecometrix Incorporated (Ecometrix). The IMPACT model was originally developed by BEAK Consultants Ltd. in 1993 as part of a research initiative partially funded by the Atomic Energy Control Board (now the Canadian Nuclear Safety Commission, CNSC). Since the initial development, IMPACT has been continuously updated to improve the interface, to integrate various operating systems, and most importantly, to embody an up-to-date understanding of the fate, transport, and toxicity of metals, radionuclides, and other constituents released to the environment.

IMPACT updates have been funded by the CANDU Owners Group in order to ensure that it is continually aligned with the Canadian Standards Association (CSA) relevant standards. The CSA standard N288.1 presents analytical equations and parameters for radiological pathways analysis, developed with the participation of experts from the nuclear industry, government agencies and consultants, including the CNSC and Ecometrix. The CSA standard N288.6 provides guidance for environmental risk assessment at nuclear facilities, and references N288.1 for radiological risk assessment.

The IMPACT 5.6.1 is a dynamic version of the model and was tailored to align with the guidance in CSA standards N288.6-22 (CSA Group, 2022) and N288.1-20 (CSA Group, 2020). It contains differential equations for COPC transport, allowing for non-steady-state conditions, whereas N288.1 contains the corresponding steady-state equations.

The IMPACT model is a customizable tool that allows the user to assess the transport and fate of COPCs through a user-specified environment. The model is used to estimate concentrations of COPCs in a range of environmental media, based on releases and environmental features. The IMPACT model enables the quantification of potential doses and hazard quotients (HQs) for aquatic and terrestrial ecological receptors, as well as humans. The graphical user interface features make it possible to create or modify scenarios quickly and without the need to change the programming code. Thus, users can construct complex models to predict potential environmental effects in a wide variety of natural environments without the need for programming skills or the use of multiple, complex model interfaces.

The IMPACT model has been applied to ecological and human health risk assessments at several proposed and operating uranium mines and mills including Cigar Lake Mine (Cameco, 2004), Key Lake Mine (Ecometrix 2005, 2012, 2013a) and Millennium Mine (Ecometrix 2013b, c). This model has also been used extensively by other nuclear facilities for ecological and human health risk assessments, to support preparation of derived release limits and for annual public dose calculations. The extensive environmental database developed for northern Saskatchewan since the 1970s to present day has gone through numerous updates as a larger available database of environmental data became available over time. This has helped develop more statistically rigorous relationships for COPC transfer among various environmental compartments. Substantial effort was made during this assessment to review parameters in the model and compare to the existing baseline dataset for the Rook I Project, which is further discussed in Section 3.0, Development of Model Parameters.

1.2 Objective of this Document

The objective of this document is to present the structure and functioning of the IMPACT model as implemented for the Rook I Project (the Project). The Project is a proposed new uranium mining and milling operation located adjacent to Patterson Lake in the southwestern Athabasca Basin in northern Saskatchewan. The Project site and surrounding local and regional study areas are shown in Figure 1-1.




Figure 1-1: Rook I Project Site and Study Areas

This document discusses the inputs and assumptions used in the IMPACT model, including receptor characteristics, exposure pathways, and the derivation and identification of site-specific information.

1.3 Structure of this Document

The document contains the following sections and content:

- Section 2.0: Describes the model structure for ecological and human receptor assessment, specific assumptions made for the Project, and the generic equations used to calculate the transfer of constituents between environmental media and the receptors.
- Section 3.0: Presents the development of project-specific input parameters and describes the approach used for calibration and validation based on regional monitoring data.

2.0 The IMPACT Model

The IMPACT model simulates the transport of constituents from sources through various environmental media such as air, water, soil, and sediment, to receptors. The model estimates the resulting concentration of constituents in environmental media, potential uptake (i.e., absorption) by aquatic and terrestrial vegetation and animals, and potential intake (i.e., ingestion) by and dose to animals and humans.

Environmental pathways are the transport, transformation, and transfer mechanisms by which COPCs travel from sources to receptors. A pathway exists when there is a point at which COPC uptake or external exposure by an ecological or human receptor may occur, following COPC transport from the source through environmental media. If a complete transport pathway does not exist between the source of a COPC and an ecological or human receptor, then no exposure is expected to occur.

The links within a pathway represent different processes of COPC transfer, which depend on the receptors and environmental conditions. These processes can include intake, transfer, and accumulation of constituents.

IMPACT is consistent with the calculations outlined in CSA N288.1-20 and CSA N288.6-22. As such the default model parameters are consistent with these publications. All environmental parameters including bioaccumulation factors, transfer factors and other constants can be adjusted to site or regional data as these are available. In general, it is preferable to use regional values, and this approach was followed in this assessment. Values for water-sediment transfer as well as aquatic animals and plants were based on regional datasets, which have been developed using extensive environmental data collected over many years (Ecometrix 2005, 2012, 2013c). These values will be identified as regional data. For the remaining factors, literature values were selected. In this selection process, the CSA standards were preferred to other available literature.

This IMPACT model was set up to be representative of the environment it models and the transfer processes between environmental components. Where deemed applicable, field measurements were used either as inputs or as a validation of modelled values. Some model parameters were selected based on regional monitoring data and previous model calibrations to regional data for northern Saskatchewan. Model-predicted COPC concentrations for the Rook I baseline condition were compared to measured Rook I baseline data so that model predictions would not be underestimating as compared to measurements.

2.1 The IMPACT Model Structure

The basic spatial units of assessment in IMPACT are "polygons". IMPACT supports two types of polygons, aquatic and terrestrial, which represent aquatic and terrestrial environments respectively. Ecological and human receptors inhabit polygons. Polygons represent zones of surface water or land that have similar physical, chemical, biological, and/or hydrological characteristics. Individual polygons are given specific attributes, such as topography for land polygons and water depth and flow for water polygons, which can be based on site-specific

information. Each polygon is given a specific spatial extent that is defined by a centroid point (with X and Y coordinates) and a surface area. Polygons can be connected by water or air pathways. A number of receptors may reside within each polygon and may have exposure connections to other polygons.

The transfer of constituents between environmental media, receptors, and polygons is conceptualized by links (i.e., arrows indicating direction of transfer). The links represent different transfer processes depending on the context. For example, a link between two waterbodies may represent flow of water, and a link between water and sediment may represent sedimentation. Each transfer process is represented by a differential equation.

Transfer of constituents is modelled iteratively with a constant user-selected timestep. IMPACT uses the Euler Method to numerically solve the set of ordinary differential equations at each timestep. The modelling timestep for this Project was selected to optimize computing time and produce a convergent solution in a dynamic system. The modelling timestep chosen was 0.02 years, which roughly equates to one week. Results were monitored at intervals of 0.02 years to 2 years, depending on the change rate for the scenario.

2.2 Water Polygons

Lakes and streams are defined within IMPACT as water polygons, which are distinct from land polygons. Water polygons can be inhabited by aquatic receptors and provide exposure pathways for terrestrial and human receptors.

The IMPACT model includes flow and mass balance in lakes and streams. Constituents enter the aquatic environment from a source and travel through various waterbodies. As constituents travel through a series of connected waterbodies such as lakes, concentrations in water can decrease as a result of mixing with natural inflows from the surrounding watershed and interactions with lake sediment. The sediment-water exchange of constituents is estimated using chemical-specific partitioning coefficients. The water and sediment pathways involve the exchange of constituents between surface water and sediment through the following processes:

- sorption and desorption between dissolved and particulate forms in water and sediment;
- settling of particulates from water to sediment;
- diffusive exchange between sediment porewater and the water column; and
- loss to deeper sediments through accumulation and burial.

The model estimates concentrations in the water and sediment using the advection dispersion equation (detailed below), which is essentially a mass conservation equation. The partial differential equations are solved iteratively. The model estimates changes in concentrations in water and sediment through each downstream waterbody, over time.



Aquatic receptors reside in water polygons. Terrestrial receptors (e.g., moose, humans) reside in land polygons and can be linked to aquatic receptors from water polygons.

The IMPACT model included the following distinct water polygons: Broach Lake, Patterson Lake North Arm – West Basin, Patterson Lake South Arm, Beet Lake, Lloyd Lake. Forrest Lake North Basin was considered as a flow-through channel, in the same way as the Clearwater River. This is discussed in more detail in Section 3.1.1, Lake Morphometry.

The Clearwater River has short sections in between these waterbodies, so Clearwater River chemical characteristics would be similar to the outflow of the upstream waterbody at any given location within the study area. The Clearwater River between Beet Lake and Lloyd Lake is modelled in sections to reflect increased flow rates at various points along the way.

2.2.1 Ecological Receptors Residing in Water Polygons

Aquatic plants and animals are assigned to water polygons. The aquatic ecological receptors that were considered in this iteration of the IMPACT model include aquatic macrophytes (i.e., plants), phytoplankton, zooplankton, benthic invertebrates, and fish.

Macrophytes

Aquatic macrophytes are primary producers that occupy the lowest level in the food chain and are exposed to constituents in surface water. Macrophytes can potentially uptake metals in their roots and shoots and are modelled accordingly. Macrophytes provide a pathway for the introduction of bioavailable constituents and their compounds into the food chain through direct consumption by terrestrial herbivores (e.g., moose [Alces alces] and woodland caribou [Rangifer tarandus caribou]).

Phytoplankton

Phytoplankton are primary producers that occupy the lowest level in the food chain and are assessed for exposure to constituents in surface water.

Zooplankton

Zooplankton are primary consumers that occupy the second lowest level in the food chain and are assessed for exposure to constituents in surface water.

Benthic Invertebrates

Benthic invertebrates are primary consumers. These organisms are important food sources for aquatic and semi-aquatic animals. In the IMPACT model, benthic invertebrates are assumed to be exposed to COPCs in the aquatic environment directly through contact with water and sediment.




Fish

Fish were collected from waterbodies within the vicinity of the Project local and regional study areas to provide data on baseline COPC concentrations in muscle and bone tissues in large-bodied fish (Annex V.1, Aquatic Environment Baseline Report). Fish species observed and sampled include northern pike (Esox lucius) and lake whitefish (Coregonus clupeaformis). Northern pike was selected for IMPACT modelling purposes to represent a piscivorous (i.e., fish-eating) top carnivore. Lake whitefish was selected for **IMPACT** modelling represent bottom-feeding fish. Northern pike and lake whitefish represent different trophic levels. Although there may be other fish species that are not included explicitly in the model, northern pike and lake whitefish are assumed to be representative of other similar species. Since baseline data are available for these species, model predictions can be validated against measured data (Section 3.6.2, Model Validation).

Fish species were used as surrogates for amphibians (e.g., frogs) because the sensitive life stages of amphibians (i.e., egg and tadpole) are aquatic and similar to the sensitive life stages of fish. During the tadpole stage, tadpoles and fish have similar exposure pathways (i.e., gills for breathing, absorption through skin, and similar feeding habits). Toxicological data for amphibians are limited, but the available data for cadmium, cobalt, copper, lead, nickel, and zinc for sensitive life stages indicate that the selected toxicity reference values for fish are protective of amphibians. Therefore, the fish species in the model are expected to provide a reasonable surrogate for, and protection of, amphibians during the sensitive tadpole stage.

2.2.2 Aquatic Pathways

Aquatic pathways include the transfer of COPCs between water, sediment, aquatic animals, and aquatic plants. Environmental media (i.e., water and sediment) and aquatic ecological receptors (i.e., aquatic plants, phytoplankton, zooplankton, benthic invertebrates, and fish) are connected by arrows in the IMPACT model (Figure 2-1). The arrows represent equations listed below the figure. Model outputs include concentrations in environmental media and receptors, and dose and risk values for aquatic ecological receptors.



DRM = Dose-Risk Monitor; ConcM = Concentration Monitor.

Figure 2-1: Representation of Transfer between Aquatic Media in the IMPACT Model



The equations for concentrations in water and sediment are partial differential equations that are solved numerically within IMPACT. Each of these equations is characterized by a series of parameters that describe the physical and biochemical environment of each lake and stream represented by the model. These parameters can be divided into general categories to represent the following components and processes:

- physical environment;
- natural background conditions;
- project-related constituent loadings; and
- biochemical exchange processes.

There are four basic equations that describe the concentrations in the aquatic environment for water and sediment.

Water column concentration (C_{wc}) :

$$\frac{dC_{wc}}{dt} = \frac{W_w + Q_{in} \cdot C_{in} + Q_{gw} \cdot C_{gw}}{V_w} + \lambda_{parent} \cdot C_{parent} - \frac{k_s}{Z_w} \left[(1 - f_w) \cdot C_{wc} - C_{pw} \right]$$
$$-C_{wc} \cdot \left[\lambda_T + \frac{g_w \cdot f_w}{Z_w} + \frac{Q_{out} + (1 - f_w) \cdot Q_{gw \to out}}{V_w} \right]$$

Sediment layer concentration (C_s):

$$\frac{dC_{s}}{dt} = \frac{g_{w} \cdot f_{w} \cdot C_{wc}}{z_{s}} + \lambda_{parent} \cdot C_{parent} + \frac{k_{s}}{z_{s}} \left[(1 - f_{w}) \cdot C_{wc} - C_{pw} \right] + \frac{Q_{gw} \cdot C_{gw}}{V_{s}}$$
$$-C_{s} \cdot \left[\lambda_{T} + \frac{g_{b}}{z_{s}} + \frac{(1 - f_{s}) \cdot Q_{gw \mapsto out}}{\varepsilon_{s} \cdot V_{s}} \right]$$

The fraction of a constituent that is particulate in the water column (fw) defined as:

$$f_{w} = \frac{K_{d} \cdot \frac{\rho_{s}}{\varepsilon_{s}}}{1 + K_{d} \cdot \frac{\rho_{s}}{\varepsilon_{s}}}$$

The sediment-water transport coefficient (k_s) defined as:

$$k_{S} = \frac{D^{*}}{z_{i}}$$

The burial rate of sediments (g_b) defined as:

$$g_b = \frac{g_w \cdot SS}{\rho_s}$$

Aquatic animal concentrations (Caa) are calculated as:

$$C_{aa} = C_{wc} \cdot BAF_{aa} \cdot \alpha \cdot (1 - OF_s) + C_{pw} \cdot BAF_{aa} \cdot \alpha \cdot OF_s$$

Aquatic plant concentrations (Cap) are calculated as:

$$C_{ap} = C_{wc} \cdot BAF_{ap}$$

where:

α = food web multiplier(s) (unitless)
 BAF_{aa} = bioaccumulation factor for aquatic animals L/kg)
 BAF_{ap} = bioaccumulation factor for aquatic plants L/kg)

 C_{in} = concentration of constituent entering water column (mg/L, Bq/L)

 C_{qw} = concentration of constituent in seepage (input) groundwater (mg/L, Bq/L)

 C_{parent} = concentration of parent constituent (mg/L, Bg/L)

 C_{pw} = concentration in the surficial sediment pore water (mg/L, Bq/L) C_s = concentration of constituent in surficial sediments (mg/kg, Bq/kg)

 C_{wc} = concentration of constituent in water column (mg/L, Bq/L)

D* = sediment-water column diffusion coefficient (m²/s)

 ε_s = porosity of surficial sediment (unitless)

 f_s = fraction of a constituent that is particulate in the sediment layer (unitless) f_w = fraction of a constituent that is particulate in the water column (unitless)

 g_b = burial rate of sediments (m/s)

 g_w = settling rate of particulates in water column (m/s) K_d = water-to-sediment partitioning coefficient (L/kg) k_s = sediment-water transport coefficient (m/s)

 λ_{parent} = first-order decay constant for parent constituent (1/s)

 λ_T = total first-order decay constant for constituent (1/s), which is the sum of the universal

decay constant and the media-specific decay constant

OF_S = sediment occupancy factor (unitless)

Q_{in} = inflow rate from upstream surface water (L/s)

 Q_{gw} = inflow rate from groundwater (L/s)

 Q_{out} = net outflow rate to downstream surface water (L/s)

 $Q_{gw \rightarrow out}$ = outflow rate to groundwater (L/s)

 ho_s = bulk density of surficial sediment (kg dw/L) SS = suspended solids concentration (mg/L) V_s = volume of surficial sediment layer (L) V_w = volume of surface waterbody (L)

Ww = total effluent emission rate from all sub-sources (mg/s, Bq/s)
 z_i = sediment-water column diffusion interface thickness (mm)

 z_s = thickness of sediment layer (m)

 $z_w = mean lake depth (m)$

Many parameters representing the physical environment were derived from baseline studies (Annex IV.2, Hydrometric Monitoring Characterization Report) including waterbody surface areas and volumes. Information from published literature and from experience with similar environments was used to quantify physical parameters that are conceptual or that were not observed directly (e.g., sediment interface thickness). A summary of these parameters used as inputs for the above transport equations is presented in Table 2-1.

It should be noted that the underlying assumption for water-sediment transport in IMPACT is that minimal suspended solids will be introduced through the effluent.

Table 2-1: Water-Sediment Transport Modelling Parameters

Model Parameter		Value	Unit
Mixing depth (thickness of sediment layer)	Z _s	0.03	m
Dry bulk density	ρ_s	0.11	kg dw/L
Water content (porosity	٤s	0.96	unitless
Diffusion coefficient	D*	3.16 × 10 ⁻¹⁰	m²/s
Interface thickness	Zi	0.01	m
Settling rate	gw	2	mm/yr
Suspended solids	SS	2	mg/L

Note: Values calibrated based on regional data.

dw = dry weight.

The natural background conditions represent the quality of water and sediment within the watershed prior to mining. Where possible, field data were used to quantify natural background conditions (Section 3.2.1, Background Water Quality and Section, 3.3.1, Background Sediment Quality). Monitoring data indicated that concentrations of some constituents in water were below analytical detection limits. Concentrations in the sediments from lakes were generally measurable. Water-sediment partitioning coefficients were selected based on the available data (Section 3.3.2, Water to Sediment Partitioning Coefficients).

The source loads of constituents from natural and Project-related sources represent the boundary conditions of the model. Natural sources are represented by the chemical influx from natural groundwater discharge, overland runoff, and stream inflows from the surrounding landscape. These natural loadings were estimated from natural background water quality and inflow rates for site-specific conditions.

2.2.3 Radiological Dose to Aquatic Ecological Receptors

Radiological dose to an aquatic receptor is the radiation energy absorbed due to radiation emissions from radionuclides present in the environment (water or sediment) or in the tissues of the organism. Through comparison with dose benchmarks, the risk to the organism can be quantified. This model characterizes dose and risk resulting from both background and project exposure.

The total dose (D_{total}) to an aquatic ecological receptor from each constituent is:

$$D_{total} = D_{int} + D_{ext}$$

The dose of COPCs to an aquatic ecological receptor due to incorporated radioactivity (D_{int}) is:

$$D_{int} = C_t \cdot DCF_{int}$$

The external dose to the aquatic ecological receptor from radioactivity in water and sediment (D_{ext}) is:

$$D_{ext} = [C_w \cdot (OF_w + 0.50F_{ws} + 0.50F_{ss}) + C_s \cdot (OF_s + 0.50F_{ss})] \cdot DCF_{ext}$$

where:

C_t = whole body tissue concentration from water (becquerels per kilogram fresh weight (Bq/kg fw))

 C_w = water concentration (becquerels per litre (Bq/L))

C_s = sediment concentration (Bq/kg fw)

 D_{int} = internal radiation dose (micrograys per hour (μ Gy/h))

 D_{ext} = external radiation dose (μ Gy/h)

 DCF_{int} = dose coefficient for radionuclide in tissue ([μ Gy/h])/[Bq/kg fw])

 DCF_{ext} = external dose coefficient for water ([μ Gy/h]/[Bq/kg fw sediment] or [μ Gy/h]/[Bq/L

water])

OF_s = fraction of time spent immersed in sediment (unitless). Assumption based on characteristics of the organism; see Table 2-5

OF_{ss} = fraction of time spent on the sediment surface (unitless). Assumption based on characteristics of the organism; see Table 2-5

 OF_w = fraction of time spent immersed in the water column (unitless). Assumption based on

characteristics of the organism; see Table 2-5

 OF_{ws} = fraction of time spent on the water surface (unitless). Assumption based on

characteristics of the organism; see Table 2-5

2.2.4 Non-radiological Risk to Aquatic Receptors

Risk to aquatic receptors from exposure to non-radionuclide COPCS is expressed as a Hazard Quotient (HQ). For aquatic receptors, it is characterized through comparison of the exposure concentration with a reference toxicity concentration. This model calculates the risk from both background and Project exposure. The HQ for aquatic ecological receptors due to exposure to COPCs in water is estimated as:

$$HQ = \frac{1}{RC_{wc}} \cdot \left[C_{wc} \cdot (1 - OF_s) + C_{pw} \cdot OF_s \right]$$

where:

 C_{pw} = concentration in the surficial sediment pore water (mg/L) C_{wc} = concentration of constituent in water column (mg/L)

OF_s = sediment occupancy factor (unitless). Assumption based on characteristics of the

organism; see Table 2-5

RC_{wc} = reference toxic concentration – water column (mg/L)

2.3 Land Polygons

Terrestrial receptors are modelled as residing in land polygons. Land polygons provide exposure pathways for terrestrial and human receptors. Land polygons are populated by one or more terrestrial receptors that are expected to occupy the habitat represented by the prevailing vegetation community and physical characteristics of the surrounding environment.

IMPACT models both terrestrial and aquatic dietary components for terrestrial receptors. Terrestrial pathways include the transfer of constituents between air, soil, water, plants, and animals.

2.3.1 Terrestrial Ecological Receptors

Terrestrial receptors are divided into ecological receptors and human receptors. Ecological receptors typically are selected to include representative species of terrestrial plants, small and large mammals, invertebrates, birds, and riparian animals. Human receptors can be modelled to represent the habits of population groups and ages that are expected to reside in the area of interest (Section 2.4, Exposure of Human Receptors).

Terrestrial ecological receptors are assigned to a land polygon. The terrestrial ecological receptors that were selected include primary producers (plants) and consumers (invertebrates and animals). Primary consumers (herbivores) and secondary consumers (omnivores and carnivores) were selected from among mammals and birds that are known to be present in the vicinity of the Project and are known to be of value to Indigenous Groups and local communities. Documented rationale for receptor selection is provided in TSD XXI, Section 6.1.1, Selected Ecological Receptors. The terrestrial ecological receptors considered in the IMPACT model include terrestrial plants, terrestrial invertebrates, terrestrial and riparian mammals, and birds, including waterfowl.

Terrestrial Vegetation

Terrestrial vegetation types are dietary components for terrestrial animals and humans. They are represented by browse (shrubs and grasses), lichen (*Cladonia* spp., *Cladina* spp.), blueberry (*Vaccinium mytilloides*), and Labrador tea (*Ledum groenlandicum*). Plants are exposed to constituents in soil through contact with soil and contaminant uptake from soil via bioaccumulation. Lichen is exposed to constituents depositing from air. Berries and Labrador tea represent plants used in the Traditional Foods diet (i.e., diet made up of plants and animals fished, hunted, or gathered from the land).

Terrestrial Invertebrates

Terrestrial invertebrates are considered to be primary consumers. They are represented by earthworms, which live in soil and are therefore exposed to constituents in soil layers through direct contact. Earthworms acquire nutrition through organic matter in soil and decomposing remains of other animals. Terrestrial invertebrates are part of the diet for the red fox (*Vulpes vulpes*).

Terrestrial and Riparian Mammals

Mammalian herbivores are represented by woodland caribou, snowshoe hare (*Lepus americanus*), moose, southern red-backed vole (*Myodes gapperi*), and beaver (*Castor canadensis*). Herbivores convert vegetable matter to animal protein, and in turn are consumed by omnivores and carnivores. Snowshoe hare and southern red-backed vole are small mammals that consume berries and browse. These species are important prey for larger terrestrial predators such as red fox and grey wolf (*Canis lupus*). The southern red-backed vole is abundant in the vicinity of the Project (Annex VIII.1, Wildlife Baseline Report 1 [Mammals, Waterfowl, and Raptors]). Woodland caribou and moose are large ungulates with distinct home ranges and linkages to aquatic environments during the summer period. The primary food sources for Woodland caribou are lichen in the winter and terrestrial vegetation (browse) and aquatic vegetation (macrophytes) in the summer. Moose consume terrestrial and aquatic vegetation. Moose is part of the Traditional Foods diet and is assumed to be representative for other ungulate species such as white-tailed deer (*Odocoileus virginianus*).

Black bear (*Ursus americanus*) and red fox are terrestrial omnivores that are opportunistic and rely on readily available and easily accessed foods. Black bears consume berries and fish, and therefore have a dietary link to the aquatic environment. Red foxes consume a combination of terrestrial mammals and birds, terrestrial invertebrates, and berries.

Little brown myotis (the bat species *Myotis lucifugus*) represents terrestrial mammals that have a primarily insectivorous diet. In the IMPACT model, the diet of little brown myotis is composed of benthic invertebrates, representing the aquatic larval life stage of aerial insects. Although this diet originates from a water polygon, little brown myotis have negligible sediment intake exposures, because they are assumed to feed aerially.

Beaver and muskrat (*Ondatra zibethicus*) are riparian mammals that are exposed to constituents in both aquatic and terrestrial environments. Beavers are herbivores, with a major component of their diet composed of terrestrial vegetation and a minor component as aquatic vegetation. Beavers are also a component of the Traditional Foods diet. The muskrat's diet is fully aquatic, and muskrats are assumed to consume a combination of aquatic plants and benthic invertebrates.

Terrestrial predators representing the top level of the food chain include grey wolf and mink (*Neovison vison*). Grey wolf consumes terrestrial herbivores including moose, caribou, and hare. The mink's diet has linkages to both land and water polygons through the consumption of fish, benthic invertebrates, riparian mammals, and birds.

Terrestrial and Riparian Birds

Grouse (*Tetraoninae* sp.) and Canada goose (*Branta canadensis*) are terrestrial herbivores. Grouse is part of the Traditional Foods diet and is assumed to be representative for other birds with similar dietary characteristics, such as the ptarmigan.

Rusty blackbird (*Euphagus carolinus*) has a varied diet but feeds primarily on aquatic invertebrates. The rusty blackbird is a Species of Special Concern under the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) and the *Species at Risk Act*, and suitable habitat for this species is common and widespread (Annex VIII.2, Wildlife Baseline Report 2 [Amphibians, Birds, and Bats]). Rusty blackbird can be considered representative for other insectivorous birds that have similar diets and exposure pathways.

Mallard (*Anas platyrhynchos*) and common loon (*Gavia immer*) are migratory water birds that are common to northern Saskatchewan. These riparian birds have a primarily aquatic diet; the mallard consumes macrophytes and benthic invertebrates, and the loon consumes fish and benthic invertebrates.

2.3.2 Home Ranges and Residency Factors

Home range is defined as the geographic area encompassed by an animal's activities, excluding migration, over a specific time (USEPA 1993). The home range size is used to determine the portion of time that an individual animal is expected to be exposed to COPCs through contact with environmental media. Small mammals such as muskrat and vole have small home ranges that are represented by small polygons with similar vegetation communities.

The residency factor represents the fraction of time an animal is expected to be exposed to a location and the proportion of the animal's intake (i.e., food, water) that is taken from each location. Animals are given a residency factor of one if they receive 100% of their exposures from one polygon location. The IMPACT model looked at four polygon locations where potential exposures to Project emissions could occur (Patterson Lake North Arm – West Basin, Patterson Lake South Arm, Beet Lake, Lloyd Lake) and one reference location that is unexposed to Project emissions (Reference Lake represented by Broach Lake). Each polygon location includes both

water and land polygons, as animals will be exposed by both aquatic (drinking) as well as terrestrial pathways at each location.

Animals with large home ranges (i.e., greater than 10 km²), such as the grey wolf, woodland caribou, and black bear, interact with a number of different exposed polygon locations for feeding and water intake. All other receptors have relatively small home ranges (i.e., less than 10 km²) and were assumed to reside in the same polygon year-round. The estimated home ranges of selected mammalian receptors with potentially larger home ranges are presented in Table 2-2. The grey wolf has a home range of approximately 1,300 km², while the woodland caribou and black bear have home ranges of approximately 80 km² and therefore have the potential to be present at more than one polygon location to obtain their food. Home ranges are generally conservatively selected based on the expected home range during sensitive life stages.

Animals that are migratory are unlikely to have 100% residency at one polygon location for the entire year. However, because toxicity reference values that are used to establish potential risk are typically based on sensitive life stages such as reproduction, a 100% residency can be conservatively assumed for a specific life stage to capture the risk from exposure for the life stage of interest. For instance, mallards are not present in the vicinity of the Project for over half the year due to migration but have a small home range during the nesting and rearing of their young. Therefore, it is conservatively assumed that the mallard has a 100% residency factor at one polygon location where the young are hatched and reared.

Similarly, moose can move to different geographical locations between seasons, but generally remain within small seasonal home ranges of 5 km² to 10 km² during the summer and winter. For modelling purposes, these animals (with local home ranges or migratory patterns) are associated with one polygon location (100% residency).

Residency factors for receptors with large home ranges (i.e., caribou, bear, and grey wolf) were estimated based on the relative sizes of waterbodies (surface areas) that are located within an area consistent with the receptor's expected home range size. The objective is to determine the time spent at exposure locations and reference locations.

Table 2-2: Estimated Home Ranges of Selected Terrestrial Ecological Receptors

Ecological	Н	ome Range (k	m²)	Community	C
Receptor	Expected	Minimum	Maximum	Comments	Source
Woodland caribou	Female home ranges from GPS tracking data in the Saskatchewan Boreal Shield. The expected home range used for assessment is based on an average female range during the most sensitive life stage of the woodland caribou (i.e., calving/post-calving).				(McLoughlin et al. 2016)
Moose	10	5	10	Moose are essentially solitary and have one or several distinct home ranges to which they are strongly attached. Movements between seasonal home ranges may be extensive but home ranges are generally small. The same home ranges are generally returned to year after year.	(Franzmann 1981; Wilson and Ruff 1999)
Black bear	80	40	130	Female home ranges from GPS tracking data in the Saskatchewan Boreal Shield.	(McLoughlin et al. 2019)
Red fox	4	2.8	34	Red foxes tend to concentrate their activities near dens, preferred hunting areas with abundant food supplies, although young fox may travel great distances in search of new territory.	(Hinterland Who's Who 1993; Jones and Theberge 1981)
Grey wolf	y wolf 1,300 100 2,451 Home ranges for wolves in Minnesota, Alaska, southern Yukon, and British Columbia.		(RIC 1998)		
Mink	7.7 0.06 16.3 Expected value based on a study in southern Manitoba.			(Arnold and Fritzell 1987; FCSAP 2012)	

The caribou or bear receptor at Patterson Lake (Table 2-3) was assumed to spend time at Patterson Lake North Arm – West Basin and Patterson Lake South Arm (i.e., exposed locations). The caribou or bear receptor at either Patterson Lake location is likely to overlap with the other due to the size of their expected home range of 80 km² and therefore exposures to these locations were assessed by one caribou or bear visiting both locations. It was also assumed that the caribou or bear spends a proportion of time at unexposed locations within its home range (i.e., lakes upstream of future water discharge points). One small lake northeast of Patterson Lake North Arm – West Basin (~0.59 km²) was included as an unexposed location for the Patterson Lake caribou or bear receptor. As a result, the Patterson Lake caribou or bear receptor was assumed to spend 43% of its time at Patterson Lake North Arm – West Basin (12.56 km²), 55% of its time at Patterson Lake South Arm (15.95 km²), and 2% of its time at an unexposed location (0.59 km²).

Similarly, a caribou or bear living near Beet Lake and within areas south of Beet Lake was considered. Within the 80 km² range, the Beet Lake caribou or bear receptor was assumed to spend 48% of its time at exposed locations and 52% of its time at unexposed locations. The exposed locations comprise Forrest Lake – North Basin (1.47 km²), which is hydrologically connected to discharges via Patterson Lake and Beet Lake (8.84 km²). Because food sources for the caribou or bear receptor have not been modelled at Forrest Lake, the Forrest Lake caribou or bear receptor was assumed to obtain their food from Beet Lake and the land adjacent to Beet Lake. Unexposed locations include a portion of Forrest Lake – South Basin (7.54 km²) and other smaller, isolated lakes (0.78 km² and 2.87 km²) within the 80 km² range that are located to the east of Forrest Lake and South of Beet Lake.

Due to its large home range of 1,300 km², the grey wolf located near Patterson Lake was also assumed to spend only a portion of its time at exposed locations. The exposed locations include Patterson Lake North Arm – West Basin and East Basin (21.84 km² combined), Patterson Lake South Arm (15.95 km²), and Beet Lake, Forrest Lake – North Basin, and Naomi Lake (12.76 km² combined). For waterbody locations where food sources have not been modelled (i.e., Patterson Lake North Arm - East Basin), the grey wolf was assumed to obtain food from the nearest modelled polygon (i.e., Patterson Lake North Arm – West Basin). The unexposed locations include Broach Lake; Lake C, Lake E, Lake G, Lake H, Unnamed Lake 1 (near Dennis Lake), and Unnamed Lake 2 (Near Naomi Lake). These locations are included in the hydrology report (Annex IV.2). The sum of the surface areas of these waterbodies was used to determine the proportion of time the grey wolf near Patterson Lake spends at the reference (unexposed) locations, adding up to 55.64 km². Therefore, the grey wolf located near Patterson Lake was assumed to spend 47.6% of its time at exposed locations and 52.4% of its time at unexposed locations. Since the unexposed areas only include a fraction of the total surface area of unexposed waterbodies in the area, this approach likely overestimates the relative amount of time spent at exposed locations and therefore provides a conservative estimate of risk.

Due to the large size of Lloyd Lake relative to the terrestrial ecological receptors' home ranges, and its greater distance from the Project, the wolf, caribou, and bear at this location were assumed to have 100% residency at Lloyd Lake (i.e., residency factor of 1).

Table 2-3 presents the residency factors that were applied in the model for each terrestrial ecological receptor. The exposed lakes part of the LSA and RSA are shown on Figure 1-1.

Table 2-3: Residency Factors for Terrestrial Ecological Receptors

Terrestrial Ecological Receptor	Polygon	Surface Area of Waterbody (ha)	Total Waterbody (ha)	Residency Factor
Woodland caribou at Reference Lake	Reference Lake	-	-	1
Woodland caribou at Patterson Lake	Reference Lake Patterson Lake North Arm – West Basin Patterson Lake South Arm	59 1,256 1,595	2,910	0.02 0.43 0.55
Woodland caribou at Beet Lake	Reference Lake Beet Lake ^(a)	1,119 1,031 ^(a)	2,150	0.52 0.48
Woodland caribou at Lloyd Lake	Lloyd Lake	-	-	1
Snowshoe hare	Reference Lake Patterson Lake North Arm – West Basin Patterson Lake South Arm Beet Lake Lloyd Lake	- - - -	- - - -	1 1 1 1
Moose	Reference Lake Patterson Lake North Arm – West Basin Patterson Lake South Arm Beet Lake Lloyd Lake	- - - -	- - - -	1 1 1 1
Little brown myotis	Reference Lake Patterson Lake North Arm – West Basin Patterson Lake South Arm Beet Lake Lloyd Lake	- - - -	- - - -	1 1 1 1
Southern red- backed vole	Reference Lake Patterson Lake North Arm – West Basin Patterson Lake South Arm Beet Lake Lloyd Lake	- - - -	- - - -	1 1 1 1
Black bear at Reference Lake	Reference Lake	-	-	1
Black bear at Patterson Lake	Reference Lake Patterson Lake North Arm – West Basin Patterson Lake South Arm	59 1,256 1,595	2,910	0.02 0.43 0.55
Black bear at Beet Lake	Reference Lake Beet Lake ^(a)	1,119 1,031 ^(a)	2,150	0.52 0.48
Black bear at Lloyd Lake	Lloyd Lake	-	-	1
Red fox	Reference Lake Patterson Lake North Arm – West Basin Patterson Lake South Arm Beet Lake Lloyd Lake	- - - -	- - - -	1 1 1 1

Terrestrial Ecological Receptor	Polygon	Surface Area of Waterbody (ha)	Total Waterbody (ha)	Residency Factor
Grey wolf at Reference Lake	Reference Lake	-	-	1
Grey wolf at Patterson Lake	Reference Lake ^(b) Patterson Lake North Arm – West Basin ^(c) Patterson Lake South Arm Beet Lake ^(d)	5,564 ^(b) 2,184 ^(c) 1,595 1,276 ^(d)	10,619	0.524 0.206 0.150 0.120
Grey wolf at Lloyd Lake	Lloyd Lake	-	-	1
Beaver	Reference Lake Patterson Lake North Arm – West Basin Patterson Lake South Arm Beet Lake Lloyd Lake	- - - -	- - - -	1 1 1 1
Muskrat	Reference Lake Patterson Lake North Arm – West Basin Patterson Lake South Arm Beet Lake Lloyd Lake	- - - -	- - - -	1 1 1 1
Mink	Reference Lake Patterson Lake North Arm – West Basin Patterson Lake South Arm Beet Lake Lloyd Lake	- - - -	- - - -	1 1 1 1
Rusty blackbird	Reference Lake Patterson Lake North Arm – West Basin Patterson Lake South Arm Beet Lake Lloyd Lake	- - - -	- - - -	1 1 1 1
Canada goose	Reference Lake Patterson Lake North Arm – West Basin Patterson Lake South Arm Beet Lake Lloyd Lake	- - - -	- - - -	1 1 1 1
Grouse	Reference Lake Patterson Lake North Arm – West Basin Patterson Lake South Arm Beet Lake Lloyd Lake	- - - -	- - - -	1 1 1 1
Mallard	Reference Lake Patterson Lake North Arm – West Basin Patterson Lake South Arm Beet Lake Lloyd Lake	- - - -	- - - -	1 1 1 1

Terrestrial Ecological Receptor	Polygon	Surface Area of Waterbody (ha)	Total Waterbody (ha)	Residency Factor
	Reference Lake	-	-	1
	Patterson Lake North Arm – West Basin	-	-	1
Common loon	Patterson Lake South Arm	-	-	1
	Beet Lake	-	-	1
	Lloyd Lake	-	-	1

a) Includes Forrest Lake - North Basin.

2.3.3 Exposure Assumptions for Terrestrial Ecological Receptors

2.3.3.1 Body Weight and Intake Rates

The body weight and intake rates are required for the calculation of exposure to birds and mammals (Table 2-4). Body weights and intake rates were obtained, in order of preference, from CSA N288.1-20, the US Environmental Protection Agency (USEPA) Wildlife Exposure Factors Handbook (USEPA 1993), the Federal Contaminated Sites Action Plan Module 3: Standardization of Wildlife Receptor Characteristics (FCSAP 2012), and from Oak Ridge National Laboratory (Sample and Suter 1994). For species not represented in the above sources, additional sources as identified in Table 2-4 were consulted to identify representative body weights, and feed intake rates were calculated using allometric equations from the USEPA (1993).

Water intake and inhalation rates were determined using allometric equations from the USEPA (1993) for birds and mammals. The incidental ingestion of soil and/or sediment was estimated from feed intake. Incidental ingestion varied from 2% to 10.4% of food intake as dry weight, depending on the biota (Beyer et al. 1994). However, no incidental soil or sediment ingestion was assumed for the little brown myotis, which is an aerial insectivore; nor was any sediment ingestion assumed for the common loon, which feeds from the water column.

b) Includes Broach Lake; Lake C, Lake E, Lake G, Lake H, Unnamed Lake 1 (near Dennis Lake), and Unnamed Lake 2 (near Naomi Lake); and Forrest Lake – South Basin.

c) Includes Patterson Lake North Arm - East Basin.

d) Includes Forrest Lake - North Basin and Naomi Lake.

^{- =} Not required for the determination of residency factor; the receptor resides at a single location.

Table 2-4: Bird and Mammal Body Weights and Intake Rates

Receptor	Body	Weight		Total Fe	ed Intake		Feed Type Dietary Components Fraction		Feed Int	Feed Intake Rate Moist		Intake of Soil ^(r)	Intake of Sediment ^(r)	Intake of Soil and Sediment ^(k)	Basis of the Soil and Sediment Intake Value	Total Soil/ Sediment Intake Rate ⁽¹⁾	Water Intake Rate ^(m)	Inhalati	ion Rate	
	kg	Source	kg dw/d	Source	kg fw/d	Source		fw	dw	kg dw/d	kg fw/d	Unitless	%	%	%		kg dw/d	L/d	m³/d	Source
Woodland caribou	180	е	3.82	m	12.35	n/a	Browse	0.5	0.32	1.24	6.18	0.80	5.15	1.65	6.8	Bison	0.3	10.6	34.8	m
							Lichen	0.2	0.43	1.66	2.47	0.33								
							Macrophytes	0.3	0.24	0.927	3.71	0.75		_						
Snowshoe hare	1.8	С	0.110	c	0.564	n/a	Browse	0.9	0.92	0.102	0.508	0.80	3.7	0	3.7	Average of small	0.004	0.168	0.9	c,m
							Berries (blueberry)	0.1	0.08	0.008	0.056	0.85				mammals				
Moose	400	b	8.0	b	38.1	n/a	Browse	0.8	0.76	6.10	30.5	0.80	1.5	0.5	2.0	Moose	0.2	21.8	65.9	m
							Macrophytes	0.2	0.24	1.90	7.62	0.75								
Little brown myotis	0.0075	d	0.001	n/a	0.0037	d	Benthic invertebrates	1	1	0.0009	0.0037	0.75	0	0	0	Assumed negligible (aerial)	0	0.00121	0.011	m
Southern red-	0.0329	a, j	0.00196	n/a	0.012	a, j	Berries (blueberry)	0.6	0.53	0.001	0.007	0.85	2.4	0	2.4	Meadow vole	0.000047	0.00458	0.048	а
backed vole							Browse	0.4	0.47	0.001	0.005	0.80								
Black bear	102.5	0	3.1	b	17.1	n/a	Berries (blueberry)	0.7	0.58	1.79	12.0	0.85	2.9	0	2.9	Average of	0.09	6.39	22.2	m
							Fish (northern pike)	0.3	0.42	1.28	5.13	0.75				large mammals				
Red fox	4.54	a	0.105	n/a	0.429	a	Small mammals (vole)	0.4	0.490	0.052	0.172	0.70	2.8	0	2.8	Red fox	0.00295	0.386	1.83	m
							Terrestrial birds (goose)	0.2	0.245	0.026	0.086	0.70								
							Terrestrial invertebrates (earthworm)	0.25	0.173	0.018	0.107	0.83								
							Berries (blueberry)	0.15	0.092	0.010	0.064	0.85								
Grey wolf	45	f	1.57	m	5.23	n/a	Moose	0.45	0.45	0.707	2.356	0.70	2.8	0	2.8	Red fox	0.04	3.04	11.5	m
							Caribou	0.45	0.45	0.707	2.356	0.70								
							Small mammal (hare)	0.1	0.10	0.157	0.523	0.70								
Beaver	24	р	0.94	m	4.57	n/a	Macrophytes	0.1	0.12	0.114	0.457	0.75	1.8	0.24	2.0	Moose	0.019	1.729	6.9	m
							Browse	0.9	0.88	0.822	4.112	0.80								
Muskrat	1.17	a	0.088	n/a	0.352	a	Macrophytes	0.8	0.8	0.070	0.282	0.75	0	3.3	3.3	Mallard	0.003	0.114	0.59	а
							Benthic invertebrates	0.2	0.2	0.018	0.070	0.75								
Mink	1.02	a	0.045	n/a	0.161	а	Fish (northern pike)	0.3	0.270	0.012	0.048	0.75	2.6	0.41	3.05	Average of	0.001	0.101	0.44	а
							Benthic invertebrates	0.15	0.135	0.006	0.024	0.75				mallard and				
							Small mammals (muskrat) Birds (mallard)	0.45 0.1	0.486 0.108	0.022 0.005	0.072 0.016	0.70 0.70				red fox				
Rusty blackbird	0.064	g	0.0177	n/a	0.077	a,i	Benthic invertebrates	0.8	0.87	0.015	0.062	0.75	10.4	0	10.4	American	0.002	0.00935	0.148	m,n
							Berries (blueberry)	0.2	0.13	0.002	0.015	0.85				woodcock				
Canada goose	3.70	С	0.0229	n/a	0.115	a	Browse	1	1	0.023	0.115	0.80	8.2	0	8.2	Canada goose	0.0019	0.142	1.100	С
Grouse	0.6	b	0.042	b	0.227	n/a	Browse	0.7	0.76	0.032	0.159	0.80	8.2	0	8.2	Canada	0.0034	0.0419	0.276	m
							Berries (blueberry)	0.3	0.24	0.010	0.068	0.85				goose				

Table 2-4: Bird and Mammal Body Weights and Intake Rates

Receptor	Body \	Veight		Total Fee	ed Intake		Dietary Components	Feed Type Fraction		Feed Intake Rate		Feed Intake Rate Moisture ^(h)		Intake of Sediment ^(r)	Intake of Soil and Sediment ^(k)	Soil and	Sediment Intake Rate ^(l)	Water Intake Rate ^(m)	Inhalation Ra	
	kg	Source	kg dw/d	Source	kg fw/d	Source		fw	dw	kg dw/d	kg fw/d	Unitless	%	%	%		kg dw/d	L/d	m³/d	Source
Mallard	1.13	С	0.06	С	0.240	n/a	Macrophytes	0.25	0.25	0.015	0.060	0.75	0	3.3	3.3	Mallard	0.002	0.0640	0.02	С
							Benthic invertebrates	0.75	0.75	0.045	0.180	0.75	İ							
Common loon	5.3	b	0.159	b	0.636	n/a	Fish (northern pike)	0.9	0.9	0.143	0.572	0.75	0	0	0	Assumed	0	0.180	1.477	m
							Benthic invertebrates	0.1	0.1	0.016	0.064	0.75				negligible (sediment is washed off in water column)				

- a) (USEPA 1993) Body weights, ingestion rates and inhalation rates of adults or all groups (adults and juveniles) are an average of the listed values. If only a range is given, the upper limit of the range is used. Values for the southern red-backed vole were based on the meadow vole.
- b) (FCSAP 2012).
- c) CSA N288.1-20 (CSA Group 2020).
- d) (Sample and Suter 1994).
- e) COSEWIC Assessment and Update Status Report on the Woodland Caribou (COSEWIC 2002). Body weight calculated as a mean of the male and female upper range.
- f) (Earthroots 2020).
- g) COSEWIC Assessment and Update Status Report on the Rusty Blackbird (COSEWIC 2006).
- h) Sources of moisture/dry weight fraction values obtained from CSA N288.1-20 (CSA Group 2020) for all receptors except for earthworm (Beresford et al. 2008); and blueberries and lichen (Annex VII.3, Vegetation Chemistry Characterization Report). The blueberry value is based on the fruit only.
- i) The total feed intake for the American robin was used in the absence of a species-specific value.
- j) The body weight and total feed intake for the meadow vole was used in the absence of a species-specific value.
- k) (Beyer et al. 1994).
- l) Intake of Soil & Sediment (kg dw/d) = Total Feed Intake (kg dw/d) x Intake of Soil & Sediment (%)/100.
- m) Calculated using allometric equations in (USEPA 1993).
- n) Inhalation rate was multiplied by a factor of 3 for passerine birds as per (USEPA 1993).
- o) Average of males and females from Hinterland Who's Who (2007).
- p) Median of adult weight range of 16 to 32 kg from Hinterland Who's Who (2017).
- q^{\prime} For purposes of modelling the earthworm is counted to the terrestrial plants.
- r) The % intake of soil or sediment is calculated from the combined % intake of Soil & Sediment (%) x Feed Type Fraction Terrestrial. Intake of Sediment (%) = Total Intake of Soil & Sediment (%) x Feed Type Fraction Terrestrial. Intake of Sediment (%) = Total Intake of Soil & Sediment (%) x Feed Type Fraction Terrestrial. Intake of Sediment (%) = Total Intake of Sediment (%) x Feed Type Fraction Terrestrial.
- fw = fresh weight; dw = dry weight; n/a = not applicable.

2.3.3.2 Occupancy Factors

An occupancy factor is defined as the fraction of time the receptor species spends in or on various media. The occupancy factors are based on the experience and judgment of the risk assessor and the known behaviour of the receptor. The occupancy factors used in the IMPACT model are presented in Table 2-5.

Aquatic and Riparian Biota	OF _a	OFs	OF _{SS}	OF _W	Terrestrial Biota	OF _a	OFs	OF _{SS}
Macrophytes	-	-	0.1	0.9	Lichen	ı	-	1
Phytoplankton	-	-	-	1	Blueberry	-	0.5	0.5
Zooplankton	-	-	-	1	Earthworm	-	1	-
Benthic invertebrates	-	1	-	-	Woodland caribou	1	-	1
Lake whitefish	-	-	0.5	0.5	Snowshoe hare	1	-	1
Northern pike	-	-	-	1	Moose	1	-	1
Muskrat	1	-	0.5	-	Little brown myotis	1	-	-
Beaver	1	-	0.5	-	Southern red-backed vole	1	-	1
American mink	1	-	0.5	-	Black bear	1	-	1
Mallard	1	-	0.5	-	Red fox	1	0.2	8.0
Common loon	1	-	0.5	-	Grey wolf	1	-	1
	•		•	•	Rusty blackbird	1	-	1
					Canada goose	1	-	1
					Grouse	1	-	1

Table 2-5: Receptor Occupancy Factors

 $OF_a = occupancy factor in air; OF_s = occupancy factor in soil/sediment; OF_{ss} = occupancy factor in soil/sediment surface; OF_w = occupancy factor in water; - = not applicable.$

2.3.4 Terrestrial Pathways

Terrestrial pathways include the transfer of COPCs between air, soil, terrestrial plants, and terrestrial animals. The specific equations that consider the concentrations of COPCs in soil, terrestrial plants, and terrestrial animals along each of these pathways are listed below.

Concentration in soil (C_{soil}):

$$\frac{dC_{soil}}{dt} = \frac{C_{air} \cdot (V_g + W_r \cdot P_{pt}) \cdot D_f}{\rho_b \cdot z_{soil}} + \lambda_{parent} \cdot C_{parent} + \frac{C_{irr} \cdot Q_{irr}}{\rho_b \cdot z_{soil}} - C_{soil} \cdot \left[\lambda_T + \frac{L \cdot f_{soil}}{\rho_b \cdot z_{soil}} + \frac{v_i \cdot (1 - f_{soil})}{z_{soil} \cdot \varphi}\right]$$

where:

C_{soil} = concentration in soil (Bg/kg dw, mg/kg dw)

C_{air} = concentration in air (becquerels per cubic metre (Bq/m³, mg/m³) C_{irr} = concentration of constituent in irrigation water (Bq/L, mq/L)

 C_{parent} = concentration of parent constituent (Bg/kg, mg/kg)

D_f = deposition fraction for airborne constituents (unitless)

f_{soil} = fraction of constituent that is sorbed to solids in soil (unitless)

φ = volumetric water content (unitless)

 λ_{parent} = first order decay constant for parent constituent (1/s)

 λ_T = sum of the rate constants for all significant loss processes (1/s)

L = erosion loss rate in soil block $(kg/m^2/s)$

 P_{pt} = precipitation rate (m/s) Q_{irr} = irrigation rate (L/m²/s) ρ_b = bulk density of soil (kg/L) v_i = infiltration rate (m/s)

 V_g = atmospheric deposition velocity (m/s) W_r = wet deposition washout ratio (unitless) Z_{soil} = thickness of soil mixing layer (m)

Table 2-6: Modelling Parameters for the Soil Model

Model Parameter		Value	Unit
Mixing depth (thickness of soil layer)	Z_{soil}	0.2	m
Dry bulk density	$ ho_{ m b}$	1.5	kg dw/L
Water content (porosity)	φ	0.1	unitless
Infiltration rate	Vi	1.205E-8	$m^3/m^2/s$
Erosion rate	L	1.5E-8	kg dw/m²/s
Precipitation rate	P _{pt}	800	mm/yr
Wet deposition washout ratio (metals)	W _r	5.5E+6	unitless
Dry Deposition velocity (metals)	V _g	0.0014	m/s

Note: All values for the soil model are the default parameters specified in CSA N288.1-20, assuming sandy soil.

Concentration in terrestrial plants (C_{tp}):

$$C_{tp} = C_{int} + C_{ext}$$

$$C_{int} = C_{soil} \cdot B_{v} + C_{air} \cdot T_{f}$$

$$\frac{dC_{ext}}{dt} = \frac{C_{air} \cdot (V_{g} + W_{r} \cdot P_{pt}) \cdot f_{a}}{Y} + \lambda_{parent} \cdot C_{parent} + \frac{C_{irr} \cdot Q_{irr} \cdot f_{irr}}{Y \cdot A} - C_{ext} \cdot [\lambda_{T} + \lambda_{e}]$$

Where:

A = Area of polygon (m²)

 B_v = bioaccumulation factor for soil-to-plant transfer (dw unitless) C_{tp} = concentration in terrestrial plants (Bq/kg fw, mg/kg dw)

C_{int} = internal concentration in plant from soil uptake (Bq/kg fw, mg/kg dw)

C_{ext} = external concentration in plant from dust and irrigation (Bq/kg fw, mg/kg fw)

C_{soil} = concentration in soil (Bq/kg dw, mg/kg dw) C_{air} = concentration in air (Bg/m³, mg/m³]

C_{irr} = concentration of constituent in irrigation water (Bq/L, mg/L)

C_{parent} = concentration of parent constituent (Bq/kq)

 f_{irr} = irrigation fraction retained on irrigation (unitless)

 λ_{Parent} = first order decay constant for parent constituent (1/s)

 λ_T = sum of the rate constants for all significant loss processes (1/s)

 λ_e = effective removal decay constant (1/s)

 P_{pt} = precipitation rate (m/s) Q_{irr} = irrigation rate (L/m²/s)

 T_f = air-to-plant transfer factor (m³/kg) V_g = atmospheric deposition velocity (m/s) W_r = wet deposition washout ratio (unitless)

Y = vegetation yield (kg dw/m^2)

Concentration in terrestrial animals (C_{tq}):

$$C_{ta} = C_{air} \cdot I_a \cdot F_{inh} + F_{ing}$$

$$\begin{bmatrix} I_w \cdot \sum (C_{wc} \cdot k_w) + I_s \cdot \sum (C_s \cdot k_{sed}) + I_{soil} \cdot \sum (C_{soil} \cdot k_{soil}) + I_{tp} \cdot \sum (C_{tp} \cdot k_{tp}) + I_{ta} \cdot \sum (C_{ta} \cdot k_{ta}) + I_{tp} \cdot \sum (C_{ap} \cdot k_{ap}) + I_{aa} \cdot \sum (C_{aa} \cdot k_{aa}) \end{bmatrix}$$

Where:

C_{ta} = concentration in ingested terrestrial animals (Bq/kg fw, mg/kg fw)

 C_{wc} = concentration in water column (Bq/L, mg/L)

C_s = concentration in surficial sediment (Bq/kg dw, mg/kg dw)

C_{soil} = concentration in soil (Bg/kg dw, mg/kg dw)

C_{tp} = concentration in ingested terrestrial plants (Bq/kg dw, mg/kg dw)
C_{ap} = concentration in ingested aquatic plants (Bq/kg fw, mg/kg fw)
C_{aa} = concentration in ingested aquatic animals (Bq/kg fw, mg/kg fw)

F_{inh} = inhalation transfer factor (d/kg fw) F_{inq} = ingestion transfer factor (d/kg fw)

 I_w = water ingestion rate (L/d) I_a = inhalation rate (m³/d)

 I_s = sediment ingestion rate (kg dw/d)

 I_{soil} = soil ingestion rate (kg dw/d)

 I_{tp} = terrestrial plants ingestion rate (kg dw/d) I_{ta} = terrestrial animals ingestion rate (kg fw/d) I_{ap} = aquatic plants ingestion rate (kg fw/d)

 I_{aa} = aquatic animals ingestion rate (kg fw/d)

 k_w = fraction of water intake from contaminated sources (unitless) k_{sed} = fraction of sediment intake from contaminated sources (unitless)

 k_{soil} = fraction of soil intake from contaminated sources (unitless)

k_{ta} = fraction of terrestrial animal intake from contaminated sources (unitless)
 k_{to} = fraction of terrestrial plant intake from contaminated sources (unitless)

 k_{aa} = fraction of aquatic animal intake from contaminated sources (unitless) k_{ap} = fraction of aquatic plant intake from contaminated sources (unitless)

2.3.5 Radiological Dose to Terrestrial Ecological Receptors

Radiological dose to a terrestrial receptor is the radiation energy absorbed due to radiation emissions from radionuclides present in the environment (air or soil/sediment) or in the tissues of the organism. Through comparison with dose benchmarks, the risk to the organism can be quantified. This model characterizes dose and risk from both background and project exposure.

The total dose to terrestrial animals from each radionuclide is:

$$D_{total} = D_{int} + D_{ext}$$

The dose of COPCs to terrestrial animals due to internal radioactivity (D_{int}) is estimated using the equation shown below.

$$D_{int} = C_t \cdot DC_{int}$$

The external dose to terrestrial animals from radioactivity in sediment and soil (D_{ext}) is:

$$\begin{array}{rcl} D_{ext} &= D_{ext,s} + D_{ext,ss} \\ D_{ext,s} &= C_{soil} \cdot DC_{ext,s} \cdot OF_{s} \\ D_{ext,ss} &= C_{ss} \cdot DC_{ext,ss} \cdot OF_{ss} \end{array}$$

The external dose from noble gases (naturally occurring gases with very low chemical reactivity [e.g., helium]) to terrestrial animals is calculated in IMPACT from outdoor air; however, there are no noble gases in the Rook I model applicable to ecological receptors, so the equation is not provided.

Where:

C_t = whole body tissue concentration (Bq/kg fw)

 C_{soil} = soil concentration (Bq/kg dw) C_{ss} = surface soil concentration (Bq/m²) D_{int} = internal radiation dose (μ Gy/h)

 DC_{int} = dose coefficient for radionuclide in tissue ([µGy/h]/[Bq/kq fw])

 $D_{ext,s}$ = external radiation dose in soil (μ Gy/h)

 $D_{ext.ss}$ = external radiation dose on soil surface ($\mu Gy/h$)

 $\begin{array}{lll} DC_{ext,s} & = & dose \ coefficient \ for \ radionuclide \ in \ soil \ ([\mu Gy/h]/[Bq/kg \ dw]) \\ DC_{ext,ss} & = & dose \ coefficient \ for \ radionuclide \ on \ soil \ surface([\mu Gy/h]/[Bq/m^2]) \end{array}$

 OF_s = fraction of time spent immersed in soil (unitless)

OF_{ss} = fraction of time spent on the soil (unitless)

For animals with large home ranges (Section 2.3.2, Home Ranges and Residency Factors), the total dose is the combined exposure from several locations (Table 2-3), where the residency factors are used as weighting factors to calculate the total dose.

2.3.6 Non-radiological Risk to Terrestrial Ecological Receptors

The IMPACT model provides a means of evaluating the potential effects on environmental components by comparing the non-radiological exposures of receptors to recognized benchmarks, usually a concentration or dose-based toxicity reference values in the same units, for each COPC. This measure of potential risk is known as an HQ. This model calculates the risk from both background and project exposure.

The HQ for terrestrial animals due to ingestion of non-radiological constituents is estimated as:

$$\text{HQ} \quad = \quad \frac{1}{\text{RD} \cdot \text{BM}} \cdot \begin{bmatrix} I_w \cdot \sum (C_{\text{wc}} \cdot k_w) + I_s \cdot \sum (C_s \cdot k_{\text{sed}}) + \\ I_{\text{soil}} \cdot \sum (C_{\text{soil}} \cdot k_{\text{soil}}) + I_{\text{tp}} \cdot \sum (C_{\text{tp}} \cdot k_{\text{tp}}) + I_{\text{ta}} \cdot \sum (C_{\text{ta}} \cdot k_{\text{ta}}) + \\ I_{\text{ap}} \cdot \sum (C_{\text{ap}} \cdot k_{\text{ap}}) + I_{\text{aa}} \cdot \sum (C_{\text{aa}} \cdot k_{\text{aa}}) \end{bmatrix}$$

Where:

ВМ body mass (kg) C_{ap} = concentration in ingested aquatic plants (mg/kg fw) C_{aa} concentration in ingested aquatic animals (mg/kg fw) = C_s = concentration in surficial sediment (mg/kg dw) Csoil concentration in soil (mg/kg dw) = concentration in ingested terrestrial animals (mg/kg fw) C_{ta} = C_{tp} concentration in ingested terrestrial plants (mg/kg dw) = C_{wc} = concentration in water column (mg/L) = aquatic animals ingestion rate (kg fw/d) I_{aa} aquatic plants ingestion rate (kg fw/d) I_{ap} = Is sediment ingestion rate (kg dw/d) = I_{soil} = soil ingestion rate (kg dw/d) = terrestrial plants ingestion rate (kg dw/d) I_{tp} terrestrial animals ingestion rate (kg fw/d) I_{ta} = I_w = water ingestion rate (L/d) fraction of aquatic animal intake from contaminated sources (unitless) k_{aa} = k_{ap} = fraction of aquatic plant intake from contaminated sources (unitless) k_{sed} fraction of sediment intake from contaminated sources (unitless) = fraction of soil intake from contaminated sources (unitless) k_{soil} = fraction of terrestrial animal intake from contaminated sources (unitless) k_{ta} = k_{tp} fraction of terrestrial plant intake from contaminated sources (unitless) fraction of water intake from contaminated sources (unitless) k_{w} = RD = reference toxic dose (mg/kg/d)

2.4 Exposure of Human Receptors

2.4.1 Exposure Assumptions for Human Receptors

The human receptor groups selected for the human health risk assessment are based on the current understanding of how people use the area in the vicinity of the Project and their potential for exposure to Project-related media concentrations during one or more Project phases (i.e., Construction, Operations, and Closure). The selected human receptor groups are a camp worker, subsistence harvester, seasonal resident, and permanent resident (as shown on Figure 2-2), with rationale for receptor selection provided in TSD XXI, Section 5.1.1, Human Receptor Selection and Characterization.

Adult and one-year-old receptors were used to assess potential risk to human health. The exposure factors for these two age groups are summarized in Table 2-7. The selection process and further characteristics of the human receptors are discussed in TSD XXI, Section 4.0, Model Integration and Evaluation of Sources.

Table 2-7: Summary of Receptor Characteristics for the Human Health Risk Assessment

Receptor Characteristic	Unit	Adult (a)	One-Year- Old ^(b)	Reference
Air inhalation (mean)	m³/yr	5,950	1,830	CSA N288.1-20; Table 19
Water ingestion (mean)	L/yr	379.6	98.92	CSA N288.1-20; Table 21
Soil ingestion (mean) (c)	kg dw/d	4.00×10^{-06}	6.10×10^{-05}	CSA N288.1-20; Table 20
Sediment ingestion (c)	kg dw/d	4.00×10^{-06}	6.10×10^{-05}	CSA N288.1-20; Table 20
Exposed surface area - water (whole body)	m ²	2.19	0.72	CSA N288.1-20; Table 22
Total food ingestion (d)	kg fw/yr	706	410	CSA N288.1-20; Annex G

a) Adult applies to both male and female receptors.

The primary exposure routes for human health include ingestion (i.e., food, water, incidental amounts of soil and/or sediment), inhalation (i.e., vapours and/or particulates), dermal absorption (non-radiological), and external exposure to radiation (radiological). The potential exposure pathways are expected to be the same for all human receptors assessed.

b) The one-year-old is equivalent to the CSA N288.1-20 age class "infant".

c) Incidental ingestion rates for soil and sediment are assumed to be equal.

d) The higher total food ingestion for the adult male (from CSA N288.1) is used to represent both male and female human receptors, which is a conservative assumption for females.

dw = dry weight; fw = fresh weight.




Figure 2-2: Human Receptor Locations for the Rook I Model

2.4.1.1 Dietary Assumptions

Each of the human health receptors described in Section 2.4.1, Exposure Assumptions for Human Receptors, consumes locally sourced Traditional Foods to an extent that is consistent with either an average or a high consumer. Where possible, the total of Traditional Foods derived was obtained from pertinent Traditional Foods surveys and engagement with Indigenous Groups and local communities. Traditional Foods are those animals and plants that are fished, hunted, or gathered from the land and consumed as food (Health Canada 2018). The remainder of the total food diet is assumed to be from store-bought foods (i.e., store foods). Store foods include grain and other agricultural products of a typical diet that are not available locally.

The camp worker is assumed to work and reside at the Rook I permanent camp for half of the year and does not consume any Traditional Foods while at the site. During the other times of the year, the camp worker is assumed to be a high Traditional Foods consumer. The subsistence harvester and the permanent camp resident are assumed to be high Traditional Foods consumers, and the seasonal resident or lodge operator is assumed to be an average Traditional Foods consumer.

Adult Diet

As stated in Table 2-7, the initial assumptions for ingestion rates and components of the total foods diet for the human health risk assessment were taken from CSA N288.1-20 Adult (Table G.9b – Central) total diet.

The human diet provided in CSA N288.1-20 was derived from the 2004 food basket survey results for an adult male. The CSA N288.1-20 human diet was selected over the Health Canada (2010a) human diet. Health Canada (2010a) references Richardson (1997), which used survey results from the late 1970s. The Richardson (1997) diet was also a combined adult male/female diet. The CSA N288.1-20 total adult diet is a smaller overall diet (706 kg/yr) than the Richardson diet (808 kg/yr) by about 100 kg/yr and is based on more recent data.

The total food diet is the sum of Traditional Foods and store foods. Its dietary components, expressed in kilograms per year (kg/yr) of food consumed, are shown in Table 2-8 for non-consumers, average consumers, and high consumers of Traditional Foods.

The store food intakes in each food category were estimated from the total diet intakes provided in CSA N288.1-20, minus the Traditional Foods intakes. If the Traditional Foods intakes were higher than the total intake for the same CSA N288.1-20 food category, the store food intake was set to zero, and then the values for other store food categories were reduced as needed to obtain total dietary intakes equal to those in CSA N288.1-20.

As a final step, the amount of Traditional Foods "berries" ingested was increased from approximately 5% of the fruit and berry component of the total CSA N288.1-20 diet to 10% by reducing the store food berry component value. This adjustment was made in response to comments from local Indigenous Groups during engagement who felt that they ingest more locally sourced berries than the diets that were presented to them suggest. After adjusting the Traditional Foods diet to account for a higher ingestion rate for berries, the overall Traditional Foods diet is greater than that estimated by the First Nations Food, Nutrition and Environment Study (Chan et al. 2018). This results in a precautionary approach to modelling and assessment.

One-Year-Old Diet

Annual food consumption rates for the "one-year-old" were calculated using "adult-to-one-year-old" ratios from CSA N288.1-20 for each of the selected food categories. Annual ingestion rates for individual food items were combined into relevant food categories and the adult-to-one-year-old ratios were determined for each of these food categories.



Surrogate Ecological Receptors Used in the IMPACT Model

The concentrations of COPCs in Traditional Foods used in the exposure assessment were estimated using the IMPACT model. Based on the descriptions of the different human receptor groups and their Traditional Foods diets, surrogate ecological receptors were selected for the IMPACT model to represent each Traditional Food category. The surrogate ecological receptor selected to represent a Traditional Food category was informed and modified based on feedback received from Joint Working Group meetings and engagement with regulators. Specifically, these engagement activities informed assumptions for fish species (e.g., inclusion of northern pike and lake whitefish in the exposure assessment), large mammals (e.g., preference for moose over other large mammals such as caribou, inclusion of moose organs as a specific Traditional Food category), and small mammals (e.g., beaver identified as of value and importance to the communities). The final selection of surrogate ecological receptors for the Traditional Foods diet is shown in Table 2-9.

The local intake fractions as implemented in IMPACT for the human diet are shown in Table 2-10, and are calculated as follows:

Local Intake Fraction (%) = (Traditional Annual Food Intake of a Food Type / Total Annual Food Intake in the Corresponding Food Category) * Residency Factor * 100%

Where:

Traditional Annual Food Intake of a Food Type = traditional annual ingestion rate of foods consumed within each relevant food category, as shown in Table 2-8 (kg/yr).

Total Annual Food Intake in the Corresponding Food Category = annual ingestion rate for traditional + store food in the food category, as shown in Table 2-8 (kg/yr).

Residency Factor = the exposure area or reference area residency factor as shown in Table 2-12 (unitless).



Table 2-8: Annual Food Intakes (kg/yr) for Food Items of the Human Receptors' Diets

Food Category	Non-Traditional Foods Consumer Diet ^(a)		Average Traditional Foods Consumer (b,c,d,e,f)				High Traditional Foods Consumer (b,c,d,e,f)			
	Adult	One-Year-Old	old Adult		One-Year-Old		Adult		One-Year-Old	
	Total Diet	Total Diet	Traditional Foods	Store Foods	Traditional Foods	Store Foods	Traditional Foods	Store Foods	Traditional Foods	Store Foods
	kg/yr	kg/yr	kg/yr kg/yr		kg/yr		kg/yr			
Milk	126	243	0	126	0	243	0	32.6	0	226
Fish	8.25	1.83	26.8	0	5.94	0	84.8	0	18.76	0
Northern pike	=	-	13.4	0	2.97	0	42.4	0	9.38	0
Lake whitefish	-	-	13.4	0	2.97	0	42.4	0	9.38	0
Meat - large mammals	67.9	8.18	6.52	24.1	0.785	2.90	20.6	0	2.48	0
Meat - large mammal organs	2.04	0.51	3.59	0	0.897	0	9.16	0	2.29	0
Meat - small mammals	0.230	0.02	17.4	0	1.66	0	56.9	0	5.42	0
Meat - birds and eggs	58.1	10.4	-	53.1	-	9.48	-	42.3	-	7.55
Aquatic bird	-	-	1.71	0	0.304	0	3.46	0	0.618	0
Terrestrial bird	-	-	3.29	0	0.587	0	12.3	0	2.19	0
Root vegetables	47.9	8.7	0	47.9	0	8.65	0	47.9	0	8.65
Other vegetables	297	82.5	0.844	296	0.225	82.3	2.45	294	0.681	81.9
Fruits and berries	99.4	54.6	9.94	89.5	5.46	49.2	9.94	89.5	5.46	49.2
Total diet - subtotal	-	-	70.1	636	15.9	395	200	506	37.9	373
Total diet ^(a)	706	410	706 411			706		411		

Values less than 0 are rounded to 3 significant figures.

^{- =} value not available, or not applicable.



Ref. 19-2574 APRIL 2024

a) Total diet is derived from CSA N288.1-20 central dietary intakes Table G.9b.

b) Traditional Foods annual consumption rates for adults are taken from the *First Nations Food, Nutrition and Environment Study* (Chan et al. 2018) Boreal Shield diet for Saskatchewan First Nations communities for either "average" or "high" Traditional Foods consumers (male) for the Traditional Foods diet.

c) Annual food consumption rates for "one-year-old" were calculated using "Adult to one-year-old" ratios (CSA N288.1 Adult (Central): one-year-old) derived for each food category. Annual consumption rates for individual food items were combined into relevant food categories and the ratios for each food categories were derived based on the total kg/yr for each group.

d) The store food intakes in each food category were estimated from the non-Traditional Foods intakes provided in CSA N288.1-20, minus the Traditional Foods intakes. If the Traditional Foods intakes were higher than the total intake for the food category, the store food intake was set to zero, and then the values for other store food categories were reduced as needed to obtain total dietary intakes equal to those in CSA N288.1-20.

e) The fractions of northern pike (predator species) and lake whitefish (forage species) were set at 50% each of the total fish diet based on input received from regulatory bodies to include both predator and forage species for a conservative assessment.

f) The fraction of Traditional Foods "fruits and berries" was increased from approximately 5% (Chan et al. 2018) to 10% of the total fruits and berries based on input received from local Indigenous communities. The total food diet was adjusted by reducing the store foods and increasing Traditional Foods fruits and Berries fractions, respectively.

Table 2-9: Surrogate Ecological Receptors Used in the Traditional Foods Diet

Traditional Food Category	Representative Ecological Receptor	Traditional Food Items	Rationale ^(a)
Fish			
Fish	Northern pike	Walleye Northern pike	 Predator species and forage species represent two ecological trophic levels Northern pike and lake whitefish are present in all lakes in the vicinity of the Project according to baseline studies
	Lake whitefish	Lake whitefish	 Important Traditional Food items for Boreal Shield and Boreal Plains ecozones Both are known to be fished in the vicinity of the Project
Game Meat/Orga	ins		
	Moose meat Moose meat	Moose meat	
Large mammal Moose organs	Moose kidney Moose liver Caribou kidney	 Large herbivore with linkages to the aquatic environment Generally present in the vicinity of the Project according to baseline studies Important Traditional Food items for Boreal Shield and Boreal Plains ecozones 	
Small mammal	Beaver	Beaver meat Rabbit meat	 Small herbivore with linkages to the aquatic environment Generally present in the vicinity of the Project according to baseline studies Important Traditional Food items for Boreal Shield ecozone
Birds		•	
Aquatic bird	Mallard	Mallard	 Omnivorous bird with linkages to the aquatic environment Seasonally present in the vicinity of the Project according to baseline studies Important Traditional Food items for Boreal Shield and Boreal Plains ecozones
Terrestrial bird	Grouse	Grouse Goose	 Upland herbivore bird Generally present in the vicinity of the Project according to baseline studies Important Traditional Food items for Boreal Shield and Boreal Plains ecozones
Berries/Plants		_	
Berries	Blueberry	Blueberry	 Generally present in the vicinity of the Project according to baseline studies Important Traditional Food items for Boreal Shield and Boreal Plains ecozones
Plants	Labrador tea	Mint Rat root	 Generally present in the vicinity of the Project according to baseline studies Important Traditional Food items for Boreal Shield and Boreal Plains ecozones

a) The selection of surrogate ecological receptors to represent Traditional Food items was informed by feedback during engagement activities.



Table 2-10: Local Intake Fractions for Human Diet

			Worker dult)		al Resident Year-Old)		Resident lult)	Subsistence Harvester (One-Year-Old)		Subsistence Harvester (Adult)		Permanent Resident (One- Year-Old)	Permanent Resident (Adult)
Food Category	Food Type	Local Intake Reference % ^(a)	Local Intake Exposure %	Local Intake Reference %	Local Intake Exposure %	Local Intake Reference %	Local Intake Exposure %	Local Intake Reference %	Local Intake Exposure %	Local Intake Reference %	Local Intake Exposure %	Local Intake Exposure %	Local Intake Exposure %
A quatic Animals	Northern pike	25.00	25.00	35.00	15.00	35.00	15.00	25.00	25.00	25.00	25.00	50.00	50.00
Aquatic Animals	Lake whitefish	25.00	25.00	35.00	15.00	35.00	15.00	25.00	25.00	25.00	25.00	50.00	50.00
	Store foods	92.21	-	98.94	-	95.14	-	96.63	-	83.17	-	96.63	83.17
	Moose	0.78	0.78	0.14	0.06	0.68	0.29	0.32	0.32	1.69	1.69	0.64	3.38
Townstrial Animals	Moose organs	0.35	0.35	0.16	0.07	0.38	0.16	0.30	0.30	0.75	0.75	0.59	1.50
Terrestrial Animals	Beaver	2.17	2.17	0.29	0.12	1.82	0.78	0.70	0.70	4.68	4.68	1.40	9.35
	Mallard	0.13	0.13	0.05	0.02	0.18	0.08	0.08	0.08	0.28	0.28	0.16	0.57
	Grouse	0.47	0.47	0.10	0.04	0.34	0.15	0.28	0.28	1.01	1.01	0.57	2.02
Torroctrial Dlants	Labrador tea	9.88	9.88	2.88	1.24	5.48	2.35	5.54	5.54	9.88	9.88	11.08	19.75
Terrestrial Plants	Fruits and berries	40.12	40.12	67.12	28.76	64.52	27.65	44.46	44.46	40.12	40.12	88.92	80.25

a) The % values were derived from the annual food intakes for components of the human receptors' diets as provided in Table 2-8 and the residency factors as provided in Table 2-12. Local Intake Fraction (%) = (Traditional Annual Food Intake of a Food Type / Total Annual Food Intake in the Corresponding Food Category) * Residency Factor * 100%.

^{- =} value not available, or not applicable.

The concentrations of COPCs in store foods (Table 2-11) used in the exposure assessment are consistent with other recent human health risk assessments conducted for uranium mine and mill projects in Saskatchewan. The proportions of different food items in store foods for the Hatchet Lake Band (CanNorth 2000) were used to derive weighted average concentrations of constituents in the store food diet of human receptors based on Health Canada data (2000 to 2011) for non-radiological constituents in foods for all Canadian cities (Health Canada 2011) and United Nations Scientific Committee on the Effects of Atomic Radiation (UNSCEAR 2000) for radiological constituents.

Table 2-11: Concentrations of Constituents of Potential Concern in Store Foods

Constituents of Potential Concern	Concentration (a)
Non-radionuclides (b)	mg/kg fw
Arsenic	0.0066
Cobalt	0.0055
Copper	0.57
Molybdenum	0.097
Uranium	0.0014
Radionuclides (c)	Bq/kg fw
Lead-210	0.039
Polonium-210	0.048
Radium-226	0.042
Thorium-230	0.0058
Uranium-234	0.0097
Uranium-238	0.0097

a) For the estimation of constituents of potential concern in store foods, proportions of food types in the store food diet were aligned with regional data from CanNorth (2000) for the Hatchet Lake First Nation.

2.4.2 Exposure Duration and Frequency

The residency assumptions for human receptors used for the human health risk assessment are summarized in Table 2-12. This includes the fraction of time the receptor spends at a given location as well as the exposure frequency (which refers to the frequency at which the activity causing the exposure occurs). The exposure duration (which refers to the length of time the receptor engages in the activity causing the exposure) was either the duration of the Project phases (43 years) for non-carcinogens, or the lifetime of the receptor for carcinogens.

b) Non-radionuclides in store foods were calculated from Health Canada data on trace elements in foods from the Total Diet Study (2000 to 2011) for all Canadian cities (Health Canada 2011).

c) Calculated from UNSCEAR (2000).

Bq = becquerel; fw = fresh weight.

Table 2-12: Summary of Residency Assumptions for Human Health Receptor Groups

Human Health Receptor	115 5(1) (2)	Builting (a)		Residency ^(a)	Exposure Frequency		
Group	Life Stage(s)	Residence(s)	Residency Factor	Fraction of Time at Location	(Months/Year)		
All Project Phases: Constr	ruction, Operations, Clo	osure			•		
Camp worker	Adult	Patterson Lake Camp/ Patterson Lake South Arm/ reference location	0.5 / 0.25 / 0.25	6 months/year 3 months/year 3 months/year	Yes, while away from the Camp		
		Patterson Lake South Arm / reference location	0.5 / 0.5	6 months/year & 6 months/year			
Subsistence harvester	Adult and one-year- old Beet Lake/ reference location		0.5 / 0.5	6 months/year & 6 months/year	Yes, year-round		
		Lloyd Lake/ reference location	0.5 / 0.5	6 months/year & 6 months/year			
Constant and and	Adult and one-year-	Patterson Lake South Arm / reference location	0.3 / 0.7	4 months/year & 8 months/year	Voc. while in the study area		
Seasonal resident	old	Lloyd Lake/ reference location	0.3 / 0.7	4 months/year & 8 months/year	Yes, while in the study area		
Far Future							
Permanent camp resident	Adult and one-year- old	Patterson Lake North Arm – West Basin	1	12 months/ year	Yes, year-round		
		Patterson Lake South Arm / reference location	0.5 / 0.5	6 months/year & 6 months/year			
Subsistence harvester	Adult and one-year- old	Beet Lake/ reference location	0.5 / 0.5	6 months/year & 6 months/year	Yes, year-round		
		Lloyd Lake/ reference location	0.5 / 0.5	6 months/year & 6 months/year			
Seasonal resident	Adult and one-year-	Patterson Lake South Arm / reference location	0.3 / 0.7	4 months/year & 8 months/year	Voc. while in the study area		
Seasonal resident	old	Lloyd Lake/ reference location	0.3 / 0.7	4 months/year & 8 months/year	Yes, while in the study area		

a) The exposure duration for subsistence harvesters, seasonal residents and permanent residents is considered to be over a lifetime; the exposure duration for the camp worker is considered to be the duration of the Project phases (43 years).



Ref. 19-2574 APRIL 2024

2.4.3 Radiological Dose

The dose to human receptors from all radioactive isotopes except radon-222 is estimated using the equations listed below.

Radiological dose due to ingestion (D_f):

$$D_{f} = DCF_{f} \left[\cdot I_{w} \cdot \sum C_{wc} \cdot k_{w} \cdot rf \cdot + I_{s} \cdot \sum C_{soil} \cdot EF_{soil} + I_{sed} \cdot \sum C_{s} \cdot EF_{s} \cdot DF_{s} + I_{tp} \cdot \sum C_{tp} \cdot k_{tp} \cdot \rho_{f} + I_{ta} \cdot \sum C_{ta} \cdot k_{ta} \cdot \rho_{f} + I_{ap} \cdot \sum C_{ap} \cdot k_{ap} \cdot \rho_{f} + I_{aa} \cdot \sum C_{aa} \cdot k_{aa} \cdot \rho_{f} \right]$$

Radiological dose due to external exposure to water (Dw):

$$D_w = C_{wc} \cdot [k_w \cdot OF_w + k_w^S \cdot rf \cdot OF_w^S + k_w^B \cdot rf \cdot D_c \cdot OF_w^B] \cdot DCF_w$$

Radiological dose due to external exposure to soil (D_a):

$$D_g = C_{soil} \cdot f_o \cdot f_r \cdot [f_u + (1 - f_u) \cdot S_g] \cdot DCF_g \cdot \rho_b \cdot z_{soil}$$

Radiological dose due to external exposure to shoreline sediment (D_s):

$$D_s = C_s \cdot OF_s \cdot W \cdot DF_s \cdot DCF_s$$

Radiological dose due to inhalation in air (D_i):

$$D_i = I_a \cdot OF_i \cdot C_a \cdot DCF_{inh}$$

Radiological dose due to immersion in air (Da):

$$D_a = f_0 \cdot [f_u + (1 - f_u) \cdot S_b] \cdot C_a \cdot DCF_a$$

Total dose (D_{total}) to a human receptor from all pathways of exposure:

$$D_{total} = D_f + D_w + D_g + D_s + D_i + D_a$$

where:

 $\begin{array}{lll} C_{aa} & = & \text{aquatic animal concentration (Bq/kg fw)} \\ C_{ap} & = & \text{aquatic plant concentration (Bq/kg fw)} \\ C_{s} & = & \text{sediment concentration (Bq/kg dw)} \\ C_{soil} & = & \text{surficial soil concentration (Bq/kg dw)} \\ C_{tp} & = & \text{terrestrial plant concentration (Bq/kg dw)} \\ C_{ta} & = & \text{terrestrial animal concentration (Bq/kg fw)} \\ \end{array}$

 C_{wc} = water column concentration (Bq/L)

 D_a = dose due to external exposure in air (mSv/yr)

 D_c = bathtub correction factor (unitless)

 D_f = dose due to ingestion (millisieverts per year (mSv/yr) D_g = dose due to external exposure to soil/sediment (mSv/yr)

 D_i = dose due to inhalation (mSv/yr)

 D_{w} dose due to external exposure in water (mSv/yr) D_{total} total dose (mSv/yr) = DCF_a = dose coefficient for external exposure in air ([Sv/vr]/[Bg/m³]) **DCF**_f dose coefficient for ingestion (sieverts per becauerel [Sv/Ba]) = DCF_{α} = dose coefficient for external exposure to soil ([Sv/yr]/[Bg/m²]) **DCF**_{inh} dose coefficient for inhalation (Sv/Bg) = DCF_s dose coefficient for external exposure to sediment ([Sv/yr]/[Bq/kq]) = DCF_w dose coefficient for external exposure in water ([Sv/yr]/[Bq/L]) = DFs dilution factor for shoreline deposits (unitless) = **EF**soil number of days per year in which soil ingestion occurs (d/yr) = EF_s number of days per year in which sediment ingestion occurs (d/yr) = f_{\circ} fraction of total time spent by the individual at the particular location (unitless) = f_r dose reduction factor for non-uniformity of ground surface (unitless) f_{u} = time spent outdoors at a particular location as a fraction of total time spent at that location (unitless) I_a inhalation rate (m³/yr) aquatic animals ingestion rate (kg fw/yr) = I_{aa} I_{ap} aquatic plants ingestion rate (kg fw/yr) = ls sediment ingestion rate (kg dw/d) = soil ingestion rate (kg dw/d) = I_{soil} terrestrial animal ingestion rate (kg fw/yr) I_{ta} = I_{tp} = terrestrial plant ingestion rate (kg dw/yr) = water indestion rate (L/vr) I_{w} fraction of air inhalation intake from contaminated source (unitless) k_a = fraction of aquatic animal intake from contaminated source (unitless) = fraction of aquatic plant intake from contaminated source (unitless) k_{ap} = $k_{\text{ta}} \\$ = fraction of terrestrial animal intake from contaminated source (unitless) fraction of terrestrial plant intake from contaminated source (unitless) = k_w = fraction of water intake from contaminated source (unitless) fraction of bathing in contaminated water (unitless) = k_{w}^{S} fraction of beach swimming in contaminated water (unitless) = OF: occupancy factor for inhalation (unitless) = OF_s occupancy factor for sediment (unitless) OFw occupancy factor for swimming in a surface water body (unitless) = OF_w^B occupancy factor in bath water (unitless) = OFw^S occupancy factor in swimming water (unitless) = dry bulk density of the soil (kg dry soil per m³) = ρ_b adjustment factor for food processing (unitless) ρf = dose reduction factor for water treatment (unitless) rf = S_h = building shielding factor (unitless) groundshine dose reduction factor for indoor shielding (unitless) S_{q} = W shore-width factor that describes the shoreline exposure geometry (unitless) =



=

Zsoil

soil mixing depth (m)

Dose from Radon

Consistent with recommendations in CSA N288.6-22 and Health Canada (2010b), the dose from radon in air is calculated according to the following equation:

$$Dose_{Rn} = \left[\frac{C_{Rn}}{3700 \frac{Bq}{m^3}} \times F \times \frac{ET}{170h} \right] \times 4mSv$$

Radon typically dissipates quickly outdoors and is usually not a human health issue. However, dose from exposure to radon indoors should be quantified. The outdoor equilibrium fraction is needed to estimate the indoor equilibrium fraction. The equilibrium fraction for radon in outdoor air was estimated as a function of the travel time from the radon source using guidance from Health Canada and the USEPA according to the following equation (Health Canada 2010b; USEPA 1986):

$$F_{out} = 1.0 - 0.0479e^{-\frac{t}{4.39}} - 2.1963e^{-\frac{t}{38.6}} + 1.2442e^{-\frac{t}{28.4}}$$

The equilibrium fraction for radon in indoor air (F_{in}) was estimated as a function of the outdoor equilibrium fraction (F_{out}) using guidance from Health Canada (2010b) and the USEPA (1986) according to the following equation:

$$F_{in} = 0.35(1 + F_{out})$$

where:

 C_{Rn} = concentration of radon in air (Bq/m³) $Dose_{Rn}$ = incremental dose from radon (mSv/yr)

ET = exposure time (h/yr)

F = degree of equilibrium between radon and decay products (unitless)

F_{in} = indoor equilibrium fraction (unitless) F_{out} = outdoors equilibrium fraction (unitless)

t = travel time from source to receptor (minutes) (distance from source to receptor divided

by mean wind speed)

2.4.4 Non-radiological Dose

The human receptor exposure to non-radiological COPCs is estimated from the following equations:

Non-radiological intake due to inhalation (D_i):

$$D_i = \frac{C_{air} \cdot I_a \cdot k_a \cdot OF_i}{BM}$$

Non-radiological dose due to ingestion (D_f):

$$D_{f} = \frac{1}{BM \cdot C} \cdot \left[I_{w} \cdot \sum (C_{wc} \cdot k_{w} \cdot rf) + I_{s} \right]$$

$$\cdot \sum (C_{s} \cdot EF_{s} \cdot DF_{s} \cdot RAF_{ing}) + I_{soil}$$

$$\cdot \sum (C_{soil} \cdot EF_{soil} \cdot RAF_{ing}) + I_{tp}$$

$$\cdot \sum (C_{tp} \cdot k_{tp} \cdot \rho_{f}) + I_{ta}$$

$$\cdot \sum (C_{ta} \cdot k_{ta} \cdot \rho_{f}) + I_{ap} \cdot \sum (C_{ap} \cdot k_{ap} \cdot \rho_{f}) + I_{aa} \cdot \sum (C_{aa} \cdot k_{aa} \cdot \rho_{f}) \right]$$

Non-radiological external exposure to water (dermal absorption) (D_w):

$$D_{w} = \frac{k_{p} \cdot SA_{w}}{BM} \cdot \left[C_{w}^{S} \cdot OF_{w}^{S} + C_{w}^{B} \cdot rf \cdot OF_{w}^{B} + C_{w}^{P} \cdot rf \cdot OF_{w}^{P}\right]$$

Non-radiological external exposure to soil and/or sediment (dermal absorption) ($D_{s,d}$):

$$D_{s,d} = \frac{[C_{soil} * (SA_{hands} * AF_{hands} + SA_{other} * AF_{other}) * RAF_{derm} * EF_{soil}]}{BM * C}$$

where:

 AF_{hands} soil adherence factor to hands (kg/m² d) AF_{other} soil adherence factor to skin other than hands (kg/m² d) ВМ body mass (kg) = C conversion factor (365 days per year) = C_{aa} aquatic animal concentration (mg/kg fw) C_{air} air concentration (mg/m³) C_{ap} = aquatic plant concentration (mg/kg fw) C_s sediment concentration (mg/kg dw) = C_{soil} = surficial soil concentration (mg/kg dw) C_{tp} = terrestrial plant concentration (mg/kg dw) terrestrial animal concentration (mg/kg fw) C_{ta} = C_w^s = water concentration at the beach (mg/L) C_{w}^{B} water concentration in the bath (mg/L) C_w^P water concentration in the pool (mg/L) = C_{wc} water column concentration (mg/L) = D_{f} dose due to ingestion (mg/kg d) Di = dose due to inhalation (mg/kg d) $D_{s,d}$ dose due to dermal contact with soil (mg/kg d) = days per year exposed to sediment (d/yr) EF_s days per year exposed to soil (d/yr) **EF**_{soil} I_a inhalation rate (m³/yr) = aquatic animals ingestion rate (kg fw/yr) I_{aa} aquatic plants ingestion rate (kg fw/yr) = I_{ap} ľ = sediment ingestion rate (kg dw/d)

 I_{soil} = soil ingestion rate (kg dw/d)

 I_{ta} = terrestrial animal ingestion rate (kg fw/yr) I_{tp} = terrestrial plant ingestion rate (kg dw/yr)

 I_w = water ingestion rate (L/yr)

k_a = fraction of air inhalation intake from contaminated source (unitless)
 k_{aa} = fraction of aquatic animal intake from contaminated source (unitless)
 k_{ap} = fraction of aquatic plant intake from contaminated source (unitless)

k_p = chemical specific permeability coefficient (cm/h)

k_{ta} = fraction of terrestrial animal intake from contaminated source (unitless)
 k_{tb} = fraction of terrestrial plant intake from contaminated source (unitless)

 OF_i = occupancy factor for inhalation (unitless) OF_W^B = occupancy factor in bath water (unitless) OF_W^P = occupancy factor in pool water (unitless) OF_W^S = occupancy factor at the beach (unitless)

 ρ_f = adjustment factor for food processing (unitless)

RAF_{derm} = relative absorption factor for soil dermal contact (unitless)

RAF_{ing} = relative absorption factor for ingestion (unitless) rf = dose reduction factor for water treatment (unitless)

 SA_w = surface area of exposed skin (m²)

 SA_{hands} = surface area of hands (m²)

 SA_{other} = surface area of exposed skin other than hands (m²)

Potential non-cancer effects are evaluated using HQ values estimated as described in Section 2.3.6, Non-radiological Risk to Terrestrial Ecological Receptors. Potential cancer risks from exposure to carcinogenic constituents are evaluated as incremental lifetime cancer risks.

3.0 Development of Model Parameters

3.1 Ambient Hydrology

A baseline sampling program is fundamental to the development and application of an ecological model for Environmental Assessment. In addition to data on environmental media, many other parameters are required to quantify the transport and fate of constituents in the environment. Many of those parameters are not typically measured and are therefore estimated in the modelling process.

3.1.1 Lake Morphometry

Waterbodies from Patterson Lake to the Lloyd Lake inlet were modelled to assess the effects of the Project on the downstream environment. Broach Lake, upstream of Patterson Lake, was modelled as a reference location. The lakes were modelled based on characteristics consistent with measured values as presented in Table 3-1. All values are based on the hydrological modelling inputs presented in Annex IV.2.

Table 3-1: Lake Morphometry for all Modelled Lakes in the IMPACT Model

Lake Modelled	Average Depth (m)	Area (km²)
Broach Lake	25.9	9.3
Patterson Lake North Arm – East Basin	7.0	9.3
Patterson Lake North Arm – West Basin	18.7	12.6
Patterson Lake South Arm	14.8	16.0
Beet Lake	10.7	8.9
Naomi Lake	1.8	2.4
Lloyd Lake inlet	3.8	6.4

The stream model in IMPACT was used to model rivers and streams to connect Broach Lake and Patterson Lake North Arm East Basin, and Naomi Lake and Lloyd Lake. The three basins of Patterson Lake were modelled as separate waterbodies: North Arm – East Basin, North Arm – West Basin, and South Arm. Forrest Lake was split into Forrest Lake – North Basin and Forrest Lake – South Basin, with the direction of flow from the South Basin to the North Basin. For the Forrest Lake – North Basin, the stream model in IMPACT was used. As such, the Forrest Lake – North Basin acts as a flow-through channel, and no sediment interaction was modelled.

3.1.2 Surface Water Flows

Watersheds and yields were implemented consistently with the hydrological baseline model (Appendix 9A, Hydrological Modelling Summary Report). Average monthly flows from the Hydrological Modelling Summary Report are shown in Table 3-2 at the model nodes, and Figure 3-1 shows the flow direction through the modelled environment as implemented in IMPACT. The outflow in IMPACT is calculated as the sum of the upstream flow entering a body of water and the local inflow, defined as any inflow from the contributing watershed and tributaries. The local inflow was calculated based on watershed yield and area. Based on this, inputs for the local inflows were calculated for input into the IMPACT model (Table 3-3).






Figure 3-1: Waterbodies and Flow Direction as Implemented in the IMPACT Model

Table 3-2: Average Monthly Outflows at the Model Nodes

						Outflow	(m³/s)						Average (m³/s)
Node Name	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	
Broach Lake	0.1	0.1	0.1	0.3	0.5	0.4	0.3	0.2	0.2	0.2	0.2	0.2	0.3
Clearwater River above Patterson Lake	0.2	0.2	0.3	0.8	1.2	0.9	0.7	0.6	0.6	0.6	0.5	0.3	0.6
Clearwater River below Patterson Lake	1.1	1.0	1.1	1.2	1.7	1.7	1.6	1.5	1.5	1.4	1.2	1.1	1.4
Clearwater River below Forrest Lake	1.9	1.7	1.7	1.8	2.7	2.8	2.6	2.4	2.3	2.3	2.3	2.1	2.2
Clearwater River below Beet Lake	2.0	1.8	1.8	2.1	3.3	3.0	2.7	2.6	2.5	2.5	2.4	2.2	2.4
Clearwater River below Naomi Lake	2.4	2.2	2.1	2.6	4.6	4.0	3.5	3.3	3.2	3.2	3.0	2.7	3.1
Clearwater River above Mirror River Confluence	3.5	3.5	3.5	6.0	10.0	8.0	6.6	6.0	6.0	6.0	5.4	4.0	5.7
Clearwater River below Mirror River Confluence	10.8	10.7	10.8	18.5	30.6	24.5	20.4	18.5	18.4	18.4	16.4	12.2	17.5
Clearwater River above Lloyd Lake	11.7	11.6	11.7	20.0	33.1	26.5	22.1	20.0	19.9	20.0	17.8	13.2	19.0
Clearwater River at Outlet of Lloyd Lake	17.6	16.1	15.5	19.4	34.8	33.2	28.4	24.2	22.6	23.5	21.3	19.5	23.0



Table 3-3: Mean Monthly Local Inflows for All Waterbodies Modelled in IMPACT

Laka Nawa		Mean Monthly Local Inflow (m³/s)											
Lake Name	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	m³/s
Broach Lake	0.13	0.12	0.13	0.29	0.54	0.43	0.31	0.24	0.23	0.23	0.21	0.17	0.25
Clearwater River above Patterson Lake	0.07	0.12	0.16	0.53	0.61	0.43	0.35	0.32	0.32	0.33	0.26	0.09	0.30
Patterson Lake North Arm – East Basin	0.23	0.21	0.20	0.11	0.13	0.23	0.25	0.25	0.23	0.22	0.20	0.23	0.21
Patterson Lake North Arm – West Basin	0.25	0.22	0.22	0.12	0.15	0.25	0.27	0.27	0.25	0.24	0.22	0.25	0.23
Patterson Lake South Arm	0.40	0.36	0.35	0.19	0.24	0.39	0.44	0.43	0.41	0.39	0.35	0.40	0.36
Forrest Lake – North Basin	0.80	0.71	0.60	0.61	1.00	1.07	1.00	0.90	0.88	0.93	1.05	0.94	0.87
Beet Lake	0.10	0.09	0.12	0.23	0.63	0.20	0.11	0.13	0.17	0.16	0.06	0.08	0.17
Naomi Lake	0.46	0.40	0.35	0.51	1.28	1.03	0.78	0.69	0.68	0.69	0.65	0.53	0.67
Clearwater River above Mirror River confluence	1.07	1.25	1.38	3.45	5.38	3.95	3.10	2.78	2.81	2.82	2.35	1.26	2.63
Clearwater River above Lloyd Lake	8.18	8.08	8.17	14.00	23.15	18.56	15.42	14.00	13.93	13.95	12.44	9.22	13.26
Lloyd Lake	5.22	5.22	1.61	1.61	4.14	4.14	5.27	5.27	3.11	3.11	4.90	4.90	4.04



3.2 Water Quality

3.2.1 Background Water Quality

To characterize regional background water quality, a database of baseline water quality data was compiled from baseline studies performed between 2018 and 2020 (Annex V.1). The waterbodies included in the calculation of a regional baseline were Patterson Lake (i.e., North Arm – East Basin, North Arm – West Basin, and South Arm), all measurement stations along the Clearwater River, Lake H, Lake G, Forrest Lake, Beet Lake, Naomi Lake, Mirror River, and Lloyd Lake. The geometric mean was selected as the baseline concentration for constituents that had measured data over the detection limit. In the case of constituents for which measured values in water were under the detection limit but sediment concentration measurements were over the detection limit, the baseline water concentration was calculated from the geometric mean of the sediment measurements using the partitioning coefficients (Kd; Section 3.3.2).

Lead-210 and its daughter polonium-210 were considered to be in transient equilibrium. Activity concentrations of a parent and daughter nuclide in transient equilibrium are governed by the following relationship:

$$Q_B = \frac{\lambda_B}{\lambda_B - \lambda_A} \cdot Q_A$$

where:

 Q_A = activity of the parent Q_B = activity of the daughter λ_A = half life of the parent λ_B = half life of the daughter

Polonium-210 and lead-210 were balanced to remain in transient equilibrium. Sediment concentration of Po-210 was calculated from measured values of water concentration using the K_d. The transient equilibrium assumption was used to calculate sediment concentrations of lead-210 from Po-210 sediment concentrations and, using the distribution coefficient (K_d), lead-210 sediment concentrations were used to model water concentrations of lead-210.

The inflow concentrations for each waterbody were assigned slightly higher values than the background concentrations since sedimentation processes remove metals and radionuclides from the inflow. Inflow concentrations were calculated from the background concentration, the residence time within each respective waterbody, the sedimentation rate, and the COPC-specific partitioning coefficient. The slightly higher inflow values were used in the model to predict stable concentrations in water and sediment within the range of observed values of background conditions and are presented in Table 3-4.

Table 3-4: Calibrated Local Inflow Concentrations for Each Water Polygon Modelled in IMPACT


СОРС	Water Baseline	Clearwater River upstream of Patterson Lake	Broach Lake	Patterson Lake North Arm East Basin	Patterson Lake North Arm West Basin	Patterson Lake South Arm	Forrest Lake North	Beet Lake	Naomi Lake	Clearwater river upstream of Mirror River Confluence	Clearwater river downstream of Mirror River Confluence	Lloyd Lake Inlet
	mg/L	mg/L	mg/L	mg/L	mg/L	mg/L	mg/L	mg/L	mg/L	mg/L	mg/L	mg/L
Arsenic	1.24×10^{-04}	1.24×10^{-04}	3.19×10^{-03}	3.85 × 10 ⁻⁰³	4.75 × 10 ⁻⁰³	3.79 × 10 ⁻⁰³	1.24×10^{-04}	4.37×10^{-03}	4.23×10^{-04}	1.24×10^{-04}	1.24×10^{-04}	2.55 × 10 ⁻⁰⁴
Chloride	4.22 × 10 ⁻⁰¹	4.22 × 10 ⁻⁰¹	4.22×10^{-01}	4.22 × 10 ⁻⁰¹	4.22 × 10 ⁻⁰¹	4.22 × 10 ⁻⁰¹	4.22×10^{-01}	4.22 × 10 ⁻⁰¹	4.22 × 10 ⁻⁰¹	4.22×10^{-01}	4.22 × 10 ⁻⁰¹	4.22 × 10 ⁻⁰¹
Cobalt	5.83 × 10 ⁻⁰⁴	5.83 × 10 ⁻⁰⁴	9.58 × 10 ⁻⁰⁴	1.04 × 10 ⁻⁰³	1.15 × 10 ⁻⁰³	1.03 × 10 ⁻⁰³	5.83 × 10 ⁻⁰⁴	1.10 × 10 ⁻⁰³	6.20 × 10 ⁻⁰⁴	5.83×10^{-04}	5.83 × 10 ⁻⁰⁴	5.99 × 10 ⁻⁰⁴
Copper	8.43 × 10 ⁻⁰⁴	8.43 × 10 ⁻⁰⁴	1.49 × 10 ⁻⁰³	1.63 × 10 ⁻⁰³	1.82 × 10 ⁻⁰³	1.62 × 10 ⁻⁰³	8.43 × 10 ⁻⁰⁴	1.74 × 10 ⁻⁰³	9.06 × 10 ⁻⁰⁴	8.43 × 10 ⁻⁰⁴	8.43 × 10 ⁻⁰⁴	8.71 × 10 ⁻⁰⁴
Molybdenum	2.06 × 10 ⁻⁰⁴	2.06 × 10 ⁻⁰⁴	3.75 × 10 ⁻⁰⁴	4.11 × 10 ⁻⁰⁴	4.60 × 10 ⁻⁰⁴	4.07 × 10 ⁻⁰⁴	2.06 × 10 ⁻⁰⁴	4.39 × 10 ⁻⁰⁴	2.23 × 10 ⁻⁰⁴	2.06 × 10 ⁻⁰⁴	2.06 × 10 ⁻⁰⁴	2.14 × 10 ⁻⁰⁴
Sulphate	1.72 × 10 ⁻⁰⁵	1.03 × 10 ⁻⁰²	1.06 × 10 ⁺⁰⁰	1.06 × 10 ⁺⁰⁰	1.06 × 10 ⁺⁰⁰	1.06 × 10 ⁺⁰⁰	1.03 × 10 ⁻⁰²	1.06 × 10 ⁺⁰⁰	1.06 × 10 ⁺⁰⁰	1.03 × 10 ⁻⁰²	1.03 × 10 ⁻⁰²	1.06 × 10 ⁺⁰⁰
Phosphorus	1.03 × 10 ⁻⁰²	1.06 × 10 ⁺⁰⁰	1.05 × 10 ⁻⁰²	1.06 × 10 ⁻⁰²	1.06 × 10 ⁻⁰²	1.06 × 10 ⁻⁰²	1.06 × 10 ⁺⁰⁰	1.06 × 10 ⁻⁰²	1.03 × 10 ⁻⁰²	1.06 × 10 ⁺⁰⁰	1.06 × 10 ⁺⁰⁰	1.03 × 10 ⁻⁰²
Uranium	1.00 × 10 ⁻⁰⁴	1.00 × 10 ⁻⁰⁴	6.16 × 10 ⁻⁰⁴	7.26 × 10 ⁻⁰⁴	8.78 × 10 ⁻⁰⁴	7.16 × 10 ⁻⁰⁴	1.00 × 10 ⁻⁰⁴	8.14 × 10 ⁻⁰⁴	1.51 × 10 ⁻⁰⁴	1.00×10^{-04}	1.00 × 10 ⁻⁰⁴	1.22 × 10 ⁻⁰⁴
	Bq/L	Bq/L	Bq/L	Bq/L	Bq/L	Bq/L	Bq/L	Bq/L	Bq/L	Bq/L	Bq/L	Bq/L
Lead-210	1.03 × 10 ⁻⁰²	5.07 × 10 ⁻⁰³	1.67 × 10 ⁻⁰¹	1.98 × 10 ⁻⁰¹	2.48 × 10 ⁻⁰¹	1.96 × 10 ⁻⁰¹	5.07 × 10 ⁻⁰³	2.26 × 10 ⁻⁰¹	2.05 × 10 ⁻⁰²	5.07×10^{-03}	5.07 × 10 ⁻⁰³	1.18 × 10 ⁻⁰²
Polonium-210	1.03 × 10 ⁻⁰²	5.16 × 10 ⁻⁰³	1.70 × 10 ⁻⁰¹	2.01 × 10 ⁻⁰¹	2.52 × 10 ⁻⁰¹	2.00 × 10 ⁻⁰¹	5.15 × 10 ⁻⁰³	2.30 × 10 ⁻⁰¹	2.08 × 10 ⁻⁰²	5.20 × 10 ⁻⁰³	5.16 × 10 ⁻⁰³	1.20 × 10 ⁻⁰²
Radium-226	1.03 × 10 ⁻⁰²	5.54 × 10 ⁻⁰³	2.26 × 10 ⁻⁰²	2.63 × 10 ⁻⁰²	3.13 × 10 ⁻⁰²	2.59 × 10 ⁻⁰²	5.54 × 10 ⁻⁰³	2.92 × 10 ⁻⁰²	7.21 × 10 ⁻⁰³	5.54×10^{-03}	5.54 × 10 ⁻⁰³	6.27 × 10 ⁻⁰³
Thorium-230	9.09 × 10 ⁻⁰³	9.09 × 10 ⁻⁰³	1.45 × 10 ⁻⁰²	1.56 × 10 ⁻⁰²	1.72 × 10 ⁻⁰²	1.55 × 10 ⁻⁰²	9.09 × 10 ⁻⁰³	1.65 × 10 ⁻⁰²	9.61 × 10 ⁻⁰³	9.09 × 10 ⁻⁰³	9.09 × 10 ⁻⁰³	9.32 × 10 ⁻⁰³
Uranium-234	3.56 × 10 ⁻⁰⁴	3.56 × 10 ⁻⁰⁴	2.18 × 10 ⁻⁰³	2.58 × 10 ⁻⁰³	3.12 × 10 ⁻⁰³	2.54 × 10 ⁻⁰³	3.56 × 10 ⁻⁰⁴	2.89 × 10 ⁻⁰³	5.34 × 10 ⁻⁰⁴	3.56 × 10 ⁻⁰⁴	3.56 × 10 ⁻⁰⁴	4.34 × 10 ⁻⁰⁴
Uranium-238	3.56 × 10 ⁻⁰⁴	3.56 × 10 ⁻⁰⁴	2.18 × 10 ⁻⁰³	2.58 × 10 ⁻⁰³	3.12 × 10 ⁻⁰³	2.54 × 10 ⁻⁰³	3.56 × 10 ⁻⁰⁴	2.89 × 10 ⁻⁰³	5.34 × 10 ⁻⁰⁴	3.56 × 10 ⁻⁰⁴	3.56 × 10 ⁻⁰⁴	4.34 × 10 ⁻⁰⁴


ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT - APPENDIX A - IMPACT MODEL


Development of Model Parameters


Modelled average baseline concentrations are presented in Table 3-5 and a comparison to measured values is shown in Figure 3-2 and Figure 3-3. Parameters include arsenic, chloride, phosphorus, copper, cobalt, sulphate, and uranium, as well as the considered radionuclides (i.e., lead-210, polonium-210, uranium-234, uranium-238, radium-226, thorium-230). The purpose of these plots is to show trends over time for selected COPCs and the generally good agreement between measured and predicted values that resulted from the calibration. Water concentrations for cobalt and copper are likely overestimates, since the modelled concentration is over the detection limit, with most or all measured values being under the detection limit for copper and cobalt, respectively. This overestimation is a result of the method for derivation of water concentrations based on measured sediment values (Section 3.3, Sediment Quality). Unlike chloride and sulphate, metals and radionuclides are expected to interact with sediment. In the case of radionuclides, ingrowth and decay are other factors that may influence activities in the receiving environment.






































Note: Blank outline symbol (no-fill) indicates that the measurement was under the detection limit. Sampling stations summarized in Appendix A.

WQO = Water Quality Objective.

Figure 3-2: Modelled and Measured Baseline Concentrations in Water

3.3 Sediment Quality

3.3.1 Background Sediment Quality

To characterize regional background sediment quality, baseline water quality data were used from baseline studies performed between 2018 and 2020 (Annex V.1). The waterbodies included in the calculation of a regional sediment baseline were Patterson Lake (i.e., North Arm – East Basin and West Basin, South Arm), all measurement stations along the Clearwater River, Lake H, Lake G, Forrest Lake, Beet Lake, Naomi Lake, Mirror River, and Lloyd Lake.

Concentrations in sediment were modelled based on concentrations in water (Section 3.3.2). If water concentration measurements for a constituent were under the detection limit, the baseline concentration in sediment was set to the geometric mean of the measured sediment concentrations. This was the case for copper, cobalt, and thorium-230. The lead-210 sediment concentration was based on its transient equilibrium with polonium-210.

Average baseline concentrations are presented in Table 3-5, and a comparison to measured values is shown in Figure 3-2 and Figure 3-3. Parameters include arsenic, copper, cobalt, uranium, and the considered radionuclides (i.e., lead-210, polonium-210, uranium-234, uranium-238, radium-226, thorium-230). These plots indicate generally good agreement between measured and predicted values that resulted from the calibration. Metals and radionuclides are expected to interact with water and, in the case of radionuclides, ingrowth and decay are other factors that may influence activities in the receiving environment.

Table 3-5: Baseline Water and Sediment Concentrations Used in the IMPACT Model

conc	Water Baseline Concentration	Sediment Baseline Concentration
СОРС	mg/L	mg/kg dw
Arsenic	1.24 × 10 ⁻⁰⁴	$1.00 \times 10^{+01}$
Chloride	4.22 × 10 ⁻⁰¹	-
Cobalt	5.83 × 10 ⁻⁰⁴	1.45 × 10 ⁺⁰⁰
Copper	8.43 × 10 ⁻⁰⁴	2.52 × 10 ⁺⁰⁰
Molybdenum	2.06 × 10 ⁻⁰⁴	6.51 × 10 ⁻⁰¹
Sulphate	1.72 × 10 ⁻⁰⁵	-
Phosphorus	1.03 × 10 ⁻⁰²	$1.01 \times 10^{+00}$
Uranium	1.00 × 10 ⁻⁰⁴	1.93 × 10 ⁺⁰⁰
	Bq/L	Bq/kg dw
Lead-210 ⁽¹⁾	1.03 × 10 ⁻⁰²	$1.01 \times 10^{+00}$
Polonium-210	1.03 × 10 ⁻⁰²	$1.01 \times 10^{+00}$
Radium-226	1.03 × 10 ⁻⁰²	$1.01 \times 10^{+00}$
Thorium-230	9.09 × 10 ⁻⁰³	2.09 × 10 ⁺⁰¹
Uranium-234 ⁽²⁾	3.56 × 10 ⁻⁰⁴	6.84 × 10 ⁺⁰⁰
Uranium-238 ⁽²⁾	3.56 × 10 ⁻⁰⁴	$6.84 \times 10^{+00}$


Dash (-) represents no value calculated as COPC does not partition to sediment.


Bold = measured values. Values that are not bolded indicate modelled values.


- (1) Calculated to be in transient equilibrium with polonium-210.
- (2) Calculated based on the specific activity of uranium.


COPC = constituent of potential concern; dw = dry weight.


























Blank outline symbol (no-fill) indicates that the measurement was under the detection limit. Sampling stations summarized in Appendix A.

SQO = Sediment Quality Objective.

Figure 3-3: Modelled and Measured Baseline Concentrations in Sediment

3.3.2 Water to Sediment Partitioning Coefficients

The water-to-sediment partitioning coefficients (K_d) are regional parameters governing the exchange between sediment and water in lakes. The K_d values are used to estimate the fraction of a constituent that is associated with the particulate fraction in the shallow sediment layer (f_s). The fraction of COPC in the solid phase is estimated using the following equation:

$$f_{s} = \frac{K_{d} \cdot \frac{\rho_{s}}{\varepsilon_{s}}}{1 + K_{d} \cdot \frac{\rho_{s}}{\varepsilon_{s}}}$$

where:

 ε_s = porosity of surficial sediment (unitless)

K_d = distribution coefficient between water and sediment (L/kg)

 ρ_s = bulk density of surficial sediment (kg/L)

The K_d values used in this model are presented in Table 3-6. They consist of regional published values that have been calibrated on similar sites in northern Saskatchewan.

Table 3-6: Distribution Coefficients Used in the IMPACT Model

СОРС	Distribution Coefficient
	L/kg dw
Arsenic	9.64 × 10 ⁺⁰⁴
Cobalt	2.50 × 10 ⁺⁰³
Copper	$3.00 \times 10^{+03}$
Molybdenum	$3.17 \times 10^{+03}$
Uranium	2.00 × 10 ⁺⁰⁴
Lead-210	1.20 × 10 ⁺⁰⁵
Polonium-210	1.20 × 10 ⁺⁰⁵
Radium-226	1.20 × 10 ⁺⁰⁴
Thorium-230	2.30 × 10 ⁺⁰³
Uranium-234	2.00 × 10 ⁺⁰⁴
Uranium-238	2.00 × 10 ⁺⁰⁴

COPC = constituent of potential concern; dw = dry weight.

3.4 Air Quality

Risk through air exposure pathways is considered during Operations. For the baseline, it is assumed that none of the considered COPC is present in air. Consequently, any risk increase through the air pathway is due to the operation of the mine. An exception is radon-222, which is discussed in detail in Section 2.4.3, Radiological Dose. A constant background concentration of 2.9 Bq/m³ for radon-222 was used for the evaluation of incremental radon concentrations and incremental dose due to radon.

Modelled air quality during Operations was provided at the locations of interest for human and ecological receptors from the atmospheric environment model (Appendix 7A, Air Dispersion Modelling Report).

3.5 Soil Quality

3.5.1 Background Soil Quality

Regional background soil chemistry was derived from baseline data collected in 2019 (Annex VI, Terrain and Soils Baseline Report). The geometric mean of 18 soil samples at 6 locations was used to characterize background concentrations of metals and radionuclides in soil. Table 3-7 presents the selected background soil concentrations used in the IMPACT model. Sandy soil was selected for the IMPACT model based on baseline studies identifying the soils of the region are dominantly brunisolic soils that have been developed on sandy glacial till deposits (Annex VI).

COPC Unit **Soil Baseline Concentration** 5.89×10^{-01} Arsenic mg/kg dw Cobalt mg/kg dw 4.32×10^{-01} 6.01×10^{-01} Copper mg/kg dw 1.42×10^{-01} Molybdenum mg/kg dw 2.47×10^{-01} Uranium mg/kg dw Lead-210 $4.00 \times 10^{+01}$ Bq/kq dw $1.12 \times 10^{+01}$ Polonium-210 Bq/kg dw Radium-226 Bq/kg dw $1.79 \times 10^{+01}$ $2.00 \times 10^{+01}$ Thorium-230 Bq/kg dw Uranium-234^(a) Bq/kg dw $3.06 \times 10^{+00}$ Uranium-238(a) $3.06 \times 10^{+00}$ Bq/kq dw

Table 3-7: Soil Baseline Concentrations

a) Uranium-234 and Uranium-238 concentration calculated using a specific activity of 12.356 Bq/mg.

COPC = constituent of potential concern; Bq = becquerel; dw = dry weight.

3.5.2 Soil Quality during Operations

Soil quality during Operations was estimated based on the calculated air quality at each location of interest for human and ecological receptors. IMPACT calculates COPC concentrations in terms

of concentration per kg of dry soil, following the soil model equations in CSA N288.1-20 (Section 2.3.4, Terrestrial Pathways).

For wet deposition, the fraction of the time that precipitation falls when the wind blows from a specific sector was set conservatively to 0.36 for all wind directions (CSA Group 2020).

Concentrations calculated from deposition of air on soil during Operations were added to baseline concentrations to represent the total exposure concentrations in soil.

3.6 Transfer of Constituents to Aquatic Receptors

A fundamental premise of pathways analysis is that chemical uptake by receptors is related to the receptors' level of exposure. A linear relationship is usually assumed and is represented by a bioaccumulation factor (BAF). For aquatic receptors, the BAF is the concentration in the organism (e.g., Cfish) divided by the concentration in water (Cwater). The BAF may be estimated from organism and water data by fitting a regression line that generally is assumed to pass through the origin, such as "Cfish = Slope · Cwater". The slope of the regression line is the BAF. It represents the equilibrium ratio of Cfish / Cwater.

The IMPACT model uses the BAF model as a simplification of the more complex multi-media uptake process. While several media (e.g., water and food) may contribute to COPC uptake into the animal tissue, at steady state, all media concentrations will have a fixed ratio to the tissue concentration. Thus, the BAF based on water and tissue measurements will reflect all the multi-media contributions to uptake. If animals and their prey are in equilibrium with the COPC in the environment, their concentrations can be estimated using an overall BAF between the water and the organism (Thomann and Mueller 1987). If animals have not reached equilibrium because they have not spent enough time in the exposure situation, because they are migratory or the project effects are transient, the BAF provides a conservative estimate of tissue concentrations.

As a result of physiological control, intracellular storage, and different excretion mechanisms, biota have an ability to actively regulate the body burden of many metals, including selenium, and maintain homeostatic control over a range of exposures (Chapman et al. 1996; Hamilton and Mehrle 1986; Wood and Port 2000). These homeostatic controls can produce non-linear relationships between the steady-state tissue concentrations and the environmental exposure concentrations (Newman and Unger 2002). However, these complicating issues do not diminish the importance, or negate the practical application, of BAFs in the assessment of environmental risks associated with COPCs.

Because of non-linearity and other factors, BAF values cited in the literature vary over a considerable range. Bioaccumulation factor values based on site-specific data are preferred. The values of BAFs based on background concentrations tend to be higher than those based on higher exposure concentrations in affected environments. In this model BAFs representative of regional conditions have been used. The predictions are compared to measured baseline values in Section 3.6.2.

3.6.1 Aquatic Bioaccumulation Factors

Bioaccumulation factors relate the COPCs in the environmental media to the concentration in the receptor. Water-based BAFs were used to calculate COPC concentrations in aquatic and terrestrial plant, invertebrate, and fish tissues. These factors were generally obtained from CSA N288.1-20 and the International Atomic Energy Agency (IAEA 2010), and from publicly available regional data from other uranium mine sites in northern Saskatchewan. Table 3-8 shows all aquatic BAFs used in the model.

The water-based BAF values for aquatic organisms were based on measured ratios of COPC concentration in tissue vs. water, under multi-media equilibrium conditions.

BAF (L/kg wet weight) =
$$\frac{\text{Concentration in Tissue (mg/kg wet weight)}}{\text{Concentration in Water (mg/L)}}$$

While expressed relative to water, BAFs represent all pathways of COPC uptake into the organism, because all pathways were operating under the environmental conditions of BAF measurement. These pathways include food ingestion, dermal absorption, and uptake across the gills.

СОРС	Zooplankton ⁽¹⁾	Benthic Invertebrates ⁽¹⁾	Macrophytes ⁽²⁾	Northern Pike	Whitefish
Arsenic	5.40 × 10 ⁺⁰²	$5.40 \times 10^{+02}$	$1.72 \times 10^{+02}$	$3.00 \times 10^{+02}$	$3.00 \times 10^{+02}$
Cobalt	2.00 × 10 ⁺⁰²	$2.00 \times 10^{+02}$	$7.90 \times 10^{+02}$	$1.20 \times 10^{+01}$	$1.20 \times 10^{+01}$
Copper	7.80 × 10 ⁺⁰³	$7.80 \times 10^{+03}$	$2.17 \times 10^{+02}$	$5.00 \times 10^{+02}$	5.00 × 10 ⁺⁰¹
Molybdenum	1.62 × 10 ⁺⁰²	1.62 × 10 ⁺⁰²	$6.80 \times 10^{+00}$	1.00 × 10- ⁰¹	1.00 × 10- ⁰¹
Uranium	1.00 × 10 ⁺⁰²	$1.00 \times 10^{+02}$	$5.04 \times 10^{+01}$	$2.00 \times 10^{+01}$	5.00 × 10 ⁺⁰⁰
Lead-210	2.30 × 10 ⁺⁰³	$2.30 \times 10^{+03}$	1.90 × 10 ⁺⁰³	$6.00 \times 10^{+01}$	2.20 × 10 ⁺⁰¹
Polonium-210	1.58 × 10 ⁺⁰⁴	$1.58 \times 10^{+04}$	$2.93 \times 10^{+02}$	$1.50 \times 10^{+02}$	1.50 × 10 ⁺⁰²
Radium-226	2.50 × 10 ⁺⁰²	$2.50 \times 10^{+02}$	1.63 × 10 ⁺⁰²	$1.20 \times 10^{+01}$	1.20 × 10 ⁺⁰¹
Thorium-230	5.00 × 10 ⁺⁰²	5.00 × 10 ⁺⁰²	$2.32 \times 10^{+02}$	$6.00 \times 10^{+00}$	$6.00 \times 10^{+00}$
Uranium-234	1.00 × 10 ⁺⁰²	$1.00 \times 10^{+02}$	$5.04 \times 10^{+01}$	$2.00 \times 10^{+01}$	5.00 × 10 ⁺⁰⁰
Uranium-238	$1.00 \times 10^{+02}$	$1.00 \times 10^{+02}$	$5.04 \times 10^{+01}$	$2.00 \times 10^{+01}$	$5.00 \times 10^{+00}$

Table 3-8: Aquatic Bioaccumulation Factors (L/kg fw)

Regional data was used for aquatic BAFs, except for:

COPC = constituent of potential concern; fw = fresh weight.

3.6.2 Model Validation

Figure 3-4 presents the predicted and observed concentrations of COPCs in aquatic macrophytes from baseline sampling in 2019 (Annex V.1). The predicted concentrations fall within the spread of measurements for macrophytes, except for thorium-230, where the model overpredicts macrophyte measurements that are under the detection limit. This comparison suggests that the parameters used in the model are suitable for describing the transport processes in the

⁽¹⁾ Th-230, Ra-226, U and Co BAFs for Benthic Invertebrates are from Thompson (1972).


⁽²⁾ Pb BAF for Macrophytes is from IAEA TRS 472 (IAEA 2010).

ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT - APPENDIX A - IMPACT MODEL


Development of Model Parameters

environment for aquatic plants. In the instance of thorium-230, the model produces conservative results. Refinement of the BAFs for aquatic plants can be completed as new data are available during future phases of the Project.

Figure 3-5 and Figure 3-6 present the predicted and observed concentrations of COPCs in northern pike and lake whitefish from baseline sampling in 2018 and 2019 (Annex V.1). Measured concentrations of radionuclides are mostly under the detection limit, which agrees with the modelled concentrations. In the case of northern pike, the data suggest that the model may be overpredicting concentrations in arsenic, cobalt, and copper. For lake whitefish, the model may be underpredicting copper; however, only data from one sampling event were available, and the BAFs used are consistent with regionally calibrated models representing exposure conditions. The predicted concentrations of the remaining COPCs are in good agreement with the measured values. The generally good agreement of modelled results with measured data validates the use of selected transfer factors and modelled processes. However, refinement of the BAFs for aquatic animals can be completed as new data become available during future phases of the Project.



Blank outline symbol (no-fill) indicates that the measurement was under the detection limit.



Blank outline symbol (no-fill) indicates that the measurement was under the detection limit.

Figure 3-4: Modelled and Measured Baseline Concentrations (fresh weight) of Radionuclides in Macrophytes





Figure 3-5: Modelled and Measured Baseline Concentrations (fresh weight) of Radionuclides in Northern Pike






Figure 3-6: Modelled and Measured Baseline Concentrations (fresh weight) of Radionuclides in Lake Whitefish

3.6.3 Dose Coefficients for Aquatic Receptors

Dose coefficients (DCFs) for all external and internal exposure routes considered are presented in Table 3-9 and Table 3-10, respectively. Dose coefficients for internal and external exposure pathways are based on values published by the International Commission on Radiological Protection (ICRP) for aquatic plants and northern pike (ICRP 2008). For benthic invertebrates, zooplankton, and whitefish, DCFs were calculated with the ERICA Tool (Brown et al. 2008).

The DCF values from the ICRP (2008) and the ERICA Tool 1.3.1 (Brown et al. 2008) do not incorporate radiation quality factors for relative biological effectiveness for low beta and alpha components. The relative biological effectiveness is a radioecological weighting factor that represents the ratio of doses from different types of radiation needed to produce the same biological effect. Therefore, the "low beta" components of the DCFs were multiplied by 2, and the alpha components were multiplied by 10 (as per CSA N288.6-22), to represent their greater relative effectiveness.

Table 3-9: External Dose Coefficients for Aquatic Receptors

	DCF External (μGy/h)/(Bq/kg[fw sediment] or Bq/L[water])										
СОРС	Macrophytes ^(a)	Northern Pike ^(b)	Whitefish ^(c)	Zooplankton ^(d)	Benthic Invertebrates ^(e)						
Lead-210	3.38 × 10 ⁻⁰⁵	4.00×10^{-06}	3.80×10^{-06}	1.50 × 10 ⁻⁰⁴	1.10 × 10 ⁻⁰⁴						
Polonium-210	4.58×10^{-09}	4.17×10^{-09}	4.30×10^{-09}	4.90×10^{-09}	4.90×10^{-09}						
Radium-226	1.13×10^{-03}	9.17 × 10 ⁻⁰⁴	9.10×10^{-04}	1.40×10^{-03}	1.30×10^{-03}						
Thorium-230	4.58×10^{-07}	2.50×10^{-07}	2.40×10^{-07}	1.00×10^{-06}	9.00 × 10 ⁻⁰⁷						
Uranium-234	4.17×10^{-07}	1.63×10^{-07}	1.50×10^{-07}	1.00×10^{-06}	9.40×10^{-07}						
Uranium-238	3.17×10^{-07}	1.00×10^{-07}	9.50×10^{-08}	7.80×10^{-07}	7.20×10^{-07}						

a) Dose coefficients are based on seaweed (ICRP 2008).



b) Dose coefficients are based on trout (ICRP 2008).

c) Dose coefficients are the value for benthic fish from the ERICA Database (Brown et al. 2008).

d) Dose coefficients are from the ERICA Database (Brown et al. 2008).

e) Dose coefficients are the value for insect larvae from the ERICA Database (Brown et al. 2008).

μGy = microgray, Bq = becquerel; fw = fresh weight; COPC = constituent of potential concern; DCF = dose coefficient.

DCF Internal (µGy/h)/(Bq/kg fw) Northern **Benthic COPC** Macrophytes^(a) Whitefish(c) Zooplankton^(d) Pike(b) Invertebrate^(e) Lead-210 2.17×10^{-04} 2.46×10^{-04} 2.50×10^{-04} 1.00×10^{-04} 1.40×10^{-04} Polonium-210 3.04×10^{-02} 3.04×10^{-02} 3.10×10^{-03} 3.10×10^{-02} 3.10×10^{-03} Radium-226 1.38×10^{-01} 1.39×10^{-01} 1.43×10^{-01} 1.37×10^{-01} 1.37×10^{-01} 2.71×10^{-02} Thorium-230 2.71×10^{-02} 2.70×10^{-02} 2.70×10^{-02} 2.70×10^{-02} 2.75×10^{-02} 2.75×10^{-02} 2.80×10^{-02} 2.80×10^{-02} 2.80×10^{-02} Uranium-234 2.42×10^{-02} 2.42×10^{-02} 2.40×10^{-02} 2.40×10^{-02} 2.40×10^{-02} Uranium-238

Table 3-10: Internal Dose Coefficients for Aquatic Receptors

3.7 Transfer of Constituents to Terrestrial Receptors and Humans

The COPCs may be transferred from environmental media to terrestrial receptors through food, water, and air. Ingestion and inhalation transfer factors (TFs) are defined as the ratio between the COPC concentrations in the receptor and the COPC intake rates. The TF may be determined for specific organs or other body tissues (e.g., food to liver; food to eggs). Soil to terrestrial plant transfers may include deposition from air to plant, as well as soil-to-plant bioaccumulation, and are defined as the ratio between the concentration of a COPC in the plant and that in the soil, both in dry weight.

3.7.1 Soil-to-Plant Transfer

A plant may take up COPCs by dry and wet deposition of COPCs from air and raindrop splash onto the plant, as well as from uptake of COPCs from soil to the plant via the root system. In the case of most COPCs, transfer from deposition on soil and then uptake from soil into the plant is considered to be the dominant pathway. The terrestrial plants include browse, blueberries, lichen, and Labrador tea. The soil-to-plant pathway is not considered applicable to lichen; therefore, the soil-to-plant BAFs are zero for lichen. Lichens are considered to take up COPCs via wet and dry deposition directly from air only (IAEA, 2010).

Uptake of COPCs by browse, blueberries, and Labrador tea is represented by soil-to-plant BAF values derived from regional data from northern Saskatchewan, as shown in Table 3-11. Uptake of COPCs by terrestrial invertebrates from soil is represented by the soil-to-invertebrate BAFs, as shown in Table 3-10.

a) Dose coefficients are based on seaweed (ICRP 2008).

b) Dose coefficients are based on trout (ICRP 2008).

c) Dose coefficients are the value for benthic fish from the ERICA Database (Brown et al. 2008).

d) Dose coefficients are from the ERICA Database (Brown et al. 2008).

e) Dose coefficients are the value for insect larvae from the ERICA Database (Brown et al. 2008).

µGy = microgray; Bq = becquerel; fw = fresh weight; COPC = constituent of potential concern; DCF = dose coefficient.

Table 3-11: Bioaccumulation Factors for Terrestrial Plants and Invertebrates

Bio	accumulation Factor kg(dw soil)/kg	g(dw plant)
СОРС	Terrestrial Plants ^(a)	Terrestrial Invertebrates ^(b)
Arsenic	4.26 × 10 ⁻⁰¹	2.24 × 10 ⁻⁰¹
Cobalt	5.80 × 10 ⁻⁰¹	3.58 × 10 ⁻⁰²
Copper	6.64 × 10 ⁺⁰⁰	5.15 × 10 ⁻⁰¹
Molybdenum	1.00 × 10 ⁺⁰⁰	$1.54 \times 10^{+00}$
Uranium	1.62 × 10 ⁺⁰⁰	5.20 × 10 ⁻⁰²
Lead-210	8.00 × 10 ⁻⁰²	1.68 × 10 ⁻⁰¹
Polonium-210	3.86 × 10 ⁻⁰¹	1.64 × 10 ⁻⁰²
Radium-226	6.00 × 10 ⁻⁰¹	5.29 × 10 ⁻⁰¹
Thorium-230	3.12 × 10 ⁻⁰²	5.20 × 10 ⁻⁰²
Uranium-234	1.62 × 10 ⁺⁰⁰	5.20 × 10 ⁻⁰²
Uranium-238	1.62 × 10 ⁺⁰⁰	5.20 × 10 ⁻⁰²


a) BAFs for terrestrial plants are compiled from regional data.

3.7.2 Model Validation

Figure 3-7 presents the predicted and observed concentrations of COPCs in blueberries from baseline sampling in 2019 (Annex VII.3). Modelled concentrations generally fall within the range of measured values from the two sampling events. Based on available data, the model is overpredicting arsenic (which is non-detect in blueberry samples), cobalt, and uranium. This comparison suggests a good agreement of available data with the modelling approach. For constituents for which the model predictions fall outside of the observed range, concentrations are conservatively overestimated. Refinement of the BAFs for aquatic plants can be completed as new data are available.

b) BAFs for terrestrial invertebrates from Sample et al (1998).

dw = dry weight; COPC = constituent of potential concern.






Figure 3-7: Modelled and Measured Concentrations (dry weight) of Radionuclides in Blueberries

3.7.3 Dose Coefficients for Terrestrial Plants and Invertebrates

Dose coefficients for all external and internal exposure routes to terrestrial plants and invertebrates are presented in Table 3-12 and Table 3-13, respectively. Dose coefficients for internal and external exposure pathways are based on values published by the ICRP (ICRP 2008).

Table 3-12: External Dose Coefficients for Terrestrial Plants and Invertebrates

СОРС		ternal Soil Su Gy/h)/(Bq/m²)	DCF External Soil (μGy/h)/(Bq/kg[dw soil])		
COPC	Blueberries ^(a)	Browse ^(b)	Labrador Tea ^(b)	Terrestrial Invertebrates ^(c)	
Lead-210	7.50 × 10 ⁻⁰⁹	1.13×10^{-06}	1.13×10^{-06}	5.83 × 10 ⁻⁰⁷	
Polonium-210	1.96 × 10 ⁻¹¹	6.67×10^{-11}	6.67×10^{-11}	4.58×10^{-09}	
Radium-226	3.88 × 10 ⁻⁰⁶	1.33×10^{-05}	1.33×10^{-05}	9.17 × 10 ⁻⁰⁴	
Thorium-230	2.67 × 10 ⁻⁰⁹	3.04×10^{-07}	3.04×10^{-07}	2.08×10^{-07}	
Uranium-234	3.38 × 10 ⁻⁰⁹	3.67×10^{-07}	3.67×10^{-07}	1.75 × 10 ⁻⁰⁷	
Uranium-238	2.63 × 10 ⁻⁰⁹	3.04×10^{-07}	3.04×10^{-07}	1.25 × 10 ⁻⁰⁷	

a) Dose coefficients are based on pine tree (ICRP 2008).

 μ Gy = microgray; Bq = becquerel; dw = dry weight; COPC = constituent of potential concern; DCF = dose coefficient.

Table 3-13: Internal Dose Coefficients for Terrestrial Plants and Invertebrates

	DCF Internal (μGy/h)/(Bq/kg(fw plant))								
СОРС	Blueberries ^(a)	Browse ^(b)	Labrador Tea ^(b)	Terrestrial Invertebrate ^(c)					
Lead-210	2.50×10^{-04}	2.25×10^{-04}	2.25 × 10 ⁻⁰⁴	2.25 × 10 ⁻⁰⁴					
Polonium-210	3.04×10^{-02}	3.04×10^{-02}	3.04×10^{-02}	3.04×10^{-02}					
Radium-226	1.41×10^{-01}	1.38×10^{-01}	1.38×10^{-01}	1.38 × 10 ⁻⁰¹					
Thorium-230	2.71 × 10 ⁻⁰²	2.71×10^{-02}	2.71×10^{-02}	2.71 × 10 ⁻⁰²					
Uranium-234	2.75×10^{-02}	2.75×10^{-02}	2.75×10^{-02}	2.75×10^{-02}					
Uranium-238	2.42×10^{-02}	2.42×10^{-02}	2.42×10^{-02}	2.42 × 10 ⁻⁰²					

a) Dose coefficients are based on pine tree (ICRP 2008).

μGy = microgray; Bq = becquerel; fw = fresh weight; COPC = constituent of potential concern; DCF = dose coefficient.



b) Dose coefficients are based on grass (ICRP 2008).

c) Dose coefficients from ICRP (2008).

b) Dose coefficients are based on grass (ICRP 2008).

c) Dose coefficients from ICRP (2008).

3.7.4 Ingestion Transfer Factors for Terrestrial Receptors

Ingestion transfer factors are COPC- and biota-specific. Ingestion TFs from forage to tissue for agricultural livestock are available in CSA N288.1-20. An allometric equation (i.e., transfer proportional to a -3/4 power of body weight) (CSA N288.6-22) was applied to TFs available for beef and poultry from CSA N288.1-20, the IAEA (2010), or the National Council on Radiation Protection and Measurements (NCRP 1996) to estimate the TFs for the mammal and bird receptors, respectively. The derived ingestion TFs are presented in Table 3-14 and Table 3-15.

3.7.5 Inhalation Transfer Factors for Terrestrial Receptors

Inhalation TFs were calculated from the ingestion TF, by adjusting the ingestion TF by a COPC-specific inhalation/ingestion ratio (II) from CSA N288.1-20. The derived inhalation TFs are presented in Table 3-16.



Table 3-14: Ingestion Transfer Factors for Mammals (d/kg fw)

СОРС	Beef Transfer Factor	Beef Transfer Factor Reference	Woodland Caribou	Snowshoe Hare	Moose	Little Brown Myotis	Southern Red-Backed Vole	Black Bear	Red Fox	Grey Wolf	Beaver	Muskrat	Mink	Beef Organs Transfer Factor	Beef Organs Transfer Factor Reference	Moose Organs ^(d)
Non-radionuclides																
Arsenic	2.00 × 10 ⁻⁰²	(a)	4.93 × 10 ⁻⁰²	1.56 × 10 ⁺⁰⁰	2.71 × 10 ⁻⁰²	9.51 × 10 ⁺⁰¹	3.14 × 10 ⁺⁰¹	7.53 × 10 ⁻⁰²	7.80 × 10 ⁻⁰¹	1.40 × 10 ⁻⁰¹	2.24 × 10 ⁻⁰¹	2.15 × 10 ⁺⁰⁰	2.39 × 10 ⁺⁰⁰	1.70 × 10 ⁻⁰¹	(a)	2.30 × 10 ⁻⁰¹
Cobalt	4.30 × 10 ⁻⁰⁴	(a)	1.06 × 10 ⁻⁰³	3.35 × 10 ⁻⁰²	5.83 × 10 ⁻⁰⁴	2.05 × 10 ⁺⁰⁰	6.75 × 10 ⁻⁰¹	1.62 × 10 ⁻⁰³	1.68 × 10 ⁻⁰²	3.00 × 10 ⁻⁰³	4.81 × 10 ⁻⁰³	4.62 × 10 ⁻⁰²	5.14 × 10 ⁻⁰²	1.00 × 10 ⁻⁰¹	(a)	1.36 × 10 ⁻⁰¹
Copper	1.00 × 10 ⁻⁰²	(b)	2.47 × 10 ⁻⁰²	7.80 × 10 ⁻⁰¹	1.36 × 10 ⁻⁰²	4.76 × 10 ⁺⁰¹	1.57 × 10 ⁺⁰¹	3.76 × 10 ⁻⁰²	3.90 × 10 ⁻⁰¹	6.98 × 10 ⁻⁰²	1.12 × 10 ⁻⁰¹	1.07 × 10 ⁺⁰⁰	1.19 × 10 ⁺⁰⁰	4.00 × 10 ⁻⁰¹	(a,d)	5.42 × 10 ⁻⁰¹
Molybdenum	1.00 × 10 ⁻⁰³	(a)	1.23 × 10 ⁻⁰²	3.90 × 10 ⁻⁰¹	6.78 × 10 ⁻⁰³	2.38 × 10 ⁺⁰¹	7.85 × 10 ⁺⁰⁰	1.88 × 10 ⁻⁰²	1.95 × 10 ⁻⁰¹	3.49 × 10 ⁻⁰²	5.59 × 10 ⁻⁰²	5.37 × 10 ⁻⁰¹	5.97 × 10 ⁻⁰¹	5.00 × 10 ⁻⁰³	(a)	6.78 × 10 ⁻⁰³
Uranium	3.90 × 10 ⁻⁰⁴	(a)	9.62 × 10 ⁻⁰⁴	3.04 × 10 ⁻⁰²	5.29 × 10 ⁻⁰⁴	1.86 × 10 ⁺⁰⁰	6.13 × 10 ⁻⁰¹	1.47 × 10 ⁻⁰³	1.52 × 10 ⁻⁰²	2.72 × 10 ⁻⁰³	4.36 × 10 ⁻⁰³	4.19 × 10 ⁻⁰²	4.66 × 10 ⁻⁰²	6.90 × 10 ⁻⁰⁴	(a)	9.35 × 10 ⁻⁰⁴
Radionuclides																
Lead-210	7.00 × 10 ⁻⁰⁴	(b)	1.73 × 10 ⁻⁰³	5.46 × 10 ⁻⁰²	9.49 × 10 ⁻⁰⁴	3.33 × 10 ⁺⁰⁰	1.10 × 10 ⁺⁰⁰	2.63 × 10 ⁻⁰³	2.73 × 10 ⁻⁰²	4.88 × 10 ⁻⁰³	7.83 × 10 ⁻⁰³	7.52 × 10 ⁻⁰²	8.36 × 10 ⁻⁰²	2.20 × 10 ⁻⁰²	(a,d)	2.98 × 10 ⁻⁰²
Polonium-210	5.00 × 10 ⁻⁰³	(c)	1.23 × 10 ⁻⁰²	3.90 × 10 ⁻⁰¹	6.78 × 10 ⁻⁰³	2.38 × 10 ⁺⁰¹	7.85 × 10 ⁺⁰⁰	1.88 × 10 ⁻⁰²	1.95 × 10 ⁻⁰¹	3.49 × 10 ⁻⁰²	5.59 × 10 ⁻⁰²	5.37 × 10 ⁻⁰¹	5.97 × 10 ⁻⁰¹	5.00 × 10 ⁻⁰⁵	(a,d)	6.78 × 10 ⁻⁰⁵
Radium-226	1.70 × 10 ⁻⁰³	(a)	4.19 × 10 ⁻⁰³	1.33 × 10 ⁻⁰¹	2.30 × 10 ⁻⁰³	8.09 × 10 ⁺⁰⁰	2.67 × 10 ⁺⁰⁰	6.40 × 10 ⁻⁰³	6.63 × 10 ⁻⁰²	1.19 × 10 ⁻⁰²	1.90 × 10 ⁻⁰²	1.83 × 10 ⁻⁰¹	2.03 × 10 ⁻⁰¹	9.50 × 10 ⁻⁰⁴	(a)	1.29 × 10 ⁻⁰³
Thorium-230	2.30 × 10 ⁻⁰⁴	(a)	5.67 × 10 ⁻⁰⁴	1.79 × 10 ⁻⁰²	3.12 × 10 ⁻⁰⁴	1.09 × 10 ⁺⁰⁰	3.61 × 10 ⁻⁰¹	8.66 × 10 ⁻⁰⁴	8.97 × 10 ⁻⁰³	1.60 × 10 ⁻⁰³	2.57 × 10 ⁻⁰³	2.47 × 10 ⁻⁰²	2.75 × 10 ⁻⁰²	6.30 × 10 ⁻⁰²	(a)	8.54 × 10 ⁻⁰²

Transfer factors in this table were calculated using the following allometric equation: TF (d/kg fw) = aBW^{-0.75}, where BW = body weight, TF = transfer factor, d/kg fw = days per kilogram fresh weight, "a" is a constant calculated using published transfer factor and body weight values for beef or beef or gans as per CSA N288.1 (CSA Group 2020), and applied to calculate species-specific transfer factors.

a) CSA N288.1-20 (CSA Group 2020).

b) Handbook of Parameter Values for the Prediction of Radionuclide Transfer in Terrestrial and Freshwater Environments (IAEA 2010).

c) NCRP No. 123 (NCRP 1996).

d) For elements that are not listed in Table G.3 (CSA Group 2020), a surrogate element was used as follows (surrogate in parentheses): copper (silver), lead (tin), polonium (tellurium).

COPC = constituent of potential concern; TF = transfer factor; BW = body weight.

Table 3-15: Ingestion Transfer Factors for Birds (d/kg fw)

СОРС	Poultry Transfer Factor	Poultry Transfer Factor Reference	Rusty Blackbird	Canada Goose	Grouse	Mallard	Loon
Non-radionuclides							
Arsenic	1.20 × 10 ⁺⁰⁰	(a)	1.59 × 10 ⁺⁰¹	7.56 × 10 ⁻⁰¹	2.96 × 10 ⁺⁰⁰	1.84 × 10 ⁺⁰⁰	5.78 × 10 ⁻⁰¹
Cobalt	9.70 × 10 ⁻⁰¹	(a)	1.28 × 10 ⁺⁰¹	6.11 × 10 ⁻⁰¹	2.39 × 10 ⁺⁰⁰	1.49 × 10 ⁺⁰⁰	4.67 × 10 ⁻⁰¹
Copper	6.20 × 10 ⁻⁰¹	(c)	8.19 × 10 ⁺⁰⁰	3.91 × 10 ⁻⁰¹	1.53 × 10 ⁺⁰⁰	9.51 × 10 ⁻⁰¹	2.99 × 10 ⁻⁰¹
Molybdenum	1.80 × 10 ⁻⁰¹	(a)	2.38 × 10 ⁺⁰⁰	1.13 × 10 ⁻⁰¹	4.44 × 10 ⁻⁰¹	2.76 × 10 ⁻⁰¹	8.67 × 10 ⁻⁰²
Uranium	7.50 × 10 ⁻⁰¹	(a)	9.91 × 10 ⁺⁰⁰	4.73 × 10 ⁻⁰¹	1.85 × 10 ⁺⁰⁰	1.15 × 10 ⁺⁰⁰	3.61 × 10 ⁻⁰¹
Radionuclides							
Lead-210	5.00 × 10 ⁻⁰²	(c)	6.61 × 10 ⁻⁰¹	3.15 × 10 ⁻⁰²	1.23 × 10 ⁻⁰¹	7.67×10^{-02}	2.41 × 10 ⁻⁰²
Polonium-210	2.40 × 10 ⁺⁰⁰	(b)	3.17 × 10 ⁺⁰¹	1.51 × 10 ⁺⁰⁰	5.92 × 10 ⁺⁰⁰	$3.68 \times 10^{+00}$	1.16 × 10 ⁺⁰⁰
Radium-226	3.00 × 10 ⁻⁰²	(a)	3.97 × 10 ⁻⁰¹	1.89 × 10 ⁻⁰²	7.40 × 10 ⁻⁰²	4.60 × 10 ⁻⁰²	1.44 × 10 ⁻⁰²
Thorium-230	1.00 × 10 ⁻⁰²	(a)	1.32 × 10 ⁻⁰¹	6.30 × 10 ⁻⁰³	2.47 × 10 ⁻⁰²	1.53 × 10 ⁻⁰²	4.81 × 10 ⁻⁰³

Transfer factors in this table were calculated using the following allometric equation: TF (d/kg FW) = aBW^{-0.75}, where "a" is a constant calculated using published TF and BW values for poultry, and applied to calculate species-specific TFs. a) CSA N288.1-20 (CSA Group 2020).

b) Handbook of Parameter Values for the Prediction of Radionuclide Transfer in Terrestrial and Freshwater Environments (IAEA 2010)

c) Transfer factor was calculated based on the beef TF published in NCRP No. 123 (NCRP 1996), adjusted to poultry using a ratio of 62, as per the CANDU Owners Group DRL Guidance (Hart 2013).

COPC = constituent of potential concern; TF = transfer factor; FW = fresh weight; BW = body weight; DRL = derived release limit.

Table 3-16: Inhalation Transfer Factors for Mammals and Birds (d/kg fw)

СОРС	Inhalation/ Ingestion Ratio (II)	Woodland Caribou	Snowshoe Hare	Moose	Little Brown Myotis	Southern Red-Backed Vole	Black Bear	Red Fox	Grey Wolf	Beaver	Muskrat	Mink	Moose Organs	Rusty Blackbird	Canada Goose	Grouse	Mallard	Loon
Non-radionucl	ides																	
Arsenic	0.75	3.70×10^{-02}	1.17 × 10 ⁺⁰⁰	2.03 × 10 ⁻⁰²	7.14 × 10 ⁺⁰¹	2.36 × 10 ⁺⁰¹	5.64 × 10 ⁻⁰²	5.85 × 10 ⁻⁰¹	1.05 × 10 ⁻⁰¹	1.68 × 10 ⁻⁰¹	1.61 × 10 ⁺⁰⁰	1.79 × 10 ⁺⁰⁰	1.73 × 10 ⁻⁰¹	1.19 × 10 ⁺⁰¹	5.67 × 10 ⁻⁰¹	2.22 × 10 ⁺⁰⁰	$1.38 \times 10^{+00}$	4.33 × 10 ⁻⁰¹
Cobalt	1.71	1.81×10^{-03}	5.74 × 10 ⁻⁰²	9.97 × 10 ⁻⁰⁴	3.50 × 10 ⁺⁰⁰	1.15 × 10 ⁺⁰⁰	2.77 × 10 ⁻⁰³	2.87 × 10 ⁻⁰²	5.13 × 10 ⁻⁰³	8.22 × 10 ⁻⁰³	7.90 × 10 ⁻⁰²	8.79 × 10 ⁻⁰²	2.32 × 10 ⁻⁰¹	2.19 × 10 ⁺⁰¹	$1.05 \times 10^{+00}$	4.09 × 10 ⁺⁰⁰	2.55 × 10 ⁺⁰⁰	7.99 × 10 ⁻⁰¹
Copper	2.91	7.18×10^{-02}	2.27 × 10 ⁺⁰⁰	3.94 × 10 ⁻⁰²	1.38 × 10 ⁺⁰²	4.57 × 10 ⁺⁰¹	1.10 × 10 ⁻⁰¹	1.14 × 10 ⁺⁰⁰	2.03 × 10 ⁻⁰¹	3.25 × 10 ⁻⁰¹	3.13 × 10 ⁺⁰⁰	3.48 × 10 ⁺⁰⁰	1.58 × 10 ⁺⁰⁰	2.38 × 10 ⁺⁰¹	$1.14 \times 10^{+00}$	4.45 × 10 ⁺⁰⁰	$2.77 \times 10^{+00}$	8.69 × 10 ⁻⁰¹
Molybdenum	0.63	1.55 × 10 ⁻⁰³	4.91 × 10 ⁻⁰²	8.54 × 10 ⁻⁰⁴	3.00 × 10 ⁺⁰⁰	9.90 × 10 ⁻⁰¹	2.37 × 10 ⁻⁰³	2.46 × 10 ⁻⁰²	4.40 × 10 ⁻⁰³	7.04 × 10 ⁻⁰³	6.77 × 10 ⁻⁰²	7.53 × 10 ⁻⁰²	8.54 × 10 ⁻⁰²	1.50 × 10 ⁺⁰⁰	7.15 × 10 ⁻⁰²	2.80 × 10 ⁻⁰¹	1.74 × 10 ⁻⁰¹	5.46 × 10 ⁻⁰²
Uranium	6.51	6.26 × 10 ⁻⁰³	1.98 × 10 ⁻⁰¹	3.44 × 10 ⁻⁰³	1.21 × 10 ⁺⁰¹	3.99 × 10 ⁺⁰⁰	9.55 × 10 ⁻⁰³	9.90 × 10 ⁻⁰²	1.77 × 10 ⁻⁰²	2.84 × 10 ⁻⁰²	2.73 × 10 ⁻⁰¹	3.03 × 10 ⁻⁰¹	6.09 × 10 ⁻⁰³	6.45 × 10 ⁺⁰¹	3.08 × 10 ⁺⁰⁰	1.20 × 10 ⁺⁰¹	7.49 × 10 ⁺⁰⁰	2.35 × 10 ⁺⁰⁰
Radionuclides																		
Lead-210	24.2	4.18 × 10 ⁻⁰²	1.32 × 10 ⁺⁰⁰	2.30 × 10 ⁻⁰²	8.06 × 10 ⁺⁰¹	2.66 × 10 ⁺⁰¹	6.38 × 10 ⁻⁰²	6.61 × 10 ⁻⁰¹	1.18 × 10 ⁻⁰¹	1.89 × 10 ⁻⁰¹	1.82 × 10 ⁺⁰⁰	2.02 × 10 ⁺⁰⁰	7.22 × 10 ⁻⁰¹	1.60 × 10 ⁺⁰¹	7.63 × 10 ⁻⁰¹	2.99 × 10 ⁺⁰⁰	1.86 × 10 ⁺⁰⁰	5.83 × 10 ⁻⁰¹
Polonium-210	0.91	1.12 × 10 ⁻⁰²	3.55 × 10 ⁻⁰¹	6.17 × 10 ⁻⁰³	2.16 × 10 ⁺⁰¹	7.15 × 10 ⁺⁰⁰	1.71 × 10 ⁻⁰²	1.77 × 10 ⁻⁰¹	3.17 × 10 ⁻⁰²	5.09 × 10 ⁻⁰²	4.89 × 10 ⁻⁰¹	5.44 × 10 ⁻⁰¹	6.17 × 10 ⁻⁰⁵	2.89 × 10 ⁺⁰¹	1.38 × 10 ⁺⁰⁰	5.39 × 10 ⁺⁰⁰	3.35 × 10 ⁺⁰⁰	1.05 × 10 ⁺⁰⁰
Radium-226	1.11	4.66 × 10 ⁻⁰³	1.47 × 10 ⁻⁰¹	2.56 × 10 ⁻⁰³	8.98 × 10 ⁺⁰⁰	2.96 × 10 ⁺⁰⁰	7.10 × 10 ⁻⁰³	7.36 × 10 ⁻⁰²	1.32 × 10 ⁻⁰²	2.11 × 10 ⁻⁰²	2.03 × 10 ⁻⁰¹	2.25 × 10 ⁻⁰¹	1.43 × 10 ⁻⁰³	4.40 × 10 ⁻⁰¹	2.10 × 10 ⁻⁰²	8.21 × 10 ⁻⁰²	5.11 × 10 ⁻⁰²	1.60 × 10 ⁻⁰²
Thorium-230	101	5.73 × 10 ⁻⁰²	1.81 × 10 ⁺⁰⁰	3.15 × 10 ⁻⁰²	1.11 × 10 ⁺⁰²	3.65 × 10 ⁺⁰¹	8.74 × 10 ⁻⁰²	9.06 × 10 ⁻⁰¹	1.62 × 10 ⁻⁰¹	2.60 × 10 ⁻⁰¹	2.50 × 10 ⁺⁰⁰	2.78 × 10 ⁺⁰⁰	8.62 × 10 ⁺⁰⁰	1.33 × 10 ⁺⁰¹	6.37 × 10 ⁻⁰¹	2.49 × 10 ⁺⁰⁰	1.55 × 10 ⁺⁰⁰	4.86 × 10 ⁻⁰¹

a) Table G.8 (CSA Group 2020).

b) For elements that are not listed in Table G.8, a surrogate element was used as follows (surrogate in parentheses): copper (silver), lead (tin), polonium (tellurium).

COPC = constituent of potential concern.

3.7.6 Dose Coefficients for Terrestrial Animals, Birds, and Humans

Dose coefficients for all internal and external exposure routes to terrestrial animals and birds are presented in Table 3-17 and Table 3-18, respectively. The DCFs for terrestrial animals and birds follow the approach of the ICRP (2008). Progeny with half lives shorter than 10 days are included in each DCF.

Table 3-17: Internal Dose coefficients for Terrestrial Animals and Birds

		DC	F internal (μG	y/h)/(Bq/kg	(fw animal))			
СОРС	Beaver ^(a)	Black Bear ^(b)	Canada Goose ^(c)	Grey Wolf ^(a)	Grouse ^(c)	Little Brown Myotis ^(a)	Loon ^(c)	Mallard ^(c)
Lead-210	2.4 × 10 ⁻⁰⁴	2.5 × 10 ⁻⁰⁴	2.5 × 10 ⁻⁰⁴	2.4 × 10 ⁻⁰⁴	2.5 × 10 ⁻⁰⁴	2.4 × 10 ⁻⁰⁴	2.5 × 10 ⁻⁰⁴	2.5 × 10 ⁻⁰⁴
Polonium-210	3.0 × 10 ⁻⁰²	3.0 × 10 ⁻⁰²	3.0 × 10 ⁻⁰²	3.0 × 10 ⁻⁰²	3.0 × 10 ⁻⁰²	3.0 × 10 ⁻⁰²	3.0 × 10 ⁻⁰²	3.0 × 10 ⁻⁰²
Radium-226	1.4 × 10 ⁻⁰¹	1.4 × 10 ⁻⁰¹	1.4 × 10 ⁻⁰¹	1.4 × 10 ⁻⁰¹	1.4 × 10 ⁻⁰¹	1.4 × 10 ⁻⁰¹	1.4 × 10 ⁻⁰¹	1.4 × 10 ⁻⁰¹
Thorium-230	2.7 × 10 ⁻⁰²	2.7 × 10 ⁻⁰²	2.7 × 10 ⁻⁰²	2.7 × 10 ⁻⁰²	2.7 × 10 ⁻⁰²	2.7 × 10 ⁻⁰²	2.7 × 10 ⁻⁰²	2.7 × 10 ⁻⁰²
Uranium-234	2.8 × 10 ⁻⁰²	2.8 × 10 ⁻⁰²	2.8 × 10 ⁻⁰²	2.8 × 10 ⁻⁰²	2.8 × 10 ⁻⁰²	2.8 × 10 ⁻⁰²	2.8 × 10 ⁻⁰²	2.8 × 10 ⁻⁰²
Uranium-238	2.4 × 10 ⁻⁰²	2.4 × 10 ⁻⁰²	2.4 × 10 ⁻⁰²	2.4 × 10 ⁻⁰²	2.4 × 10 ⁻⁰²	2.4 × 10 ⁻⁰²	2.4 × 10 ⁻⁰²	2.4 × 10 ⁻⁰²
СОРС	Mink ^(a)	Moose ^(b)	Muskrat ^(a)	Red Fox ^(a)	Rusty Black- bird ^(c)	Snowshoe Hare ^(a)	Southern Red-Backed Vole ^(a)	Woodland Caribou ^(b)
Lead-210	2.4 × 10 ⁻⁰⁴	2.5 × 10 ⁻⁰⁴	2.4 × 10 ⁻⁰⁴	2.4 × 10 ⁻⁰⁴	2.5 × 10 ⁻⁰⁴	2.4 × 10 ⁻⁰⁴	2.4 × 10 ⁻⁰⁴	2.5 × 10 ⁻⁰⁴
Polonium-210	3.0 × 10 ⁻⁰²	3.0 × 10 ⁻⁰²	3.0 × 10 ⁻⁰²	3.0 × 10 ⁻⁰²	3.0 × 10 ⁻⁰²	3.0 × 10 ⁻⁰²	3.0 × 10 ⁻⁰²	3.0 × 10 ⁻⁰²
Radium-226	1.4 × 10 ⁻⁰¹	1.4 × 10 ⁻⁰¹	1.4 × 10 ⁻⁰¹	1.4 × 10 ⁻⁰¹	1.4 × 10 ⁻⁰¹	1.4 × 10 ⁻⁰¹	1.4 × 10 ⁻⁰¹	1.4 × 10 ⁻⁰¹
Thorium-230	2.7 × 10 ⁻⁰²	2.7 × 10 ⁻⁰²	2.7 × 10 ⁻⁰²	2.7 × 10 ⁻⁰²	2.7 × 10 ⁻⁰²	2.7 × 10 ⁻⁰²	2.7 × 10 ⁻⁰²	2.7 × 10 ⁻⁰²
Uranium-234	2.8 × 10 ⁻⁰²	2.8 × 10 ⁻⁰²	2.8 × 10 ⁻⁰²	2.8 × 10 ⁻⁰²	2.8 × 10 ⁻⁰²	2.8 × 10 ⁻⁰²	2.8 × 10 ⁻⁰²	2.8 × 10 ⁻⁰²
Uranium-238	2.4 × 10 ⁻⁰²	2.4 × 10 ⁻⁰²	2.4 × 10 ⁻⁰²	2.4 × 10 ⁻⁰²	2.4 × 10 ⁻⁰²	2.4 × 10 ⁻⁰²	2.4 × 10 ⁻⁰²	2.4 × 10 ⁻⁰²

a) Dose coefficients are based on rat (ICRP 2008).

b) Dose coefficients are based on deer (ICRP 2008).

c) Dose coefficients are based on duck (ICRP 2008).

COPC = constituent of potential concern; DCF = dose coefficient.

Table 3-18: External Dose Coefficients for Terrestrial Animals and Birds

			DCF exteri	nal (µGy/h)/((Bq/m²)			
СОРС	Beaver ^(a)	Black Bear ^(b)	Canada Goose ^(c)	Grey Wolf ^(a)	Grouse ^(c)	Little Brown Myotis ^(a)	Loon ^(c)	Mallard ^(c)
Lead-210	6.3 × 10 ⁻⁰⁹	2.3 × 10 ⁻⁰⁹	5.8 × 10 ⁻⁰⁹	6.3 × 10 ⁻⁰⁹	5.8 × 10 ⁻⁰⁹	6.3 × 10 ⁻⁰⁹	5.8 × 10 ⁻⁰⁹	5.8 × 10 ⁻⁰⁹
Polonium-210	2.8 × 10 ⁻¹¹	1.4 × 10 ⁻¹¹	2.6 × 10 ⁻¹¹	2.8 × 10 ⁻¹¹	2.6 × 10 ⁻¹¹	2.8 × 10 ⁻¹¹	2.6 × 10 ⁻¹¹	2.6 × 10 ⁻¹¹
Radium-226	5.4 × 10 ⁻⁰⁶	2.8 × 10 ⁻⁰⁶	5.0 × 10 ⁻⁰⁶	5.4 × 10 ⁻⁰⁶	5.0 × 10 ⁻⁰⁶	5.4 × 10 ⁻⁰⁶	5.0 × 10 ⁻⁰⁶	5.0 × 10 ⁻⁰⁶
Thorium-230	1.3 × 10 ⁻⁰⁹	5.4 × 10 ⁻¹⁰	1.3 × 10 ⁻⁰⁹	1.3 × 10 ⁻⁰⁹	1.3 × 10 ⁻⁰⁹	1.3 × 10 ⁻⁰⁹	1.3 × 10 ⁻⁰⁹	1.3 × 10 ⁻⁰⁹
Uranium-234	7.5 × 10 ⁻¹⁰	2.8 × 10 ⁻¹⁰	7.1 × 10 ⁻¹⁰	7.5 × 10 ⁻¹⁰	7.1 × 10 ⁻¹⁰	7.5 × 10 ⁻¹⁰	7.1 × 10 ⁻¹⁰	7.1 × 10 ⁻¹⁰
Uranium-238	4.1 × 10 ⁻¹⁰	1.5 × 10 ⁻¹⁰	3.9 × 10 ⁻¹⁰	4.1 × 10 ⁻¹⁰	3.9 × 10 ⁻¹⁰	4.1 × 10 ⁻¹⁰	3.9 × 10 ⁻¹⁰	3.9 × 10 ⁻¹⁰
СОРС	Mink ^(a)	Moose ^(b)	Muskrat ^(a)	Red Fox ^(a)	Rusty Black- bird ^(c)	Snowshoe Hare ^(a)	Southern Red-Backed Vole ^(a)	Woodland Caribou ^(b)
Lead-210	6.3 × 10 ⁻⁰⁹	2.3 × 10 ⁻⁰⁹	6.3 × 10 ⁻⁰⁹	6.3 × 10 ⁻⁰⁹	5.8 × 10 ⁻⁰⁹	6.3 × 10 ⁻⁰⁹	6.3 × 10 ⁻⁰⁹	2.3 × 10 ⁻⁰⁹
Polonium-210	2.8 × 10 ⁻¹¹	1.4 × 10 ⁻¹¹	2.8 × 10 ⁻¹¹	2.8 × 10 ⁻¹¹	2.6 × 10 ⁻¹¹	2.8 × 10 ⁻¹¹	2.8 × 10 ⁻¹¹	1.4 × 10 ⁻¹¹
Radium-226	5.4 × 10 ⁻⁰⁶	2.8 × 10 ⁻⁰⁶	5.4 × 10 ⁻⁰⁶	5.4 × 10 ⁻⁰⁶	5.0 × 10 ⁻⁰⁶	5.4 × 10 ⁻⁰⁶	5.4 × 10 ⁻⁰⁶	2.8 × 10 ⁻⁰⁶
Thorium-230	1.3 × 10 ⁻⁰⁹	5.4 × 10 ⁻¹⁰	1.3 × 10 ⁻⁰⁹	1.3 × 10 ⁻⁰⁹	1.3 × 10 ⁻⁰⁹	1.3 × 10 ⁻⁰⁹	1.3 × 10 ⁻⁰⁹	5.4 × 10 ⁻¹⁰
Uranium-234	7.5 × 10 ⁻¹⁰	2.8 × 10 ⁻¹⁰	7.5 × 10 ⁻¹⁰	7.5 × 10 ⁻¹⁰	7.1 × 10 ⁻¹⁰	7.5 × 10 ⁻¹⁰	7.5 × 10 ⁻¹⁰	2.8 × 10 ⁻¹⁰
Uranium-238	4.1 × 10 ⁻¹⁰	1.5 × 10 ⁻¹⁰	4.1 × 10 ⁻¹⁰	4.1 × 10 ⁻¹⁰	3.9 × 10 ⁻¹⁰	4.1 × 10 ⁻¹⁰	4.1 × 10 ⁻¹⁰	1.5 × 10 ⁻¹⁰

a) Dose coefficients are based on rat (ICRP 2008).

COPC = constituent of potential concern; DCF = dose coefficient.

Dose coefficients for all internal and external exposure routes for humans are presented in Table 3-19. The DCFs for ingestion and inhalation by human receptors were taken from CSA N288.1-20 and are progeny inclusive.

The external DCFs used in this IMPACT model update were derived based on the methodology described in CSA N288.1-20. Uranium-238 progeny associated with each of the radionuclides, as required, were included in the external DCFs as follows:

- uranium-238 with its progeny thorium-234 and protactinium-234m;
- uranium-234 was modelled explicitly and no progeny were included in the external DCF;
- thorium-230 was modelled explicitly and no progeny were included in the external DCF;
- radium-226 with its progeny radon-222 (except for air immersion), polonium-218, lead-214, bismuth-214, and polonium-214;
- lead-210 with its progeny bismuth-210; and



b) Dose coefficients are based on deer (ICRP 2008).

c) Dose coefficients are based on duck (ICRP 2008).

• radon-222 (air immersion only) with its progeny polonium-218, lead-214, and bismuth-214.

The achievement of transient or secular equilibrium by ingrowth of progeny takes time. Therefore, the available time for ingrowth in each exposure medium must be considered in deciding whether it is reasonable to assume equilibrium. The timeframes for ingrowth of the progeny assumed for each of the external pathways were two hours for air immersion, one day for water immersion, and 40 years for exposure to soils and sediment. Details for the calculation of the external DCFs are described in CSA N288.1-20.

Table 3-19: Human Dose Coefficients for Internal and External Exposure

СОРС	Ingestion DCF	Inhalation DCF	External Soil DCF	External Water Immersion DCF	External Sediment DCF	External Air Immersion DCF
	(Sv/Bq)	(Sv/Bq)	(Sv/yr)/ (Bq/m²)	(Sv/yr)/ (Bq/L)	(Sv/yr)/ (Bq/kg)	(Sv/yr)/ (Bq/m³)
			Adult			
Lead-210	6.9 × 10 ⁻⁰⁷	1.1 × 10 ⁻⁰⁶	1.2×10^{-09}	4.5 × 10 ⁻⁰⁹	1.8×10^{-09}	1.4×10^{-09}
Polonium-210	1.2 × 10 ⁻⁰⁶	3.3 × 10 ⁻⁰⁶	2.6×10^{-13}	2.7 × 10 ⁻¹¹	7.4 × 10 ⁻¹²	1.2 × 10 ⁻¹¹
Radium-226	2.8 × 10 ⁻⁰⁷	3.5 × 10 ⁻⁰⁶	5.3 × 10 ⁻⁰⁸	9.2 × 10 ⁻⁰⁷	1.5 × 10 ⁻⁰⁶	9.0 × 10 ⁻⁰⁹
Thorium-230	2.1 × 10 ⁻⁰⁷	1.4 × 10 ⁻⁰⁵	2.4 × 10 ⁻¹¹	1.1 × 10 ⁻⁰⁹	2.3 × 10 ⁻¹⁰	4.7 × 10 ⁻¹⁰
Uranium-234	4.9 × 10 ⁻⁰⁸	3.5 × 10 ⁻⁰⁶	1.9 × 10 ⁻¹¹	4.4 × 10 ⁻¹⁰	7.8 × 10 ⁻¹¹	1.9 × 10 ⁻¹⁰
Uranium-238	4.5 × 10 ⁻⁰⁸	2.9 × 10 ⁻⁰⁶	3.7 × 10 ⁻⁰⁹	2.5 × 10 ⁻⁰⁹	2.2 × 10 ⁻⁰⁸	7.9 × 10 ⁻¹¹
		1-	year old			
Lead-210	3.6 × 10 ⁻⁰⁶	3.7 × 10 ⁻⁰⁶	1.5×10^{-09}	4.8 × 10 ⁻⁰⁹	2.3×10^{-09}	1.8×10^{-09}
Polonium-210	8.8 × 10 ⁻⁰⁶	1.1 × 10 ⁻⁰⁵	3.3×10^{-13}	3.5 × 10 ⁻¹¹	9.6 × 10 ⁻¹²	1.6 × 10 ⁻¹¹
Radium-226	9.6 × 10 ⁻⁰⁷	1.1 × 10 ⁻⁰⁵	6.9 × 10 ⁻⁰⁸	1.2 × 10 ⁻⁰⁶	2.0 × 10 ⁻⁰⁶	1.2 × 10 ⁻⁰⁸
Thorium-230	4.1 × 10 ⁻⁰⁷	3.5 × 10 ⁻⁰⁵	2.6 × 10 ⁻¹¹	1.4 × 10 ⁻⁰⁹	3.0 × 10 ⁻¹⁰	6.1 × 10 ⁻¹⁰
Uranium-234	1.3 × 10 ⁻⁰⁷	1.1 × 10 ⁻⁰⁵	2.4 × 10 ⁻¹¹	5.7 × 10 ⁻¹⁰	1.0 × 10 ⁻¹⁰	2.5 × 10 ⁻¹⁰
Uranium-238	1.2 × 10 ⁻⁰⁷	9.4 × 10 ⁻⁰⁶	4.8 × 10 ⁻⁰⁹	3.3 × 10 ⁻⁰⁹	2.8 × 10 ⁻⁰⁸	1.0 × 10 ⁻¹⁰

COPC = constituent of potential concern; DCF = dose coefficient; Sv/yr = sieverts per year; Bq = becquerel.

4.0 References

- Arnold TW, Fritzell EK. 1987. Food habits of prairie mink during the waterfowl breeding season. Canadian Journal of Zoology. 65, 2322–2324.
- Beresford NA, Barnett CL, Howard BJ, Scott WA, Brown JE, Copplestone D. 2008. Derivation of transfer parameters for use within the ERICA Tool and the default concentration ratios for terrestrial biota. Journal of Environmental Radioactivity. 99 1393-1407.
- Beyer WN, Connor E, Gerould S. 1994. Estimates of Soil Ingestion by Wildlife. Journal of Wildlife Management 58.
- Brown JE, Alfonso B, Avila R, Beresford NA, Copplestone D, Prohl G, Ulanovsky A. 2008. The ERICA Tool. Journal of Environmental Radioactivity. 99 1393-1407 99, 1371–1383. September.
- Cameco. 2004. Cigar Lake Project: Construction, Operation, and Decommissioning Environmental Assessment Study Report.
- CanNorth. 2000. Hatchet Lake Dietary Survey. Prepared for: Atomic Energy Control Board, Ottawa ON. March 2000.
- Chan L, Receveur O, Sadik T, Schwartz H, Ing A, Fediuk K, Tikhonov C (University of Ottawa). 2018. First Nations Food, Nutrition and Environment Study (FNFNES): Results from Saskatchewan (2015).
- Chapman PM, Allen HE, Z'Graggen MN. 1996. Evaluation of bioaccumulation factors in regulating metals. Environmental Science and Technology. 30, 448A-452A.
- COSEWIC (Committee on the Status of Endangered Wildlife in Canada). 2002. COSEWIC assessment and update status report on the woodland caribou *Rangifer tarandus* caribou in Canada.
- COSEWIC. 2006. Rusty blackbird (*Euphagus carolinus*): COSEWIC assessment and status report 2017.
- CSA Group (Canadian Standards Association Group). 2022. CSA N288.6-22: Environmental Risk Assessments at Nuclear Facilities and Uranium Mines and Mills.
- CSA Group. 2020. CSA N288.1-20: Guidelines for Calculating Derived Release Limits for Radioactive Material in Airborne or Liquid Effluents for Normal Operation of Nuclear Facilities.
- Earthroots. 2020. Wolves in Ontario. Available at http://wolvesontario.org/wolves-ontario/.
- Ecometrix (Ecometrix Incorporated). 2005. Key Lake Project: Environmental Risk Assessment 2005. Report prepared for Cameco Corporation.
- Ecometrix. 2012. Key Lake IMPACT model 2011 Update. Report prepared for Cameco Corporation.



- Ecometrix. 2013a. Key Lake Extension Project Ecological and Human Health Risk Assessment. Report prepared for Cameco Corporation. In: Cameco. 2013. Key Lake Extension Project Environmental Impact Statement, Annex VII.
- Ecometrix. 2013b. Millennium Project Ecological and Human Health Risk Assessment. Report prepared for Cameco Corporation. In: Cameco. 2013. Millennium Project Environmental Impact Statement, Annex J.
- Ecometrix. 2013c. Millennium IMPACT model 2012. Report prepared for Cameco Corporation. In: Cameco. 2013. Millennium Project Environmental Impact Statement, Annex K.
- FCSAP (Federal Contaminated Sites Action Plan). 2012. Ecological Risk Assessment Guidance, Module 3: Standardization of Wildlife Receptor Characteristics. Prepared for: Environment Canada (Government of Canada), Vancouver, BC. March.
- Franzmann AW. 1981. Alces alces. Mammalian Species. 154:1-7. May.
- Hamilton SJ, Mehrle PM. 1986. Metallothionein in fish: review of its importance in assessing stress from metal contaminants. Transactions of the American Fisheries Society 115, 596–609.
- Hart D. 2013. Derived Release Limits. Report No. COG-06-3090-R3-I.
- Health Canada. 2010a. Federal Contaminated Site Risk Assessment in Canada, Part I: Guidance on Human Health Preliminary Quantitative Risk Assessment (PQRA), Version 2.0. Contaminated Sites Program. September.
- Health Canada. 2010b. Federal Contaminated Site Risk Assessment in Canada. Part VI: Guidance on Human Health Detailed Quantitative Radiological Risk Assessment (DQRARAD). September.
- Health Canada. 2011. Food and Nutrition. Dietary Intakes of Contaminants and other Chemicals for Different Age-sex Groups of Canadians.
- Health Canada. 2018. Guidance for Evaluating Human Health Impacts in Environmental Assessment: Country Foods. Report No. H129- 54/5-2018. May.
- Hinterland Who's Who. 1993. Profile on the Red Fox. Available at http://www.hww.ca/en/wildlife/mammals/red-fox.html.
- Hinterland Who's Who. 2007. Profile on the Black Bear. Available at http://www.hww.ca/en/wildlife/mammals/black-bear.html.
- Hinterland Who's Who. 2017. Profile on the Beaver. Available at http://www.hww.ca/en/wildlife/mammals/beaver.html.
- IAEA (International Atomic Energy Agency). 2010. Handbook of Parameter Values for the Prediction of Radionuclide Transfer in Terrestrial and Freshwater Environments. Technical Report Series No. 472.
- ICRP (International Commission on Radiological Protection). 2008. Environmental Protection the Concept and Use of Reference Animals and Plants. ICRP Publication 108.



- Jones DM, Theberge JB. 1981. Summer home range and habitat utilisation of the red fox (*Vulpes vulpes*) in a tundra habitat, northwest British Columbia. Canadian Journal of Zoology. 804–812.
- McLoughlin PD, Stewart K, Superbie C, Perry T, Tomchuk P, Greuel R, Singh K, Truchon-Savard A, Henkelman J, Johnstone JF (Department of Biology, University of Saskatchewan). 2016. Population dynamics and critical habitat of woodland caribou in the Saskatchewan Boreal Shield. Interim Project Report, 2013-2016.
- McLoughlin PD, Superbie CM, Stewart K, Tomchuk P, Neufeld B, Barks D, Perry T, Greuel R, Regan C, Truchon-Savard A, Hart S, Henkelman J, Johnstone JF (Department of Biology, University of Saskatchewan). 2019. Population and habitat ecology of boreal caribou and their predators in the Saskatchewan Boreal Shield. Final Report.
- NCRP (National Council on Radiation Protection and Measurements). 1996. Screening Models for Releases of Radionuclides to Atmosphere, Surface Water, and Ground Vol. I and Vol. II. Report No. 123.
- Newman MC, Unger MA. 2002. Fundamentals of Ecotoxicology, 2nd ed. CRC/Lewis Press, Boca Raton, FL.
- RIC (Ministry of the Environment, Lands and Parks Research for the Terrestrial Ecosystems Task Force Resources Inventory Committee). 1998. Inventory Methods for Wolf and Cougar: Standards for Components of British Columbia's Biodiversity. Report No. 34.
- Richardson GM. 1997. Compendium of Canadian Human Exposure Factors for Risk Assessment. O'Connor Associates.
- Sample BE, Suter GWI (Oak Ridges National Laboratory). 1994. Estimating Exposure of Terrestrial Wildlife to Contaminants. Report No. ES/ER/TM-125. September.
- Sample BE, Beauchamp JJ, Efroymson RA, Suter GW, Ashwood TL. 1998. Development and Validation of Bioaccumulation Models for Earthworms. Report No. ES/ER/TM-220.
- Species at Risk Act. SC. 2002, c 29. Last amended 12 August 2021. Available at: https://laws-lois.justice.gc.ca/eng/acts/s-15.3/.
- Thomann RV, Mueller JA. 1987. Principles of Surface Water Quality Modeling and Control. Harper & Row, Publishers, New York.
- Thompson, S.E., C.A. Burton, D.J. Quinn and Y.C. Ng. 1972. Concentration Factors of Chemical Elements in Edible Aquatic Organisms. Lawrence Livermore Laboratory, University of California. Rep. UCRL-50564 Rev. 1.
- UNSCEAR (United Nations Scientific Committee on the Effects of Atomic Radiation). 2000. Volume 1, Annex B: Exposures from Natural Radiation Sources and Effects of Ionizing Radiation, Report to the General Assembly with Scientific Annexes.
- USEPA (United States Environmental Protection Agency). 1986. Standard for Radon-222 Emissions from Licensed Uranium Mill Tailings. Report No. EPA 520/1-86-009.
- USEPA. 1993. Wildlife Exposure Factors Handbook. Report No. EPA/600/R-93/187.



- Wilson D, Ruff S. 1999. The Smithsonian Book of North American Mammals. Smithsonian Institution Press, Washington.
- Wood CM, Port P. 2000. Intracellular pH regulation and buffer capacity in CO₂/HCO₃-buffered media in cultured epithelial cells from rainbow trout gills. Journal of Comparative Physiology. 170, 175–184.

Appendix A Water and Sediment Sampling Stations

Label ID	· · · · · · · · · · · · · · · · · · ·
BeetC	Beet Creek
BeetL	Beet Lake
CR	Clearwater River
FL	Forrest Lake
JedC	Jed Creek
LG	Lake G
LH	Lake H
LLI	Lloyd Lake Inlet
LLO	Lloyd Lake Outlet

Name

Label ID

Mid Mid-field MR Mirror River

PLNW Patterson Lake North Arm West Basin
PLNE Patterson Lake North Arm East Basin

PLS Patterson Lake South Arm

The location of sampling stations is described in detail in Annex V.1, which describes the baseline studies performed between 2018 and 2020.

Appendix B Ecological Receptor Profiles



APPENDIX B – ECOLOGICAL RECEPTOR PROFILES

One of the key considerations, which defines the scope of a risk assessment is the selection of ecological receptors. In selecting ecological receptors, it is important to identify plants and animals that are likely to be most exposed to the effects of the Project. As it is not possible to evaluate all ecological species at a site, representative species were selected based on the selection process provided in Table 7.1 of CSA N288.6 (2022).

This appendix describes the aquatic and terrestrial ecological receptors (groups or species) selected for the assessment. The general characteristics of each receptor including any status under COSEWIC, SARA or provincial ranking, its distribution and presence in the Regional Study Area (RSA), its home range and diet, as well as how it is incorporated into the ecological and/or human health models and the protection level that it is afforded in the risk assessment.

1.0 Aquatic and Riparian Biota

1.1 Aquatic Vegetation

1.1.1 Macrophytes

Macrophytes are aquatic plants growing in or near water and can be either emergent, submergent or floating. Emergent macrophytes are rooted in shallow water whereas submergent macrophytes are typically rooted in sediment and the entire plant is submerged. Macrophytes are primary producers that provide food, cover, and shelter for wildlife, such as spawning and nursery habitats for fish and nesting habitats for waterfowl, improve water quality and clarity, and help to stabilize shorelines and bottom sediments.

Dense amounts of submergent macrophytes including various pondweeds were found in the Clearwater River upstream of Mirror River, while downstream of Mirror River, emergent macrophytes were limited to areas close to the shoreline. Submergent macrophytes were also sampled at Lloyd Lake as part of baseline studies. A general lack of emergent macrophytes were noted in Patterson Lake, and for this reason the lake has been characterized as only marginally suitable for northern pike spawning (CanNorth 2021a).

In the ecological model, macrophytes are exposed to aquatic release through surface water, and they provide food for woodland caribou, moose, muskrat, and mallard. Because of their importance as a potential link between contaminants in sediments and surface water to other aquatic and terrestrial receptors, macrophytes were conservatively assumed to be present in every surface water body within the RSA.

In the ERA, macrophytes are assessed as a group at a community level.



1.1.2 Phytoplankton

Phytoplankton are small primary producers that use sunlight to produce oxygen and nutrients for other organisms. Phytoplankton are an important food sources for organisms in an aquatic environment.

Phytoplankton are present in all surface water bodies in the RSA. The phytoplankton communities in Patterson Lake North, Forrest Lake, Beet Lake, and Naomi Lake, Broach Lake, and Hodge Lake were sampled in fall 2018 (CanNorth 2021a). The phytoplankton communities were dominated by blue-green algae (Cyanophycota) primarily of the families Synechococcaceae and Chroococcaeae. Diatoms (Bacillariophyta) were also an important component of the phytoplankton communities of Beet and Naomi lakes.

In the model, phytoplankton are exposed to aquatic release through surface water. Phytoplankton communities represent a potential link between contaminants in surface water and other aquatic and terrestrial receptors.

In the ERA, phytoplankton are assessed as a group at a community level.

1.2 Aquatic Invertebrates

1.2.1 Epifaunal and Infaunal Benthic Invertebrates

Benthic invertebrates or "benthos" live on or within the sediments of surface water bodies. Benthic invertebrates play an integral role in the fresh water ecosystem through nutrient cycling and they function as an important food source for wildlife such as the diving (e.g. Bufflehead) and dabbling (e.g. Mallard) ducks, and fish (e.g. White Sucker). Benthic invertebrates represent a relatively large fraction of biomass in Patterson Lake and other lakes in the RSA (CanNorth 2021a).

Epifaunal benthic invertebrates live in water bodies on or attached to submerged substrates such as the bottom sediment or hard surfaces and feed at the substrate surface. Epifaunal benthic invertebrates primarily include amphipods, bivalves, shrimps, crabs, snails, and some aquatic insects. Amphipoda (*Gammarus* and *Hyalella*, also called scuds and fresh water shrimp) are an important group of epifaunal invertebrates in the RSA depending on the area (CanNorth 2021a). Ephemeroptera (mayfly nymphs) form a representative group of epifaunal benthic invertebrates generally found throughout the RSA that represent a relatively large fraction of the biomass of aquatic invertebrates (CanNorth 2021a).

Infaunal benthic invertebrates live and burrow in the sediments and feed within sediments. Infaunal benthic invertebrates primarily include bivalves, worms, and some aquatic insects. Chiromomids (non-biting midge larvae) was the most abundant group of infaunal benthic invertebrates found in the RSA, followed by Oligochaeta (aquatic earthworms and sludge worms), and Mollusca (other than Unionidae [river mussels]), depending on the area (CanNorth 2021a).

In the model, benthic invertebrates are exposed to aquatic release through surface water and sediment. Benthic invertebrates provide water to fish and sediment to fish pathway links and a



link between aquatic and terrestrial ecosystems. Many benthic invertebrate species feed on decaying organic matter and thereby form an important link between the decomposer and primary consumer levels.

In the ERA, benthic invertebrates are assessed as a group at a community level.

1.2.2 Zooplankton

Zooplankton are small aquatic animals that are intermediaries in the aquatic food chain. Zooplankton are primary or secondary consumers that are also prey of other larger zooplankton predators or fish. Exposed to aquatic release through surface water.

Zooplankton are present in all surface water bodies in the RSA. The zooplankton communities in Patterson Lake North, Forrest Lake, Beet Lake, and Naomi Lake, Broach Lake, and Hodge Lake were sampled in fall 2018 (CanNorth 2021a). Zooplankton communities consisted of mainly crustaceans (Cyclopoida) and rotifers (Ploima). Other crustaceans (Cladocera) were also abundant in some areas.

In the model, zooplankton are exposed to aquatic release through surface water. Zooplankton communities represent a potential link between contaminants in surface water and phytoplankton, and other aquatic and terrestrial receptors.

In the ERA, zooplankton are assessed as a group at a community level.

1.3 Fish

1.3.1 Lake Whitefish

Lake whitefish (*Coregonus culpeaformis*) is a benthic cool water fish species of the family Salmonidae, that is found in fresh water lakes throughout much of Canada. It is a species of ecological and economic value. Regionally, the Other Land and Resource Use Memorandum (InterGroup 2018b) identified lake whitefish as a target of commercial fishing in Patterson Lake, Lloyd Lake, and Lake La Loche. Lake whitefish are also an important component of the traditional foods diet for First Nations in Saskatchewan (Chan et al. 2018, 2019).

Fish community surveys that targeted lake whitefish were conducted in 2018 at Broach, Hodge, G, H, and Patterson lakes, Lloyd Lake Inlet and the midfield area of the Clearwater River, and in 2019 at Broach Lake, Jed Creek, Patterson Lake, Patterson Creek, Forrest Lake, Beet Channel, Beet Lake, Beet Creek, Naomi Lake, Naomi Creek, Clearwater Creek, and Clearwater Nearfield. Lake whitefish were found to be present in many of these waterbodies, and were found to spawn in Patterson, Forrest and Beet lakes and Patterson Creek (CanNorth 2021a). Lake whitefish tissue samples were retained for chemical analysis. It is assumed that the lake whitefish spends 100% of its time in the Project area waterbody where it is assessed.

Lake whitefish ingest plankton during their larval and post larval stages then switch to benthic invertebrates, including snails, bivalves and insect larvae, and fish eggs. In the model, lake



whitefish are exposed to aquatic release through surface water and sediments. In the human health model, they are a component of the Traditional Foods diet.

In the ERA, lake whitefish are assessed as a representative benthopelagic fish species at a population level.

1.3.2 Northern Pike

Northern pike (*Esox lucius*) is a pelagic top predator fish, of the family Esicidae, that is widely distributed in fresh water environments across Canada. It is of ecological and economic importance. Regionally, the Other Land and Resource Use Memorandum (InterGroup 2018b) identified northern pike as a target of commercial fishing in Patterson Lake, Lloyd Lake, and Lake La Loche. Northern pike is also an important component of the traditional foods diet for First Nations in Saskatchewan (Chan et al. 2018, 2019).

Fish community surveys that targeted northern pike were conducted in 2018 at Broach, Hodge, G, H, and Patterson lakes, Lloyd Lake Inlet and the midfield area of the Clearwater River, and in 2019 at Broach Lake, Jed Creek, Patterson Lake, Patterson Creek, Forrest Lake, Beet Channel, Beet Lake, Beet Creek, Naomi Lake, Naomi Creek, Clearwater Creek, and Clearwater Nearfield. Northern Pike were found to be present in all the waterbodies. Northern pike were found to spawn in Patterson, Forrest, Naomi lakes, Patterson, Jed, Beet, Naomi, Clearwater creeks, and Clearwater River (CanNorth 2021a). Northern pike tissue samples were retained for chemical analysis. It is assumed that the northern pike spends 100% of its time in the Project area waterbody where it is assessed.

Northern pike are exposed to aquatic release through surface water and consumption of prey (aquatic vertebrates and/or aquatic invertebrates). In the model, northern pike are exposed to aquatic release through surface water. Northern pike represent a potential link between contaminants in surface water, other aquatic and terrestrial receptors, and human receptor groups. In the ecological model, northern pike are consumed by bear, mink, and loon. In the human health model, they are a component of the human Traditional Foods diet and are used as a surrogate for other common predatory fish, such as walleye, which are also items in the human Traditional Foods diet.

In the ERA, northern pike are assessed as a representative pelagic predator fish species at a population level.



1.4 Riparian Mammals

1.4.1 Beaver

North American beaver (*Castor canadensis*) is a large rodent of the family Castoridae, measuring approximately 1.3 m including its large tail, and weighing about 20 kg. They are found throughout much of North America from southwest Alaska to northeast Mexico except for western California (EC & CWF 2017). Beavers are present in the RSA and are trapped for fur and meat and harvested by Indigenous people (Omnia 2020a). Beavers are an important component of the traditional foods diet for First Nations living in the Boreal Shield ecoregion of Saskatchewan (Chan et al. 2018, 2019).

Beavers live in wetland areas engineering their habitat through the construction of dams and channels to create suitable habitat with water deep enough not to freeze to the bottom in winter. Beavers mate in winter and produce 3 or 4 kits by May or June after a 100-day gestation period. For the ecological model, it is assumed that the beaver spends 100% of its time at the location where it is assessed.

Beavers consume a woody diet (aspen, poplar, willow, birch) during the winter from submerged caches prepared during the open water season. If the area is relatively free of predators, beavers may travel 125 m away in search of suitable winter food. During the other seasons beaver switch to an herbaceous diet of grasses, herbs, twigs, fruits, and aquatic plants. For the ecological model it is assumed that the beaver's diet consists of browse (90%) and macrophytes (10%).

In the model, beavers are exposed to aquatic release through water, food (terrestrial and aquatic vegetation) and sediment. In the ecological model, beavers are prey for larger predators such as wolves. In the human health model, beavers are a component of the First Nation's traditional foods diet and are used as a surrogate for other small mammals, such as hare.

In the ERA, beavers are assessed as a representative riparian herbivore species at a population level.

1.4.2 Muskrat

The muskrat (*Ondatra zibethicus*) is a large rodent (Muridae), measuring approximately 50 cm from tip of the nose to tail, and weighing on average 1 kg. Muskrats exist all over North America, from the Arctic Ocean in the north to the Gulf of Mexico in the south, from the Pacific Ocean in the west to the Atlantic Ocean in the east. Their mean home range size in the summer is between 0.048 to 0.17 hectares (USEPA 1993). Muskrats are present in the RSA and are trapped for fur and meat and harvested by Indigenous people (Omnia 2020a).

Muskrats prefer fresh water marshes, marshy areas of lakes, and slow-moving streams. The preferred water depth in these areas is 1 to 2 m, deep enough not to freeze fully during the winter but shallow enough to allow aquatic vegetation to grow. Muskrats nest in compact mounds of partially dried and decayed plant material such as cattails and bulrushes. In winter, muskrats generally occupy lodges that they build through burrowing underneath their mounds, and spends



most of its time eating and sleeping. They are capable of extended dives, allowing them to dig channels and burrows underwater, cut submerged stems and roots with their teeth, and travel long distances beneath the ice (EC & CWF 1986). Breeding generally occurs in March, April, or May. Birth of the litter usually occurs within 1 month of mating and usually contains 5 to 10 young. Breeding can occur multiple times throughout the season (EC & CWF 1986). For the ecological model, it is assumed that the muskrat spends 100% of its time at the location where it is assessed.

Muskrats mainly feed on aquatic plants such as cattails, bulrushes, horsetails, or pondweeds; although they prefer cattails. When aquatic plants are unavailable, muskrats are also known to feed on fish, frogs, and clams. In the model, muskrats are exposed to aquatic release through water, food, and sediment. For the ecological model it is assumed that the muskrat's diet consists of aquatic plants (80%) and benthic invertebrates (20%).

In the ERA, muskrats are assessed as a representative riparian herbivore species at a population level.

1.4.3 American Mink

American mink (*Neovison vison*) is a semi-aquatic, carnivorous mammal belonging to the weasel family (Mustelidae), measuring approximately 30 to 40 cm not including the tail and weighing on average 1 kg. Minks are widely distributed throughout North America, except in the far north, arid areas in southwestern United States, and Mexico. Minks are present in the RSA and are trapped for fur and harvested by Indigenous people (Omnia 2020a).

Preferred habitats for minks generally consist of aquatic habitats such as rivers, lakes, streams, marshes, and swamps, with wooded cover. Mink typically use bank burrows of other animals such as muskrats, cavities in tree roots, rock or brush piles, log jams, and beaver lodges for denning instead of constructing their own (USEPA 1993). Minks have been known to travel up to 200 m away from a water body and males have the largest home ranges of up to 10 km² (USEPA 1993). For the ecological model, it is assumed that the mink spends 100% of its time at the location where it is assessed.

Minks are nocturnal opportunistic hunters, consuming a variety of prey around shorelines or emergent vegetation, preferring small mammals such as voles, muskrats, and shrews. Other secondary prey items include aquatic birds, fish, amphibians, and crustaceans (USEPA 1993). For the ecological model it is assumed that the muskrat's diet consists of small mammals, represented by muskrat (45%); fish, represented by northern pike (30%); benthic invertebrates (15%) and aquatic birds; represented by mallard (10%).

In the ERA, minks are assessed as a representative riparian carnivore species at a population level.



1.5 Riparian Birds

1.5.1 Mallard

The mallard (*Anas platyrhynchos*) is an omnivorous migratory duck (Anatidae) that has a summer breeding range in Canada (USEPA 1993; FCSAP 2012). Male mallards average 1.1 kg in weight and females about 1.2 kg in weight (FCSAP 2012). Mallards are common to northern Saskatchewan and are known to breed in the RSA (Omnia 2020a).

Mallards are an important component of the traditional foods diet for First Nations living in the Boreal Shield and Boreal Plains ecozones of Saskatchewan (Chan et al. 2018, 2019).

The general habitat of the mallard is wetlands. The mean home range of a mallard in spring is between 111 and 620 hectares (USEPA 1993). Mallard typically nest on the ground in thick vegetation away from a waterbody. Mating pairs may produce more than one clutch per season, with the first clutch generally being finished by in late Spring, in the northern United States (USEPA 1993). Females remain with their brood until fledging (USEPA 1993). Although mallards are not present in the vicinity of the Project for over half the year due to migration, they have a small home range while nesting and rearing of their young. For the ecological model, it is conservatively assumed that the mallard spends 100% of its time at the location where it is assessed.

The mallard forages by dabbling and filtering though sediment (USEPA 1993). The bulk of the mallard's diet is plant material, mostly aquatic plants and seeds, with the remaining portions of the diet consisting of aquatic invertebrates, especially during the breeding season (FCSAP 2012). For the ecological model it is assumed that the mallard's diet while in the RSA consists of benthic invertebrates (75%) and aquatic plants (25%).

In the model, mallards are exposed to aquatic release through water, food, and sediment. In the ecological model, mallards are prey for mink. In the human health model, mallards are a component of the First Nation's traditional foods diet and are used as a surrogate for other water birds with a similar diet.

In the ERA, mallards are assessed as a representative riparian avian species at a population level.

1.5.2 Common Loon

The loon (*Gavia immer*) is a piscivorous migratory waterfowl of the family Gaviidae, that breeds in Canada during the summer (FCSAP 2012). Male loons average 5.3 kg in weight and females about 4.7 kg in weight (FCSAP 2012). Loons are common to northern Saskatchewan and are known to breed in the RSA (Omnia 2020a).

Loons commonly inhabit open water areas of lakes and rivers during the summer (FCSAP 2012). Loons do not walk well on land so they spend their time on water. They prefer to nest offshore, on islands, islets, or floating mounds of vegetation in shallow water. Loon pairs may reuse the same nesting site year after year. They have one brood per year with 1 or 2 chicks being produced. Eggs hatch after an incubation period of 26-29 days (Cornell Lab (no date)). For the ecological



model, it is conservatively assumed that the loon spends 100% of its time at the location where it is assessed.

Loons surface dive to the water column to catch fish. Loons are opportunistic feeders but appear to prefer perch to salmon (FCSAP 2012). Aquatic invertebrates and crayfish are significant secondary components of their diet especially if the water is not clear. For the ecological model, loons are assumed to eat 90% fish (northern pike) and 10% aquatic invertebrates.

In the model, loons are exposed to aquatic release through water and food.

In the ERA, loons are assessed as a representative avian piscivorous species at a population level.

2.0 Terrestrial Biota

2.1 Terrestrial Invertebrates

Terrestrial invertebrates are considered to be primary consumers. They are represented by earthworms, which live in soil and are therefore exposed to constituents in soil layers through direct contact. Earthworms live in soil and depending on the species they either move vertically or horizontally indifferent soil layers. Earthworms acquire their nutrition through the organic matter in soil as well as the decomposing remains of other animals. They can devour one third of their own body weight per day.

For the ecological model, terrestrial invertebrates represented by the earthworm are assumed to spend 100% of their time at the location where they are assessed.

In the model, terrestrial invertebrates are exposed to atmospheric release through deposition to soil. Terrestrial invertebrate communities represent a potential link between contaminants in air and other terrestrial receptors. In the ecological model, terrestrial invertebrates are part of the diet for the red fox (*Vulpes vulpes*).

In the ERA, terrestrial invertebrates are assessed as a group at a community level.

2.2 Terrestrial Vegetation

Terrestrial vegetation types are dietary components for terrestrial animals and humans. They are represented by browse (various shrubs and grasses), lichen (various species), blueberries (*Vaccinium myrtilloides*), and Labrador tea (*Rhododendron groenlandicum*).

Individual plant species are included as food for other ecological receptors and for human receptors in the ecological and human health models. Three individual species are assessed in the model as representative species for lichen, fruits, and berries (blueberry), and leafy plants and browse (Labrador tea).

Plants are immobile and therefore are assumed to spend 100% of their time at the location where they are assessed.



In the ERA terrestrial vegetation is assessed as a group at a community level.

2.2.1 Lichen

Lichens are not plants, but rather they are composite organisms and symbiotic partnerships between fungus and algae. Some species are arboreal and grow on branches of higher plants, while others are terrestrial and grow on rocks and soil. The fungus provides structure and protection to the algae, while the algae which has chlorophyll, photosynthesizes and provides nutrition to the fungus. Lichens are found around the world in many different habitats.

Lichens are present in the RSA, especially in regenerating and recently burned jackpine stands that are common in the SSA (CanNorth 2021c). The forest floors of these communities have a greater proportion of lichens than mosses compared to the areas of mature jackpine stands that are also present in the RSA. Due to recurrent burn cycles in the area the dominant lichens are terrestrial species. Reindeer lichen is a dominant groundcover in regenerating coniferous forests in the area (Omnia 2020b).

In the model, lichen is exposed to atmospheric release depositing from air. In the ecological model lichens are food for woodland caribou.

2.2.2 Blueberry

Blueberry is a low spreading deciduous shrub that is common to Canada and northern United States. It often grows in the form of small thickets in open areas of coniferous forests in loose acidic soil, and in burn areas. Blueberries produce clusters of small sweet fruits. Blueberries are generally present in the RSA (CanNorth 2021c). Blueberries are an important Traditional Foods component for First Nations living in the Boreal Shield and Boreal Plains ecozones of Saskatchewan (Chan et al. 2018, 2019).

In the model, plants like blueberry are exposed to atmospheric release through deposition to soil and uptake from soil. In the ecological model blueberries are part of the diet for a number of mammals and birds including: snowshoe hare; Southern red-backed vole; black bear; red fox; rusty blackbird; and grouse. In the human health model blueberries are a component of the Traditional Foods diet, and is used as a surrogate for other fruits and berries in the human Traditional Foods diet, such as rosehips, cranberries, mooseberry, and squashberry (Chan et al. 2018, 2019).

2.2.3 Labrador Tea

Labrador tea is a common name for three closely related plant species in the genus Rhododendron, including *Rhododendron groenlandicum*, which is present in the RSA (CanNorth 2021c). Labrador tea is an important Traditional Foods component for First Nations living in the Boreal Shield and Boreal Plains ecozones of Saskatchewan (Chan et al. 2018, 2019), and is infused as a tea.

In the model, plants like Labrador tea are exposed to atmospheric release through deposition to soil. In the human health model Labrador tea is a component of the Traditional Foods diet, and is



used as a surrogate for other plants in the human Traditional Foods diet, such as mint and rat root (Chan et al. 2018, 2019).

2.3 Terrestrial Mammals

2.3.1 Woodland Caribou

Woodland caribou (*Rangifer tarandus caribou*) is a medium sized member of the deer family (Cervidae), weighing between 100 and 250 kg (COSEWIC 2002). A body weight of 180 kg, representing the upper range mean for adult male and female woodland caribou (COSEWIC 2002) was selected for the ecological model. The boreal population of woodland caribou is designated as Threatened by COSEWIC and SARA and is provincially-ranked S3, meaning that the species is vulnerable/rare to uncommon in Saskatchewan (CanNorth 2021b). Woodland caribou were observed on two occasions in the RSA during baseline studies (CanNorth 2021b).

Woodland caribou have a relatively large home range that varies from 67 km² to 267 km² (McLoughlin 2016). Caribou seek mature jack pine- and black spruce-dominated forest, black spruce bog, and open muskeg habitats common to the Boreal Shield during most of the year, and inaccessible rough terrain such as muskeg or islands on lakes for calving (McLoughlin 2016). For the ecological model, a home rage of 80 km² was considered appropriate based on the average female range during the most sensitive life stage of the caribou (calving and post calving). For the ecological model, the caribou is assumed to interact with more than one exposure location in the RSA for feeding and water intake, due to the large home range.

The food source for the woodland caribou in the winter is terrestrial or arboreal lichens; terrestrial and aquatic vegetation are also food sources in the remainder of the year. Although the current suitability of the site for woodland caribou is relatively low due to an abundance of recently burned or harvested forest in the vicinity of the RSA, the habitat potential over the timeframe of the Project is moderate to high (Omnia 2020a). For the ecological model a diet comprised of 50% browse, 20% lichen and 30% macrophytes is assumed for the woodland caribou.

In the model, caribou are exposed to aquatic and atmospheric release through water and food. In the ecological model, caribou are part of the diet for the grey wolf. Caribou are represented by moose in the human traditional foods diet.

In the ERA, woodland caribou are assessed as a species of concern at an individual level and as a representative mammalian herbivore species at a population level.

2.3.2 Snowshoe Hare

The snowshoe hare (*Lepus americanus*) is a North American hare of the family Leporidae. Adult snowshoe hares typically weigh between 0.9 kg to 1.9 kg (FCSAP 2012) and females are often slightly larger than males (EC & CWF 2005). For the ecological model a bodyweight of 1.8 kg was selected for snowshoe hare. Snowshoe hares are one of the most common woodland mammals of every province and territory of Canada, and are present in the RSA (Omnia 2020a).



Snowshoe hares prefer forested areas with a heavy understory such as conifer-dominated habitats and deciduous riparian forests. The size of their home range can reach up to 10 ha (EC & CWF 2005), but their foraging range is relatively small (1.6 ha) especially for females during the spring and summer reproductive season (FCSAP 2012). In the ecological model snowshoe hare are assumed to reside 100% of the time at the location where they are assessed.

During the summer, the snowshoe hare's diet consists of grasses, sedges, and forbs while during the winter they eat stems and branches of woody plants (FCSAP 2012). They are also known to consume fruits and berries and some animal protein. In the ecological model, snowshoe hares are assumed to eat 90% browse and 10% berries represented by blueberries.

In the model, snowshoe hares are exposed to aquatic and atmospheric release through water and food. In the ecological model, snowshoe hare is part of the diet for the grey wolf. Snowshoe hare is represented by beaver in the human traditional foods diet in the human health model.

In the ERA, snowshoe hare is assessed as a representative upland mammalian herbivore species at a population level.

2.3.3 Moose

Moose (*Alces alces*) is a large ungulate of the family Cervidae, with an average weight for males and females of 453 kg and 350 kg, respectively (FCSAP 2012). For the ecological model a body weight of 400 kg was selected. Moose are found in Canadian forests from the Alaska boundary to the eastern tip of Newfoundland and Labrador (EC & CWF 1997). Moose meat and organs are important components of the traditional foods diet for First Nations living in the Boreal Shield and Boreal Plains ecozones of Saskatchewan (Chan et al. 2018, 2019).

Moose are essentially solitary animals that have one or several distinct seasonal home ranges to which they are strongly attached. Movements between seasonal home ranges may be extensive but home ranges are generally small and moose usually return to the same home ranges year after year (Franzmann 1981; Wilson and Ruff 1999). A seasonal home range of 10 km² was selected for the ecological model.

Breeding season, or rut, begins in mid-September. Cows usually bear one or two calves in the Spring, and the calves stay with the cow until she calves again the following spring. The number of calves born to a cow reflects the availability of nutrition during the rut. During the winter months, moose live almost solely on twigs and shrubs such as balsam fir, poplar, red osier dogwood, birch, willow, and red and striped maples (EC & CWF 1997). In summer the moose's diet includes leaves, some upland plants, and water plants in great quantity where available (EC & CWF 1997). The diet used in the ecological model reflects a spring/summer diet at the end of gestation and when calves are nursing. It assumes 80% browse and 20% macrophytes.

In the model, moose are exposed to aquatic and atmospheric releases through water and food. In the ecological model moose are part of the grey wolf diet. The moose is part of the Traditional

Foods diet in the human health model, and is used as a surrogate for other ungulate species such as white-tailed deer (*Odocoileus virginianus*) and woodland caribou.

In the ERA, moose are assessed as a representative ungulate species at a population level.

2.3.4 Little Brown Myotis

Little brown myotis (*Myotis lucifugus*) is a mouse-eared microbat of the family Vespertilionidae, with a body weight between 0.007 kg and 0.009 kg (EC & CWF 2013). Its range extends from Alaska, across most of North America and into central Mexico. It was, until recently, the most common bat species in Canada but its population is adversely affected by a disease called White Nose Syndrome. The little brown myotis is designated as an Endangered species under COSEWIC and SARA and is provincially-ranked S4. Two bat species, little brown myotis and northern myotis (*Myotis septentrionalis*), have breeding ranges overlapping the RSA (CanNorth 2021b). The northern myotis is also designated as an Endangered species under COSEWIC and SARA and is provincially-ranked S3 (CanNorth 2021b).

Little brown myotis does not migrate, but its summer roosting range can be up to 1000 km from its winter roosting range, also known as hibernacula, where it hibernates without eating. Bats mate in late summer, between August and early October, prior to moving to their hibernacula. Pregnant female bats leave the hibernacula in mid-April and move to their summer maternity or nursery roosts. While pregnant, females forage over a large area but stay closer to their maternity roost once the pups are born. The ecological model assumes a small home range such as during the sensitive reproductive period, and little brown myotis are assumed to spend 100% of their time at the location where they are assessed.

Little brown myotis are nocturnal opportunistic feeders. While at their summer roosts, their diet consists of a great variety of small, flying insects, typically moths, flies, mosquitoes, mayflies, beetles, and midges. In the ecological model, the diet of little brown myotis is composed exclusively of aerial insects with an aquatic larval life stage, represented by benthic invertebrates.

In the model, little brown myotis are exposed to aquatic release through water and food.

In the ERA, little brown myotis are assessed as a species of special concern on an individual level and as a representative insectivorous mammal or other mammals with a similar diet on a population level.

2.3.5 Southern Red-Backed Vole

Southern red-backed vole (*Myodes gapperi*), is a woodland vole of the family Cricetidae, found in Canada and the northern United States, that weighs up to 0.042 kg. Red-backed voles are present in the RSA and were among the most abundant small mammals trapped during baseline studies (Omnia 2020a). Similar small mammals trapped during baseline studies included meadow voles (*Microtus pennsylvanicus*), deer mice (*Peromyscus maniculatus*), masked shrew (*Sorex cinerus*), and water shrew (*Sorex palustris*) (Omnia 2020a). Red-backed vole tissue samples were retained for chemical analysis. For the model, the body weight and total feed intake for the meadow vole were



used because it is a similar species that is well characterized and commonly used in risk assessments (FCSAP 2012).

Baseline studies indicate that red-backed voles are present in all ecosites and vegetation cover types in the RSA. Coarse woody debris and shrub cover are two major components of red-backed vole habitat (Omnia 2020a). They often reside in burrows created by other small mammals, and use runways they created through the vegetation understory or snow. They remain active year-round and generally feed during the night. Females may have two to four litters a year. In the ecological model, voles are assumed to reside 100% of the time at the location where they are assessed.

Southern red-backed voles are omnivorous. They generally feed on green plants, underground fungi, seeds, nuts, roots, and berries, but also eat small terrestrial invertebrates such as insects and snails. They are known to store food for later use. In the ecological model, voles are assumed to eat 60% berries (blueberries) and 40% browse.

In the model, voles are exposed to aquatic and atmospheric releases through water and food. In the ecological model voles are part of the diet for red fox.

In the ERA, voles are assessed as a representative mammalian herbivore at a population level.

2.3.6 Black Bear

Black bear (*Ursus americanus*) is a large terrestrial omnivore. The body weight of black bears varies seasonally and is at its greatest in the fall before winter hibernation. The summer (June/July) weights for males and females are approximately 90 kg and 46 kg, respectively, resulting in an average weight of 68 kg (FCSAP 2012). Although its natural range is across North America, black bears have been extirpated from much of their territory in the continental United States. Currently, black bears are commonly found across most of Canada and Alaska (EC & CWF 2007). Black bears are widespread within the RSA (CanNorth 2021b; Omnia 2020a).

Black bears are found in a variety of habitats but appear to prefer heavily wooded areas and dense bushland. Black bears mate in June or early July, and the cubs, generally one or two, are born the following January or February while the mother is still in her winter den (EC & CWF 2007). The availability of food in the foraging range leading up to denning is important to reproductive success because the bear's fetuses do not "implant" into the uterus until late fall when she enters her winter den (EC & CWF 2007). Home ranges from for female black bears in the Saskatchewan Boreal Shield range from 40 km² to 130 km², as determined from GPS tracking data (McLoughlin et al. 2019). In the ecological model an average female home range of 80 km² is used. For the ecological model, the black bear is assumed to interact with more than one exposure location in the RSA for feeding and water intake, due to the large home range.

Black bears are omnivores and eat seasonally available food. Most of their diet is composed of plant material, including many berries (blueberries, buffalo berries, strawberries, elderberries, Saskatoon berries), fruits (black cherries, and apples) and nuts (acorns, hazelnuts, and beechnuts)



(EC & CWF 2007; FCSAP 2012). Black bears also consume animals, especially invertebrates (ants, bee larvae), fish, small mammals and birds, and carrion (EC & CWF 2007; FCSAP 2012). In the ecological model, black bears are assumed to eat 70% berries (blueberries) and 30% fish (northern pike).

In the model, black bears are exposed to aquatic and atmospheric releases through water and food.

In the ERA, black bears are assessed as a representative terrestrial mammalian omnivorous species at a population level.

2.3.7 Red Fox

The red fox (*Vulpes vulpes*) is a small mammal that ranges in length between 90 cm and 112 cm, and weighs approximately 4.54 kg (USEPA 1993). They are the most widely distributed carnivore in the world (USEPA 1993). Red foxes are found throughout Canada in all provinces and territories. Red foxes are common to northern Saskatchewan and are known to be present in the RSA (Omnia 2020a).

Red foxes occupy many habitats, but in the RSA the highest densities were associated with anthropogenically modified habitats (Omnia 2020a). They generally occupy a home range between 4 to 8 km² and reside in a main underground den and one or more other burrows within their home range. Foxes tend to concentrate their activities near to their dens (USEPA 1993). Foxes breed between late December to mid-March, and pups are born from March through May, with litter sizes ranging from 1 to 10 pups. Pup-rearing is the primary focus of the red fox during spring and early summer. The red fox is assumed to spend 100% of its time at the location where it is assessed.

Red foxes are primarily carnivorous but also eat insects and plant material (fruits, berries, seeds and nuts) (USEPA 1993). Their diet varies with the seasons, eating mainly small mammals in fall and winter, nesting waterfowl in the spring, and insects and berries in the summer (EC & CWF 2019). For the ecological model it is assumed that the red fox's diet consists of 70% small mammals (48% hares and 32% voles) and 30% terrestrial plants represented by browse.

In the model, foxes are exposed to aquatic and atmospheric releases through water and food.

In the ERA, foxes are assessed as a representative mammalian carnivore species at a population level.

2.3.8 Grey Wolf

The grey wolf (*Canis lupus*) is a terrestrial carnivore representing the top level of the food chain. Grey wolves generally weigh between 27 kg and 45 kg, but can be larger (Earthroots 2020). The grey wolf's historical range was across North America, Europe and Asia but they have been extirpated from much of this range. Currently, they are still common in lightly settled portions of Canada from Labrador to British Columbia and in the Yukon Territory and the Northwest



Territories (EC & CWF 1993). The northern grey wolf (Canis lupus occidentalis) has been observed in the RSA during baseline studies (CanNorth 2021b). Wolves have been trapped and harvested for fur in the northern Saskatchewan, including in the Fur Conservation Area (N-19) of La Loche (Omnia 2020b).

Wolves are territorial. The size of their territory varies depending on the availability of prey (EC & CWF 1993). They live and hunt in packs generally composed on a male, female and their offspring. They generally breed once a year in the wild and pups are born in a den that is commonly dug into the soil or in old stumps or rock crevasses (EC & CWF 1993). Home ranges for grey wolves in Minnesota, Alaska, southern Yukon, and British Columbia range from 100 km² to 2,451 km² (RIC 1998). An expected home range of 1,300 km² was selected for the ecological model. For the ecological model, the grey wolf is assumed to interact with more than one exposure location in the RSA for feeding and water intake, due to the large home range.

The wolf's diet consumes terrestrial herbivores including large prey such as moose, caribou, and small prey such as hare (Earthroots 2020). Small prey generally makes up only a small portion of the overall diet (EC & CWF 1993). For the ecological model a diet comprised of 45% moose, 45% caribou and 10% small mammals (snowshoe hare).

In the model, grey wolves are exposed to aquatic and atmospheric releases through water and food.

In the ERA, grey wolves are assessed as a representative avian piscivorous species at a population level.

2.4 Terrestrial Birds

2.4.1 Canada Goose

Canada goose (*Branta canadensis*) is a large terrestrial herbivore of the family Anatidae. Canada geese breed in open forested areas near lake shores and coastal marshes from the arctic tundra through temperate climates (USEPA 1993). Canada goose was the most commonly detected bird species in the RSA during baseline studies (CanNorth 2021b) and are known to breed in the RSA (Omnia 2020a). Geese, including Canada geese and brant geese (*Branta bernicla*), are part of the Traditional Foods diet for the Boreal Plaines and Boreal Shield ecozones of Saskatchewan (Chan et al. 2018, 2019). The USEPA (1993) considers brant geese to be similar to Canada geese.

Body weights for Canada geese vary seasonally and are at their maximum in spring prior to migrating to their northern breeding grounds. Their body weights decline during reproduction and rearing of young (USEPA 1993). For the ecological model, a body weight of 3.7 kg is used for the Canada goose.

The foraging areas for Canada geese vary seasonally. After hatching, goose families tend to move away from their nesting site to other areas, generally on land in riparian areas and near water, with adequate cover and forage to rear their broods (USEPA 1993). Canada geese are almost exclusively vegetarian. They are primarily grazers but consume some grit to aid digestion (USEPA

1993). In the ecological model, Canada geese are assumed to eat 100% browse, and they are assumed to spend 100% of their time at the location where they are assessed.

In the model, Canada geese are exposed to aquatic and atmospheric releases through water and food.

In the ERA, Canada geese are assessed as a representative avian herbivore species at a population level. Although geese are part of the human traditional foods diet, grouse is used as a surrogate for geese in the human health model.

2.4.2 Spruce Grouse

Grouse are avian ground foraging herbivores. Spruce grouse (*Dendragapus canadensis*) have been observed in the RSA during baseline wildlife studies (CanNorth 2021b). Adult spruce grouse average 0.6 kg in weight (FCSAP 2012). Grouse are an important component of the traditional foods diet for First Nations living in the Boreal Shield and Boreal Plains ecozones of Saskatchewan (Chan et al. 2018, 2019).

Grouse are sedentary resident species with a foraging range of 3 to 24 hectares (FCSAP, 2012). Spruce grouse inhabit evergreen forests, especially spruce, jack pine, and lodgepole pine, and appear to prefer young, regenerating tracts that have a dense understory (Cornell Lab n.d.). In the ecological model, grouse are assumed to spend 100% of their time at the location where they are assessed.

Spruce grouse eat pine needles as the main component of their diet, with berries as a secondary food and insects and fungus comprise a small portion of their overall diet (FCSAP 2012). In the ecological model, grouse are assumed to eat 70% browse and 30% berries (blueberry).

In the model, grouse are exposed to aquatic and atmospheric releases through water and food. In the human health model, grouse is part of the human traditional foods diet, and is a surrogate for other terrestrial birds, such as geese and ptarmigan, that have similar diets.

In the ERA, grouse are assessed as a representative avian herbivore species at a population level.

2.4.3 Rusty Blackbird

The Rusty Blackbird (*Euphagus carolinus*) is a medium sized migratory passerine weighing about 64 g. Wintering areas occur primarily within the southeastern United States (EC 2015). It is designated as a Species of Special Concern under COSEWIC and SARA.

The rusty blackbird breeds throughout the boreal forest. Their preferred habitat includes forest wetlands, peat bogs, sedge meadows, marshes, swamps, beaver ponds, and pasture edges (COSEWIC 2006). Suitable summer habitat for the rusty blackbird was found to be common and widespread within the RSA, and rusty blackbirds were observed in the RSA during baseline studies (CanNorth 2021b).



The breeding season for the rusty blackbird begins in April or May, and ends between August to October. Females typically build nests placed in thickets of small conifers or deciduous shrubs, or in dead trees close to water. Females lay a clutch between 3 to 6 eggs and the nesting period lasts between 11 and 13 days (COSEWIC 2016). For the ecological risk assessment, it is assumed that the rusty blackbird spends 100% of its time at the location where it is assessed.

The main dietary components of the Rusty Blackbird include invertebrates associated with aquatic environments, supplanting its diet with seeds and small fruits in the fall and winter (COSEWIC 2006). For the ecological model, it is assumed that the Rusty Blackbird's diet consists of 80% aquatic invertebrates and 20% berries.

In the model, rusty blackbirds are exposed to aquatic and atmospheric releases through water and food.

In the ERA, rusty blackbird is assessed as a species of special concern on an individual level and is considered a suitable surrogate species for other species at risk with similar characteristics and diets that have been observed in the RSA during baseline studies, including common nighthawk, olive-sided flycatcher, and barn swallow (CanNorth 2021b). The rusty blackbird is assessed as a representative insectivorous avian species for other birds with a similar diet on a population level.



3.0 References

- CanNorth. 2021a. Rook I Project Aquatic Environment Baseline Report. Revised Draft Report. Prepared for NexGen Energy Ltd. January.
- CanNorth. 2021b. Rook I Project Wildlife Environment Baseline Report. Revised Draft Report. Prepared for NexGen Energy Ltd. January.
- CanNorth 2021c. Rook I Project Vegetation Inventory Environment Baseline Report. Revised Draft Report. Prepared for NexGen Energy Ltd. January.
- Chan L, Receveur O, Batal M, Sadik T, Schwartz H, Ing A, Fediuk K, Tikhonov C. 2018. First Nations Food, Nutrition and Environment Study (FNFNES): Results from Saskatchewan (2015). Ottawa: University of Ottawa.
- Chan L, Receveur O, Batal M, Sadik T, Schwartz H, Ing A, Fediuk K, Tikhonov C. 2019. Erratum to the First Nations Food, Nutrition and Environment Study (FNFNES): Results from Saskatchewan 2015. Ottawa: University of Ottawa, 2018. July.
- Cornell Lab. (no date) All about Birds. Common Loon. Retrieved September 2021 from: https://www.allaboutbirds.org/quide/Common Loon/lifehistory#.
- COSEWIC (Committee on the Status of Endangered Wildlife in Canada). 2002. COSEWIC assessment and update status report on the woodland caribou *Rangifer tarandus caribou* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. xi + 98 pp.
- COSEWIC. 2006. Rusty blackbird (Euphagus carolinus): COSEWIC assessment and status report 2017.
- CSA (Canadian Standards Association Group). 2022. CSA N288.6-22: Environmental Risk Assessments at Nuclear Facilities and Uranium Mines and Mills.
- Drilling, N., R. Titman and F. McKinney. 2002. Mallard (*Anas platyrhynchos*). In the Birds of North America, No. 658 (A. Poole and F. Gill, eds). The Birds of North America Online, Ithaca, New York.
- Earthroots. 2020. Wolves in Ontario. Available at http://wolvesontario.org/wolves-ontario/.
- EC (Environment Canada). 2015. Management Plan for the Rusty Blackbird (*Euphagus carolinus*) in Canada. *Species at Risk Act* Management Plan Series. Environment Canada, Ottawa. iv + 26 pp.
- EC & CWF (Environment Canada and Canada Wildlife Federation). 1986 (last revised by M. Aleksuik). Muskrat. Retrieved in September 2021 from https://www.hww.ca/en/wildlife/mammals/muskrat.html.
- EC & CWF. 1993 (last revised by L. Carbyn). Wolf. Retrieved in September 2021 from https://www.hww.ca/en/wildlife/mammals/wolf.html.
- EC & CWF. 1997. Moose. Retrieved in September 2021 from https://www.hww.ca/en/wildlife/mammals/moose.html.



- EC & CWF. 2005 (last revised by K.E. Hodges). Snowshoe Hare. Retrieved in September 2021 from https://www.hww.ca/en/wildlife/mammals/snowshoe-hare.html.
- EC & CWF. 2007 (last revised by C. Samson). Black Bear. Retrieved in September 2021 from https://www.hww.ca/en/wildlife/mammals/black-bear.html.
- EC & CWF. 2013 (text by A. Langlois) Little Brown Bat. Retrieved in September 2021 from https://www.hww.ca/en/wildlife/mammals/little-brown-bat.html.
- EC & CWF. 2017 (last revised by G.A. Hood). Beaver. Retrieved in September 2021 from https://www.hww.ca/en/wildlife/mammals/beaver.html.
- FCSAP (Federal Contaminated Sites Action Plan). 2012. Ecological Risk Assessment Guidance, Module 3: Standardization of Wildlife Receptor Characteristics. Prepared for: Environment Canada (Government of Canada), Vancouver, B.C., Prepared by Azimuth Consulting Group, Vancouver, B.C. March 2012.
- Franzmann AW. 1981. Alces alces. Mammalian Species. 154:1–7. May.
- InterGroup (InterGroup Consultants). 2018a. NexGen-Rook I Traditional Land Use Desktop Study. Draft. December.
- InterGroup. 2018b. Desktop Review of Other Land and Resources Use in the Vicinity of the Rook I Project. Draft. November.
- McLoughlin PD, Stewart K, Superbie C, Perry T, Tomchuk P, Greuel R, Singh K, Truchon-Savard A, Henkelman J, Johnstone JF (Department of Biology, University of Saskatchewan). 2016. Population dynamics and critical habitat of woodland caribou in the Saskatchewan Boreal Shield. Interim Project Report, 2013-2016.
- NLFLR (Newfoundland and Labrador Fisheries and Land Resources). 2019. American Mink. Retrieved from,
 - https://www.flr.gov.nl.ca/wildlife/snp/programs/education/animal facts/mammals/mink.html.
- Omnia (Omnia Ecological Services). 2020a. Terrestrial Environment Wildlife and Vegetation Baseline Inventory. Draft. December.
- Omnia. 2020b. Terrestrial Environment Vegetation Baseline Inventory. Draft. December.
- RIC (Ministry of the Environment, Lands and Parks Research for the Terrestrial Ecosystems Task Force Resources Inventory Committee). 1998. Inventory Methods for Wolf and Cougar: Standards for Components of British Columbia's Biodiversity. Report No. 34.
- USEPA (United States Environmental Protection Agency). 1993. Wildlife Exposure Factors Handbook. Document No. EPA/600/R-93/187. December.
- Wilson D, and Ruff S. 1999. The Smithsonian Book of North American Mammals. Smithsonian Institution Press, Washington.



ENVIRONMENTAL RISK ASSESSMENT FOR THE ROOK I PROJECT

Appendix C – Model Results in Support of the Environmental Risk Assessment

APPENDIX C – MODEL RESULTS IN SUPPORT OF THE ENVIRONMENTAL RISK ASSESSMENT



Table C.1: Maximum Concentrations of non-radiological COPCs for the Application Case and Upper Bound Scenario - Project Lifespan

							Maxim	um Concentra	tion during O	perations	•				
	Location				Application (Case				<u> </u>	U	pper Bound S	Scenario		
		Chloride	Sulphate	Arsenic	Cobalt	Copper	Molybdenum	Uranium	Chloride	Sulphate	Arsenic	Cobalt	Copper	Molybdenum	Uranium
	Reference (Broach Lake)	4.22E-01	1.06E+00	1.24E-04	5.83E-04	8.43E-04	2.06E-04	1.00E-04	4.22E-01	1.06E+00	1.24E-04	5.83E-04	8.43E-04	2.06E-04	1.00E-04
	Clearwater River downstream of Broach Lake	4.22E-01	1.06E+00	1.24E-04	5.83E-04	8.43E-04	2.06E-04	1.00E-04	4.22E-01	1.06E+00	1.24E-04	5.83E-04	8.43E-04	2.06E-04	1.00E-04
	Patterson Lake North Arm - East Basin	4.32E-01	1.07E+00	1.25E-04	5.88E-04	8.44E-04	2.07E-04	1.00E-04	4.33E-01	1.07E+00	1.25E-04	5.89E-04	8.44E-04	2.07E-04	1.01E-04
	Patterson Lake North Arm - West Basin	4.35E+00	2.14E+02	4.36E-04	7.71E-04	9.95E-04	5.98E-04	3.84E-04	1.00E+02	2.16E+02	4.46E-04	7.78E-04	9.99E-04	1.39E-03	9.25E-04
	Patterson Lake South Arm	3.17E+00	1.45E+02	1.50E-04	6.85E-04	9.23E-04	4.07E-04	1.76E-04	7.16E+01	1.47E+02	1.50E-04	6.89E-04	9.25E-04	8.15E-04	3.09E-04
Material (magell)	Forrest Lake North Basin	2.10E+00	9.01E+01	1.39E-04	6.45E-04	8.92E-04	3.30E-04	1.47E-04	4.40E+01	9.08E+01	1.40E-04	6.48E-04	8.93E-04	5.82E-04	2.28E-04
Water (mg/L)	Beet Lake	1.97E+00	8.25E+01	1.28E-04	6.36E-04	8.84E-04	3.09E-04	1.28E-04	4.05E+01	8.33E+01	1.28E-04	6.38E-04	8.85E-04	5.20E-04	1.77E-04
	Naomi Lake	1.64E+00	6.49E+01	1.26E-04	6.24E-04	8.74E-04	2.85E-04	1.20E-04	3.18E+01	6.56E+01	1.26E-04	6.25E-04	8.75E-04	4.48E-04	1.55E-04
	Clearwater River upstream Mirror River Confluence	1.08E+00	3.57E+01	1.25E-04	6.05E-04	8.60E-04	2.49E-04	1.11E-04	1.74E+01	3.60E+01	1.25E-04	6.06E-04	8.60E-04	3.37E-04	1.30E-04
	Clearwater River downstream Mirror River Confluence	6.20E-01	1.16E+01	1.24E-04	5.90E-04	8.48E-04	2.19E-04	1.04E-04	5.55E+00	1.17E+01	1.24E-04	5.90E-04	8.48E-04	2.46E-04	1.09E-04
	Lloyd Lake Inlet	5.86E-01	9.71E+00	1.24E-04	5.89E-04	8.47E-04	2.17E-04	1.03E-04	4.65E+00	9.80E+00	1.24E-04	5.89E-04	8.47E-04	2.39E-04	1.07E-04
	Reference (Broach Lake)	-	-	1.00E+01	1.45E+00	2.52E+00	6.51E-01	1.93E+00	-	-	1.00E+01	1.45E+00	2.52E+00	6.51E-01	1.93E+00
	Patterson Lake North Arm - West Basin - East Basin	_	-	1.00E+01	1.46E+00	2.52E+00	6.52E-01	1.93E+00	-	-	1.00E+01	1.46E+00	2.52E+00	6.52E-01	1.93E+00
Sediment (mg/kg dw)	Patterson Lake North Arm - West Basin - West Basin	_	-	3.05E+01	1.88E+00	2.93E+00	1.74E+00	6.33E+00	-	-	3.10E+01	1.90E+00	2.94E+00	3.97E+00	1.42E+01
Seament (mg/kg aw/	Patterson Lake South Arm	_	-	1.17E+01	1.68E+00	2.73E+00	1.21E+00	3.09E+00	-	-	1.17E+01	1.69E+00	2.73E+00	2.39E+00	5.19E+00
	Beet Lake	-	-	1.03E+01	1.57E+00	2.62E+00	9.39E-01	2.36E+00	-	-	1.03E+01	1.58E+00	2.63E+00	1.55E+00	3.12E+00
	Naomi Lake	-	-	1.01E+01	1.54E+00	2.60E+00	8.72E-01	2.23E+00	-	-	1.01E+01	1.55E+00	2.60E+00	1.34E+00	2.77E+00
	Lloyd Lake Inlet	-	-	1.00E+01	1.47E+00	2.53E+00	6.81E-01	1.97E+00	-	-	1.00E+01	1.47E+00	2.53E+00	7.43E-01	2.04E+00
	Reference (Broach Lake)	-	-	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	-	-	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
	Patterson Lake North Arm - West Basin	_	-	3.00E-08	2.00E-08	1.70E-07	1.20E-07	9.38E-06	-	-	3.00E-08	2.00E-08	1.70E-07	1.20E-07	9.38E-06
	Patterson Lake South Arm	-	-	1.00E-08	1.00E-08	3.00E-08	2.00E-08	1.34E-06	-	-	1.00E-08	1.00E-08	3.00E-08	2.00E-08	1.34E-06
Air (mg/m³)	Beet Lake	-	-	1.00E-08	1.00E-08	2.00E-08	1.00E-08	7.90E-07	-	-	1.00E-08	1.00E-08	2.00E-08	1.00E-08	7.90E-07
	Clearwater River and Mirror River Confluence	_	-	1.00E-08	1.00E-08	1.00E-08	1.00E-08	5.00E-08	-	-	1.00E-08	1.00E-08	1.00E-08	1.00E-08	5.00E-08
	Lloyd Lake	-	-	1.00E-08	1.00E-08	1.00E-08	1.00E-08	3.00E-08	-	-	1.00E-08	1.00E-08	1.00E-08	1.00E-08	3.00E-08
	Reference (Broach Lake)	-	-	5.89E-01	4.32E-01	6.01E-01	1.42E-01	2.47E-01	-	-	5.89E-01	4.32E-01	6.01E-01	1.42E-01	2.47E-01
	Patterson Lake North Arm - West Basin	-	-	5.89E-01	4.34E-01	6.19E-01	1.47E-01	1.30E+00	=	-	5.89E-01	4.34E-01	6.19E-01	1.47E-01	1.30E+00
	Patterson Lake South Arm	-	-	5.89E-01	4.33E-01	6.04E-01	1.43E-01	3.97E-01	=	-	5.89E-01	4.33E-01	6.04E-01	1.43E-01	3.97E-01
Soil (mg/kg dw)	Beet Lake	_	_	5.89E-01	4.33E-01	6.03E-01	1.43E-01	3.36E-01	-	_	5.89E-01	4.33E-01	6.03E-01	1.43E-01	3.36E-01
	Clearwater River and Mirror River Confluence	-	-	5.89E-01	4.33E-01	6.02E-01	1.43E-01	2.53E-01	-	-	5.89E-01	4.33E-01	6.02E-01	1.43E-01	2.53E-01
	Lloyd Lake	-	-	5.89E-01	4.33E-01	6.02E-01	1.43E-01	2.51E-01	-	-	5.89E-01	4.33E-01	6.02E-01	1.43E-01	2.51E-01

Table C.2: Maximum Concentrations of non-radiological COPCs for the Application Case and Upper Bound Scenario – Far-Future

							Maximum C	oncentratio	n Far-Futur	e Scenario					
	Location			А	pplication Cas	ie						Upper Bound	Scenario		
		Chloride	Sulphate	Arsenic	Cobalt	Copper	Molybdenum	Uranium	Chloride	Sulphate	Arsenic	Cobalt	Copper	Molybdenum	Uranium
	Reference (Broach Lake)	4.22E-01	1.06E+00	1.24E-04	5.83E-04	8.43E-04	2.06E-04	1.00E-04	4.22E-01	1.06E+00	1.24E-04	5.83E-04	8.43E-04	2.06E-04	1.00E-04
	Clearwater River downstream of Broach Lake	4.22E-01	1.06E+00	1.24E-04	5.83E-04	8.43E-04	2.06E-04	1.00E-04	4.22E-01	1.06E+00	1.24E-04	5.83E-04	8.43E-04	2.06E-04	1.00E-04
	Patterson Lake North Arm - East Basin	4.22E-01	1.06E+00	1.24E-04	5.83E-04	8.43E-04	2.06E-04	1.00E-04	4.22E-01	1.06E+00	1.24E-04	5.83E-04	8.43E-04	2.06E-04	1.00E-04
	Patterson Lake North Arm - West Basin	8.39E-01	2.16E+00	1.31E-04	1.43E-03	2.18E-03	4.60E-03	1.00E-03	1.04E+00	2.79E+00	1.35E-04	1.88E-03	2.85E-03	1.71E-02	1.00E-03
	Patterson Lake South Arm	7.27E-01	1.86E+00	1.24E-04	1.10E-03	1.63E-03	2.76E-03	3.50E-04	8.76E-01	2.33E+00	1.25E-04	1.37E-03	2.02E-03	1.00E-02	3.51E-04
	Forrest Lake North Basin	6.07E-01	1.55E+00	1.24E-04	8.94E-04	1.32E-03	1.76E-03	2.52E-04	6.98E-01	1.83E+00	1.24E-04	1.06E-03	1.56E-03	6.16E-03	2.52E-04
Water (mg/L)	Beet Lake	5.94E-01	1.51E+00	1.24E-04	8.54E-04	1.25E-03	1.54E-03	1.93E-04	6.78E-01	1.77E+00	1.24E-04	9.98E-04	1.46E-03	5.31E-03	1.93E-04
	Naomi Lake	5.56E-01	1.41E+00	1.24E-04	7.92E-04	1.16E-03	1.23E-03	1.66E-04	6.22E-01	1.62E+00	1.24E-04	9.03E-04	1.32E-03	4.12E-03	1.66E-04
	Clearwater River upstream Mirror River Confluence	4.94E-01	1.25E+00	1.24E-04	6.96E-04	1.01E-03	7.56E-04	1.36E-04	5.30E-01	1.36E+00	1.24E-04	7.55E-04	1.10E-03	2.32E-03	1.36E-04
	Clearwater River downstream Mirror River Confluence	4.44E-01	1.12E+00	1.24E-04	6.17E-04	8.94E-04	3.72E-04	1.11E-04	4.54E-01	1.15E+00	1.24E-04	6.35E-04	9.20E-04	8.41E-04	1.11E-04
	Lloyd Lake Inlet	4.40E-01	1.11E+00	1.24E-04	6.11E-04	8.85E-04	3.42E-04	1.09E-04	4.49E-01	1.14E+00	1.24E-04	6.26E-04	9.06E-04	7.27E-04	1.09E-04
	Reference (Broach Lake)	-	-	1.00E+01	1.45E+00	2.52E+00	6.51E-01	1.93E+00	-	-	1.00E+01	1.45E+00	2.52E+00	6.51E-01	1.93E+00
	Patterson Lake North Arm - West Basin - East Basin	_	_	1.00E+01	1.45E+00	2.52E+00	6.51E-01	1.93E+00	-	_	1.00E+01	1.45E+00	2.52E+00	6.51E-01	1.93E+00
Sediment (mg/kg dw)	Patterson Lake North Arm - West Basin - West Basin	-	-	1.06E+01	3.56E+00	6.51E+00	1.45E+01	1.93E+01	-	_	1.09E+01	4.68E+00	8.52E+00	5.39E+01	1.93E+01
	Patterson Lake South Arm	-	-	1.00E+01	2.73E+00	4.86E+00	8.71E+00	6.74E+00	-	-	1.01E+01	3.41E+00	6.04E+00	3.16E+01	6.74E+00
	Beet Lake	-	-	1.00E+01	2.13E+00	3.74E+00	4.85E+00	3.72E+00	-	-	1.00E+01	2.49E+00	4.36E+00	1.68E+01	3.72E+00
	Naomi Lake	-	-	1.00E+01	1.97E+00	3.46E+00	3.87E+00	3.19E+00	-	-	1.00E+01	2.25E+00	3.93E+00	1.30E+01	3.19E+00
	Lloyd Lake Inlet	-	-	1.00E+01	1.52E+00	2.64E+00	1.08E+00	2.09E+00	-	-	1.00E+01	1.56E+00	2.71E+00	2.29E+00	2.09E+00
	Reference (Broach Lake)	-	-	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	-	-	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
	Patterson Lake North Arm - West Basin	-	-	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	-	-	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
A: , (, , , , , , , , , 3)	Patterson Lake South Arm	-	-	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	-	-	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Air (mg/m³)	Beet Lake	-	-	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	-	-	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
	Clearwater River and Mirror River Confluence	-	-	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	-	-	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
	Lloyd Lake	-	_	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	-	-	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
	Reference (Broach Lake)	-	-	5.89E-01	4.32E-01	6.01E-01	1.42E-01	2.47E-01	-	-	5.89E-01	4.32E-01	6.01E-01	1.42E-01	2.47E-01
	Patterson Lake North Arm - West Basin	-	-	5.89E-01	4.33E-01	6.02E-01	1.42E-01	3.57E-01	-	-	5.89E-01	4.33E-01	6.02E-01	1.42E-01	3.57E-01
Soil (ma/ka dw)	Patterson Lake South Arm	-	-	5.89E-01	4.33E-01	6.01E-01	1.42E-01	2.63E-01	-	-	5.89E-01	4.33E-01	6.01E-01	1.42E-01	2.63E-01
Soil (mg/kg dw)	Beet Lake	-	-	5.89E-01	4.33E-01	6.01E-01	1.42E-01	2.57E-01	-	_	5.89E-01	4.33E-01	6.01E-01	1.42E-01	2.57E-01
	Clearwater River and Mirror River Confluence	-	-	5.89E-01	4.33E-01	6.01E-01	1.42E-01	2.48E-01	-	_	5.89E-01	4.33E-01	6.01E-01	1.42E-01	2.48E-01
	Lloyd Lake		-	5.89E-01	4.33E-01	6.01E-01	1.42E-01	2.48E-01	-	-	5.89E-01	4.33E-01	6.01E-01	1.42E-01	2.48E-01

Table C.3: Maximum Concentrations of non-radiological COPCs for the RFD Scenario - Project Lifespan and Far-Future

							Ma	aximum Conc	entration RFI	D case					
	Location			F	Project Lifes	pan						Far-Futu	re		
		Chloride	Sulphate	Arsenic	Cobalt	Copper	Molybdenum	Uranium	Chloride	Sulphate	Arsenic	Cobalt	Copper	Molybdenum	Uranium
	Reference (Broach Lake)	4.22E-01	1.06E+00	1.24E-04	5.83E-04	8.43E-04	2.06E-04	1.00E-04	4.22E-01	1.06E+00	1.24E-04	5.83E-04	8.43E-04	2.06E-04	1.00E-04
	Clearwater River downstream of Broach Lake	4.22E-01	1.06E+00	1.24E-04	5.83E-04	8.43E-04	2.06E-04	1.00E-04	4.22E-01	1.06E+00	1.24E-04	5.83E-04	8.43E-04	2.06E-04	1.00E-04
	Patterson Lake North Arm - East Basin	4.32E-01	1.07E+00	1.25E-04	5.88E-04	8.44E-04	2.07E-04	1.00E-04	4.22E-01	1.06E+00	1.24E-04	5.83E-04	8.43E-04	2.06E-04	1.00E-04
	Patterson Lake North Arm - West Basin	4.35E+00	2.14E+02	4.36E-04	7.71E-04	9.95E-04	5.98E-04	3.84E-04	8.39E-01	2.16E+00	1.31E-04	1.43E-03	2.18E-03	4.60E-03	1.00E-03
	Patterson Lake South Arm	4.27E+00	2.14E+02	6.93E-04	7.82E-04	1.01E-03	5.72E-04	5.37E-04	7.27E-01	1.86E+00	1.24E-04	1.10E-03	1.63E-03	2.76E-03	3.50E-04
Motor (mg/l)	Forrest Lake North Basin	2.85E+00	1.36E+02	4.84E-04	7.08E-04	9.47E-04	4.37E-04	3.76E-04	6.07E-01	1.55E+00	1.24E-04	8.94E-04	1.32E-03	1.76E-03	2.52E-04
Water (mg/L)	Beet Lake	2.37E+00	1.09E+02	2.21E-04	6.71E-04	9.15E-04	3.69E-04	2.52E-04	5.94E-01	1.51E+00	1.24E-04	8.54E-04	1.25E-03	1.54E-03	1.93E-04
	Naomi Lake	1.95E+00	8.57E+01	1.74E-04	6.50E-04	8.97E-04	3.30E-04	2.06E-04	5.56E-01	1.41E+00	1.24E-04	7.92E-04	1.16E-03	1.23E-03	1.66E-04
	Clearwater River upstream Mirror River Confluence	1.25E+00	4.69E+01	1.52E-04	6.19E-04	8.72E-04	2.73E-04	1.59E-04	4.94E-01	1.25E+00	1.24E-04	6.96E-04	1.01E-03	7.56E-04	1.36E-04
	Clearwater River downstream Mirror River Confluence	6.72E-01	1.49E+01	1.32E-04	5.94E-04	8.52E-04	2.26E-04	1.18E-04	4.44E-01	1.12E+00	1.24E-04	6.17E-04	8.94E-04	3.72E-04	1.11E-04
	Lloyd Lake Inlet	6.28E-01	1.25E+01	1.30E-04	5.92E-04	8.50E-04	2.23E-04	1.14E-04	4.40E-01	1.11E+00	1.24E-04	6.11E-04	8.85E-04	3.42E-04	1.09E-04
	Reference (Broach Lake)	-	-	1.00E+01	1.45E+00	2.52E+00	6.51E-01	1.93E+00	-	-	1.00E+01	1.45E+00	2.52E+00	6.51E-01	1.93E+00
	Patterson Lake North Arm - West Basin - East Basin	-	-	1.00E+01	1.46E+00	2.52E+00	6.52E-01	1.93E+00	-	-	1.00E+01	1.45E+00	2.52E+00	6.51E-01	1.93E+00
	Patterson Lake North Arm - West Basin - West Basin	-	-	3.05E+01	1.88E+00	2.93E+00	1.74E+00	6.33E+00	-	-	1.06E+01	3.56E+00	6.51E+00	1.45E+01	1.93E+01
Sediment (mg/kg dw)	Patterson Lake South Arm	-	-	2.29E+01	1.77E+00	2.82E+00	1.39E+00	4.46E+00	-	-	1.00E+01	2.73E+00	4.86E+00	8.71E+00	6.74E+00
	Beet Lake	-	-	1.22E+01	1.62E+00	2.68E+00	1.03E+00	2.86E+00	-	-	1.00E+01	2.13E+00	3.74E+00	4.85E+00	3.72E+00
	Naomi Lake	-	-	1.11E+01	1.58E+00	2.64E+00	9.46E-01	2.58E+00	-	-	1.00E+01	1.97E+00	3.46E+00	3.87E+00	3.19E+00
	Lloyd Lake Inlet	-	-	1.01E+01	1.47E+00	2.53E+00	6.91E-01	2.01E+00	-	-	1.00E+01	1.52E+00	2.64E+00	1.08E+00	2.09E+00
	Reference (Broach Lake)	-	-	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	-	-	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
	Patterson Lake North Arm - West Basin	-	-	1.00E-08	1.00E-08	1.72E-07	1.23E-07	9.47E-06	-	-	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Air (mg/m³)	Patterson Lake South Arm	-	-	1.00E-08	1.00E-08	1.00E-08	1.00E-08	1.51E-06	-	-	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Air (mg/m²)	Beet Lake	-	-	1.00E-08	1.00E-08	1.00E-08	1.00E-08	8.45E-07	-	-	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
	Clearwater River and Mirror River Confluence	-	-	1.00E-08	1.00E-08	1.00E-08	1.00E-08	6.36E-08	-	-	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
	Lloyd Lake	-	-	1.00E-08	1.00E-08	1.00E-08	1.00E-08	1.00E-08	-	-	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
	Reference (Broach Lake)	-	-	5.89E-01	4.32E-01	6.01E-01	1.42E-01	2.47E-01	-	-	5.89E-01	4.32E-01	6.01E-01	1.42E-01	2.47E-01
	Patterson Lake North Arm - West Basin	-	-	5.89E-01	4.33E-01	6.19E-01	1.47E-01	1.31E+00	-	-	5.89E-01	4.33E-01	6.02E-01	1.42E-01	3.59E-01
Soil (ma/ka du)	Patterson Lake South Arm	-	-	5.89E-01	4.33E-01	6.02E-01	1.43E-01	4.16E-01	-	-	5.89E-01	4.33E-01	6.01E-01	1.42E-01	2.65E-01
Soil (mg/kg dw)	Beet Lake	-	-	5.89E-01	4.33E-01	6.02E-01	1.43E-01	3.42E-01	-	-	5.89E-01	4.33E-01	6.01E-01	1.42E-01	2.57E-01
	Clearwater River and Mirror River Confluence	-	-	5.89E-01	4.33E-01	6.02E-01	1.43E-01	2.54E-01	-	-	5.89E-01	4.33E-01	6.01E-01	1.42E-01	2.48E-01
	Lloyd Lake	-	-	5.89E-01	4.33E-01	6.02E-01	1.43E-01	2.48E-01	-	-	5.89E-01	4.33E-01	6.01E-01	1.42E-01	2.47E-01

Table C.4: Maximum Concentrations of Radiological COPCs for the Application Case and Upper Bound Scenario - Project Lifespan

						Maximu	ım Concentra	ntion Project	t Lifespan				
	Location			Applicat	ion Case					Upper Bou	nd Scenario)	
		Uranium-	Uranium-	Thorium-	Radium-	Lead-	Polonium-	Uranium-	Uranium-	Thorium-	Radium-	Lead-	Polonium-
		238	234	230	226	210	210	238	234	230	226	210	210
	Reference (Broach Lake)	3.56E-04	3.56E-04	9.09E-03	5.54E-03	5.07E-03	5.15E-03	3.56E-04	3.56E-04	9.09E-03	5.54E-03	5.07E-03	5.15E-03
	Clearwater River downstream of Broach Lake	3.56E-04	3.56E-04	9.09E-03	5.54E-03	5.07E-03	5.15E-03	3.56E-04	3.56E-04	9.09E-03	5.54E-03	5.07E-03	5.15E-03
	Patterson Lake North Arm - East Basin	3.58E-04	3.58E-04	9.12E-03	5.55E-03	5.08E-03	5.16E-03	3.58E-04	3.58E-04	9.12E-03	5.55E-03	5.08E-03	5.16E-03
	Patterson Lake North Arm - West Basin	3.88E-03	3.89E-03	5.17E-02	9.63E-03	7.74E-03	7.82E-03	1.06E-02	1.06E-02	5.22E-02	9.82E-03	1.94E-02	1.99E-02
	Patterson Lake South Arm	1.30E-03	1.31E-03	3.41E-02	6.99E-03	5.22E-03	5.31E-03	2.95E-03	2.96E-03	3.44E-02	7.05E-03	5.86E-03	5.96E-03
Water (Bq/L)	Forrest Lake North Basin	9.38E-04	9.40E-04	2.44E-02	6.43E-03	5.16E-03	5.25E-03	1.94E-03	1.95E-03	2.46E-02	6.47E-03	5.55E-03	5.65E-03
	Beet Lake	7.04E-04	7.06E-04	2.22E-02	6.15E-03	5.09E-03	5.17E-03	1.31E-03	1.31E-03	2.24E-02	6.18E-03	5.17E-03	5.26E-03
	Naomi Lake	6.01E-04	6.02E-04	1.92E-02	5.99E-03	5.08E-03	5.16E-03	1.03E-03	1.03E-03	1.94E-02	6.01E-03	5.11E-03	5.20E-03
	Clearwater River upstream Mirror River Confluence	4.89E-04	4.89E-04	1.46E-02	5.78E-03	5.07E-03	5.16E-03	7.20E-04	7.22E-04	1.46E-02	5.79E-03	5.09E-03	5.18E-03
	Clearwater River downstream Mirror River Confluence	3.96E-04	3.96E-04	1.07E-02	5.61E-03	5.07E-03	5.16E-03	4.66E-04	4.67E-04	1.08E-02	5.62E-03	5.07E-03	5.16E-03
	Lloyd Lake Inlet	3.88E-04	3.88E-04	1.04E-02	5.60E-03	5.07E-03	5.15E-03	4.43E-04	4.44E-04	1.05E-02	5.60E-03	5.07E-03	5.16E-03
	Reference (Broach Lake)	6.84E+00	6.84E+00	2.09E+01	6.47E+01	3.59E+02	3.65E+02	6.84E+00	6.84E+00	2.09E+01	6.47E+01	3.59E+02	3.65E+02
	Patterson Lake North Arm - East Basin	6.86E+00	6.86E+00	2.09E+01	6.48E+01	3.59E+02	3.66E+02	6.86E+00	6.86E+00	2.09E+01	6.48E+01	3.59E+02	3.66E+02
	Patterson Lake North Arm - West Basin	6.16E+01	6.18E+01	1.16E+02	1.04E+02	4.92E+02	5.00E+02	1.59E+02	1.59E+02	1.17E+02	1.06E+02	9.84E+02	1.00E+03
Sediment (Bq/kg dw)	Patterson Lake South Arm	2.13E+01	2.14E+01	7.49E+01	7.82E+01	3.68E+02	3.74E+02	4.72E+01	4.74E+01	7.57E+01	7.88E+01	4.01E+02	4.08E+02
	Beet Lake	1.22E+01	1.22E+01	4.93E+01	7.04E+01	3.60E+02	3.67E+02	2.17E+01	2.17E+01	4.97E+01	7.07E+01	3.65E+02	3.71E+02
	Naomi Lake	1.06E+01	1.06E+01	4.28E+01	6.89E+01	3.60E+02	3.66E+02	1.72E+01	1.73E+01	4.31E+01	6.91E+01	3.62E+02	3.68E+02
	Lloyd Lake Inlet	7.33E+00	7.33E+00	2.38E+01	6.53E+01	3.59E+02	3.65E+02	8.19E+00	8.19E+00	2.38E+01	6.53E+01	3.59E+02	3.66E+02
	Reference (Broach Lake)	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
	Patterson Lake North Arm	9.26E-05	9.26E-05	9.26E-05	9.26E-05	9.26E-05	9.26E-05	9.26E-05	9.26E-05	9.26E-05	9.26E-05	9.26E-05	9.26E-05
Air (Bq/m³)	Patterson Lake South Arm	1.51E-05	1.51E-05	1.51E-05	1.51E-05	1.51E-05	1.51E-05	1.51E-05	1.51E-05	1.51E-05	1.51E-05	1.51E-05	1.51E-05
All (bq/III)	Beet Lake	8.80E-06	8.80E-06	8.80E-06	8.80E-06	8.80E-06	8.80E-06	8.80E-06	8.80E-06	8.80E-06	8.80E-06	8.80E-06	8.80E-06
	Clearwater River and Mirror River Confluence	5.48E-07	5.48E-07	5.48E-07	5.48E-07	5.48E-07	5.48E-07	5.48E-07	5.48E-07	5.48E-07	5.48E-07	5.48E-07	5.48E-07
	Lloyd Lake	2.94E-07	2.94E-07	2.94E-07	2.94E-07	2.94E-07	2.94E-07	2.94E-07	2.94E-07	2.94E-07	2.94E-07	2.94E-07	2.94E-07
	Reference (Broach Lake)	3.06E+00	3.06E+00	2.00E+01	1.79E+01	4.00E+01	1.12E+01	3.06E+00	3.06E+00	2.00E+01	1.79E+01	4.00E+01	1.12E+01
	Patterson Lake North Arm	1.34E+01	1.34E+01	3.16E+01	2.97E+01	4.76E+01	1.89E+01	1.34E+01	1.34E+01	3.16E+01	2.97E+01	4.76E+01	1.89E+01
Soil (Bq/kg dw)	Patterson Lake South Arm	4.74E+00	4.74E+00	2.19E+01	1.98E+01	4.12E+01	1.25E+01	4.74E+00	4.74E+00	2.19E+01	1.98E+01	4.12E+01	1.25E+01
Soli (bq/kg dw)	Beet Lake	4.04E+00	4.04E+00	2.11E+01	1.91E+01	4.07E+01	1.20E+01	4.04E+00	4.04E+00	2.11E+01	1.91E+01	4.07E+01	1.20E+01
	Clearwater River and Mirror River Confluence	3.12E+00	3.12E+00	2.01E+01	1.80E+01	4.00E+01	1.13E+01	3.12E+00	3.12E+00	2.01E+01	1.80E+01	4.00E+01	1.13E+01
	Lloyd Lake	3.09E+00	3.09E+00	2.00E+01	1.80E+01	4.00E+01	1.12E+01	3.09E+00	3.09E+00	2.00E+01	1.80E+01	4.00E+01	1.12E+01

Table C.5: Maximum Concentrations of Radiological COPCs for the Application Case and Upper Bound Scenario - Far-Future

						Maxi	imum Concer	tration Far	Future				
	Location			Applicat	ion Case					Upper Bou	nd Scenaric)	
		Uranium-	Uranium-	Thorium-	Radium-	Lead-	Polonium-	Uranium-	Uranium-	Thorium-	Radium-	Lead-	Polonium-
		238	234	230	226	210	210	238	234	230	226	210	210
	Reference (Broach Lake)	3.56E-04	3.56E-04	9.09E-03	5.54E-03	5.07E-03	5.15E-03	3.56E-04	3.56E-04	9.09E-03	5.54E-03	5.07E-03	5.15E-03
	Clearwater River downstream of Broach Lake	3.56E-04	3.56E-04	9.09E-03	5.54E-03	5.07E-03	5.15E-03	3.56E-04	3.56E-04	9.09E-03	5.54E-03	5.07E-03	5.15E-03
	Patterson Lake North Arm - East Basin	3.56E-04	3.56E-04	9.09E-03	5.54E-03	5.07E-03	5.15E-03	3.56E-04	3.56E-04	9.09E-03	5.54E-03	5.07E-03	5.15E-03
	Patterson Lake North Arm - West Basin	1.15E-02	1.16E-02	9.72E-03	7.14E-03	5.30E-03	5.39E-03	1.15E-02	1.16E-02	9.72E-03	7.15E-03	5.30E-03	5.39E-03
	Patterson Lake South Arm	3.44E-03	3.46E-03	9.48E-03	6.13E-03	5.08E-03	5.17E-03	3.45E-03	3.46E-03	9.48E-03	6.13E-03	5.08E-03	5.17E-03
Water (Bq/L)	Forrest Lake North Basin	2.23E-03	2.24E-03	9.32E-03	5.90E-03	5.08E-03	5.16E-03	2.23E-03	2.24E-03	9.32E-03	5.90E-03	5.08E-03	5.16E-03
	Beet Lake	1.50E-03	1.51E-03	9.29E-03	5.79E-03	5.07E-03	5.16E-03	1.51E-03	1.51E-03	9.29E-03	5.79E-03	5.07E-03	5.16E-03
	Naomi Lake	1.16E-03	1.17E-03	9.25E-03	5.72E-03	5.07E-03	5.15E-03	1.16E-03	1.17E-03	9.25E-03	5.72E-03	5.07E-03	5.15E-03
	Clearwater River upstream Mirror River Confluence	7.91E-04	7.93E-04	9.17E-03	5.64E-03	5.07E-03	5.15E-03	7.92E-04	7.93E-04	9.17E-03	5.64E-03	5.07E-03	5.15E-03
	Clearwater River downstream Mirror River Confluence	4.87E-04	4.87E-04	9.11E-03	5.57E-03	5.07E-03	5.15E-03	4.87E-04	4.87E-04	9.11E-03	5.57E-03	5.07E-03	5.15E-03
	Lloyd Lake Inlet	4.60E-04	4.60E-04	9.11E-03	5.56E-03	5.07E-03	5.15E-03	4.60E-04	4.60E-04	9.11E-03	5.56E-03	5.07E-03	5.15E-03
	Reference (Broach Lake)	6.84E+00	6.84E+00	2.09E+01	6.47E+01	3.59E+02	3.65E+02	6.84E+00	6.84E+00	2.09E+01	6.47E+01	3.59E+02	3.65E+02
	Patterson Lake North Arm - East Basin	6.84E+00	6.84E+00	2.09E+01	6.47E+01	3.59E+02	3.65E+02	6.84E+00	6.84E+00	2.09E+01	6.47E+01	3.59E+02	3.65E+02
	Patterson Lake North Arm - West Basin	2.22E+02	2.22E+02	2.23E+01	8.34E+01	3.76E+02	3.82E+02	2.22E+02	2.23E+02	2.23E+01	8.35E+01	3.76E+02	3.82E+02
Sediment (Bq/kg dw)	Patterson Lake South Arm	6.63E+01	6.65E+01	2.18E+01	7.16E+01	3.60E+02	3.66E+02	6.63E+01	6.65E+01	2.18E+01	7.16E+01	3.60E+02	3.66E+02
	Beet Lake	2.89E+01	2.90E+01	2.13E+01	6.77E+01	3.59E+02	3.66E+02	2.89E+01	2.90E+01	2.13E+01	6.77E+01	3.59E+02	3.66E+02
	Naomi Lake	2.24E+01	2.24E+01	2.12E+01	6.69E+01	3.59E+02	3.65E+02	2.24E+01	2.25E+01	2.12E+01	6.69E+01	3.59E+02	3.65E+02
	Lloyd Lake Inlet	8.84E+00	8.85E+00	2.09E+01	6.50E+01	3.59E+02	3.65E+02	8.84E+00	8.85E+00	2.09E+01	6.50E+01	3.59E+02	3.65E+02
	Reference (Broach Lake)	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
	Patterson Lake North Arm	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Air (Dar/ma3)	Patterson Lake South Arm	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Air (Bq/m³)	Beet Lake	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
	Clearwater River and Mirror River Confluence	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
	Lloyd Lake	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
	Reference (Broach Lake)	3.06E+00	3.06E+00	2.00E+01	1.79E+01	4.00E+01	1.12E+01	3.06E+00	3.06E+00	2.00E+01	1.79E+01	4.00E+01	1.12E+01
	Patterson Lake North Arm	4.14E+00	4.14E+00	2.65E+01	2.58E+01	4.00E+01	1.12E+01	4.14E+00	4.14E+00	2.65E+01	2.58E+01	4.00E+01	1.12E+01
Soil (Bq/kg dw)	Patterson Lake South Arm	3.23E+00	3.23E+00	2.11E+01	1.92E+01	4.00E+01	1.12E+01	3.23E+00	3.23E+00	2.11E+01	1.92E+01	4.00E+01	1.12E+01
Soli (Bq/kg aw)	Beet Lake	3.16E+00	3.16E+00	2.06E+01	1.87E+01	4.00E+01	1.12E+01	3.16E+00	3.16E+00	2.06E+01	1.87E+01	4.00E+01	1.12E+01
	Clearwater River and Mirror River Confluence	3.06E+00	3.06E+00	2.00E+01	1.80E+01	4.00E+01	1.12E+01	3.06E+00	3.06E+00	2.00E+01	1.80E+01	4.00E+01	1.12E+01
	Lloyd Lake	3.06E+00	3.06E+00	2.00E+01	1.80E+01	4.00E+01	1.12E+01	3.06E+00	3.06E+00	2.00E+01	1.80E+01	4.00E+01	1.12E+01

Table C.6: Maximum Concentrations of Radiological COPCs for the RFD Scenario - Project Lifespan and Far-Future

						Max	rimum Conce	ntration RFD) Case				
	Location			Project	Lifespan					Far f	uture		
		Uranium-	Uranium-	Thorium-	Radium-	Lead-	Polonium-	Uranium-	Uranium-	Thorium-	Radium-	Lead-	Polonium-
		238	234	230	226	210	210	238	234	230	226	210	210
	Reference (Broach Lake)	3.56E-04	3.56E-04	9.09E-03	5.54E-03	5.07E-03	5.15E-03	3.56E-04	3.56E-04	9.09E-03	5.54E-03	5.07E-03	5.15E-03
	Clearwater River downstream of Broach Lake	3.56E-04	3.56E-04	9.09E-03	5.54E-03	5.07E-03	5.15E-03	3.56E-04	3.56E-04	9.09E-03	5.54E-03	5.07E-03	5.15E-03
	Patterson Lake North Arm - East Basin	3.58E-04	3.58E-04	9.12E-03	5.55E-03	5.08E-03	5.16E-03	3.56E-04	3.56E-04	9.09E-03	5.54E-03	5.07E-03	5.15E-03
	Patterson Lake North Arm - West Basin	3.88E-03	3.89E-03	5.17E-02	9.63E-03	7.74E-03	7.82E-03	1.15E-02	1.16E-02	9.72E-03	7.14E-03	5.30E-03	5.39E-03
	Patterson Lake South Arm	5.79E-03	5.81E-03	5.50E-02	1.13E-02	7.87E-03	7.90E-03	3.44E-03	3.46E-03	9.48E-03	6.13E-03	5.08E-03	5.17E-03
Water (Bq/L)	Forrest Lake North Basin	3.79E-03	3.80E-03	3.81E-02	9.18E-03	6.85E-03	6.89E-03	2.23E-03	2.24E-03	9.32E-03	5.90E-03	5.08E-03	5.16E-03
	Beet Lake	2.24E-03	2.25E-03	2.97E-02	7.76E-03	5.47E-03	5.56E-03	1.50E-03	1.51E-03	9.29E-03	5.79E-03	5.07E-03	5.16E-03
	Naomi Lake	1.68E-03	1.68E-03	2.48E-02	7.14E-03	5.26E-03	5.35E-03	1.16E-03	1.17E-03	9.25E-03	5.72E-03	5.07E-03	5.15E-03
	Clearwater River upstream Mirror River Confluence	1.08E-03	1.08E-03	1.76E-02	6.41E-03	5.17E-03	5.26E-03	7.91E-04	7.93E-04	9.17E-03	5.64E-03	5.07E-03	5.15E-03
	Clearwater River downstream Mirror River Confluence	5.78E-04	5.79E-04	1.17E-02	5.81E-03	5.10E-03	5.19E-03	4.87E-04	4.87E-04	9.11E-03	5.57E-03	5.07E-03	5.15E-03
	Lloyd Lake Inlet	5.31E-04	5.32E-04	1.12E-02	5.75E-03	5.09E-03	5.18E-03	4.60E-04	4.60E-04	9.11E-03	5.56E-03	5.07E-03	5.15E-03
	Reference (Broach Lake)	6.84E+00	6.84E+00	2.09E+01	6.47E+01	3.59E+02	3.65E+02	6.84E+00	6.84E+00	2.09E+01	6.47E+01	3.59E+02	3.65E+02
	Patterson Lake North Arm - East Basin	6.86E+00	6.86E+00	2.09E+01	6.48E+01	3.59E+02	3.66E+02	6.84E+00	6.84E+00	2.09E+01	6.47E+01	3.59E+02	3.65E+02
C 1: /D //	Patterson Lake North Arm - West Basin	6.16E+01	6.18E+01	1.16E+02	1.04E+02	4.92E+02	5.00E+02	2.22E+02	2.22E+02	2.23E+01	8.34E+01	3.76E+02	3.82E+02
Sediment (Bq/kg dw)	Patterson Lake South Arm	3.84E+01	3.85E+01	8.94E+01	8.82E+01	4.41E+02	4.48E+02	6.63E+01	6.65E+01	2.18E+01	7.16E+01	3.60E+02	3.66E+02
uw)	Beet Lake	1.85E+01	1.85E+01	5.74E+01	7.49E+01	3.71E+02	3.77E+02	2.89E+01	2.90E+01	2.13E+01	6.77E+01	3.59E+02	3.66E+02
	Naomi Lake	1.50E+01	1.50E+01	4.90E+01	7.22E+01	3.65E+02	3.71E+02	2.24E+01	2.24E+01	2.12E+01	6.69E+01	3.59E+02	3.65E+02
	Lloyd Lake Inlet	7.91E+00	7.91E+00	2.46E+01	6.57E+01	3.60E+02	3.66E+02	8.84E+00	8.85E+00	2.09E+01	6.50E+01	3.59E+02	3.65E+02
	Reference (Broach Lake)	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
	Patterson Lake North Arm	9.43E-05	9.43E-05	9.43E-05	9.43E-05	9.43E-05	9.43E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
A:u (Dar (ma3)	Patterson Lake South Arm	1.81E-05	1.81E-05	1.81E-05	1.81E-05	1.81E-05	1.81E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Air (Bq/m³)	Beet Lake	9.92E-06	9.92E-06	9.92E-06	9.92E-06	9.92E-06	9.92E-06	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
	Clearwater River and Mirror River Confluence	7.23E-07	7.23E-07	7.23E-07	7.23E-07	7.23E-07	7.23E-07	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
	Lloyd Lake	4.73E-07	4.73E-07	4.73E-07	4.73E-07	4.73E-07	4.73E-07	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
	Reference (Broach Lake)	3.06E+00	3.06E+00	2.00E+01	1.79E+01	4.00E+01	1.12E+01	3.06E+00	3.06E+00	2.00E+01	1.79E+01	4.00E+01	1.12E+01
	Patterson Lake North Arm	1.36E+01	1.36E+01	3.18E+01	2.99E+01	4.77E+01	1.91E+01	4.16E+00	4.16E+00	2.66E+01	2.60E+01	4.00E+01	1.12E+01
Soil (Bq/kg dw)	Patterson Lake South Arm	5.08E+00	5.08E+00	2.23E+01	2.02E+01	4.15E+01	1.27E+01	3.27E+00	3.27E+00	2.13E+01	1.95E+01	4.00E+01	1.12E+01
Son (bq/kg uw)	Beet Lake	4.16E+00	4.16E+00	2.12E+01	1.92E+01	4.08E+01	1.20E+01	3.17E+00	3.17E+00	2.07E+01	1.88E+01	4.00E+01	1.12E+01
	Clearwater River and Mirror River Confluence	3.14E+00	3.14E+00	2.01E+01	1.80E+01	4.01E+01	1.13E+01	3.06E+00	3.06E+00	2.01E+01	1.80E+01	4.00E+01	1.12E+01
	Lloyd Lake	3.11E+00	3.11E+00	2.01E+01	1.80E+01	4.00E+01	1.13E+01	3.06E+00	3.06E+00	2.00E+01	1.80E+01	4.00E+01	1.12E+01

Table C.7: Maximum Concentrations of non-radiological COPCs in Biota for the Application Case and Upper Bound Scenario - Project Lifespan

						Maximum Con	centration P	roject Lifes	pan (mg/kg	fw)		
	Biota	Location			Application	ı Case			U	pper Bound	Scenario	
			Arsenic	Cobalt	Copper	Molybdenum	Uranium	Arsenic	Cobalt	Copper	Molybdenum	Uranium
		Reference (Broach Lake)	2.13E-02	9.21E-02	1.83E-01	1.40E-03	5.06E-03	2.13E-02	9.21E-02	1.83E-01	1.40E-03	5.06E-03
		Patterson Lake North Arm - West Basin	7.50E-02	1.22E-01	2.16E-01	4.07E-03	1.93E-02	7.68E-02	1.23E-01	2.17E-01	9.45E-03	4.66E-02
	Macrophytes	Patterson Lake South Arm	2.57E-02	1.08E-01	2.00E-01	2.77E-03	8.89E-03	2.58E-02	1.09E-01	2.01E-01	5.54E-03	1.56E-02
	. ,	Beet Lake	2.20E-02	1.00E-01	1.92E-01	2.10E-03	6.47E-03	2.20E-02	1.01E-01	1.92E-01	3.54E-03	8.93E-03
		Lloyd Lake	2.13E-02	9.30E-02	1.84E-01	1.48E-03	5.19E-03	2.13E-02	9.30E-02	1.84E-01	1.62E-03	5.41E-03
Aquatic Plants		Reference (Broach Lake)	2.13E-02	9.21E-02	1.83E-01	1.40E-03	5.06E-03	2.13E-02	9.21E-02	1.83E-01	1.40E-03	5.06E-03
		Patterson Lake North Arm - West Basin	7.50E-02	1.22E-01	2.16E-01	4.07E-03	1.93E-02	7.68E-02	1.23E-01	2.17E-01	9.45E-03	4.66E-02
	Phytoplankton	Patterson Lake South Arm	2.57E-02	1.08E-01	2.00E-01	2.77E-03	8.89E-03	2.58E-02	1.09E-01	2.01E-01	5.54E-03	1.56E-02
		Beet Lake	2.20E-02	1.00E-01	1.92E-01	2.10E-03	6.47E-03	2.20E-02	1.01E-01	1.92E-01	3.54E-03	8.93E-03
		Lloyd Lake	2.13E-02	9.30E-02	1.84E-01	1.48E-03	5.19E-03	2.13E-02	9.30E-02	1.84E-01	1.62E-03	5.41E-03
		Reference (Broach Lake)	5.60E-02	1.16E-01	6.53E+00	3.32E-02	9.64E-03	5.60E-02	1.16E-01	6.53E+00	3.32E-02	9.64E-03
		Patterson Lake North Arm - West Basin	1.71E-01	1.50E-01	7.59E+00	8.87E-02	3.16E-02	1.74E-01	1.51E-01	7.62E+00	2.02E-01	7.10E-02
	Benthic	Patterson Lake South Arm	6.55E-02	1.34E-01	7.07E+00	6.18E-02	1.55E-02	6.57E-02	1.35E-01	7.09E+00	1.22E-01	2.59E-02
	Invertebrates	Beet Lake	5.75E-02	1.25E-01	6.80E+00	4.78E-02	1.18E-02	5.76E-02	1.26E-01	6.81E+00	7.88E-02	1.56E-02
		Lloyd Lake	5.61E-02	1.17E-01	6.56E+00	3.47E-02	9.84E-03	5.61E-02	1.17E-01	6.56E+00	3.78E-02	1.02E-02
		Reference (Broach Lake)	6.68E-02	1.17E-01	6.57E+00	3.34E-02	1.00E-02	6.68E-02	1.17E-01	6.57E+00	3.34E-02	1.00E-02
		Patterson Lake North Arm - West Basin	2.35E-01	1.54E-01	7.76E+00	9.69E-02	3.84E-02	2.41E-01	1.56E-01	7.79E+00	2.25E-01	9.25E-02
	Zooplankton	Patterson Lake South Arm	8.07E-02	1.37E-01	7.20E+00	6.59E-02	1.76E-02	8.11E-02	1.38E-01	7.21E+00	1.32E-01	3.09E-02
	•	Beet Lake	6.91E-02	1.27E-01	6.89E+00	5.01E-02	1.28E-02	6.91E-02	1.28E-01	6.90E+00	8.43E-02	1.77E-02
		Lloyd Lake	6.69E-02	1.18E-01	6.61E+00	3.51E-02	1.03E-02	6.69E-02	1.18E-01	6.61E+00	3.87E-02	1.07E-02
Aquatic Animals		Reference (Broach Lake)	3.71E-02	7.00E-03	4.21E-01	2.06E-05	2.01E-03	3.71E-02	7.00E-03	4.21E-01	2.06E-05	2.01E-03
		Patterson Lake North Arm - West Basin	1.31E-01	9.25E-03	4.97E-01	5.98E-05	7.67E-03	1.34E-01	9.34E-03	5.00E-01	1.39E-04	1.85E-02
	Northern Pike	Patterson Lake South Arm	4.49E-02	8.22E-03	4.61E-01	4.07E-05	3.53E-03	4.51E-02	8.27E-03	4.62E-01	8.15E-05	6.19E-03
		Beet Lake	3.84E-02	7.63E-03	4.42E-01	3.09E-05	2.57E-03	3.84E-02	7.66E-03	4.42E-01	5.20E-05	3.54E-03
		Lloyd Lake	3.72E-02	7.06E-03	4.23E-01	2.17E-05	2.06E-03	3.72E-02	7.06E-03	4.24E-01	2.39E-05	2.15E-03
		Reference (Broach Lake)	3.56E-02	6.99E-03	4.21E-02	2.06E-05	4.97E-04	3.56E-02	6.99E-03	4.21E-02	2.06E-05	4.97E-04
		Patterson Lake North Arm - West Basin	1.22E-01	9.18E-03	4.95E-02	5.83E-05	1.83E-03	1.24E-01	9.27E-03	4.97E-02	1.34E-04	4.30E-03
	Whitefish	Patterson Lake South Arm	4.27E-02	8.16E-03	4.59E-02	3.98E-05	8.50E-04	4.29E-02	8.21E-03	4.60E-02	7.95E-05	1.47E-03
		Beet Lake	3.68E-02	7.60E-03	4.40E-02	3.04E-05	6.26E-04	3.68E-02	7.62E-03	4.41E-02	5.09E-05	8.55E-04
		Lloyd Lake	3.57E-02	7.05E-03	4.23E-02	2.16E-05	5.09E-04	3.57E-02	7.05E-03	4.23E-02	2.37E-05	5.30E-04
		Reference (Broach Lake)	2.51E-02	2.51E-02	3.99E-01	1.42E-02	4.01E-02	2.51E-02	2.51E-02	3.99E-01	1.42E-02	4.01E-02
		Patterson Lake North Arm - West Basin	2.54E-02	2.54E-02	4.12E-01	1.58E-02	2.91E-01	2.54E-02	2.54E-02	4.12E-01	1.58E-02	2.91E-01
	Blueberries	Patterson Lake South Arm	2.52E-02	2.52E-02	4.01E-01	1.45E-02	7.59E-02	2.52E-02	2.52E-02	4.01E-01	1.45E-02	7.59E-02
		Beet Lake	2.52E-02	2.52E-02	4.00E-01	1.44E-02	6.12E-02	2.52E-02	2.52E-02	4.00E-01	1.44E-02	6.12E-02
		Lloyd Lake	2.52E-02	2.52E-02	4.00E-01	1.44E-02	4.09E-02	2.52E-02	2.52E-02	4.00E-01	1.44E-02	4.09E-02
Terrestrial Plants		Reference (Broach Lake)	5.02E-02	5.01E-02	7.98E-01	2.85E-02	8.01E-02	5.02E-02	5.01E-02	7.98E-01	2.85E-02	8.01E-02
		Patterson Lake North Arm - West Basin	5.10E-02	5.09E-02	8.27E-01	3.27E-02	6.73E-01	5.10E-02	5.09E-02	8.27E-01	3.27E-02	6.73E-01
	Browse	Patterson Lake South Arm	5.05E-02	5.05E-02	8.03E-01	2.92E-02	1.65E-01	5.05E-02	5.05E-02	8.03E-01	2.92E-02	1.65E-01
		Beet Lake	5.05E-02	5.05E-02	8.01E-01	2.88E-02	1.30E-01	5.05E-02	5.05E-02	8.01E-01	2.88E-02	1.30E-01
		Lloyd Lake	5.05E-02	5.05E-02	7.99E-01	2.88E-02	8.20E-02	5.05E-02	•	7.99E-01	2.88E-02	8.20E-02

Table C.7: Maximum Concentrations of non-radiological COPCs in Biota for the Application Case and Upper Bound Scenario - Project Lifespan

						Maximum Con	centrati <u>on P</u>	roject Lifes	pan (mg/ko	fw)		
	Biota	Location			Application					pper Bound	Scenario	
			Arsenic	Cobalt	Copper	Molybdenum	Uranium	Arsenic	Cobalt	Copper	Molybdenum	Uranium
		Reference (Broach Lake)	5.02E-02	5.01E-02	7.98E-01	2.85E-02	8.01E-02	5.02E-02	5.01E-02	7.98E-01	2.85E-02	8.01E-02
		Patterson Lake North Arm - West Basin	5.80E-02	5.56E-02	8.66E-01	6.06E-02	2.85E+00	5.80E-02	5.56E-02	8.66E-01	6.06E-02	2.85E+00
	Labrador Tea	Patterson Lake South Arm	5.28E-02	5.28E-02	8.10E-01	3.38E-02	4.76E-01	5.28E-02	5.28E-02	8.10E-01	3.38E-02	4.76E-01
		Beet Lake	5.28E-02	5.28E-02	8.06E-01	3.11E-02	3.14E-01	5.28E-02	5.28E-02	8.06E-01	3.11E-02	3.14E-01
		Lloyd Lake	5.28E-02	5.28E-02	8.02E-01	3.11E-02	8.90E-02	5.28E-02	5.28E-02	8.02E-01	3.11E-02	8.90E-02
		Reference (Broach Lake)	5.58E-02	7.33E-02	7.58E-01	6.70E-02	1.68E-02	5.58E-02	7.33E-02	7.58E-01	6.70E-02	1.68E-02
		Patterson Lake North Arm - West Basin	1.00E-01	1.03E-01	1.01E+00	2.46E-01	1.40E+01	1.00E-01	1.03E-01	1.01E+00	2.46E-01	1.40E+01
	Lichen	Patterson Lake South Arm	7.07E-02	8.82E-02	8.03E-01	9.68E-02	2.01E+00	7.07E-02	8.82E-02	8.03E-01	9.68E-02	2.01E+00
		Beet Lake	7.07E-02	8.82E-02	7.88E-01	8.19E-02	1.19E+00	7.07E-02	8.82E-02	7.88E-01	8.19E-02	1.19E+00
		Lloyd Lake	7.07E-02	8.82E-02	7.73E-01	8.19E-02	6.15E-02	7.07E-02	8.82E-02	7.73E-01	8.19E-02	6.15E-02
		Reference (Broach Lake)	2.24E-02	2.63E-03	5.26E-02	3.73E-02	2.19E-03	2.24E-02	2.63E-03	5.26E-02	3.73E-02	2.19E-03
		Patterson Lake North Arm - West Basin	2.33E-02	3.19E-03	5.88E-02	4.18E-02	2.67E-01	2.33E-02	3.19E-03	5.88E-02	4.18E-02	2.67E-01
	Terrestrial	Patterson Lake South Arm	2.27E-02	2.91E-03	5.37E-02	3.80E-02	4.00E-02	2.27E-02	2.91E-03	5.37E-02	3.80E-02	4.00E-02
	Invertebrate	Beet Lake	2.27E-02	2.91E-03	5.33E-02	3.76E-02	2.45E-02	2.27E-02	2.91E-03	5.33E-02	3.76E-02	2.45E-02
		Lloyd Lake	2.27E-02	2.91E-03	5.29E-02	3.76E-02	3.03E-03	2.27E-02	2.91E-03	5.29E-02	3.76E-02	3.03E-03
		Reference (Broach Lake)	5.56E-02	1.25E-03	3.78E-01	1.36E-03	1.48E-03	5.56E-02	1.25E-03	3.78E-01	1.36E-03	1.48E-03
		Patterson Lake North Arm - West Basin	7.25E-02	1.33E-03	3.93E-01	1.61E-03	1.23E-02	7.29E-02	1.34E-03	3.93E-01	1.70E-03	1.24E-02
	Beaver	Patterson Lake South Arm	5.72E-02	1.29E-03	3.81E-01	1.42E-03	3.03E-03	5.72E-02	1.29E-03	3.81E-01	1.47E-03	3.07E-03
		Beet Lake	5.61E-02	1.27E-03	3.80E-01	1.39E-03	2.39E-03	5.61E-02	1.28E-03	3.80E-01	1.41E-03	2.40E-03
		Lloyd Lake	5.59E-02	1.26E-03	3.79E-01	1.38E-03	1.52E-03	5.59E-02	1.26E-03	3.79E-01	1.38E-03	1.52E-03
		Reference (Broach Lake)	5.21E-02	8.47E-04	3.52E-01	1.01E-03	1.10E-03	5.21E-02	8.47E-04	3.52E-01	1.01E-03	1.10E-03
	Diagle Dage	Patterson Lake North Arm - West Basin	6.97E-02	8.74E-04	3.67E-01	1.07E-03	4.55E-03	7.03E-02	8.75E-04	3.68E-01	1.09E-03	4.60E-03
	Black Bear	Beet Lake	5.46E-02	8.91E-04	3.55E-01	1.04E-03	1.40E-03	5.46E-02	8.91E-04	3.55E-01	1.05E-03	1.40E-03
		Lloyd Lake	5.23E-02	8.58E-04	3.53E-01	1.02E-03	1.12E-03	5.23E-02	8.58E-04	3.53E-01	1.02E-03	1.12E-03
Terrestrial		Reference (Broach Lake)	4.71E-02	8.12E-05	1.04E-01	9.04E-05	4.57E-05	4.71E-02	8.12E-05	1.04E-01	9.04E-05	4.57E-05
Animals	Grey Wolf	Patterson Lake North-West Basin	5.57E-02	8.27E-05	1.06E-01	9.96E-05	1.47E-04	5.59E-02	8.28E-05	1.06E-01	1.08E-04	1.49E-04
		Lloyd Lake	4.78E-02	8.18E-05	1.04E-01	9.27E-05	4.71E-05	4.78E-02	8.18E-05	1.04E-01	9.34E-05	4.72E-05
		Reference (Broach Lake)	2.80E-02	2.42E-04	2.78E-01	1.40E-04	5.15E-05	2.80E-02	2.42E-04	2.78E-01	1.40E-04	5.15E-05
		Patterson Lake North Arm - West Basin	8.43E-02	3.06E-04	3.24E-01	3.40E-04	1.98E-04	8.56E-02	3.08E-04	3.25E-01	7.49E-04	3.53E-04
	Mink	Patterson Lake South Arm	3.27E-02	2.76E-04	3.02E-01	2.43E-04	8.27E-05	3.28E-02	2.77E-04	3.02E-01	4.59E-04	1.24E-04
		Beet Lake	2.88E-02	2.59E-04	2.90E-01	1.92E-04	6.47E-05	2.88E-02	2.60E-04	2.91E-01	3.04E-04	7.99E-05
		Lloyd Lake	2.81E-02	2.44E-04	2.80E-01	1.45E-04	5.24E-05	2.81E-02	2.44E-04	2.80E-01	1.56E-04	5.38E-05
		Reference (Broach Lake)	5.87E-02	1.37E-03	3.51E-01	1.26E-03	1.37E-03	5.87E-02	1.37E-03	3.51E-01	1.26E-03	1.37E-03
		Patterson Lake North Arm - West Basin	9.28E-02	1.53E-03	3.67E-01	1.53E-03	1.11E-02	9.36E-02	1.54E-03	3.67E-01	1.73E-03	1.14E-02
	Moose	Patterson Lake South Arm	9.26E-02	1.52E-03	3.66E-01	1.51E-03	1.11E-02	9.34E-02	1.53E-03	3.66E-01	1.68E-03	1.13E-02
		Beet Lake	5.94E-02	1.42E-03	3.54E-01	1.29E-03	2.19E-03	5.94E-02	1.42E-03	3.54E-01	1.34E-03	2.22E-03
		Lloyd Lake	5.90E-02	1.38E-03	3.52E-01	1.27E-03	1.40E-03	5.90E-02	1.38E-03	3.52E-01	1.28E-03	1.40E-03
		Reference (Broach Lake)	4.99E-01	3.19E-01	1.40E+01	1.26E-01	2.42E-03	4.99E-01	3.19E-01	1.40E+01	1.26E-01	2.42E-03
	Moose Organs	Patterson Lake North Arm - West Basin	7.89E-01	3.56E-01	1.47E+01	1.53E-01	1.97E-02	7.96E-01	3.57E-01	1.47E+01	1.73E-01	2.02E-02
		Patterson Lake South Arm	7.87E-01	3.53E-01	1.46E+01	1.51E-01	1.96E-02	7.94E-01	3.55E-01	1.47E+01	1.68E-01	2.01E-02

Table C.7: Maximum Concentrations of non-radiological COPCs in Biota for the Application Case and Upper Bound Scenario - Project Lifespan

					Maximum Con	centration P	roject Life <u>s</u>	pan (mg/ko	g fw)		
Biota	Location			Application					pper Bound	Scenario	
		Arsenic	Cobalt	Copper	Molybdenum	Uranium	Arsenic	Cobalt	Copper	Molybdenum	Uranium
	Beet Lake	5.05E-01	3.30E-01	1.41E+01	1.29E-01	3.88E-03	5.05E-01	3.30E-01	1.41E+01	1.34E-01	3.93E-03
	Lloyd Lake	5.01E-01	3.22E-01	1.41E+01	1.27E-01	2.48E-03	5.01E-01	3.22E-01	1.41E+01	1.28E-01	2.48E-03
	Reference (Broach Lake)	7.19E-02	8.64E-04	5.12E-01	4.59E-04	2.70E-04	7.19E-02	8.64E-04	5.12E-01	4.59E-04	2.70E-04
	Patterson Lake North Arm - West Basin	2.20E-01	1.13E-03	5.96E-01	1.23E-03	8.89E-04	2.24E-01	1.14E-03	5.98E-01	2.80E-03	1.99E-03
Muskrat	Patterson Lake South Arm	8.41E-02	1.00E-03	5.55E-01	8.54E-04	4.33E-04	8.44E-02	1.01E-03	5.56E-01	1.69E-03	7.27E-04
	Beet Lake	7.38E-02	9.36E-04	5.34E-01	6.61E-04	3.30E-04	7.39E-02	9.39E-04	5.35E-01	1.09E-03	4.38E-04
	Lloyd Lake	7.20E-02	8.71E-04	5.15E-01	4.79E-04	2.75E-04	7.20E-02	8.72E-04	5.15E-01	5.23E-04	2.85E-04
	Reference (Broach Lake)	7.78E-03	7.79E-05	2.78E-02	2.35E-04	8.18E-05	7.78E-03	7.79E-05	2.78E-02	2.35E-04	8.18E-05
	Patterson Lake North Arm - West Basin	8.01E-03	8.08E-05	2.89E-02	2.67E-04	9.88E-04	8.01E-03	8.09E-05	2.89E-02	2.79E-04	9.91E-04
Red Fox	Patterson Lake South Arm	7.83E-03	7.94E-05	2.80E-02	2.42E-04	2.12E-04	7.83E-03	7.94E-05	2.80E-02	2.48E-04	2.12E-04
	Beet Lake	7.83E-03	7.91E-05	2.79E-02	2.38E-04	1.58E-04	7.83E-03	7.91E-05	2.79E-02	2.41E-04	1.58E-04
	Lloyd Lake	7.82E-03	7.88E-05	2.78E-02	2.37E-04	8.47E-05	7.82E-03	7.88E-05	2.78E-02	2.37E-04	8.47E-05
	Reference (Broach Lake)	4.69E-02	9.87E-04	3.44E-01	1.27E-03	1.37E-03	4.69E-02	9.87E-04	3.44E-01	1.27E-03	1.37E-03
	Patterson Lake North Arm - West Basin	4.76E-02	1.00E-03	3.57E-01	1.45E-03	1.13E-02	4.76E-02	1.00E-03	3.57E-01	1.46E-03	1.13E-02
Snowshoe Hare	Patterson Lake South Arm	4.71E-02	9.95E-04	3.46E-01	1.30E-03	2.79E-03	4.71E-02	9.95E-04	3.46E-01	1.31E-03	2.79E-03
	Beet Lake	4.71E-02	9.95E-04	3.46E-01	1.29E-03	2.21E-03	4.71E-02	9.95E-04	3.46E-01	1.29E-03	2.21E-03
	Lloyd Lake	4.71E-02	9.95E-04	3.45E-01	1.28E-03	1.40E-03	4.71E-02	9.95E-04	3.45E-01	1.29E-03	1.40E-03
	Reference (Broach Lake)	1.67E-02	3.54E-04	1.26E-01	4.59E-04	4.98E-04	1.67E-02	3.54E-04	1.26E-01	4.59E-04	4.98E-04
	Patterson Lake North Arm - West Basin	1.69E-02	3.59E-04	1.30E-01	5.19E-04	3.87E-03	1.69E-02	3.59E-04	1.30E-01	5.25E-04	3.87E-03
Southern Red- Backed Vole	Patterson Lake South Arm	1.67E-02	3.57E-04	1.27E-01	4.70E-04	9.79E-04	1.67E-02	3.57E-04	1.27E-01	4.73E-04	9.80E-04
Backed Voie	Beet Lake	1.67E-02	3.57E-04	1.26E-01	4.65E-04	7.82E-04	1.67E-02	3.57E-04	1.26E-01	4.66E-04	7.82E-04
	Lloyd Lake	1.67E-02	3.56E-04	1.26E-01	4.64E-04	5.09E-04	1.67E-02	3.56E-04	1.26E-01	4.64E-04	5.09E-04
	Reference (Broach Lake)	6.30E-02	1.08E-03	1.92E-01	1.03E-03	7.02E-04	6.30E-02	1.08E-03	1.92E-01	1.03E-03	7.02E-04
Woodland	Patterson Lake North Arm - West Basin	1.01E-01	1.24E-03	2.05E-01	1.78E-03	1.97E-02	1.02E-01	1.25E-03	2.05E-01	2.07E-03	2.00E-02
Caribou	Beet Lake	6.43E-02	1.12E-03	1.93E-01	1.10E-03	2.21E-03	6.43E-02	1.12E-03	1.93E-01	1.15E-03	2.23E-03
	Lloyd Lake	6.49E-02	1.12E-03	1.94E-01	1.13E-03	8.23E-04	6.49E-02	1.12E-03	1.94E-01	1.14E-03	8.27E-04
	Reference (Broach Lake)	3.71E-02	2.88E-02	2.59E-01	2.87E-03	3.27E-02	3.71E-02	2.88E-02	2.59E-01	2.87E-03	3.27E-02
	Patterson Lake North Arm - West Basin	3.76E-02	2.92E-02	2.69E-01	3.26E-03	2.61E-01	3.77E-02	2.92E-02	2.69E-01	3.27E-03	2.61E-01
Grouse	Patterson Lake South Arm	3.73E-02	2.90E-02	2.61E-01	2.94E-03	6.53E-02	3.73E-02	2.90E-02	2.61E-01	2.95E-03	6.53E-02
	Beet Lake	3.73E-02	2.90E-02	2.61E-01	2.91E-03	5.19E-02	3.73E-02	2.90E-02	2.61E-01	2.91E-03	5.19E-02
	Lloyd Lake	3.73E-02	2.90E-02	2.60E-01	2.91E-03	3.34E-02	3.73E-02	2.90E-02	2.60E-01	2.91E-03	3.34E-02
	Reference (Broach Lake)	5.23E-03	4.08E-03	3.63E-02	4.06E-04	4.59E-03	5.23E-03	4.08E-03	3.63E-02	4.06E-04	4.59E-03
	Patterson Lake North Arm - West Basin	5.33E-03	4.15E-03	3.77E-02	4.68E-04	3.78E-02	5.33E-03	4.15E-03	3.77E-02	4.81E-04	3.78E-02
Canada Goose	Patterson Lake South Arm	5.25E-03	4.12E-03	3.66E-02	4.18E-04	9.33E-03	5.25E-03	4.12E-03	3.66E-02	4.25E-04	9.34E-03
	Beet Lake	5.25E-03	4.11E-03	3.65E-02	4.12E-04	7.38E-03	5.25E-03	4.11E-03	3.65E-02	4.15E-04	7.38E-03
	Lloyd Lake	5.25E-03	4.11E-03	3.64E-02	4.10E-04	4.69E-03	5.25E-03	4.11E-03	3.64E-02	4.11E-04	4.69E-03
	Reference (Broach Lake)	1.97E-02	8.78E-04	1.15E+00	5.85E-04	6.64E-05	1.97E-02	8.78E-04	1.15E+00	5.85E-04	6.64E-05
Little Brown	Patterson Lake North Arm - West Basin	6.02E-02	1.14E-03	1.34E+00	1.56E-03	2.18E-04	6.12E-02	1.15E-03	1.34E+00	3.57E-03	4.89E-04
Myotis	Patterson Lake South Arm	2.31E-02	1.02E-03	1.24E+00	1.09E-03	1.06E-04	2.32E-02	1.02E-03	1.25E+00	2.15E-03	1.79E-04
	Beet Lake	2.03E-02	9.50E-04	1.20E+00	8.44E-04	8.11E-05	2 025 02	9.53E-04	1.20E+00	1.39E-03	1.08E-04

Table C.7: Maximum Concentrations of non-radiological COPCs in Biota for the Application Case and Upper Bound Scenario - Project Lifespan

					Maximum Con	centration P	roject Lifes	pan (mg/kg	j fw)		
Biota	Location			Application	Case			U	pper Bound	Scenario	
		Arsenic	Cobalt	Copper	Molybdenum	Uranium	Arsenic	Cobalt	Copper	Molybdenum	Uranium
	Lloyd Lake	1.98E-02	8.86E-04	1.15E+00	6.12E-04	6.77E-05	1.98E-02	8.86E-04	1.15E+00	6.68E-04	7.01E-05
	Reference (Broach Lake)	1.44E-02	5.39E-03	1.97E-01	1.88E-04	6.47E-04	1.44E-02	5.39E-03	1.97E-01	1.88E-04	6.47E-04
	Patterson Lake North Arm - West Basin	4.98E-02	7.03E-03	2.30E-01	5.03E-04	2.37E-03	5.09E-02	7.09E-03	2.31E-01	1.15E-03	5.49E-03
Loon	Patterson Lake South Arm	1.74E-02	6.26E-03	2.14E-01	3.51E-04	1.10E-03	1.74E-02	6.29E-03	2.15E-01	6.92E-04	1.89E-03
	Beet Lake	1.49E-02	5.84E-03	2.06E-01	2.71E-04	8.13E-04	1.49E-02	5.86E-03	2.06E-01	4.47E-04	1.10E-03
	Lloyd Lake	1.45E-02	5.44E-03	1.98E-01	1.97E-04	6.62E-04	1.45E-02	5.44E-03	1.98E-01	2.15E-04	6.88E-04
	Reference (Broach Lake)	5.77E-02	4.36E-02	1.13E+00	2.04E-03	6.80E-03	5.77E-02	4.36E-02	1.13E+00	2.04E-03	6.80E-03
	Patterson Lake North Arm - West Basin	1.77E-01	5.66E-02	1.32E+00	5.44E-03	2.23E-02	1.80E-01	5.71E-02	1.32E+00	1.24E-02	5.02E-02
Mallard	Patterson Lake South Arm	6.76E-02	5.05E-02	1.23E+00	3.79E-03	1.09E-02	6.78E-02	5.08E-02	1.23E+00	7.48E-03	1.83E-02
	Beet Lake	5.93E-02	4.72E-02	1.18E+00	2.94E-03	8.30E-03	5.94E-02	4.74E-02	1.18E+00	4.83E-03	1.10E-02
	Lloyd Lake	5.78E-02	4.40E-02	1.14E+00	2.13E-03	6.93E-03	5.78E-02	4.40E-02	1.14E+00	2.32E-03	7.18E-03
	Reference (Broach Lake)	7.99E-02	1.09E-01	3.39E+00	6.19E-03	1.83E-02	7.99E-02	1.09E-01	3.39E+00	6.19E-03	1.83E-02
	Patterson Lake North Arm - West Basin	1.93E-01	1.36E-01	3.93E+00	1.44E-02	1.00E-01	1.96E-01	1.37E-01	3.95E+00	3.12E-02	1.24E-01
Rusty Blackbird	Patterson Lake South Arm	8.92E-02	1.23E-01	3.67E+00	1.04E-02	3.15E-02	8.95E-02	1.24E-01	3.67E+00	1.93E-02	3.75E-02
	Beet Lake	8.14E-02	1.16E-01	3.53E+00	8.35E-03	2.53E-02	8.14E-02	1.16E-01	3.54E+00	1.29E-02	2.74E-02
	Lloyd Lake	8.00E-02	1.09E-01	3.40E+00	6.41E-03	1.86E-02	8.00E-02	1.09E-01	3.41E+00	6.88E-03	1.88E-02

Table C.8: Maximum Concentrations of non-radiological COPCs in Biota for the Application Case and Upper Bound Scenario – Far-Future

						Maximum Co	oncentration	n Far-Future	(mg/ka fw)		
	Biota	Location			Application					per Bound	Scenario	
	210111		Arsenic	Cobalt	Copper	Molybdenum	Uranium	Arsenic	Cobalt	Copper	Molybdenum	Uranium
		Reference (Broach Lake)	2.13E-02	9.21E-02	1.83E-01	1.40E-03	5.06E-03	2.13E-02	9.21E-02	1.83E-01	1.40E-03	5.06E-03
		Patterson Lake North Arm - West Basin	2.26E-02	2.26E-01	4.73E-01	3.13E-02	5.06E-02	2.33E-02	2.96E-01	6.19E-01	1.16E-01	5.06E-02
	Macrophytes	Patterson Lake South Arm	2.14E-02	1.73E-01	3.53E-01	1.88E-02	1.77E-02	2.14E-02	2.16E-01	4.39E-01	6.80E-02	1.77E-02
		Beet Lake	2.13E-02	1.35E-01	2.72E-01	1.04E-02	9.74E-03	2.13E-02	1.58E-01	3.17E-01	3.61E-02	9.74E-03
		Lloyd Lake	2.13E-02	9.65E-02	1.92E-01	2.32E-03	5.48E-03	2.13E-02	9.88E-02	1.97E-01	4.94E-03	5.48E-03
Aquatic Plants		Reference (Broach Lake)	2.13E-02	9.21E-02	1.83E-01	1.40E-03	5.06E-03	2.13E-02	9.21E-02	1.83E-01	1.40E-03	5.06E-03
		Patterson Lake North Arm - West Basin	2.26E-02	2.26E-01	4.73E-01	3.13E-02	5.06E-02	2.33E-02	2.96E-01	6.19E-01	1.16E-01	5.06E-02
	Phytoplankton	Patterson Lake South Arm	2.14E-02	1.73E-01	3.53E-01	1.88E-02	1.77E-02	2.14E-02	2.16E-01	4.39E-01	6.80E-02	1.77E-02
		Beet Lake	2.13E-02	1.35E-01	2.72E-01	1.04E-02	9.74E-03	2.13E-02	1.58E-01	3.17E-01	3.61E-02	9.74E-03
		Lloyd Lake	2.13E-02	9.65E-02	1.92E-01	2.32E-03	5.48E-03	2.13E-02	9.88E-02	1.97E-01	4.94E-03	5.48E-03
		Reference (Broach Lake)	5.60E-02	1.16E-01	6.53E+00	3.32E-02	9.64E-03	5.60E-02	1.16E-01	6.53E+00	3.32E-02	9.64E-03
	B 41:	Patterson Lake North Arm - West Basin	5.94E-02	2.84E-01	1.69E+01	7.40E-01	9.65E-02	6.12E-02	3.73E-01	2.21E+01	2.75E+00	9.66E-02
	Benthic Invertebrates	Patterson Lake South Arm	5.63E-02	2.18E-01	1.26E+01	4.44E-01	3.37E-02	5.64E-02	2.72E-01	1.57E+01	1.61E+00	3.37E-02
	ilivertebrates	Beet Lake	5.61E-02	1.70E-01	9.71E+00	2.47E-01	1.86E-02	5.61E-02	1.98E-01	1.13E+01	8.54E-01	1.86E-02
		Lloyd Lake	5.60E-02	1.21E-01	6.85E+00	5.50E-02	1.05E-02	5.60E-02	1.24E-01	7.01E+00	1.17E-01	1.05E-02
		Reference (Broach Lake)	6.68E-02	1.17E-01	6.57E+00	3.34E-02	1.00E-02	6.68E-02	1.17E-01	6.57E+00	3.34E-02	1.00E-02
		Patterson Lake North Arm - West Basin	7.08E-02	2.85E-01	1.70E+01	7.46E-01	1.00E-01	7.30E-02	3.75E-01	2.23E+01	2.77E+00	1.00E-01
	Zooplankton	Patterson Lake South Arm	6.71E-02	2.19E-01	1.27E+01	4.47E-01	3.50E-02	6.73E-02	2.74E-01	1.58E+01	1.62E+00	3.51E-02
		Beet Lake	6.69E-02	1.71E-01	9.78E+00	2.49E-01	1.93E-02	6.69E-02	2.00E-01	1.14E+01	8.60E-01	1.93E-02
Aquatic Animals		Lloyd Lake	6.68E-02	1.22E-01	6.90E+00	5.54E-02	1.09E-02	6.68E-02	1.25E-01	7.06E+00	1.18E-01	1.09E-02
Aquatic Animais		Reference (Broach Lake)	3.71E-02	7.00E-03	4.21E-01	2.06E-05	2.01E-03	3.71E-02	7.00E-03	4.21E-01	2.06E-05	2.01E-03
		Patterson Lake North Arm - West Basin	3.94E-02	1.71E-02	1.09E+00	4.60E-04	2.01E-02	4.06E-02	2.25E-02	1.43E+00	1.71E-03	2.01E-02
	Northern Pike	Patterson Lake South Arm	3.73E-02	1.31E-02	8.14E-01	2.76E-04	7.01E-03	3.74E-02	1.64E-02	1.01E+00	1.00E-03	7.01E-03
		Beet Lake	3.71E-02	1.02E-02	6.27E-01	1.54E-04	3.87E-03	3.72E-02	1.20E-02	7.30E-01	5.31E-04	3.87E-03
		Lloyd Lake	3.71E-02	7.33E-03	4.42E-01	3.42E-05	2.17E-03	3.71E-02	7.51E-03	4.53E-01	7.27E-05	2.17E-03
		Reference (Broach Lake)	3.56E-02	6.99E-03	4.21E-02	2.06E-05	4.97E-04	3.56E-02	6.99E-03	4.21E-02	2.06E-05	4.97E-04
		Patterson Lake North Arm - West Basin	3.78E-02	1.71E-02	1.09E-01	4.60E-04	4.97E-03	3.89E-02	2.25E-02	1.42E-01	1.71E-03	4.97E-03
	Whitefish	Patterson Lake South Arm	3.58E-02	1.31E-02	8.12E-02	2.75E-04	1.73E-03	3.59E-02	1.64E-02	1.01E-01	9.99E-04	1.74E-03
		Beet Lake	3.56E-02	1.02E-02	6.25E-02	1.53E-04	9.57E-04	3.57E-02	1.20E-02	7.28E-02	5.30E-04	9.57E-04
		Lloyd Lake	3.56E-02	7.32E-03	4.42E-02	3.41E-05	5.38E-04	3.56E-02	7.50E-03	4.52E-02	7.25E-05	5.38E-04
		Reference (Broach Lake)	2.51E-02	2.51E-02	3.99E-01	1.42E-02	4.01E-02	2.51E-02	2.51E-02	3.99E-01	1.42E-02	4.01E-02
		Patterson Lake North Arm - West Basin	2.51E-02	2.51E-02	4.00E-01	1.42E-02	5.79E-02	2.51E-02	2.51E-02	4.00E-01	1.42E-02	5.79E-02
	Blueberries	Patterson Lake South Arm	2.51E-02	2.51E-02	3.99E-01	1.42E-02	4.26E-02	2.51E-02	2.51E-02	3.99E-01	1.42E-02	4.26E-02
		Beet Lake	2.51E-02	2.51E-02	3.99E-01	1.42E-02	4.16E-02	2.51E-02	2.51E-02	3.99E-01	1.42E-02	4.16E-02
Terrestrial Plants		Lloyd Lake	2.51E-02	2.51E-02	3.99E-01	1.42E-02	4.01E-02	2.51E-02	2.51E-02	3.99E-01	1.42E-02	4.01E-02
refrestrial Plants		Reference (Broach Lake)	5.02E-02	5.01E-02	7.98E-01	2.85E-02	8.01E-02	5.02E-02	5.01E-02	7.98E-01	2.85E-02	8.01E-02
		Patterson Lake North Arm - West Basin	5.02E-02	5.03E-02	7.99E-01	2.85E-02	1.16E-01	5.02E-02	5.03E-02	7.99E-01	2.85E-02	1.16E-01
	Browse	Patterson Lake South Arm	5.02E-02	5.02E-02	7.98E-01	2.85E-02	8.52E-02	5.02E-02	5.02E-02	7.98E-01	2.85E-02	8.52E-02
		Beet Lake	5.02E-02	5.02E-02	7.98E-01	2.85E-02	8.31E-02	5.02E-02	5.02E-02	7.98E-01	2.85E-02	8.31E-02
		Lloyd Lake	5.02E-02	5.02E-02	7.98E-01	2.85E-02	8.02E-02	5.02E-02	5.02E-02	7.98E-01	2.85E-02	8.02E-02

Table C.8: Maximum Concentrations of non-radiological COPCs in Biota for the Application Case and Upper Bound Scenario – Far-Future

						Maximum Co	oncentration	ո Far-Future	(mg/kg fw	r)		
	Biota	Location			Application	Case			Up	per Bound	Scenario	
			Arsenic	Cobalt	Copper	Molybdenum	Uranium	Arsenic	Cobalt	Copper	Molybdenum	Uraniun
		Reference (Broach Lake)	5.02E-02	5.01E-02	7.98E-01	2.85E-02	8.01E-02	5.02E-02	5.01E-02	7.98E-01	2.85E-02	8.01E-0
		Patterson Lake North Arm - West Basin	5.02E-02	5.03E-02	7.99E-01	2.85E-02	1.16E-01	5.02E-02	5.03E-02	7.99E-01	2.85E-02	1.16E-0
	Labrador Tea	Patterson Lake South Arm	5.02E-02	5.02E-02	7.98E-01	2.85E-02	8.52E-02	5.02E-02	5.02E-02	7.98E-01	2.85E-02	8.52E-0
		Beet Lake	5.02E-02	5.02E-02	7.98E-01	2.85E-02	8.31E-02	5.02E-02	5.02E-02	7.98E-01	2.85E-02	8.31E-0
		Lloyd Lake	5.02E-02	5.02E-02	7.98E-01	2.85E-02	8.02E-02	5.02E-02	5.02E-02	7.98E-01	2.85E-02	8.02E-0
		Reference (Broach Lake)	5.58E-02	7.33E-02	7.58E-01	6.70E-02	1.68E-02	5.58E-02	7.33E-02	7.58E-01	6.70E-02	1.68E-0
		Patterson Lake North Arm - West Basin	5.58E-02	7.33E-02	7.58E-01	6.70E-02	1.68E-02	5.58E-02	7.33E-02	7.58E-01	6.70E-02	1.68E-0
	Lichen	Patterson Lake South Arm	5.58E-02	7.33E-02	7.58E-01	6.70E-02	1.68E-02	5.58E-02	7.33E-02	7.58E-01	6.70E-02	1.68E-0
		Beet Lake	5.58E-02	7.33E-02	7.58E-01	6.70E-02	1.68E-02	5.58E-02	7.33E-02	7.58E-01	6.70E-02	1.68E-0
		Lloyd Lake	5.58E-02	7.33E-02	7.58E-01	6.70E-02	1.68E-02	5.58E-02	7.33E-02	7.58E-01	6.70E-02	1.68E-0
		Reference (Broach Lake)	2.24E-02	2.63E-03	5.26E-02	3.73E-02	2.19E-03	2.24E-02	2.63E-03	5.26E-02	3.73E-02	2.19E-0
		Patterson Lake North Arm - West Basin	2.24E-02	2.63E-03	5.27E-02	3.73E-02	3.16E-03	2.24E-02	2.63E-03	5.27E-02	3.73E-02	3.16E-0
	Terrestrial Invertebrate	Patterson Lake South Arm	2.24E-02	2.63E-03	5.26E-02	3.73E-02	2.33E-03	2.24E-02	2.63E-03	5.26E-02	3.73E-02	2.33E-
	invertebrate	Beet Lake	2.24E-02	2.63E-03	5.26E-02	3.73E-02	2.27E-03	2.24E-02	2.63E-03	5.26E-02	3.73E-02	2.27E-
		Lloyd Lake	2.24E-02	2.63E-03	5.26E-02	3.73E-02	2.19E-03	2.24E-02	2.63E-03	5.26E-02	3.73E-02	2.19E-
		Reference (Broach Lake)	5.56E-02	1.25E-03	3.78E-01	1.36E-03	1.48E-03	5.56E-02	1.25E-03	3.78E-01	1.36E-03	1.48E-
		Patterson Lake North Arm - West Basin	5.60E-02	1.57E-03	3.94E-01	1.96E-03	2.39E-03	5.63E-02	1.74E-03	4.02E-01	3.64E-03	2.39E-
	Beaver	Patterson Lake South Arm	5.56E-02	1.44E-03	3.87E-01	1.71E-03	1.65E-03	5.57E-02	1.55E-03	3.92E-01	2.69E-03	1.65E-
		Beet Lake	5.56E-02	1.35E-03	3.83E-01	1.54E-03	1.56E-03	5.56E-02	1.41E-03	3.85E-01	2.05E-03	1.56E-
		Lloyd Lake	5.56E-02	1.26E-03	3.78E-01	1.38E-03	1.49E-03	5.56E-02	1.26E-03	3.79E-01	1.43E-03	1.49E-
		Reference (Broach Lake)	5.21E-02	8.47E-04	3.52E-01	1.01E-03	1.10E-03	5.21E-02	8.47E-04	3.52E-01	1.01E-03	1.10E-
	Dia de Dage	Patterson Lake North Arm - West Basin	5.26E-02	9.25E-04	4.49E-01	1.10E-03	1.43E-03	5.28E-02	9.63E-04	4.98E-01	1.34E-03	1.43E-
	Black Bear	Beet Lake	5.43E-02	9.00E-04	3.72E-01	1.05E-03	1.15E-03	5.43E-02	9.08E-04	3.82E-01	1.10E-03	1.15E-
		Lloyd Lake	5.22E-02	8.57E-04	3.56E-01	1.02E-03	1.10E-03	5.22E-02	8.58E-04	3.58E-01	1.03E-03	1.10E-0
Terrestrial		Reference (Broach Lake)	4.71E-02	8.12E-05	1.04E-01	9.04E-05	4.57E-05	4.71E-02	8.12E-05	1.04E-01	9.04E-05	4.57E-0
Animals	Grey Wolf	Patterson Lake North-West Basin	4.73E-02	8.70E-05	1.08E-01	1.40E-04	5.54E-05	4.74E-02	9.01E-05	1.10E-01	2.83E-04	5.54E-
		Lloyd Lake	4.71E-02	8.18E-05	1.04E-01	9.51E-05	4.59E-05	4.71E-02	8.21E-05	1.04E-01	1.09E-04	4.59E-
		Reference (Broach Lake)	2.80E-02	2.42E-04	2.78E-01	1.40E-04	5.15E-05	2.80E-02	2.42E-04	2.78E-01	1.40E-04	5.15E-
		Patterson Lake North Arm - West Basin	2.96E-02	5.54E-04	7.19E-01	2.69E-03	3.95E-04	3.05E-02	7.20E-04	9.40E-01	9.92E-03	3.96E-
	Mink	Patterson Lake South Arm	2.82E-02	4.31E-04	5.37E-01	1.62E-03	1.46E-04	2.82E-02	5.32E-04	6.67E-01	5.82E-03	1.46E-
		Beet Lake	2.81E-02	3.42E-04	4.14E-01	9.09E-04	8.68E-05	2.81E-02	3.95E-04	4.82E-01	3.10E-03	8.68E-
		Lloyd Lake	2.80E-02	2.52E-04	2.92E-01	2.18E-04	5.47E-05	2.80E-02	2.58E-04	2.99E-01	4.41E-04	5.47E-
		Reference (Broach Lake)	5.87E-02	1.37E-03	3.51E-01	1.26E-03	1.37E-03	5.87E-02	1.37E-03	3.51E-01	1.26E-03	1.37E-
		Patterson Lake North Arm - West Basin	5.96E-02	2.03E-03	3.84E-01	2.45E-03	2.49E-03	6.01E-02	2.37E-03	4.00E-01	5.83E-03	2.49E-
	Moose	Patterson Lake South Arm	5.96E-02	2.01E-03	3.83E-01	2.31E-03	2.47E-03	6.01E-02	2.35E-03	4.00E-01	5.33E-03	2.47E-
		Beet Lake	5.87E-02	1.58E-03	3.61E-01	1.62E-03	1.47E-03	5.87E-02	1.69E-03	3.66E-01	2.64E-03	1.47E-
		Lloyd Lake	5.87E-02	1.39E-03	3.52E-01	1.29E-03	1.38E-03	5.87E-02	1.41E-03	3.53E-01	1.40E-03	1.38E-
		Reference (Broach Lake)	4.99E-01	3.19E-01	1.40E+01	1.26E-01	2.42E-03	4.99E-01	3.19E-01	1.40E+01	1.26E-01	2.42E-0
	Moose Organs	Patterson Lake North Arm - West Basin	5.07E-01	4.71E-01	1.54E+01	2.45E-01	4.40E-03	5.11E-01	5.52E-01	1.60E+01	5.83E-01	4.40E-0
		Patterson Lake South Arm	5.06E-01	4.67E-01	1.53E+01	2.31E-01	4.37E-03	5.11E-01	5.46E-01	1.60E+01	5.33E-01	4.37E-0

Table C.8: Maximum Concentrations of non-radiological COPCs in Biota for the Application Case and Upper Bound Scenario – Far-Future

					Maximum C	oncen <u>tratior</u>	Far-Future	e (mg/kg fw	r)		
Biota	Location			Application					per Bound	Scenario	
		Arsenic	Cobalt	Copper	Molybdenum	Uranium	Arsenic	Cobalt	Copper	Molybdenum	Uranium
	Beet Lake	4.99E-01	3.68E-01	1.44E+01	1.62E-01	2.61E-03	4.99E-01	3.94E-01	1.46E+01	2.64E-01	2.61E-03
	Lloyd Lake	4.99E-01	3.24E-01	1.41E+01	1.29E-01	2.44E-03	4.99E-01	3.27E-01	1.41E+01	1.40E-01	2.44E-03
	Reference (Broach Lake)	7.19E-02	8.64E-04	5.12E-01	4.59E-04	2.70E-04	7.19E-02	8.64E-04	5.12E-01	4.59E-04	2.70E-04
	Patterson Lake North Arm - West Basin	7.62E-02	2.12E-03	1.33E+00	1.02E-02	2.70E-03	7.85E-02	2.78E-03	1.73E+00	3.80E-02	2.70E-03
Muskrat	Patterson Lake South Arm	7.22E-02	1.62E-03	9.90E-01	6.13E-03	9.43E-04	7.24E-02	2.03E-03	1.23E+00	2.22E-02	9.44E-04
	Beet Lake	7.19E-02	1.27E-03	7.62E-01	3.41E-03	5.20E-04	7.20E-02	1.48E-03	8.88E-01	1.18E-02	5.20E-04
	Lloyd Lake	7.19E-02	9.05E-04	5.38E-01	7.60E-04	2.93E-04	7.19E-02	9.27E-04	5.51E-01	1.62E-03	2.93E-04
	Reference (Broach Lake)	7.78E-03	7.79E-05	2.78E-02	2.35E-04	8.18E-05	7.78E-03	7.79E-05	2.78E-02	2.35E-04	8.18E-05
	Patterson Lake North Arm - West Basin	7.79E-03	8.36E-05	2.80E-02	3.01E-04	1.23E-04	7.79E-03	8.66E-05	2.81E-02	4.91E-04	1.23E-04
Red Fox	Patterson Lake South Arm	7.78E-03	8.13E-05	2.79E-02	2.73E-04	8.84E-05	7.78E-03	8.31E-05	2.80E-02	3.83E-04	8.84E-05
	Beet Lake	7.78E-03	7.98E-05	2.78E-02	2.55E-04	8.54E-05	7.78E-03	8.07E-05	2.79E-02	3.12E-04	8.54E-05
	Lloyd Lake	7.78E-03	7.82E-05	2.78E-02	2.37E-04	8.20E-05	7.78E-03	7.82E-05	2.78E-02	2.42E-04	8.20E-05
	Reference (Broach Lake)	4.69E-02	9.87E-04	3.44E-01	1.27E-03	1.37E-03	4.69E-02	9.87E-04	3.44E-01	1.27E-03	1.37E-03
	Patterson Lake North Arm - West Basin	4.69E-02	9.95E-04	3.45E-01	1.33E-03	1.99E-03	4.69E-02	9.97E-04	3.45E-01	1.49E-03	1.99E-03
Snowshoe Hare	Patterson Lake South Arm	4.69E-02	9.91E-04	3.44E-01	1.30E-03	1.46E-03	4.69E-02	9.93E-04	3.45E-01	1.40E-03	1.46E-03
	Beet Lake	4.69E-02	9.90E-04	3.44E-01	1.29E-03	1.42E-03	4.69E-02	9.91E-04	3.44E-01	1.34E-03	1.42E-03
	Lloyd Lake	4.69E-02	9.89E-04	3.44E-01	1.27E-03	1.37E-03	4.69E-02	9.89E-04	3.44E-01	1.28E-03	1.37E-03
	Reference (Broach Lake)	1.67E-02	3.54E-04	1.26E-01	4.59E-04	4.98E-04	1.67E-02	3.54E-04	1.26E-01	4.59E-04	4.98E-04
	Patterson Lake North Arm - West Basin	1.67E-02	3.58E-04	1.26E-01	4.91E-04	7.22E-04	1.67E-02	3.59E-04	1.26E-01	5.81E-04	7.22E-04
Southern Red-	Patterson Lake South Arm	1.67E-02	3.56E-04	1.26E-01	4.78E-04	5.31E-04	1.67E-02	3.57E-04	1.26E-01	5.30E-04	5.31E-04
Backed Vole	Beet Lake	1.67E-02	3.55E-04	1.26E-01	4.69E-04	5.17E-04	1.67E-02	3.56E-04	1.26E-01	4.96E-04	5.17E-04
	Lloyd Lake	1.67E-02	3.55E-04	1.26E-01	4.60E-04	4.99E-04	1.67E-02	3.55E-04	1.26E-01	4.63E-04	4.99E-04
	Reference (Broach Lake)	6.30E-02	1.08E-03	1.92E-01	1.03E-03	7.02E-04	6.30E-02	1.08E-03	1.92E-01	1.03E-03	7.02E-04
	Patterson Lake North Arm - West Basin	6.39E-02	1.59E-03	2.17E-01	2.94E-03	1.50E-03	6.44E-02	1.87E-03	2.29E-01	8.37E-03	1.50E-03
Woodland Caribou	Beet Lake	6.29E-02	1.18E-03	1.97E-01	1.40E-03	7.67E-04	6.29E-02	1.24E-03	1.99E-01	2.45E-03	7.67E-04
	Lloyd Lake	6.30E-02	1.10E-03	1.93E-01	1.11E-03	7.14E-04	6.30E-02	1.11E-03	1.94E-01	1.34E-03	7.14E-04
	Reference (Broach Lake)	3.71E-02	2.88E-02	2.59E-01	2.87E-03	3.27E-02	3.71E-02	2.88E-02	2.59E-01	2.87E-03	3.27E-02
	Patterson Lake North Arm - West Basin	3.71E-02	2.89E-02	2.60E-01	2.96E-03	4.73E-02	3.71E-02	2.90E-02	2.60E-01	3.19E-03	4.73E-02
Grouse	Patterson Lake South Arm	3.71E-02	2.88E-02	2.60E-01	2.92E-03	3.48E-02	3.71E-02	2.89E-02	2.60E-01	3.06E-03	3.48E-02
	Beet Lake	3.71E-02	2.88E-02	2.60E-01	2.90E-03	3.39E-02	3.71E-02	2.88E-02	2.60E-01	2.97E-03	3.39E-02
	Lloyd Lake	3.71E-02	2.88E-02	2.59E-01	2.88E-03	3.27E-02	3.71E-02	2.88E-02	2.59E-01	2.88E-03	3.27E-02
	Reference (Broach Lake)	5.23E-03	4.08E-03	3.63E-02	4.06E-04	4.59E-03	5.23E-03	4.08E-03	3.63E-02	4.06E-04	4.59E-03
	Patterson Lake North Arm - West Basin	5.23E-03	4.16E-03	3.65E-02	4.76E-04	6.68E-03	5.23E-03	4.20E-03	3.65E-02	6.78E-04	6.68E-03
Canada Goose	Patterson Lake South Arm	5.23E-03	4.13E-03	3.64E-02	4.47E-04	4.89E-03	5.23E-03	4.15E-03	3.64E-02	5.63E-04	4.89E-03
	Beet Lake	5.23E-03	4.10E-03	3.64E-02	4.27E-04	4.76E-03	5.23E-03	4.12E-03	3.64E-02	4.88E-04	4.76E-03
	Lloyd Lake	5.23E-03	4.08E-03	3.63E-02	4.08E-04	4.59E-03	5.23E-03	4.09E-03	3.63E-02	4.14E-04	4.59E-03
	Reference (Broach Lake)	1.97E-02	8.78E-04	1.15E+00	5.85E-04	6.64E-05	1.97E-02	8.78E-04	1.15E+00	5.85E-04	6.64E-05
Little Brown	Patterson Lake North Arm - West Basin	2.09E-02	2.15E-03	2.97E+00	1.31E-02	6.64E-04	2.16E-02	2.83E-03	3.89E+00	4.85E-02	6.65E-04
Myotis	Patterson Lake South Arm	1.98E-02	1.65E-03	2.22E+00	7.82E-03	2.32E-04	1.99E-02	2.06E-03	2.76E+00	2.84E-02	2.32E-04
	Beet Lake	1.97E-02	1.29E-03	1.71E+00	4.35E-03	1.28E-04	1.98E-02	1.50E-03	1.99E+00	1.51E-02	1.28E-04

Table C.8: Maximum Concentrations of non-radiological COPCs in Biota for the Application Case and Upper Bound Scenario – Far-Future

					Maximum Co	oncentration	n Far-Future	(mg/kg fw	·)		
Biota	Location			Application	Case			Up	per Bound	Scenario	
		Arsenic	Cobalt	Copper	Molybdenum	Uranium	Arsenic	Cobalt	Copper	Molybdenum	Uranium
	Lloyd Lake	1.97E-02	9.20E-04	1.21E+00	9.69E-04	7.20E-05	1.97E-02	9.42E-04	1.23E+00	2.06E-03	7.20E-05
	Reference (Broach Lake)	1.44E-02	5.39E-03	1.97E-01	1.88E-04	6.47E-04	1.44E-02	5.39E-03	1.97E-01	1.88E-04	6.47E-04
	Patterson Lake North Arm - West Basin	1.53E-02	1.32E-02	5.10E-01	4.20E-03	6.47E-03	1.58E-02	1.74E-02	6.68E-01	1.56E-02	6.48E-03
Loon	Patterson Lake South Arm	1.45E-02	1.01E-02	3.81E-01	2.52E-03	2.26E-03	1.45E-02	1.27E-02	4.73E-01	9.13E-03	2.26E-03
	Beet Lake	1.44E-02	7.90E-03	2.93E-01	1.40E-03	1.25E-03	1.45E-02	9.24E-03	3.42E-01	4.84E-03	1.25E-03
	Lloyd Lake	1.44E-02	5.65E-03	2.07E-01	3.12E-04	7.01E-04	1.44E-02	5.79E-03	2.12E-01	6.63E-04	7.01E-04
	Reference (Broach Lake)	5.77E-02	4.36E-02	1.13E+00	2.04E-03	6.80E-03	5.77E-02	4.36E-02	1.13E+00	2.04E-03	6.80E-03
	Patterson Lake North Arm - West Basin	6.12E-02	1.07E-01	2.93E+00	4.54E-02	6.80E-02	6.31E-02	1.40E-01	3.84E+00	1.69E-01	6.81E-02
Mallard	Patterson Lake South Arm	5.80E-02	8.20E-02	2.19E+00	2.72E-02	2.37E-02	5.82E-02	1.02E-01	2.72E+00	9.87E-02	2.37E-02
	Beet Lake	5.78E-02	6.39E-02	1.68E+00	1.51E-02	1.31E-02	5.78E-02	7.47E-02	1.96E+00	5.24E-02	1.31E-02
	Lloyd Lake	5.77E-02	4.57E-02	1.19E+00	3.37E-03	7.37E-03	5.77E-02	4.68E-02	1.22E+00	7.17E-03	7.37E-03
	Reference (Broach Lake)	7.99E-02	1.09E-01	3.39E+00	6.19E-03	1.83E-02	7.99E-02	1.09E-01	3.39E+00	6.19E-03	1.83E-02
	Patterson Lake North Arm - West Basin	8.32E-02	2.42E-01	8.65E+00	1.11E-01	7.52E-02	8.50E-02	3.13E-01	1.13E+01	4.07E-01	7.52E-02
Rusty Blackbird	Patterson Lake South Arm	8.01E-02	1.90E-01	6.48E+00	6.68E-02	3.35E-02	8.03E-02	2.33E-01	8.03E+00	2.39E-01	3.35E-02
	Beet Lake	7.99E-02	1.51E-01	5.01E+00	3.77E-02	2.40E-02	7.99E-02	1.74E-01	5.82E+00	1.27E-01	2.40E-02
	Lloyd Lake	7.99E-02	1.13E-01	3.56E+00	9.40E-03	1.88E-02	7.99E-02	1.15E-01	3.64E+00	1.85E-02	1.88E-02

Table C.9: Maximum Concentrations of non-radiological COPCs in Biota for the RFD Scenario - Project Lifespan and Far-Future

						Maximum (Concentratio	on RFD Case	(mg/kg fw)			
	Biota	Location			Project Life:				(9,9)	Far-Futu	re	
	2.5.11		Arsenic	Cobalt	Copper	Molybdenum	Uranium	Arsenic	Cobalt	Copper	Molybdenum	Uranium
		Reference (Broach Lake)	2.13E-02	9.21E-02	1.83E-01	1.40E-03	5.06E-03	2.13E-02	9.21E-02	1.83E-01	1.40E-03	5.06E-03
		Patterson Lake North Arm - West Basin	7.50E-02	1.22E-01	2.16E-01	4.07E-03	1.93E-02	2.26E-02	2.26E-01	4.73E-01	3.13E-02	5.06E-02
	Macrophytes	Patterson Lake South Arm	1.19E-01	1.23E-01	2.19E-01	3.89E-03	2.70E-02	2.14E-02	1.73E-01	3.53E-01	1.88E-02	1.77E-02
		Beet Lake	3.80E-02	1.06E-01	1.98E-01	2.51E-03	1.27E-02	2.13E-02	1.35E-01	2.72E-01	1.04E-02	9.74E-03
		Lloyd Lake	2.23E-02	9.35E-02	1.84E-01	1.51E-03	5.77E-03	2.13E-02	9.65E-02	1.92E-01	2.32E-03	5.48E-03
Aquatic Plants		Reference (Broach Lake)	2.13E-02	9.21E-02	1.83E-01	1.40E-03	5.06E-03	2.13E-02	9.21E-02	1.83E-01	1.40E-03	5.06E-03
		Patterson Lake North Arm - West Basin	7.50E-02	1.22E-01	2.16E-01	4.07E-03	1.93E-02	2.26E-02	2.26E-01	4.73E-01	3.13E-02	5.06E-02
	Phytoplankton	Patterson Lake South Arm	1.19E-01	1.23E-01	2.19E-01	3.89E-03	2.70E-02	2.14E-02	1.73E-01	3.53E-01	1.88E-02	1.77E-02
		Beet Lake	3.80E-02	1.06E-01	1.98E-01	2.51E-03	1.27E-02	2.13E-02	1.35E-01	2.72E-01	1.04E-02	9.74E-03
		Lloyd Lake	2.23E-02	9.35E-02	1.84E-01	1.51E-03	5.77E-03	2.13E-02	9.65E-02	1.92E-01	2.32E-03	5.48E-03
		Reference (Broach Lake)	5.60E-02	1.16E-01	6.53E+00	3.32E-02	9.64E-03	5.60E-02	1.16E-01	6.53E+00	3.32E-02	9.64E-03
	5 di	Patterson Lake North Arm - West Basin	1.71E-01	1.50E-01	7.59E+00	8.87E-02	3.16E-02	5.94E-02	2.84E-01	1.69E+01	7.40E-01	9.65E-02
	Benthic Invertebrates	Patterson Lake South Arm	1.28E-01	1.41E-01	7.32E+00	7.07E-02	2.23E-02	5.63E-02	2.18E-01	1.26E+01	4.44E-01	3.37E-02
	invertebrates	Beet Lake	6.84E-02	1.29E-01	6.94E+00	5.27E-02	1.43E-02	5.61E-02	1.70E-01	9.71E+00	2.47E-01	1.86E-02
		Lloyd Lake	5.68E-02	1.17E-01	6.57E+00	3.52E-02	1.01E-02	5.60E-02	1.21E-01	6.85E+00	5.50E-02	1.05E-02
		Reference (Broach Lake)	6.68E-02	1.17E-01	6.57E+00	3.34E-02	1.00E-02	6.68E-02	1.17E-01	6.57E+00	3.34E-02	1.00E-02
		Patterson Lake North Arm - West Basin	2.35E-01	1.54E-01	7.76E+00	9.69E-02	3.84E-02	7.08E-02	2.85E-01	1.70E+01	7.46E-01	1.00E-01
	Zooplankton	Patterson Lake South Arm	3.74E-01	1.56E-01	7.87E+00	9.26E-02	5.37E-02	6.71E-02	2.19E-01	1.27E+01	4.47E-01	3.50E-02
		Beet Lake	1.19E-01	1.34E-01	7.13E+00	5.98E-02	2.52E-02	6.69E-02	1.71E-01	9.78E+00	2.49E-01	1.93E-02
Aquatic Animals		Lloyd Lake	7.00E-02	1.18E-01	6.63E+00	3.61E-02	1.14E-02	6.68E-02	1.22E-01	6.90E+00	5.54E-02	1.09E-02
Aquatic Allillais		Reference (Broach Lake)	3.71E-02	7.00E-03	4.21E-01	2.06E-05	2.01E-03	3.71E-02	7.00E-03	4.21E-01	2.06E-05	2.01E-03
		Patterson Lake North Arm - West Basin	1.31E-01	9.25E-03	4.97E-01	5.98E-05	7.67E-03	3.94E-02	1.71E-02	1.09E+00	4.60E-04	2.01E-02
	Northern Pike	Patterson Lake South Arm	2.08E-01	9.38E-03	5.04E-01	5.72E-05	1.07E-02	3.73E-02	1.31E-02	8.14E-01	2.76E-04	7.01E-03
		Beet Lake	6.62E-02	8.05E-03	4.57E-01	3.69E-05	5.04E-03	3.71E-02	1.02E-02	6.27E-01	1.54E-04	3.87E-03
		Lloyd Lake	3.89E-02	7.10E-03	4.25E-01	2.23E-05	2.29E-03	3.71E-02	7.33E-03	4.42E-01	3.42E-05	2.17E-03
		Reference (Broach Lake)	3.56E-02	6.99E-03	4.21E-02	2.06E-05	4.97E-04	3.56E-02	6.99E-03	4.21E-02	2.06E-05	4.97E-04
		Patterson Lake North Arm - West Basin	1.22E-01	9.18E-03	4.95E-02	5.83E-05	1.83E-03	3.78E-02	1.71E-02	1.09E-01	4.60E-04	4.97E-03
	Whitefish	Patterson Lake South Arm	1.73E-01	8.97E-03	4.89E-02	5.11E-05	2.24E-03	3.58E-02	1.31E-02	8.12E-02	2.75E-04	1.73E-03
		Beet Lake	5.89E-02	7.88E-03	4.51E-02	3.43E-05	1.10E-03	3.56E-02	1.02E-02	6.25E-02	1.53E-04	9.57E-04
		Lloyd Lake	3.70E-02	7.08E-03	4.24E-02	2.20E-05	5.53E-04	3.56E-02	7.32E-03	4.42E-02	3.41E-05	5.38E-04
		Reference (Broach Lake)	2.51E-02	2.51E-02	3.99E-01	1.42E-02	4.01E-02	2.51E-02	2.51E-02	3.99E-01	1.42E-02	4.01E-02
		Patterson Lake North Arm - West Basin	2.52E-02	2.52E-02	4.13E-01	1.58E-02	2.93E-01	2.51E-02	2.51E-02	4.00E-01	1.42E-02	5.81E-02
	Blueberries	Patterson Lake South Arm	2.52E-02	2.52E-02	4.00E-01	1.44E-02	8.05E-02	2.51E-02	2.51E-02	3.99E-01	1.42E-02	4.29E-02
		Beet Lake	2.52E-02	2.52E-02	4.00E-01	1.44E-02	6.26E-02	2.51E-02	2.51E-02	3.99E-01	1.42E-02	4.17E-02
Terrestrial Plants		Lloyd Lake	2.52E-02	2.52E-02	4.00E-01	1.44E-02	4.03E-02	2.51E-02	2.51E-02	3.99E-01	1.42E-02	4.01E-02
refrestrial Plants		Reference (Broach Lake)	5.02E-02	5.01E-02	7.98E-01	2.85E-02	8.01E-02	5.02E-02	5.01E-02	7.98E-01	2.85E-02	8.01E-02
		Patterson Lake North Arm - West Basin	5.05E-02	5.05E-02	8.27E-01	3.28E-02	6.79E-01	5.02E-02	5.02E-02	7.99E-01	2.85E-02	1.16E-01
	Browse	Patterson Lake South Arm	5.05E-02	5.05E-02	7.99E-01	2.88E-02	1.76E-01	5.02E-02	5.02E-02	7.98E-01	2.85E-02	8.59E-02
		Beet Lake	5.05E-02	5.05E-02	7.99E-01	2.88E-02	1.34E-01	5.02E-02	5.02E-02	7.98E-01	2.85E-02	8.33E-02
		Lloyd Lake	5.05E-02	5.05E-02	7.99E-01	2.88E-02	8.07E-02	5.02E-02	5.02E-02	7.98E-01	2.85E-02	8.02E-02

Table C.9: Maximum Concentrations of non-radiological COPCs in Biota for the RFD Scenario - Project Lifespan and Far-Future

						Ma <u>ximum</u>	Concentratio	n RFD Case	(mg/kg fw)			
	Biota	Location			Project Life	span				Far-Futu	re	
			Arsenic	Cobalt	Copper	Molybdenum	Uranium	Arsenic	Cobalt	Copper	Molybdenum	Uranium
		Reference (Broach Lake)	5.02E-02	5.01E-02	7.98E-01	2.85E-02	8.01E-02	5.02E-02	5.01E-02	7.98E-01	2.85E-02	8.01E-02
		Patterson Lake North Arm - West Basin	5.28E-02	5.28E-02	8.67E-01	6.14E-02	2.88E+00	5.02E-02	5.02E-02	7.99E-01	2.85E-02	1.16E-01
	Labrador Tea	Patterson Lake South Arm	5.28E-02	5.28E-02	8.02E-01	3.11E-02	5.27E-01	5.02E-02	5.02E-02	7.98E-01	2.85E-02	8.59E-02
		Beet Lake	5.28E-02	5.28E-02	8.02E-01	3.11E-02	3.30E-01	5.02E-02	5.02E-02	7.98E-01	2.85E-02	8.33E-02
		Lloyd Lake	5.28E-02	5.28E-02	8.02E-01	3.11E-02	8.31E-02	5.02E-02	5.02E-02	7.98E-01	2.85E-02	8.02E-02
		Reference (Broach Lake)	5.58E-02	7.33E-02	7.58E-01	6.70E-02	1.68E-02	5.58E-02	7.33E-02	7.58E-01	6.70E-02	1.68E-02
		Patterson Lake North Arm - West Basin	7.07E-02	8.82E-02	1.01E+00	2.50E-01	1.41E+01	5.58E-02	7.33E-02	7.58E-01	6.70E-02	1.68E-02
	Lichen	Patterson Lake South Arm	7.07E-02	8.82E-02	7.73E-01	8.19E-02	2.27E+00	5.58E-02	7.33E-02	7.58E-01	6.70E-02	1.68E-02
		Beet Lake	7.07E-02	8.82E-02	7.73E-01	8.19E-02	1.27E+00	5.58E-02	7.33E-02	7.58E-01	6.70E-02	1.68E-02
		Lloyd Lake	7.07E-02	8.82E-02	7.73E-01	8.19E-02	3.17E-02	5.58E-02	7.33E-02	7.58E-01	6.70E-02	1.68E-02
		Reference (Broach Lake)	2.24E-02	2.63E-03	5.26E-02	3.73E-02	2.19E-03	2.24E-02	2.63E-03	5.26E-02	3.73E-02	2.19E-03
		Patterson Lake North Arm - West Basin	2.27E-02	2.91E-03	5.89E-02	4.19E-02	2.70E-01	2.24E-02	2.63E-03	5.27E-02	3.73E-02	3.17E-03
	Terrestrial Invertebrate	Patterson Lake South Arm	2.27E-02	2.91E-03	5.29E-02	3.76E-02	4.49E-02	2.24E-02	2.63E-03	5.26E-02	3.73E-02	2.34E-03
	invertebrate	Beet Lake	2.27E-02	2.91E-03	5.29E-02	3.76E-02	2.61E-02	2.24E-02	2.63E-03	5.26E-02	3.73E-02	2.27E-03
		Lloyd Lake	2.27E-02	2.91E-03	5.29E-02	3.76E-02	2.47E-03	2.24E-02	2.63E-03	5.26E-02	3.73E-02	2.19E-03
		Reference (Broach Lake)	5.56E-02	1.25E-03	3.78E-01	1.36E-03	1.48E-03	5.56E-02	1.25E-03	3.78E-01	1.36E-03	1.48E-03
		Patterson Lake North Arm - West Basin	7.20E-02	1.33E-03	3.93E-01	1.61E-03	1.24E-02	5.60E-02	1.57E-03	3.94E-01	1.96E-03	2.40E-03
	Beaver	Patterson Lake South Arm	7.19E-02	1.32E-03	3.80E-01	1.41E-03	3.24E-03	5.56E-02	1.44E-03	3.87E-01	1.71E-03	1.66E-03
		Beet Lake	5.86E-02	1.28E-03	3.79E-01	1.39E-03	2.46E-03	5.56E-02	1.35E-03	3.83E-01	1.54E-03	1.57E-03
		Lloyd Lake	5.60E-02	1.26E-03	3.79E-01	1.38E-03	1.50E-03	5.56E-02	1.26E-03	3.78E-01	1.38E-03	1.49E-03
		Reference (Broach Lake)	5.21E-02	8.47E-04	3.52E-01	1.01E-03	1.10E-03	5.21E-02	8.47E-04	3.52E-01	1.01E-03	1.10E-03
	Black Bear	Patterson Lake North Arm - West Basin	1.03E-01	8.74E-04	3.68E-01	1.07E-03	4.65E-03	5.26E-02	9.25E-04	4.49E-01	1.10E-03	1.44E-03
	віаск веаг	Beet Lake	5.97E-02	8.91E-04	3.56E-01	1.04E-03	1.42E-03	5.43E-02	9.00E-04	3.72E-01	1.05E-03	1.15E-03
		Lloyd Lake	5.30E-02	8.58E-04	3.53E-01	1.02E-03	1.11E-03	5.22E-02	8.57E-04	3.56E-01	1.02E-03	1.10E-03
Terrestrial		Reference (Broach Lake)	4.71E-02	8.12E-05	1.04E-01	9.04E-05	4.57E-05	4.71E-02	8.12E-05	1.04E-01	9.04E-05	4.57E-05
Animals	Grey Wolf	Patterson Lake North-West Basin	5.66E-02	8.27E-05	1.06E-01	9.97E-05	1.49E-04	4.73E-02	8.70E-05	1.08E-01	1.40E-04	5.55E-05
		Lloyd Lake	4.81E-02	8.19E-05	1.04E-01	9.28E-05	4.62E-05	4.71E-02	8.18E-05	1.04E-01	9.51E-05	4.59E-05
		Reference (Broach Lake)	2.80E-02	2.42E-04	2.78E-01	1.40E-04	5.15E-05	2.80E-02	2.42E-04	2.78E-01	1.40E-04	5.15E-05
		Patterson Lake North Arm - West Basin	8.43E-02	3.06E-04	3.24E-01	3.40E-04	1.98E-04	2.96E-02	5.54E-04	7.19E-01	2.69E-03	3.95E-04
	Mink	Patterson Lake South Arm	7.57E-02	2.89E-04	3.12E-01	2.75E-04	1.07E-04	2.82E-02	4.31E-04	5.37E-01	1.62E-03	1.46E-04
		Beet Lake	3.60E-02	2.67E-04	2.96E-01	2.10E-04	7.32E-05	2.81E-02	3.42E-04	4.14E-01	9.09E-04	8.68E-05
		Lloyd Lake	2.85E-02	2.44E-04	2.80E-01	1.47E-04	5.33E-05	2.80E-02	2.52E-04	2.92E-01	2.18E-04	5.47E-05
		Reference (Broach Lake)	5.87E-02	1.37E-03	3.51E-01	1.26E-03	1.37E-03	5.87E-02	1.37E-03	3.51E-01	1.26E-03	1.37E-03
		Patterson Lake North Arm - West Basin	9.24E-02	1.52E-03	3.67E-01	1.53E-03	1.12E-02	5.96E-02	2.02E-03	3.84E-01	2.45E-03	2.49E-03
	Moose	Patterson Lake South Arm	9.21E-02	1.51E-03	3.66E-01	1.51E-03	1.12E-02	5.96E-02	2.01E-03	3.83E-01	2.31E-03	2.48E-03
		Beet Lake	6.45E-02	1.44E-03	3.53E-01	1.30E-03	2.26E-03	5.87E-02	1.58E-03	3.61E-01	1.62E-03	1.48E-03
		Lloyd Lake	5.93E-02	1.38E-03	3.52E-01	1.27E-03	1.38E-03	5.87E-02	1.39E-03	3.52E-01	1.29E-03	1.38E-03
		Reference (Broach Lake)	4.99E-01	3.19E-01	1.40E+01	1.26E-01	2.42E-03	4.99E-01	3.19E-01	1.40E+01	1.26E-01	2.42E-03
	Moose Organs	Patterson Lake North Arm - West Basin	7.85E-01	3.54E-01	1.47E+01	1.53E-01	1.99E-02	5.07E-01	4.71E-01	1.54E+01	2.45E-01	4.41E-03
	Wioose Organis	Patterson Lake South Arm	7.83E-01	3.52E-01	1.47E+01	1.51E-01	1.98E-02	5.06E-01	4.67E-01	1.53E+01	2.31E-01	4.38E-03

Table C.9: Maximum Concentrations of non-radiological COPCs in Biota for the RFD Scenario - Project Lifespan and Far-Future

					Maximum	Concentratio	on RFD Case	(mg/kg fw)			
Biota	Location			Project Life:				· J. J	Far-Futu	re	
		Arsenic	Cobalt	Copper	Molybdenum	Uranium	Arsenic	Cobalt	Copper	Molybdenum	Uranium
	Beet Lake	5.48E-01	3.35E-01	1.41E+01	1.30E-01	4.01E-03	4.99E-01	3.68E-01	1.44E+01	1.62E-01	2.61E-03
	Lloyd Lake	5.04E-01	3.22E-01	1.41E+01	1.27E-01	2.45E-03	4.99E-01	3.24E-01	1.41E+01	1.29E-01	2.43E-03
	Reference (Broach Lake)	7.19E-02	8.64E-04	5.12E-01	4.59E-04	2.70E-04	7.19E-02	8.64E-04	5.12E-01	4.59E-04	2.70E-04
	Patterson Lake North Arm - West Basin	2.20E-01	1.13E-03	5.96E-01	1.23E-03	8.89E-04	7.62E-02	2.12E-03	1.33E+00	1.02E-02	2.70E-03
Muskrat	Patterson Lake South Arm	1.68E-01	1.05E-03	5.74E-01	9.78E-04	6.26E-04	7.22E-02	1.62E-03	9.90E-01	6.13E-03	9.43E-04
	Beet Lake	8.81E-02	9.64E-04	5.45E-01	7.29E-04	4.01E-04	7.19E-02	1.27E-03	7.62E-01	3.41E-03	5.20E-04
	Lloyd Lake	7.28E-02	8.74E-04	5.16E-01	4.87E-04	2.82E-04	7.19E-02	9.05E-04	5.38E-01	7.60E-04	2.93E-04
	Reference (Broach Lake)	7.78E-03	7.79E-05	2.78E-02	2.35E-04	8.18E-05	7.78E-03	7.79E-05	2.78E-02	2.35E-04	8.18E-05
	Patterson Lake North Arm - West Basin	7.93E-03	8.00E-05	2.89E-02	2.67E-04	9.97E-04	7.79E-03	8.35E-05	2.80E-02	3.01E-04	1.23E-04
Red Fox	Patterson Lake South Arm	8.01E-03	7.99E-05	2.79E-02	2.42E-04	2.28E-04	7.78E-03	8.13E-05	2.79E-02	2.73E-04	8.91E-05
	Beet Lake	7.86E-03	7.92E-05	2.78E-02	2.39E-04	1.64E-04	7.78E-03	7.98E-05	2.78E-02	2.55E-04	8.56E-05
	Lloyd Lake	7.83E-03	7.88E-05	2.78E-02	2.37E-04	8.28E-05	7.78E-03	7.82E-05	2.78E-02	2.37E-04	8.19E-05
	Reference (Broach Lake)	4.69E-02	9.87E-04	3.44E-01	1.27E-03	1.37E-03	4.69E-02	9.87E-04	3.44E-01	1.27E-03	1.37E-03
	Patterson Lake North Arm - West Basin	4.72E-02	9.96E-04	3.57E-01	1.46E-03	1.14E-02	4.69E-02	9.93E-04	3.45E-01	1.33E-03	1.99E-03
Snowshoe Hare	Patterson Lake South Arm	4.72E-02	9.96E-04	3.45E-01	1.29E-03	2.97E-03	4.69E-02	9.91E-04	3.44E-01	1.30E-03	1.47E-03
	Beet Lake	4.71E-02	9.95E-04	3.45E-01	1.29E-03	2.27E-03	4.69E-02	9.90E-04	3.44E-01	1.29E-03	1.43E-03
	Lloyd Lake	4.71E-02	9.95E-04	3.45E-01	1.28E-03	1.38E-03	4.69E-02	9.89E-04	3.44E-01	1.27E-03	1.37E-03
	Reference (Broach Lake)	1.67E-02	3.54E-04	1.26E-01	4.59E-04	4.98E-04	1.67E-02	3.54E-04	1.26E-01	4.59E-04	4.98E-04
	Patterson Lake North Arm - West Basin	1.68E-02	3.57E-04	1.30E-01	5.21E-04	3.90E-03	1.67E-02	3.57E-04	1.26E-01	4.91E-04	7.25E-04
Southern Red- Backed Vole	Patterson Lake South Arm	1.68E-02	3.57E-04	1.26E-01	4.66E-04	1.04E-03	1.67E-02	3.56E-04	1.26E-01	4.78E-04	5.35E-04
Backed Voie	Beet Lake	1.67E-02	3.57E-04	1.26E-01	4.65E-04	8.02E-04	1.67E-02	3.55E-04	1.26E-01	4.69E-04	5.18E-04
	Lloyd Lake	1.67E-02	3.56E-04	1.26E-01	4.64E-04	5.02E-04	1.67E-02	3.55E-04	1.26E-01	4.60E-04	4.98E-04
	Reference (Broach Lake)	6.30E-02	1.08E-03	1.92E-01	1.03E-03	7.02E-04	6.30E-02	1.08E-03	1.92E-01	1.03E-03	7.02E-04
Woodland	Patterson Lake North Arm - West Basin	1.09E-01	1.24E-03	2.04E-01	1.76E-03	2.03E-02	6.39E-02	1.59E-03	2.17E-01	2.94E-03	1.50E-03
Caribou	Beet Lake	6.81E-02	1.13E-03	1.93E-01	1.11E-03	2.32E-03	6.29E-02	1.18E-03	1.97E-01	1.40E-03	7.67E-04
	Lloyd Lake	6.54E-02	1.13E-03	1.94E-01	1.14E-03	7.47E-04	6.30E-02	1.10E-03	1.93E-01	1.11E-03	7.14E-04
	Reference (Broach Lake)	3.71E-02	2.88E-02	2.59E-01	2.87E-03	3.27E-02	3.71E-02	2.88E-02	2.59E-01	2.87E-03	3.27E-02
	Patterson Lake North Arm - West Basin	3.73E-02	2.90E-02	2.69E-01	3.27E-03	2.63E-01	3.71E-02	2.89E-02	2.60E-01	2.96E-03	4.75E-02
Grouse	Patterson Lake South Arm	3.74E-02	2.90E-02	2.60E-01	2.91E-03	6.95E-02	3.71E-02	2.88E-02	2.60E-01	2.92E-03	3.51E-02
	Beet Lake	3.73E-02	2.90E-02	2.60E-01	2.91E-03	5.33E-02	3.71E-02	2.88E-02	2.59E-01	2.90E-03	3.40E-02
	Lloyd Lake	3.73E-02	2.90E-02	2.60E-01	2.91E-03	3.29E-02	3.71E-02	2.88E-02	2.59E-01	2.88E-03	3.27E-02
	Reference (Broach Lake)	5.23E-03	4.08E-03	3.63E-02	4.06E-04	4.59E-03	5.23E-03	4.08E-03	3.63E-02	4.06E-04	4.59E-03
	Patterson Lake North Arm - West Basin	5.28E-03	4.12E-03	3.77E-02	4.70E-04	3.81E-02	5.23E-03	4.15E-03	3.65E-02	4.76E-04	6.70E-03
Canada Goose	Patterson Lake South Arm	5.31E-03	4.12E-03	3.64E-02	4.15E-04	9.95E-03	5.23E-03	4.13E-03	3.64E-02	4.47E-04	4.93E-03
	Beet Lake	5.26E-03	4.11E-03	3.64E-02	4.12E-04	7.58E-03	5.23E-03	4.10E-03	3.64E-02	4.27E-04	4.77E-03
	Lloyd Lake	5.25E-03	4.11E-03	3.64E-02	4.10E-04	4.62E-03	5.23E-03	4.08E-03	3.63E-02	4.08E-04	4.59E-03
	Reference (Broach Lake)	1.97E-02	8.78E-04	1.15E+00	5.85E-04	6.64E-05	1.97E-02	8.78E-04	1.15E+00	5.85E-04	6.64E-05
Little Brown	Patterson Lake North Arm - West Basin	6.02E-02	1.14E-03	1.34E+00	1.56E-03	2.18E-04	2.09E-02	2.15E-03	2.97E+00	1.31E-02	6.64E-04
Myotis	Patterson Lake South Arm	4.52E-02	1.07E-03	1.29E+00	1.25E-03	1.54E-04	1.98E-02	1.65E-03	2.22E+00	7.82E-03	2.32E-04
	Beet Lake	2.41E-02	9.79E-04	1.22E+00	9.30E-04	9.87E-05	1.97E-02	1.29E-03	1.71E+00	4.35E-03	1.28E-04

Table C.9: Maximum Concentrations of non-radiological COPCs in Biota for the RFD Scenario - Project Lifespan and Far-Future

					Maximum (Concentratio	n RFD Case	(mg/kg fw)			
Biota	Location			Project Lifes	pan				Far-Futur	œ .	
		Arsenic	Cobalt	Copper	Molybdenum	Uranium	Arsenic	Cobalt	Copper	Molybdenum	Uranium
	Lloyd Lake	2.00E-02	8.89E-04	1.16E+00	6.21E-04	6.94E-05	1.97E-02	9.20E-04	1.21E+00	9.69E-04	7.20E-05
	Reference (Broach Lake)	1.44E-02	5.39E-03	1.97E-01	1.88E-04	6.47E-04	1.44E-02	5.39E-03	1.97E-01	1.88E-04	6.47E-04
	Patterson Lake North Arm - West Basin	4.98E-02	7.03E-03	2.30E-01	5.03E-04	2.37E-03	1.53E-02	1.32E-02	5.10E-01	4.20E-03	6.47E-03
Loon	Patterson Lake South Arm	7.37E-02	6.58E-03	2.21E-01	4.01E-04	2.70E-03	1.45E-02	1.01E-02	3.81E-01	2.52E-03	2.26E-03
	Beet Lake	2.45E-02	6.02E-03	2.10E-01	2.99E-04	1.35E-03	1.44E-02	7.90E-03	2.93E-01	1.40E-03	1.25E-03
	Lloyd Lake	1.51E-02	5.46E-03	1.98E-01	2.00E-04	7.12E-04	1.44E-02	5.65E-03	2.07E-01	3.12E-04	7.01E-04
	Reference (Broach Lake)	5.77E-02	4.36E-02	1.13E+00	2.04E-03	6.80E-03	5.77E-02	4.36E-02	1.13E+00	2.04E-03	6.80E-03
	Patterson Lake North Arm - West Basin	1.77E-01	5.66E-02	1.32E+00	5.44E-03	2.23E-02	6.12E-02	1.07E-01	2.93E+00	4.54E-02	6.80E-02
Mallard	Patterson Lake South Arm	1.34E-01	5.32E-02	1.27E+00	4.34E-03	1.57E-02	5.80E-02	8.20E-02	2.19E+00	2.72E-02	2.37E-02
	Beet Lake	7.07E-02	4.87E-02	1.20E+00	3.24E-03	1.01E-02	5.78E-02	6.39E-02	1.68E+00	1.51E-02	1.31E-02
	Lloyd Lake	5.85E-02	4.42E-02	1.14E+00	2.16E-03	7.10E-03	5.77E-02	4.57E-02	1.19E+00	3.37E-03	7.37E-03
	Reference (Broach Lake)	7.99E-02	1.09E-01	3.39E+00	6.19E-03	1.83E-02	7.99E-02	1.09E-01	3.39E+00	6.19E-03	1.83E-02
	Patterson Lake North Arm - West Basin	1.93E-01	1.36E-01	3.93E+00	1.44E-02	1.01E-01	8.32E-02	2.42E-01	8.65E+00	1.11E-01	7.52E-02
Rusty Blackbird	Patterson Lake South Arm	1.51E-01	1.29E-01	3.79E+00	1.17E-02	3.62E-02	8.01E-02	1.90E-01	6.48E+00	6.68E-02	3.35E-02
	Beet Lake	9.21E-02	1.19E-01	3.60E+00	9.07E-03	2.71E-02	7.99E-02	1.51E-01	5.01E+00	3.77E-02	2.41E-02
	Lloyd Lake	8.06E-02	1.10E-01	3.41E+00	6.48E-03	1.86E-02	7.99E-02	1.13E-01	3.56E+00	9.40E-03	1.88E-02

Table C.10: Maximum Concentrations of Radiological COPCs in Biota for the Application Case and Upper Bound Scenario - Project Lifespan

Polymer Parterson Lake North Arm - West Basin Phytoplankton Phytoplankton Patterson Lake North Arm - West Basin Polymer Phytoplankton Patterson Lake North Arm - West Basin Phytoplankton Phytoplankton Patterson Lake North Arm - West Basin Patterson Lake North Arm - West	Maximum Concentration Project Lifespan (Bq/kg fw)													
Reference (Broach Lake) 1.79E-02 1.79E-02 2.11E+00 9.03E-01 9.03E-01 1.79E-02 1.79E-02 2.11E+00 9.03E-01 9.03E-01 1.79E-02 1.79E-02 2.11E+00 9.03E-01 1.79E-02 1.79E-02 1.79E-02 2.11E+00 9.03E-01 1.79E-02 1.79E														
Macrophytes Macrophytes Patterson Lake North Arm - West Basin 1.96E-01 1.96E-01 1.96E-01 1.96E-01 1.47E+00 9.92E+00 1.47E+00 1.49E-01 1.49E-0	Lead- 210	Polonium- 210												
Aquatic Plants Macrophytes Beet Lake Patterson Lake South Arm Beet Lake 6.57E-02 6.59E-02 7.91E+00 1.41E+00 9.92E+00 1.55E+00 1.49E-01 1.49E-01 7.99E+00 1.15E+00 Aquatic Plants Beet Lake 3.55E-02 3.55E-02 5.16E+00 1.00E+00 9.66E+00 1.51E+00 6.60E-02 6.62E-02 5.20E+00 1.01E+00 Phytoplankton Reference (Broach Lake) 1.79E-02 1.79E-02 2.11E+00 1.20E+01 1.57E+00 1.47E+01 2.23E+00 2.23E-02 2.24E-02 2.11E+00 9.93E-01 Phytoplankton Patterson Lake North Arm - West Basin 1.99E-01 1.96E-01 1.20E+01 1.57E+00 1.47E+01 2.29E+00 5.33E-01 1.59E-02 2.11E+00 9.92E+00 1.51E+00 6.60E-02 7.9E+00 1.15E+00	9.62E+00	1.51E+00												
Aquatic Plants Beet Lake 1.55E-02 1.55E-02 1.55E-02 1.55E-02 2.42E+00 1.00E+00 9.66E+00 1.51E+00 2.23E-02 2.24E-02 2.43E+00 9.13E-01	3.69E+01	5.84E+00												
Reference (Broach Lake) 1.95E-02 1.95E-02 2.42E+00 9.12E-01 9.63E+00 1.51E+00 2.23E-02 2.24E-02 2.43E+00 9.3E-01 9.03E-01 9.03E-01 1.79E-02 1.79E-	1.11E+01	1.74E+00												
Phytoplankton Phytoplankto	9.82E+00	1.54E+00												
Phytoplankton Phytoplankton Phytoplankton Phytoplankton Patterson Lake North Arm - West Basin N.9E-02 1.79E-02 1.79E	9.63E+00	1.51E+00												
Phytoplankton Patterson Lake South Arm 6.57E-02 6.59E-02 7.91E+00 1.14E+00 9.92E+00 1.55E+00 1.49E-01 7.99E+00 1.15E+00 1.01E+00 1.01E+00 1.00E+00	9.62E+00	1.51E+00												
Beet Lake 1.95E-02 1.95E-02 2.42E+00 2.10E+00 2.66E+00 1.51E+00 6.60E-02 6.62E-02 2.24E-00 2.10E+00 2.23E-02 2.24E-00 3.69E+01	5.84E+00													
Reference (Broach Lake) 1.95E-02 3.42E-02 3.42E-02 4.52E+00 9.12E-01 9.63E+00 1.51E+00 2.23E-02 2.24E-02 2.43E+00 9.13E-01	1.11E+01	1.74E+00												
Reference (Broach Lake)	9.82E+00	1.54E+00												
Patterson Lake North Arm - West Basin 1.07E-01 1.	9.63E+00	1.51E+00												
Patterson Lake South Arm 1.07E-01 1.07E-01 1.07E-01 1.07E-01 1.62E+01 1.63E+00 4.92E+01 2.36E-01 2.37E-01 1.64E+01 1.64E+01 1.64E+00 1.64	6.88E+00	4.80E+01												
Invertebrates Patterson Lake South Arm 1.07E-01 1.07E-01 1.07E-01 1.07E-01 1.07E-01 1.62E+01 1.63E+00 7.05E+00 4.92E+01 2.36E-01 2.37E-01 1.64E+01 1.64E+00 1.47E+00 1.47E+00 1.47E+00 1.47E+00 1.47E+00 1.47E+00 1.47E+00 1.48E+01 1.08E-01 1.09E-01 1.08E+01 1.08E+01 1.08E+01 1.47E+00	1.89E+01	1.32E+02												
Beet Lake 6.08E-02 6.09E-02 1.07E+01 1.47E+00 6.91E+00 4.82E+01 1.08E-01 1.09E-01 1.09E-01 1.08E+01 1.47E+00	7.69E+00	5.36E+01												
Zooplankton Reference (Broach Lake) 3.56E-02 3.56E-02 4.54E+00 1.38E+00 1.17E+01 8.13E+01 3.56E-02 4.54E+00 1.38E+00 Aquatic Animals Reference (Broach Lake) Reference (Broach Lake) 3.56E-02 3.56E-02 4.54E+00 1.38E+00 1.78E+01 1.78E+01 1.23E+02 1.06E+00 1.06E+00 2.61E+01 2.46E+00 1.30E-01 1.31E-01 1.71E+01 1.75E+00 1.20E+01 8.38E+01 2.95E-01 2.96E-01 1.72E+01 1.76E+00 1.09d Lake 4.89E-02 3.88E-02 5.22E+00 1.40E+00 1.17E+01 8.13E+01 4.43E-02 7.22E-02 7.32E+00 1.45E+00 1.09d Lake 3.88E-02 3.88E-02 5.22E+00 1.40E+00 1.17E+01 8.13E+01 4.43E-02 4.44E-02 5.23E+00 1.40E+00 1.09d Lake 7.12E-03 7.12E-03 5.45E-02 6.65E-02 3.04E-01 7.73E-01 7.12E-03 7.12E-03 5.45E-02 6.65E-02	6.99E+00	4.88E+01												
Zooplankton Patterson Lake North Arm - West Basin Patterson Lake South Arm Beet Lake Lioyd Lake 3.88E-01 1.30E-01 1.31E-01 1.31E-01 1.71E+01 1.75E+00 1.20E+01 1.75E+00 1.20E+01 1.77E+01 1.7	6.89E+00	4.81E+01												
Zooplankton Patterson Lake South Arm 1.30E-01 1.31E-01 1.71E+01 1.75E+00 1.20E+01 8.38E+01 2.95E-01 2.96E-01 1.72E+01 1.76E+00 Beet Lake 4.89E-02 4.89E-02 7.28E+00 1.45E+00 1.17E+01 8.14E+01 7.20E-02 7.22E-02 7.32E+00 1.45E+00 Lloyd Lake 3.88E-02 3.88E-02 5.22E+00 1.40E+00 1.17E+01 8.13E+01 4.43E-02 4.44E-02 5.23E+00 1.40E+00 Reference (Broach Lake) 7.12E-03 7.12E-03 5.45E-02 6.65E-02 3.04E-01 7.73E-01 7.12E-03 7.12E-03 5.45E-02 6.65E-02	1.17E+01	8.13E+01												
Beet Lake 4.89E-02 4.89E-02 7.28E+00 1.45E+00 1.17E+01 8.14E+01 7.20E-02 7.22E-02 7.32E+00 1.40E+00	4.47E+01	3.15E+02												
Aquatic Animals Lloyd Lake 3.88E-02 3.88E-02 5.22E+00 1.40E+00 1.17E+01 8.13E+01 4.43E-02 4.44E-02 5.23E+00 1.40E+00 Reference (Broach Lake) 7.12E-03 7.12E-03 5.45E-02 6.65E-02 3.04E-01 7.73E-01 7.12E-03 7.12E-03 5.45E-02 6.65E-02	1.35E+01	9.41E+01												
Aquatic Animals Reference (Broach Lake) 7.12E-03 7.12E-03 5.45E-02 6.65E-02 3.04E-01 7.73E-01 7.12E-03 7.12E-03 5.45E-02 6.65E-02	1.17E+01	8.17E+01												
Reference (Broach Lake) /.12E-03 /.12E-03 5.45E-02 6.65E-02 3.04E-01 /.73E-01 /.12E-03 7.12E-03 5.45E-02 6.65E-02	1.17E+01	8.14E+01												
Patterson Lake North Arm - West Basin 7.76F-02 7.79F-02 3.10F-01 1.16F-01 4.64F-01 1.17F+00 2.11F-01 2.12F-01 3.13F-01 1.18F-01	3.04E-01	7.73E-01												
	1.17E+00	2.99E+00												
Northern Pike Patterson Lake South Arm 2.61E-02 2.61E-02 2.05E-01 8.38E-02 3.13E-01 7.96E-01 5.89E-02 5.91E-02 2.07E-01 8.46E-02	3.51E-01	8.94E-01												
Beet Lake 1.41E-02 1.41E-02 1.33E-01 7.39E-02 3.05E-01 7.76E-01 2.62E-02 2.63E-02 1.35E-01 7.42E-02	3.10E-01	7.89E-01												
Lloyd Lake 7.75E-03 7.76E-03 6.27E-02 6.72E-02 3.04E-01 7.73E-01 8.87E-03 8.88E-03 6.28E-02 6.72E-02	3.04E-01	7.74E-01												
Reference (Broach Lake) 1.76E-03 1.76E-03 5.45E-02 6.60E-02 1.00E-01 6.94E-01 1.76E-03 1.76E-03 5.45E-02 6.60E-02	1.00E-01	6.94E-01												
Patterson Lake North Arm - West Basin 1.83E-02 1.84E-02 3.08E-01 1.12E-01 1.49E-01 1.03E+00 4.88E-02 4.90E-02 3.11E-01 1.15E-01	3.62E-01	2.53E+00												
Whitefish Patterson Lake South Arm 6.16E-03 6.18E-03 2.02E-01 8.22E-02 1.03E-01 7.14E-01 1.38E-02 1.39E-02 2.03E-01 8.29E-02	1.15E-01	7.96E-01												
Beet Lake 3.38E-03 3.38E-03 1.32E-01 7.29E-02 1.00E-01 6.97E-01 6.20E-03 6.22E-03 1.33E-01 7.32E-02	1.02E-01	7.07E-01												
Lloyd Lake 1.91E-03 1.91E-03 6.24E-02 6.67E-02 1.00E-01 6.94E-01 2.17E-03 2.17E-03 6.25E-02 6.67E-02	1.00E-01	6.95E-01												
Reference (Broach Lake) 4.95E-01 4.95E-01 6.24E-02 1.08E+00 3.20E-01 4.33E-01 4.95E-01 4.95E-01 6.24E-02 1.08E+00	3.20E-01	4.33E-01												
Patterson Lake North Arm - West Basin 2.24E+00 2.24E+00 1.06E-01 2.58E+00 1.19E+00 1.45E+00 2.24E+00 2.24E+00 2.24E+00 2.58E+00	1.19E+00	1.45E+00												
Blueberries Patterson Lake South Arm 7.80E-01 7.80E-01 6.96E-02 1.32E+00 4.62E-01 6.00E-01 7.80E-01 7.80E-01 6.96E-02 1.32E+00	4.62E-01	6.00E-01												
Beet Lake 6.61E-01 6.66E-02 1.22E+00 4.03E-01 5.30E-01 6.61E-01 6.61E-01 6.66E-02 1.22E+00	4.03E-01	5.30E-01												
Uoyd Jake 5.01F-01 5.01F-01 6.25F-02 1.08F+00 3.23F-01 4.37F-01 5.01F-01 6.25F-02 1.08F+00	3.23E-01	4.37E-01												
Terrestrial Plants Reference (Broach Lake) 9.90E-01 9.90E-01 1.25E-01 2.15E+00 6.40E-01 8.67E-01 9.90E-01 9.90E-01 1.25E-01 2.15E+00	6.40E-01	8.67E-01												
Patterson Lake North Arm - West Basin 4.34E+00 4.34E+00 2.72E+00 6.07E+00 3.28E+00 3.82E+00 4.34E+00 4.34E+00 6.07E+00	3.28E+00	3.82E+00												
Browse Patterson Lake South Arm 1.54E+00 1.54E+00 5.48E-01 2.79E+00 1.07E+00 1.35E+00 1.54E+00 5.48E-01 2.79E+00	1.07E+00	1.35E+00												
Beet Lake 1.31E+00 1.31E+00 3.72E-01 2.52E+00 8.91E-01 1.15E+00 1.31E+00 3.72E-01 2.52E+00	8.91E-01	1.15E+00												
Lloyd Lake 1.00E+00 1.33E-01 2.16E+00 6.48E-01 8.76E-01 1.00E+00 1.33E-01 2.16E+00		8.76E-01												

Table C.10: Maximum Concentrations of Radiological COPCs in Biota for the Application Case and Upper Bound Scenario - Project Lifespan

						Ma	aximum Co	ncentration I	Project Lifes	pan (Bq/kg	fw)			
	Biota	Location			Applicat	ion Case					Upper Bou	ınd Scenario	,	
			Uranium-	Uranium-	Thorium-	Radium-	Lead-	Polonium-	Uranium-	Uranium-	Thorium-	Radium-	Lead-	Polonium-
			238	234	230	226	210	210	238	234	230	226	210	210
		Reference (Broach Lake)	9.90E-01	9.90E-01	1.25E-01	2.15E+00	6.40E-01	8.67E-01	9.90E-01	9.90E-01	1.25E-01	2.15E+00	6.40E-01	8.67E-01
		Patterson Lake North Arm - West Basin	4.34E+00	4.34E+00	2.42E+01	2.76E+01	2.47E+01	2.21E+01	4.34E+00	4.34E+00	2.42E+01	2.76E+01	2.47E+01	2.21E+01
	Labrador Tea	Patterson Lake South Arm	1.54E+00	1.54E+00	4.06E+00	6.30E+00	4.57E+00	4.32E+00	1.54E+00	1.54E+00	4.06E+00	6.30E+00	4.57E+00	4.32E+00
		Beet Lake	1.31E+00	1.31E+00	2.42E+00	4.57E+00	2.93E+00	2.88E+00	1.31E+00	1.31E+00	2.42E+00	4.57E+00	2.93E+00	2.88E+00
		Lloyd Lake	1.00E+00	1.00E+00	2.01E-01	2.23E+00	7.17E-01	9.34E-01	1.00E+00	1.00E+00	2.01E-01	2.23E+00	7.17E-01	9.34E-01
		Reference (Broach Lake)	2.08E-01	2.08E-01	6.77E-01	1.33E+00	1.91E+02	1.45E+02	2.08E-01	2.08E-01	6.77E-01	1.33E+00	1.91E+02	1.45E+02
		Patterson Lake North Arm - West Basin	2.08E-01	2.08E-01	1.38E+02	1.39E+02	3.22E+02	1.72E+02	2.08E-01	2.08E-01	1.38E+02	1.39E+02	3.22E+02	1.72E+02
	Lichen	Patterson Lake South Arm	2.08E-01	2.08E-01	2.32E+01	2.38E+01	2.12E+02	1.49E+02	2.08E-01	2.08E-01	2.32E+01	2.38E+01	2.12E+02	1.49E+02
		Beet Lake	2.08E-01	2.08E-01	1.38E+01	1.44E+01	2.04E+02	1.47E+02	2.08E-01	2.08E-01	1.38E+01	1.44E+01	2.04E+02	1.47E+02
		Lloyd Lake	2.08E-01	2.08E-01	1.12E+00	1.77E+00	1.92E+02	1.45E+02	2.08E-01	2.08E-01	1.12E+00	1.77E+00	1.92E+02	1.45E+02
		Reference (Broach Lake)	2.70E-02	2.70E-02	1.77E-01	1.61E+00	1.14E+00	3.12E-02	2.70E-02	2.70E-02	1.77E-01	1.61E+00	1.14E+00	3.12E-02
	T	Patterson Lake North Arm - West Basin	1.18E-01	1.18E-01	2.80E+00	5.19E+00	3.88E+00	2.42E+00	1.18E-01	1.18E-01	2.80E+00	5.19E+00	3.88E+00	2.42E+00
	Terrestrial Invertebrate	Patterson Lake South Arm	4.19E-02	4.19E-02	6.05E-01	2.20E+00	1.59E+00	4.21E-01	4.19E-02	4.19E-02	6.05E-01	2.20E+00	1.59E+00	4.21E-01
	ilivertebrate	Beet Lake	3.57E-02	3.57E-02	4.26E-01	1.95E+00	1.40E+00	2.58E-01	3.57E-02	3.57E-02	4.26E-01	1.95E+00	1.40E+00	2.58E-01
		Lloyd Lake	2.73E-02	2.73E-02	1.85E-01	1.63E+00	1.15E+00	3.88E-02	2.73E-02	2.73E-02	1.85E-01	1.63E+00	1.15E+00	3.88E-02
		Reference (Broach Lake)	1.81E-02	1.81E-02	4.78E-03	1.84E-01	6.65E-02	2.94E-01	1.81E-02	1.81E-02	4.78E-03	1.84E-01	6.65E-02	2.94E-01
		Patterson Lake North Arm - West Basin	7.98E-02	7.98E-02	4.51E-02	5.01E-01	1.72E-01	1.01E+00	8.13E-02	8.13E-02	4.52E-02	5.02E-01	2.59E-01	1.15E+00
	Beaver	Patterson Lake South Arm	2.82E-02	2.82E-02	1.62E-02	2.38E-01	8.17E-02	4.08E-01	2.86E-02	2.86E-02	1.63E-02	2.38E-01	8.64E-02	4.16E-01
		Beet Lake	2.39E-02	2.39E-02	1.10E-02	2.15E-01	7.49E-02	3.59E-01	2.41E-02	2.41E-02	1.10E-02	2.15E-01	7.55E-02	3.61E-01
		Lloyd Lake	1.83E-02	1.83E-02	5.23E-03	1.86E-01	6.68E-02	2.96E-01	1.83E-02	1.83E-02	5.24E-03	1.86E-01	6.69E-02	2.96E-01
		Reference (Broach Lake)	1.35E-02	1.35E-02	2.77E-03	1.36E-01	2.87E-02	2.39E-01	1.35E-02	1.35E-02	2.77E-03	1.36E-01	2.87E-02	2.39E-01
	Black Bear	Patterson Lake North Arm - West Basin	3.84E-02	3.84E-02	4.73E-03	2.30E-01	5.20E-02	4.39E-01	3.89E-02	3.89E-02	4.75E-03	2.31E-01	5.59E-02	5.10E-01
	Diack Deal	Beet Lake	1.59E-02	1.59E-02	3.94E-03	1.50E-01	3.57E-02	2.66E-01	1.60E-02	1.60E-02	3.94E-03	1.50E-01	3.58E-02	2.67E-01
		Lloyd Lake	1.36E-02	1.36E-02	2.87E-03	1.37E-01	2.89E-02	2.41E-01	1.36E-02	1.36E-02	2.87E-03	1.37E-01	2.89E-02	2.41E-01
Terrestrial		Reference (Broach Lake)	5.46E-04	5.46E-04	1.52E-03	1.88E-02	2.07E-02	4.47E-01	5.46E-04	5.46E-04	1.52E-03	1.88E-02	2.07E-02	4.47E-01
Animals	Grey Wolf	Patterson Lake North-West Basin	1.04E-03	1.04E-03	2.07E-03	3.21E-02	2.31E-02	4.92E-01	1.07E-03	1.07E-03	2.07E-03	3.21E-02	2.41E-02	5.08E-01
		Lloyd Lake	5.52E-04	5.52E-04	1.54E-03	1.90E-02	2.08E-02	4.47E-01	5.53E-04	5.53E-04	1.54E-03	1.90E-02	2.08E-02	4.47E-01
		Reference (Broach Lake)	3.02E-04	3.02E-04	3.86E-03	1.50E-02	2.57E-02	1.20E+00	3.02E-04	3.02E-04	3.86E-03	1.50E-02	2.57E-02	1.20E+00
		Patterson Lake North Arm - West Basin	1.97E-03	1.97E-03	1.88E-02	2.42E-02	3.46E-02	1.64E+00	3.90E-03	3.91E-03	1.90E-02	2.45E-02	6.48E-02	3.27E+00
	Mink	Patterson Lake South Arm	6.81E-04	6.82E-04	1.22E-02	1.76E-02	2.63E-02	1.23E+00	1.20E-03	1.20E-03	1.24E-02	1.77E-02	2.83E-02	1.33E+00
		Beet Lake	4.60E-04	4.61E-04	8.27E-03	1.62E-02	2.58E-02	1.20E+00	6.49E-04	6.50E-04	8.33E-03	1.62E-02	2.61E-02	1.21E+00
		Lloyd Lake	3.14E-04	3.14E-04	4.31E-03	1.51E-02	2.57E-02	1.20E+00	3.31E-04	3.31E-04	4.32E-03	1.51E-02	2.57E-02	1.20E+00
		Reference (Broach Lake)	1.64E-02	1.64E-02	7.27E-03	1.78E-01	1.06E-01	3.65E-01	1.64E-02	1.64E-02	7.27E-03	1.78E-01	1.06E-01	3.65E-01
		Patterson Lake North Arm - West Basin	7.29E-02	7.29E-02	5.72E-02	4.71E-01	2.24E-01	1.05E+00	7.61E-02	7.61E-02	5.76E-02	4.72E-01	4.01E-01	1.34E+00
	Moose	Patterson Lake South Arm	7.23E-02	7.23E-02	5.64E-02	4.68E-01	2.23E-01	1.04E+00	7.55E-02	7.55E-02	5.67E-02	4.69E-01	4.00E-01	1.34E+00
		Beet Lake	2.18E-02	2.18E-02	1.68E-02	2.07E-01	1.14E-01	4.24E-01	2.21E-02	2.21E-02	1.69E-02	2.07E-01	1.15E-01	4.26E-01
		Lloyd Lake	1.66E-02	1.66E-02	8.07E-03	1.79E-01	1.07E-01	3.67E-01	1.66E-02	1.66E-02	8.09E-03	1.79E-01	1.07E-01	3.67E-01
		Reference (Broach Lake)	2.90E-02	2.90E-02	1.99E+00	9.96E-02	3.34E+00	3.65E-03	2.90E-02	2.90E-02	1.99E+00	9.96E-02	3.34E+00	3.65E-03
	Moose Organs	Patterson Lake North Arm - West Basin	1.29E-01	1.29E-01	1.57E+01	2.63E-01	7.04E+00	1.05E-02	1.35E-01	1.35E-01	1.58E+01	2.64E-01	1.26E+01	1.34E-02
		Patterson Lake South Arm	1.28E-01	1.28E-01	1.54E+01	2.62E-01	7.01E+00	1.04E-02	1.34E-01	1.34E-01	1.55E+01	2.62E-01	1.26E+01	1.34E-02

Table C.10: Maximum Concentrations of Radiological COPCs in Biota for the Application Case and Upper Bound Scenario - Project Lifespan

						Ma	aximum Co	ncentration F	Project Lifes	pan (Bq/kg t	fw)			
	Biota	Location			Applicat	ion Case					Upper Bou	nd Scenaric		
			Uranium-	Uranium-	Thorium-	Radium-	Lead-	Polonium-	Uranium-	Uranium-	Thorium-	Radium-	Lead-	Polonium-
			238	234	230	226	210	210	238	234	230	226	210	210
		Beet Lake	3.85E-02	3.85E-02	4.59E+00	1.16E-01	3.58E+00	4.24E-03	3.90E-02	3.90E-02	4.62E+00	1.16E-01	3.62E+00	4.26E-03
		Lloyd Lake	2.93E-02	2.93E-02	2.21E+00	1.00E-01	3.35E+00	3.67E-03	2.94E-02	2.94E-02	2.21E+00	1.00E-01	3.35E+00	3.67E-03
		Reference (Broach Lake)	9.58E-04	9.58E-04	1.29E-02	6.20E-02	1.63E-01	2.41E+00	9.58E-04	9.58E-04	1.29E-02	6.20E-02	1.63E-01	2.41E+00
		Patterson Lake North Arm - West Basin	8.65E-03	8.68E-03	7.23E-02	1.00E-01	2.28E-01	3.30E+00	2.23E-02	2.24E-02	7.30E-02	1.02E-01	4.81E-01	6.63E+00
	Muskrat	Patterson Lake South Arm	2.99E-03	3.00E-03	4.68E-02	7.52E-02	1.67E-01	2.47E+00	6.62E-03	6.65E-03	4.72E-02	7.58E-02	1.83E-01	2.70E+00
		Beet Lake	1.70E-03	1.71E-03	3.07E-02	6.76E-02	1.63E-01	2.42E+00	3.04E-03	3.04E-03	3.09E-02	6.78E-02	1.65E-01	2.45E+00
		Lloyd Lake	1.03E-03	1.03E-03	1.48E-02	6.25E-02	1.63E-01	2.41E+00	1.15E-03	1.15E-03	1.48E-02	6.26E-02	1.63E-01	2.42E+00
		Reference (Broach Lake)	1.01E-03	1.01E-03	7.96E-04	2.29E-02	7.58E-03	2.12E-02	1.01E-03	1.01E-03	7.96E-04	2.29E-02	7.58E-03	2.12E-02
		Patterson Lake North Arm - West Basin	4.54E-03	4.54E-03	3.99E-03	6.16E-02	1.88E-02	1.11E-01	4.57E-03	4.57E-03	3.99E-03	6.16E-02	1.89E-02	1.11E-01
	Red Fox	Patterson Lake South Arm	1.58E-03	1.58E-03	1.37E-03	2.92E-02	9.41E-03	3.58E-02	1.59E-03	1.59E-03	1.38E-03	2.92E-02	9.42E-03	3.58E-02
		Beet Lake	1.34E-03	1.34E-03	1.13E-03	2.66E-02	8.65E-03	2.97E-02	1.34E-03	1.34E-03	1.13E-03	2.66E-02	8.65E-03	2.97E-02
		Lloyd Lake	1.02E-03	1.02E-03	8.10E-04	2.30E-02	7.62E-03	2.15E-02	1.02E-03	1.02E-03	8.11E-04	2.30E-02	7.62E-03	2.15E-02
		Reference (Broach Lake)	1.69E-02	1.69E-02	2.73E-03	1.67E-01	2.82E-02	2.03E-01	1.69E-02	1.69E-02	2.73E-03	1.67E-01	2.82E-02	2.03E-01
		Patterson Lake North Arm - West Basin	7.44E-02	7.44E-02	2.75E-02	4.54E-01	1.07E-01	8.31E-01	7.44E-02	7.44E-02	2.75E-02	4.54E-01	1.07E-01	8.32E-01
Sı	nowshoe Hare	Patterson Lake South Arm	2.63E-02	2.63E-02	6.82E-03	2.14E-01	4.11E-02	3.06E-01	2.63E-02	2.63E-02	6.82E-03	2.14E-01	4.11E-02	3.06E-01
		Beet Lake	2.24E-02	2.24E-02	5.11E-03	1.94E-01	3.57E-02	2.63E-01	2.24E-02	2.24E-02	5.11E-03	1.94E-01	3.57E-02	2.63E-01
		Lloyd Lake	1.71E-02	1.71E-02	2.81E-03	1.68E-01	2.85E-02	2.05E-01	1.71E-02	1.71E-02	2.81E-03	1.68E-01	2.85E-02	2.05E-01
		Reference (Broach Lake)	6.15E-03	6.15E-03	8.06E-04	5.98E-02	9.13E-03	7.20E-02	6.15E-03	6.15E-03	8.06E-04	5.98E-02	9.13E-03	7.20E-02
	Sanathania Bard	Patterson Lake North Arm - West Basin	2.74E-02	2.74E-02	5.81E-03	1.53E-01	3.34E-02	2.68E-01	2.74E-02	2.74E-02	5.81E-03	1.53E-01	3.35E-02	2.68E-01
	Southern Red- Backed Vole	Patterson Lake South Arm	9.62E-03	9.62E-03	1.65E-03	7.50E-02	1.31E-02	1.04E-01	9.62E-03	9.62E-03	1.65E-03	7.50E-02	1.31E-02	1.04E-01
	backed voie	Beet Lake	8.17E-03	8.17E-03	1.29E-03	6.87E-02	1.14E-02	9.07E-02	8.17E-03	8.17E-03	1.29E-03	6.87E-02	1.14E-02	9.07E-02
		Lloyd Lake	6.22E-03	6.22E-03	8.23E-04	6.01E-02	9.21E-03	7.27E-02	6.22E-03	6.22E-03	8.23E-04	6.01E-02	9.21E-03	7.27E-02
		Reference (Broach Lake)	7.48E-03	7.48E-03	8.89E-03	1.16E-01	9.35E-01	4.81E+00	7.48E-03	7.48E-03	8.89E-03	1.16E-01	9.35E-01	4.81E+00
	Woodland	Patterson Lake North Arm - West Basin	2.11E-02	2.11E-02	1.33E-01	9.28E-01	1.27E+00	5.44E+00	2.50E-02	2.50E-02	1.33E-01	9.29E-01	1.35E+00	5.68E+00
	Caribou	Beet Lake	8.62E-03	8.62E-03	2.15E-02	1.87E-01	9.66E-01	4.88E+00	8.93E-03	8.93E-03	2.15E-02	1.87E-01	9.66E-01	4.88E+00
		Lloyd Lake	6.22E-03	6.22E-03	8.23E-04	6.01E-02	9.21E-03	7.27E-02	6.22E-03	6.22E-03	8.23E-04	6.01E-02	9.21E-03	7.27E-02
		Reference (Broach Lake)	4.04E-01	4.04E-01	2.33E-03	3.80E-02	3.34E-02	1.30E+00	4.04E-01	4.04E-01	2.33E-03	3.80E-02	3.34E-02	1.30E+00
		Patterson Lake North Arm - West Basin	1.78E+00	1.78E+00	1.37E-02	9.85E-02	9.94E-02	4.84E+00	1.78E+00	1.78E+00	1.37E-02	9.85E-02	9.95E-02	4.84E+00
	Grouse	Patterson Lake South Arm	6.28E-01	6.28E-01	4.21E-03	4.79E-02	4.42E-02	1.88E+00	6.28E-01	6.28E-01	4.21E-03	4.79E-02	4.42E-02	1.88E+00
		Beet Lake	5.34E-01	5.34E-01	3.43E-03	4.37E-02	3.97E-02	1.64E+00	5.34E-01	5.34E-01	3.43E-03	4.37E-02	3.97E-02	1.64E+00
		Lloyd Lake	4.08E-01	4.08E-01	2.37E-03	3.82E-02	3.36E-02	1.31E+00	4.08E-01	4.08E-01	2.37E-03	3.82E-02	3.36E-02	1.31E+00
		Reference (Broach Lake)	5.66E-02	5.66E-02	3.38E-04	5.34E-03	4.74E-03	1.83E-01	5.66E-02	5.66E-02	3.38E-04	5.34E-03	4.74E-03	1.83E-01
		Patterson Lake North Arm - West Basin	2.48E-01	2.48E-01	2.40E-03	1.43E-02	1.48E-02	7.18E-01	2.49E-01	2.49E-01	2.40E-03	1.43E-02	1.48E-02	7.19E-01
C	Canada Goose	Patterson Lake South Arm	8.79E-02	8.79E-02	6.89E-04	6.80E-03	6.37E-03	2.71E-01	8.80E-02	8.80E-02	6.90E-04	6.80E-03	6.38E-03	2.71E-01
		Beet Lake	7.48E-02	7.48E-02	5.41E-04	6.19E-03	5.69E-03	2.34E-01	7.49E-02	7.49E-02	5.42E-04	6.19E-03	5.69E-03	2.34E-01
		Lloyd Lake	5.72E-02	5.72E-02	3.46E-04	5.37E-03	4.77E-03	1.85E-01	5.72E-02	5.72E-02	3.46E-04	5.37E-03	4.77E-03	1.85E-01
		Reference (Broach Lake)	2.36E-04	2.36E-04	1.83E-02	4.04E-02	8.48E-02	4.23E+00	2.36E-04	2.36E-04	1.83E-02	4.04E-02	8.48E-02	4.23E+00
	Little Brown	Patterson Lake North Arm - West Basin	2.12E-03	2.13E-03	1.02E-01	6.47E-02	1.16E-01	5.79E+00	5.47E-03	5.49E-03	1.03E-01	6.59E-02	2.33E-01	1.16E+01
	Myotis	Patterson Lake South Arm	7.34E-04	7.36E-04	6.57E-02	4.88E-02	8.69E-02	4.33E+00	1.63E-03	1.63E-03	6.64E-02	4.92E-02	9.47E-02	4.72E+00
		Beet Lake	4.19E-04	4.19E-04	4.32E-02	4.39E-02		4.24E+00	7.45E-04	7.47E-04	4.36E-02	1	8.62E-02	4.30E+00

Table C.10: Maximum Concentrations of Radiological COPCs in Biota for the Application Case and Upper Bound Scenario - Project Lifespan

					Ma	aximum Co	ncentration F	Project Lifes	pan (Bq/kg f	fw)			
Biota	Location			Applicat	ion Case					Upper Bou	nd Scenario		
		Uranium- 238	Uranium- 234	Thorium- 230	Radium- 226	Lead- 210	Polonium- 210	Uranium- 238	Uranium- 234	Thorium- 230	Radium- 226	Lead- 210	Polonium- 210
	Lloyd Lake	2.52E-04	2.52E-04	2.09E-02	4.07E-02	8.48E-02	4.23E+00	2.82E-04	2.82E-04	2.09E-02	4.07E-02	8.49E-02	4.23E+00
	Reference (Broach Lake)	2.29E-03	2.29E-03	1.55E-03	1.81E-03	1.48E-02	4.07E+00	2.29E-03	2.29E-03	1.55E-03	1.81E-03	1.48E-02	4.07E+00
	Patterson Lake North Arm - West Basin	2.37E-02	2.38E-02	8.63E-03	2.97E-03	2.08E-02	5.60E+00	6.22E-02	6.25E-02	8.72E-03	3.03E-03	4.36E-02	1.14E+01
Loon	Patterson Lake South Arm	7.92E-03	7.94E-03	5.58E-03	2.21E-03	1.52E-02	4.17E+00	1.77E-02	1.77E-02	5.63E-03	2.23E-03	1.67E-02	4.55E+00
	Beet Lake	4.36E-03	4.37E-03	3.67E-03	1.98E-03	1.49E-02	4.08E+00	7.92E-03	7.95E-03	3.70E-03	1.99E-03	1.51E-02	4.14E+00
	Lloyd Lake	2.48E-03	2.48E-03	1.77E-03	1.83E-03	1.48E-02	4.07E+00	2.81E-03	2.81E-03	1.77E-03	1.83E-03	1.49E-02	4.07E+00
	Reference (Broach Lake)	2.41E-02	2.41E-02	1.51E-02	1.96E-02	1.95E-01	3.49E+01	2.41E-02	2.41E-02	1.51E-02	1.96E-02	1.95E-01	3.49E+01
	Patterson Lake North Arm - West Basin	2.17E-01	2.18E-01	8.39E-02	3.16E-02	2.71E-01	4.77E+01	5.61E-01	5.64E-01	8.48E-02	3.22E-02	5.62E-01	9.56E+01
Mallard	Patterson Lake South Arm	7.52E-02	7.54E-02	5.42E-02	2.38E-02	1.99E-01	3.57E+01	1.67E-01	1.67E-01	5.47E-02	2.39E-02	2.18E-01	3.89E+01
	Beet Lake	4.29E-02	4.29E-02	3.56E-02	2.14E-02	1.95E-01	3.50E+01	7.64E-02	7.66E-02	3.59E-02	2.15E-02	1.98E-01	3.54E+01
	Lloyd Lake	2.58E-02	2.58E-02	1.72E-02	1.98E-02	1.95E-01	3.49E+01	2.89E-02	2.89E-02	1.72E-02	1.98E-02	1.95E-01	3.49E+01
	Reference (Broach Lake)	1.74E-01	1.74E-01	4.20E-02	5.45E-02	3.34E-01	9.54E+01	1.74E-01	1.74E-01	4.20E-02	5.45E-02	3.34E-01	9.54E+01
	Patterson Lake North Arm - West Basin	8.71E-01	8.72E-01	2.13E-01	9.47E-02	4.59E-01	1.31E+02	1.16E+00	1.17E+00	2.16E-01	9.56E-02	8.45E-01	2.61E+02
Rusty Blackbird	Patterson Lake South Arm	3.03E-01	3.03E-01	1.38E-01	6.44E-02	3.44E-01	9.79E+01	3.79E-01	3.79E-01	1.40E-01	6.47E-02	3.70E-01	1.07E+02
	Beet Lake	2.39E-01	2.39E-01	9.27E-02	5.91E-02	3.37E-01	9.58E+01	2.66E-01	2.66E-01	9.34E-02	5.92E-02	3.40E-01	9.70E+01
	Lloyd Lake	1.77E-01	1.77E-01	4.72E-02	5.48E-02	3.34E-01	9.54E+01	1.79E-01	1.79E-01	4.72E-02	5.48E-02	3.34E-01	9.55E+01

Table C.11: Maximum Concentrations of Radiological COPCs in Biota for the Application Case and Upper Bound Scenario - Far-Future

							Ma <u>ximum</u>	Concentration	n Fa <u>r Futur</u>	e (Bq/kg fw)				
	Biota	Location			Applicat	ion Case				· ('P' J' /	Upper Bou	nd Scenario)	
			Uranium-	Uranium-	Thorium-	:	Lead-	Polonium-	Uranium-	Uranium-	Thorium-	<u> </u>	Lead-	Polonium-
			238	234	230	226	210	210	238	234	230	226	210	210
		Reference (Broach Lake)	1.79E-02	1.79E-02	2.11E+00	9.03E-01	9.62E+00	1.51E+00	1.79E-02	1.79E-02	2.11E+00	9.03E-01	9.62E+00	1.51E+00
		Patterson Lake North Arm - West Basin	5.81E-01	5.83E-01	2.25E+00	1.16E+00	1.01E+01	1.58E+00	5.81E-01	5.83E-01	2.25E+00	1.16E+00	1.01E+01	1.58E+00
	Macrophytes	Patterson Lake South Arm	1.74E-01	1.74E-01	2.20E+00	9.99E-01	9.65E+00	1.51E+00	1.74E-01	1.74E-01	2.20E+00	9.99E-01	9.65E+00	1.51E+00
		Beet Lake	7.58E-02	7.60E-02	2.16E+00	9.44E-01	9.63E+00	1.51E+00	7.59E-02	7.61E-02	2.16E+00	9.44E-01	9.63E+00	1.51E+00
Acustic Disute		Lloyd Lake	2.32E-02	2.32E-02	2.11E+00	9.07E-01	9.63E+00	1.51E+00	2.32E-02	2.32E-02	2.11E+00	9.07E-01	9.63E+00	1.51E+00
Aquatic Plants		Reference (Broach Lake)	1.79E-02	1.79E-02	2.11E+00	9.03E-01	9.62E+00	1.51E+00	1.79E-02	1.79E-02	2.11E+00	9.03E-01	9.62E+00	1.51E+00
		Patterson Lake North Arm - West Basin	5.81E-01	5.83E-01	2.25E+00	1.16E+00	1.01E+01	1.58E+00	5.81E-01	5.83E-01	2.25E+00	1.16E+00	1.01E+01	1.58E+00
	Phytoplankton	Patterson Lake South Arm	1.74E-01	1.74E-01	2.20E+00	9.99E-01	9.65E+00	1.51E+00	1.74E-01	1.74E-01	2.20E+00	9.99E-01	9.65E+00	1.51E+00
		Beet Lake	7.58E-02	7.60E-02	2.16E+00	9.44E-01	9.63E+00	1.51E+00	7.59E-02	7.61E-02	2.16E+00	9.44E-01	9.63E+00	1.51E+00
		Lloyd Lake	2.32E-02	2.32E-02	2.11E+00	9.07E-01	9.63E+00	1.51E+00	2.32E-02	2.32E-02	2.11E+00	9.07E-01	9.63E+00	1.51E+00
		Reference (Broach Lake)	3.42E-02	3.42E-02	4.52E+00	1.35E+00	6.88E+00	4.80E+01	3.42E-02	3.42E-02	4.52E+00	1.35E+00	6.88E+00	4.80E+01
	Donahi.	Patterson Lake North Arm - West Basin	1.11E+00	1.11E+00	4.83E+00	1.74E+00	7.20E+00	5.03E+01	1.11E+00	1.11E+00	4.83E+00	1.74E+00	7.20E+00	5.03E+01
	Benthic Invertebrates	Patterson Lake South Arm	3.31E-01	3.32E-01	4.71E+00	1.49E+00	6.90E+00	4.82E+01	3.31E-01	3.33E-01	4.71E+00	1.49E+00	6.90E+00	4.82E+01
	ilivertebrates	Beet Lake	1.45E-01	1.45E-01	4.62E+00	1.41E+00	6.89E+00	4.81E+01	1.45E-01	1.45E-01	4.62E+00	1.41E+00	6.89E+00	4.81E+01
		Lloyd Lake	4.42E-02	4.42E-02	4.53E+00	1.35E+00	6.88E+00	4.80E+01	4.42E-02	4.42E-02	4.53E+00	1.35E+00	6.88E+00	4.80E+01
		Reference (Broach Lake)	3.56E-02	3.56E-02	4.54E+00	1.38E+00	1.17E+01	8.13E+01	3.56E-02	3.56E-02	4.54E+00	1.38E+00	1.17E+01	8.13E+01
		Patterson Lake North Arm - West Basin	1.15E+00	1.16E+00	4.86E+00	1.79E+00	1.22E+01	8.51E+01	1.15E+00	1.16E+00	4.86E+00	1.79E+00	1.22E+01	8.51E+01
	Zooplankton	Patterson Lake South Arm	3.44E-01	3.46E-01	4.74E+00	1.53E+00	1.17E+01	8.16E+01	3.45E-01	3.46E-01	4.74E+00	1.53E+00	1.17E+01	8.16E+01
		Beet Lake	1.50E-01	1.51E-01	4.65E+00	1.45E+00	1.17E+01	8.14E+01	1.51E-01	1.51E-01	4.65E+00	1.45E+00	1.17E+01	8.14E+01
Aquatic Animals		Lloyd Lake	4.60E-02	4.60E-02	4.56E+00	1.39E+00	1.17E+01	8.13E+01	4.60E-02	4.60E-02	4.56E+00	1.39E+00	1.17E+01	8.13E+01
Aquatic Allillais		Reference (Broach Lake)	7.12E-03	7.12E-03	5.45E-02	6.65E-02	3.04E-01	7.73E-01	7.12E-03	7.12E-03	5.45E-02	6.65E-02	3.04E-01	7.73E-01
		Patterson Lake North Arm - West Basin	2.30E-01	2.31E-01	5.83E-02	8.57E-02	3.18E-01	8.09E-01	2.31E-01	2.31E-01	5.83E-02	8.57E-02	3.18E-01	8.09E-01
	Northern Pike	Patterson Lake South Arm	6.89E-02	6.91E-02	5.69E-02	7.35E-02	3.05E-01	7.75E-01	6.90E-02	6.92E-02	5.69E-02	7.35E-02	3.05E-01	7.75E-01
		Beet Lake	3.01E-02	3.02E-02	5.58E-02	6.95E-02	3.04E-01	7.73E-01	3.01E-02	3.02E-02	5.58E-02	6.95E-02	3.04E-01	7.73E-01
		Lloyd Lake	9.20E-03	9.20E-03	5.47E-02	6.67E-02	3.04E-01	7.73E-01	9.20E-03	9.21E-03	5.47E-02	6.67E-02	3.04E-01	7.73E-01
		Reference (Broach Lake)	1.76E-03	1.76E-03	5.45E-02	6.60E-02	1.00E-01	6.94E-01	1.76E-03	1.76E-03	5.45E-02	6.60E-02	1.00E-01	6.94E-01
		Patterson Lake North Arm - West Basin	5.70E-02	5.73E-02	5.82E-02	8.51E-02	1.05E-01	7.26E-01	5.71E-02	5.73E-02	5.82E-02	8.52E-02	1.05E-01	7.26E-01
	Whitefish	Patterson Lake South Arm	1.71E-02	1.71E-02	5.68E-02	7.30E-02	1.00E-01	6.96E-01	1.71E-02	1.71E-02	5.68E-02	7.31E-02	1.00E-01	6.96E-01
		Beet Lake	7.45E-03	7.47E-03	5.57E-02	6.90E-02	1.00E-01	6.94E-01	7.45E-03	7.48E-03	5.57E-02	6.90E-02	1.00E-01	6.94E-01
		Lloyd Lake	2.28E-03	2.28E-03	5.46E-02	6.63E-02	1.00E-01	6.94E-01	2.28E-03	2.28E-03	5.46E-02	6.63E-02	1.00E-01	6.94E-01
		Reference (Broach Lake)	4.95E-01	4.95E-01	6.24E-02	1.08E+00	3.20E-01	4.33E-01	4.95E-01	4.95E-01	6.24E-02	1.08E+00	3.20E-01	4.33E-01
		Patterson Lake North Arm - West Basin	6.71E-01	6.71E-01	8.26E-02	1.55E+00	3.20E-01	4.33E-01	6.71E-01	6.71E-01	8.26E-02	1.55E+00	3.20E-01	4.33E-01
	Blueberries	Patterson Lake South Arm	5.24E-01	5.24E-01	6.57E-02	1.15E+00	3.20E-01	4.33E-01	5.24E-01	5.24E-01	6.57E-02	1.15E+00	3.20E-01	4.33E-01
		Beet Lake	5.12E-01	5.12E-01	6.43E-02	1.12E+00	3.20E-01	4.33E-01	5.12E-01	5.12E-01	6.43E-02	1.12E+00	3.20E-01	4.33E-01
Terrestrial Plants		Lloyd Lake	4.96E-01	4.96E-01	6.25E-02	1.08E+00	3.20E-01	4.33E-01	4.96E-01	4.96E-01	6.25E-02	1.08E+00	3.20E-01	4.33E-01
refrestrat rialits		Reference (Broach Lake)	9.90E-01	9.90E-01	1.25E-01	2.15E+00	6.40E-01	8.67E-01	9.90E-01	9.90E-01	1.25E-01	2.15E+00	6.40E-01	8.67E-01
		Patterson Lake North Arm - West Basin	1.34E+00	1.34E+00	1.65E-01	3.10E+00	6.40E-01	8.67E-01	1.34E+00	1.34E+00	1.65E-01	3.10E+00	6.40E-01	8.67E-01
	Browse	Patterson Lake South Arm	1.05E+00	1.05E+00	1.31E-01	2.31E+00	6.40E-01	8.67E-01	1.05E+00	1.05E+00	1.31E-01	2.31E+00	6.40E-01	8.67E-01
		Beet Lake	1.02E+00	1.02E+00	1.29E-01	2.24E+00	6.40E-01	8.67E-01	1.02E+00	1.02E+00	1.29E-01	2.24E+00	6.40E-01	8.67E-01
		Lloyd Lake	9.91E-01	9.91E-01	1.25E-01	2.16E+00	6.40E-01	8.67E-01	9.91E-01	9.91E-01	1.25E-01	2.16E+00	6.40E-01	8.67E-01

Table C.11: Maximum Concentrations of Radiological COPCs in Biota for the Application Case and Upper Bound Scenario - Far-Future

							Maximum	Concentration	n Far <u>Futur</u>	e (Bq/kg f <u>w</u>))			
	Biota	Location			Applicat	tion Case						nd Scenario	,	
			Uranium-	Uranium-	Thorium-		Lead-	Polonium-	Uranium-	Uranium-	:	Radium-	Lead-	Polonium-
			238	234	230	226	210	210	238	234	230	226	210	210
		Reference (Broach Lake)	9.90E-01	9.90E-01	1.25E-01	2.15E+00	6.40E-01	8.67E-01	9.90E-01	9.90E-01	1.25E-01	2.15E+00	6.40E-01	8.67E-01
		Patterson Lake North Arm - West Basin	1.34E+00	1.34E+00	1.65E-01	3.10E+00	6.40E-01	8.67E-01	1.34E+00	1.34E+00	1.65E-01	3.10E+00	6.40E-01	8.67E-01
	Labrador Tea	Patterson Lake South Arm	1.05E+00	1.05E+00	1.31E-01	2.31E+00	6.40E-01	8.67E-01	1.05E+00	1.05E+00	1.31E-01	2.31E+00	6.40E-01	8.67E-01
		Beet Lake	1.02E+00	1.02E+00	1.29E-01	2.24E+00	6.40E-01	8.67E-01	1.02E+00	1.02E+00	1.29E-01	2.24E+00	6.40E-01	8.67E-01
		Lloyd Lake	9.91E-01	9.91E-01	1.25E-01	2.16E+00	6.40E-01	8.67E-01	9.91E-01	9.91E-01	1.25E-01	2.16E+00	6.40E-01	8.67E-01
		Reference (Broach Lake)	2.08E-01	2.08E-01	6.77E-01	1.33E+00	1.91E+02	1.45E+02	2.08E-01	2.08E-01	6.77E-01	1.33E+00	1.91E+02	1.45E+02
		Patterson Lake North Arm - West Basin	2.08E-01	2.08E-01	6.77E-01	1.33E+00	1.91E+02	1.45E+02	2.08E-01	2.08E-01	6.77E-01	1.33E+00	1.91E+02	1.45E+02
	Lichen	Patterson Lake South Arm	2.08E-01	2.08E-01	6.77E-01	1.33E+00	1.91E+02	1.45E+02	2.08E-01	2.08E-01	6.77E-01	1.33E+00	1.91E+02	1.45E+02
		Beet Lake	2.08E-01	2.08E-01	6.77E-01	1.33E+00	1.91E+02	1.45E+02	2.08E-01	2.08E-01	6.77E-01	1.33E+00	1.91E+02	1.45E+02
		Lloyd Lake	2.08E-01	2.08E-01	6.77E-01	1.33E+00	1.91E+02	1.45E+02	2.08E-01	2.08E-01	6.77E-01	1.33E+00	1.91E+02	1.45E+02
		Reference (Broach Lake)	2.70E-02	2.70E-02	1.77E-01	1.61E+00	1.14E+00	3.12E-02	2.70E-02	2.70E-02	1.77E-01	1.61E+00	1.14E+00	3.12E-02
		Patterson Lake North Arm - West Basin	3.66E-02	3.66E-02	2.34E-01	2.33E+00	1.14E+00	3.12E-02	3.66E-02	3.66E-02	2.34E-01	2.33E+00	1.14E+00	3.12E-02
	Terrestrial	Patterson Lake South Arm	2.86E-02	2.86E-02	1.86E-01	1.73E+00	1.14E+00	3.12E-02	2.86E-02	2.86E-02	1.86E-01	1.73E+00	1.14E+00	3.12E-02
	Invertebrate	Beet Lake	2.79E-02	2.79E-02	1.82E-01	1.68E+00	1.14E+00	3.12E-02	2.79E-02	2.79E-02	1.82E-01	1.68E+00	1.14E+00	3.12E-02
		Lloyd Lake	2.70E-02	2.70E-02	1.77E-01	1.62E+00	1.14E+00	3.12E-02	2.70E-02	2.70E-02	1.77E-01	1.62E+00	1.14E+00	3.12E-02
		Reference (Broach Lake)	1.81E-02	1.81E-02	4.78E-03	1.84E-01	6.65E-02	2.94E-01	1.81E-02	1.81E-02	4.78E-03	1.84E-01	6.65E-02	2.94E-01
		Patterson Lake North Arm - West Basin	2.77E-02	2.77E-02	5.65E-03	2.64E-01	6.84E-02	2.98E-01	2.77E-02	2.77E-02	5.65E-03	2.64E-01	6.85E-02	2.98E-01
	Beaver	Patterson Lake South Arm	2.00E-02	2.00E-02	5.00E-03	1.98E-01	6.67E-02	2.94E-01	2.00E-02	2.00E-02	5.00E-03	1.98E-01	6.67E-02	2.94E-01
		Beet Lake	1.90E-02	1.90E-02	4.91E-03	1.92E-01	6.66E-02	2.94E-01	1.90E-02	1.90E-02	4.91E-03	1.92E-01	6.66E-02	2.94E-01
		Lloyd Lake	1.81E-02	1.81E-02	4.79E-03	1.85E-01	6.66E-02	2.94E-01	1.81E-02	1.81E-02	4.79E-03	1.85E-01	6.66E-02	2.94E-01
		Reference (Broach Lake)	1.35E-02	1.35E-02	2.77E-03	1.36E-01	2.87E-02	2.39E-01	1.35E-02	1.35E-02	2.77E-03	1.36E-01	2.87E-02	2.39E-01
	Black Bear	Patterson Lake North Arm - West Basin	1.69E-02	1.69E-02	3.26E-03	1.67E-01	2.89E-02	2.41E-01	1.69E-02	1.69E-02	3.26E-03	1.67E-01	2.89E-02	2.41E-01
	ыаск веаг	Beet Lake	1.40E-02	1.40E-02	3.69E-03	1.44E-01	3.37E-02	2.49E-01	1.40E-02	1.40E-02	3.69E-03	1.44E-01	3.37E-02	2.49E-01
		Lloyd Lake	1.35E-02	1.35E-02	2.82E-03	1.36E-01	2.88E-02	2.39E-01	1.35E-02	1.35E-02	2.82E-03	1.36E-01	2.88E-02	2.39E-01
Terrestrial		Reference (Broach Lake)	5.46E-04	5.46E-04	1.52E-03	1.88E-02	2.07E-02	4.47E-01	5.46E-04	5.46E-04	1.52E-03	1.88E-02	2.07E-02	4.47E-01
Animals	Grey Wolf	Patterson Lake North-West Basin	6.54E-04	6.54E-04	1.63E-03	2.09E-02	2.08E-02	4.48E-01	6.54E-04	6.54E-04	1.63E-03	2.09E-02	2.08E-02	4.48E-01
		Lloyd Lake	5.48E-04	5.48E-04	1.52E-03	1.89E-02	2.07E-02	4.47E-01	5.48E-04	5.48E-04	1.52E-03	1.89E-02	2.07E-02	4.47E-01
		Reference (Broach Lake)	3.02E-04	3.02E-04	3.86E-03	1.50E-02	2.57E-02	1.20E+00	3.02E-04	3.02E-04	3.86E-03	1.50E-02	2.57E-02	1.20E+00
		Patterson Lake North Arm - West Basin	4.54E-03	4.56E-03	4.28E-03	1.98E-02	2.67E-02	1.25E+00	4.55E-03	4.56E-03	4.28E-03	1.98E-02	2.67E-02	1.25E+00
	Mink	Patterson Lake South Arm	1.47E-03	1.48E-03	4.02E-03	1.64E-02	2.57E-02	1.20E+00	1.47E-03	1.48E-03	4.02E-03	1.64E-02	2.57E-02	1.20E+00
		Beet Lake	7.38E-04	7.40E-04	3.95E-03	1.56E-02	2.57E-02	1.20E+00	7.39E-04	7.40E-04	3.95E-03	1.56E-02	2.57E-02	1.20E+00
		Lloyd Lake	3.42E-04	3.42E-04	3.87E-03	1.50E-02	2.57E-02	1.20E+00	3.42E-04	3.42E-04	3.87E-03	1.50E-02	2.57E-02	1.20E+00
		Reference (Broach Lake)	1.64E-02	1.64E-02	7.27E-03	1.78E-01	1.06E-01	3.65E-01	1.64E-02	1.64E-02	7.27E-03	1.78E-01	1.06E-01	3.65E-01
		Patterson Lake North Arm - West Basin	2.88E-02	2.88E-02	8.24E-03	2.53E-01	1.10E-01	3.73E-01	2.88E-02	2.88E-02	8.24E-03	2.53E-01	1.10E-01	3.73E-01
	Moose	Patterson Lake South Arm	2.86E-02	2.86E-02	7.98E-03	2.51E-01	1.10E-01	3.72E-01	2.86E-02	2.86E-02	7.98E-03	2.51E-01	1.10E-01	3.72E-01
		Beet Lake	1.76E-02	1.76E-02	7.44E-03	1.86E-01	1.06E-01	3.65E-01	1.76E-02	1.76E-02	7.44E-03	1.86E-01	1.06E-01	3.65E-01
		Lloyd Lake	1.65E-02	1.65E-02	7.28E-03	1.79E-01	1.06E-01	3.65E-01	1.65E-02	1.65E-02	7.28E-03	1.79E-01	1.06E-01	3.65E-01
		Reference (Broach Lake)	2.90E-02	2.90E-02	1.99E+00	9.96E-02	3.34E+00	3.65E-03	2.90E-02	2.90E-02	1.99E+00	9.96E-02	3.34E+00	3.65E-03
	Moose Organs	Patterson Lake North Arm - West Basin	5.09E-02	5.09E-02	2.26E+00	1.42E-01	3.47E+00	3.73E-03	5.09E-02	5.09E-02	2.26E+00	1.42E-01	3.47E+00	3.73E-03
		Patterson Lake South Arm	5.05E-02	5.06E-02	2.18E+00	1.40E-01	3.46E+00	3.72E-03	5.05E-02	5.06E-02	2.18E+00	1.40E-01	3.46E+00	3.72E-03

Table C.11: Maximum Concentrations of Radiological COPCs in Biota for the Application Case and Upper Bound Scenario - Far-Future

Beet Lake 3.11E-02 2.91E-02 2.91E-02 2.91E-02 2.91E-02 2.91E-02 2.91E-02 3.34E+00 3.65E-03 3.11E-02 2.91E-02 3.12E-02	Idium- Lead- 226 210 04E-01 3.34E+00 98E-02 3.34E+00 20E-02 1.63E-01 00E-02 1.70E-01 86E-02 1.63E-01 48E-02 1.63E-01 23E-02 1.63E-01 29E-02 7.58E-03 29E-02 7.59E-03	Polonium- 210 3.65E-03 3.65E-03 2.41E+00 2.53E+00 2.42E+00 2.42E+00 2.41E+00 2.12E-02
Beet Lake	226 210 04E-01 3.34E+00 98E-02 3.34E+00 20E-02 1.63E-01 00E-02 1.63E-01 48E-02 1.63E-01 23E-02 1.63E-01 29E-02 7.58E-03 29E-02 7.59E-03	210 3.65E-03 3.65E-03 2.41E+00 2.53E+00 2.42E+00 2.42E+00 2.41E+00 2.12E-02
Beet Lake 3.11E-02 3.11E-02 2.04E+00 1.04E-01 3.34E+00 3.65E-03 3.11E-02 3.11E-02 3.11E-02 3.0EE-04 9.98E-04	226 210 04E-01 3.34E+00 98E-02 3.34E+00 20E-02 1.63E-01 00E-02 1.63E-01 48E-02 1.63E-01 23E-02 1.63E-01 29E-02 7.58E-03 29E-02 7.59E-03	210 3.65E-03 3.65E-03 2.41E+00 2.53E+00 2.42E+00 2.42E+00 2.41E+00 2.12E-02
Lloyd Lake 2.91E-02 2.91E-02 1.99E+00 9.98E-02 3.34E+00 3.65E-03 2.91E-02 2.91E-02 1.99E+00 9.98E-02 1.99E+00 9.98E-02 1.63E-01 2.41E+00 9.58E-04 9.58E-04 1.29E-02 6.20E-02 1.63E-01 2.41E+00 9.58E-04 9.58E-04 1.29E-02 6.20E-02 1.63E-01 2.41E+00 9.58E-04 9.58E-04 1.29E-02 6.20E-02 1.38E-02 1.38E-03 1.38E-03 1.38E-03 1.38E-03 1.38E-03 1.38E-0	98E-02 3.34E+00 20E-02 1.63E-01 00E-02 1.70E-01 86E-02 1.63E-01 48E-02 1.63E-01 23E-02 1.63E-01 29E-02 7.58E-03 29E-02 7.59E-03	3.65E-03 2.41E+00 2.53E+00 2.42E+00 2.42E+00 2.41E+00 2.12E-02
Reference (Broach Lake) Patterson Lake North Arm - West Basin Patterson Lake South Arm Patterson Lake South Arm Patterson Lake South Arm Patterson Lake Patterson Lake South Arm Patterson Lake Patterson Lake South Arm Patterson Lake South Arm Patterson Lake	20E-02 1.63E-01 00E-02 1.70E-01 86E-02 1.63E-01 48E-02 1.63E-01 23E-02 1.63E-01 29E-02 7.58E-03 29E-02 7.59E-03	2.41E+00 2.53E+00 2.42E+00 2.42E+00 2.41E+00 2.12E-02
Muskrat Patterson Lake North Arm - West Basin Patterson Lake South Arm 3.10E-02 3.11E-02 1.38E-02 7.99E-02 1.70E-01 2.53E+00 3.10E-02 3.12E-02 1.38E-02 8.0 Beet Lake Lloyd Lake 4.05E-03 4.06E-03 1.32E-02 6.48E-02 1.63E-01 2.42E+00 9.28E-03 9.31E-03 1.35E-02 6.8 Reference (Broach Lake) 1.24E-03 1.24E-03 1.01E-03 7.96E-04 2.29E-02 7.59E-03 2.12E-02 1.01E-03 1.01E-03 7.96E-04 2.29E-02 7.59E-03 2.13E-02 1.43E-03 1.05E-03 3.2 Red Fox Patterson Lake South Arm 1.08E-03 1.08E-03 8.38E-04 2.45E-02 7.58E-03 2.12E-02 1.08E-03 1.05E-03 3.2	00E-02 1.70E-01 86E-02 1.63E-01 48E-02 1.63E-01 23E-02 1.63E-01 29E-02 7.58E-03 29E-02 7.59E-03	2.53E+00 2.42E+00 2.42E+00 2.41E+00 2.12E-02
Muskrat Patterson Lake South Arm 9.27E-03 9.31E-03 1.35E-02 6.86E-02 1.63E-01 2.42E+00 9.28E-03 9.31E-03 1.35E-02 6.88E-02 Beet Lake 4.05E-03 4.06E-03 1.32E-02 6.48E-02 1.63E-01 2.42E+00 4.05E-03 4.06E-03 1.32E-02 6.48E-02 Lloyd Lake 1.24E-03 1.24E-03 1.30E-02 6.23E-02 1.63E-01 2.41E+00 1.24E-03 1.24E-03 1.30E-02 6.2 Reference (Broach Lake) 1.01E-03 1.01E-03 7.96E-04 2.29E-02 7.58E-03 2.12E-02 1.01E-03 1.01E-03 7.96E-04 2.2 Patterson Lake North Arm - West Basin 1.43E-03 1.08E-03 8.38E-04 2.45E-02 7.58E-03 2.12E-02 1.08E-03 1.08E-03 8.38E-04 2.4	86E-02 1.63E-01 48E-02 1.63E-01 23E-02 1.63E-01 29E-02 7.58E-03 29E-02 7.59E-03	2.42E+00 2.42E+00 2.41E+00 2.12E-02
Beet Lake 4.05E-03 4.06E-03 1.32E-02 6.48E-02 1.63E-01 2.42E+00 4.05E-03 4.06E-03 1.32E-02 6.48E-02 Lloyd Lake 1.24E-03 1.24E-03 1.30E-02 6.23E-02 1.63E-01 2.41E+00 1.24E-03 1.24E-03 1.30E-02 6.2 Reference (Broach Lake) 1.01E-03 1.01E-03 1.05E-03 3.29E-02 7.58E-03 2.12E-02 1.01E-03 1.05E-03 3.2 Patterson Lake North Arm - West Basin 1.08E-03 1.08E-03 8.38E-04 2.45E-02 7.58E-03 2.12E-02 1.08E-03 1.08E-03 8.38E-04 2.4	48E-02 1.63E-01 23E-02 1.63E-01 29E-02 7.58E-03 29E-02 7.59E-03	2.42E+00 2.41E+00 2.12E-02
Reference (Broach Lake)1.24E-031.24E-031.30E-026.23E-021.63E-012.41E+001.24E-031.24E-031.30E-026.23E-02Reference (Broach Lake)1.01E-031.01E-031.01E-037.96E-042.29E-027.58E-032.12E-021.01E-031.01E-037.96E-042.2Patterson Lake North Arm1.08E-031.08E-031.08E-038.38E-042.45E-027.58E-032.12E-021.08E-031.08E-038.38E-042.4	23E-02 1.63E-01 29E-02 7.58E-03 29E-02 7.59E-03	2.41E+00 2.12E-02
Reference (Broach Lake) Patterson Lake North Arm - West Basin Red Fox Patterson Lake South Arm 1.01E-03 1.01E-03 1.01E-03 1.01E-03 1.01E-03 1.05E-03 29E-02 7.58E-03 29E-02 7.59E-03	2.12E-02	
Red Fox Patterson Lake North Arm - West Basin 1.43E-03 1.43E-03 1.05E-03 3.29E-02 7.59E-03 2.13E-02 1.43E-03 1.43E-03 1.05E-03 3.2 Red Fox Patterson Lake South Arm 1.08E-03 1.08E-03 8.38E-04 2.45E-02 7.58E-03 2.12E-02 1.08E-03 1.08E-03 8.38E-04 2.4	29E-02 7.59E-03	
Red Fox Patterson Lake South Arm 1.08E-03 1.08E-03 8.38E-04 2.45E-02 7.58E-03 2.12E-02 1.08E-03 1.08E-03 8.38E-04 2.4	•	
	IEE 00 3 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	2.13E-02
Beet Lake 1.05F-03 1.05F-03 8.21F-04 2.38F-02 7.58F-03 2.12F-02 1.05F-03 1.05F-03 8.21F-04 2.3	45E-02 7.58E-03	2.12E-02
1.03E 03 1.03E 03 0.21E 04 2.30E 02 2.12E 02 1.03E 03 0.21E 04 2.3	38E-02 7.58E-03	2.12E-02
Lloyd Lake 1.01E-03 1.01E-03 7.97E-04 2.29E-02 7.58E-03 2.12E-02 1.01E-03 1.01E-03 7.97E-04 2.2	29E-02 7.58E-03	2.12E-02
Reference (Broach Lake) 1.69E-02 1.69E-02 2.73E-03 1.67E-01 2.82E-02 2.03E-01 1.69E-02 1.69E-02 2.73E-03 1.6	67E-01 2.82E-02	2.03E-01
Patterson Lake North Arm - West Basin 2.30E-02 2.30E-02 3.61E-03 2.40E-01 2.82E-02 2.03E-01 2.30E-02 2.30E-02 3.61E-03 2.4	40E-01 2.82E-02	2.03E-01
Snowshoe Hare Patterson Lake South Arm 1.79E-02 1.79E-02 2.87E-03 1.79E-01 2.82E-02 2.03E-01 1.79E-02 1.79E-03 1.7	79E-01 2.82E-02	2.03E-01
Beet Lake 1.75E-02 1.75E-02 2.81E-03 1.74E-01 2.82E-02 2.03E-01 1.75E-02 1.75E-02 2.81E-03 1.7	74E-01 2.82E-02	2.03E-01
Lloyd Lake 1.70E-02 1.70E-02 2.73E-03 1.67E-01 2.82E-02 2.03E-01 1.70E-02 1.70E-02 2.73E-03 1.6	67E-01 2.82E-02	2.03E-01
Reference (Broach Lake) 6.15E-03 6.15E-03 8.06E-04 5.98E-02 9.13E-03 7.20E-02 6.15E-03 6.15E-03 8.06E-04 5.9	98E-02 9.13E-03	7.20E-02
Patterson Lake North Arm - West Basin 8.37E-03 8.37E-03 8.62E-02 9.13E-03 7.21E-02 8.37E-03 8.37E-03 1.06E-03 8.6	62E-02 9.13E-03	7.21E-02
Southern Red- Backed Vole Patterson Lake South Arm 6.52E-03 6.52E-03 8.48E-04 6.41E-02 9.13E-03 7.21E-02 6.52E-03 6.52E-03 8.48E-04 6.4	41E-02 9.13E-03	7.21E-02
Beet Lake 6.36E-03 6.36E-03 8.30E-04 6.23E-02 9.13E-03 7.21E-02 6.36E-03 6.36E-03 8.30E-04 6.2	23E-02 9.13E-03	7.21E-02
Lloyd Lake 6.16E-03 6.16E-03 8.06E-04 5.99E-02 9.13E-03 7.20E-02 6.16E-03 6.16E-03 8.06E-04 5.9	99E-02 9.13E-03	7.20E-02
Reference (Broach Lake) 7.48E-03 7.48E-03 1.16E-01 9.35E-01 4.81E+00 7.48E-03 7.48E-03 1.1	16E-01 9.35E-01	4.81E+00
Woodland Patterson Lake North Arm - West Basin 1.71E-02 1.71E-02 9.61E-03 1.38E-01 9.41E-01 4.83E+00 1.71E-02 1.71E-02 9.61E-03 1.3	38E-01 9.41E-01	4.83E+00
Caribou Beet Lake 8.26E-03 8.27E-03 8.98E-03 1.18E-01 9.39E-01 4.83E+00 8.26E-03 8.27E-03 1.1	18E-01 9.39E-01	4.83E+00
Lloyd Lake 6.16E-03 6.16E-03 8.06E-04 5.99E-02 9.13E-03 7.20E-02 6.16E-03 6.16E-03 8.06E-04 5.9	99E-02 9.13E-03	7.20E-02
Reference (Broach Lake) 4.04E-01 4.04E-01 2.33E-03 3.80E-02 3.34E-02 1.30E+00 4.04E-01 4.04E-01 2.33E-03 3.8	3.34E-02	1.30E+00
Patterson Lake North Arm - West Basin 5.48E-01 5.48E-01 3.09E-03 5.47E-02 3.34E-02 1.30E+00 5.48E-01 5.48E-01 3.09E-03 5.4	47E-02 3.34E-02	1.30E+00
Grouse Patterson Lake South Arm 4.28E-01 4.28E-01 2.46E-03 4.07E-02 3.34E-02 1.30E+00 4.28E-01 4.28E-01 2.46E-03 4.0	07E-02 3.34E-02	1.30E+00
Beet Lake 4.18E-01 4.18E-01 2.41E-03 3.96E-02 3.34E-02 1.30E+00 4.18E-01 4.18E-01 2.41E-03 3.9	96E-02 3.34E-02	1.30E+00
Lloyd Lake 4.04E-01 4.04E-01 2.34E-03 3.80E-02 3.34E-02 1.30E+00 4.04E-01 4.04E-01 2.34E-03 3.8	3.34E-02	1.30E+00
Reference (Broach Lake) 5.66E-02 5.66E-02 3.38E-04 5.34E-03 4.74E-03 1.83E-01 5.66E-02 5.66E-02 3.38E-04 5.3	34E-03 4.74E-03	1.83E-01
Patterson Lake North Arm - West Basin 7.74E-02 7.74E-02 4.45E-04 7.69E-03 4.74E-03 1.84E-01 7.74E-02 7.74E-02 4.45E-04 7.6	69E-03 4.74E-03	1.84E-01
Canada Goose Patterson Lake South Arm 6.01E-02 6.01E-02 3.56E-04 5.72E-03 4.74E-03 1.83E-01 6.01E-02 6.01E-02 3.56E-04 5.7	72E-03 4.74E-03	1.83E-01
Beet Lake 5.86E-02 5.86E-02 3.48E-04 5.56E-03 4.74E-03 1.83E-01 5.86E-02 5.86E-02 3.48E-04 5.5	56E-03 4.74E-03	1.83E-01
Lloyd Lake 5.67E-02 5.67E-02 3.39E-04 5.35E-03 4.74E-03 1.83E-01 5.67E-02 5.67E-02 3.39E-04 5.3	35E-03 4.74E-03	1.83E-01
	04E-02 8.48E-02	4.23E+00
	21E-02 8.88E-02	4.42E+00
	47E-02 8.51E-02	4.24E+00
	22E-02 8.49E-02	4.23E+00

Table C.11: Maximum Concentrations of Radiological COPCs in Biota for the Application Case and Upper Bound Scenario - Far-Future

			Maximum Concentration Far Future (Bq/kg fw)												
	Biota	Location	Application Case						Upper Bound Scenario						
			Uranium- 238	Uranium- 234	Thorium- 230	Radium- 226	Lead- 210	Polonium- 210	Uranium- 238	Uranium- 234	Thorium- 230	Radium- 226	Lead- 210	Polonium- 210	
		Lloyd Lake	3.04E-04	3.05E-04	1.83E-02	4.05E-02	8.48E-02	4.23E+00	3.04E-04	3.05E-04	1.83E-02	4.05E-02	8.48E-02	4.23E+00	
		Reference (Broach Lake)	2.29E-03	2.29E-03	1.55E-03	1.81E-03	1.48E-02	4.07E+00	2.29E-03	2.29E-03	1.55E-03	1.81E-03	1.48E-02	4.07E+00	
		Patterson Lake North Arm - West Basin	7.43E-02	7.45E-02	1.66E-03	2.34E-03	1.55E-02	4.26E+00	7.43E-02	7.46E-02	1.66E-03	2.34E-03	1.55E-02	4.26E+00	
	Loon	Patterson Lake South Arm	2.22E-02	2.23E-02	1.62E-03	2.01E-03	1.49E-02	4.08E+00	2.22E-02	2.23E-02	1.62E-03	2.01E-03	1.49E-02	4.08E+00	
		Beet Lake	9.70E-03	9.73E-03	1.59E-03	1.90E-03	1.48E-02	4.07E+00	9.70E-03	9.73E-03	1.59E-03	1.90E-03	1.48E-02	4.07E+00	
		Lloyd Lake	2.96E-03	2.97E-03	1.55E-03	1.82E-03	1.48E-02	4.07E+00	2.96E-03	2.97E-03	1.55E-03	1.82E-03	1.48E-02	4.07E+00	
		Reference (Broach Lake)	2.41E-02	2.41E-02	1.51E-02	1.96E-02	1.94E-01	3.49E+01	2.41E-02	2.41E-02	1.51E-02	1.96E-02	1.94E-01	3.49E+01	
		Patterson Lake North Arm - West Basin	7.80E-01	7.83E-01	1.61E-02	2.53E-02	2.04E-01	3.65E+01	7.81E-01	7.84E-01	1.61E-02	2.53E-02	2.04E-01	3.65E+01	
	Mallard	Patterson Lake South Arm	2.33E-01	2.34E-01	1.57E-02	2.17E-02	1.95E-01	3.50E+01	2.34E-01	2.34E-01	1.57E-02	2.17E-02	1.95E-01	3.50E+01	
		Beet Lake	1.02E-01	1.02E-01	1.54E-02	2.05E-02	1.95E-01	3.49E+01	1.02E-01	1.02E-01	1.54E-02	2.05E-02	1.95E-01	3.49E+01	
		Lloyd Lake	3.11E-02	3.12E-02	1.51E-02	1.97E-02	1.95E-01	3.49E+01	3.12E-02	3.12E-02	1.51E-02	1.97E-02	1.95E-01	3.49E+01	
		Reference (Broach Lake)	1.74E-01	1.74E-01	4.20E-02	5.45E-02	3.34E-01	9.54E+01	1.74E-01	1.74E-01	4.20E-02	5.45E-02	3.34E-01	9.54E+01	
		Patterson Lake North Arm - West Basin	8.67E-01	8.69E-01	4.60E-02	7.30E-02	3.47E-01	9.98E+01	8.67E-01	8.70E-01	4.60E-02	7.30E-02	3.47E-01	9.98E+01	
Rus	ısty Blackbird	Patterson Lake South Arm	3.61E-01	3.61E-01	4.38E-02	5.94E-02	3.35E-01	9.57E+01	3.61E-01	3.62E-01	4.38E-02	5.94E-02	3.35E-01	9.57E+01	
		Beet Lake	2.44E-01	2.45E-01	4.29E-02	5.68E-02	3.34E-01	9.54E+01	2.44E-01	2.45E-01	4.29E-02	5.68E-02	3.34E-01	9.54E+01	
		Lloyd Lake	1.80E-01	1.80E-01	4.20E-02	5.46E-02	3.34E-01	9.54E+01	1.80E-01	1.80E-01	4.20E-02	5.46E-02	3.34E-01	9.54E+01	

Table C.12: Maximum Concentrations of Radiological COPCs in Biota for the RFD Scenario - Project Lifespan and Far-Future

			Maximum Concentration RFD Case (Bq/kg fw)												
	Biota	Location			Project	Lifespan			Far Future						
			Uranium- 238	Uranium- 234	Thorium- 230	Radium- 226	Lead- 210	Polonium- 210	Uranium- 238	Uranium- 234	Thorium- 230	Radium- 226	Lead- 210	Polonium- 210	
		Reference (Broach Lake)	1.79E-02	1.79E-02	2.11E+00	9.03E-01	9.62E+00	1.51E+00	1.79E-02	1.79E-02	2.11E+00	9.03E-01	9.62E+00	1.51E+00	
		Patterson Lake North Arm - West Basin	1.96E-01	1.96E-01	1.20E+01	1.57E+00	1.47E+01	2.29E+00	5.81E-01	5.83E-01	2.25E+00	1.16E+00	1.01E+01	1.58E+00	
	Macrophytes	Patterson Lake South Arm	2.92E-01	2.93E-01	1.28E+01	1.84E+00	1.50E+01	2.31E+00	1.74E-01	1.74E-01	2.20E+00	9.99E-01	9.65E+00	1.51E+00	
		Beet Lake	1.13E-01	1.13E-01	6.90E+00	1.26E+00	1.04E+01	1.63E+00	7.58E-02	7.60E-02	2.16E+00	9.44E-01	9.63E+00	1.51E+00	
Aquatic Plants		Lloyd Lake	2.68E-02	2.68E-02	2.60E+00	9.38E-01	9.67E+00	1.51E+00	2.32E-02	2.32E-02	2.11E+00	9.07E-01	9.63E+00	1.51E+00	
Aquatic Plants		Reference (Broach Lake)	1.79E-02	1.79E-02	2.11E+00	9.03E-01	9.62E+00	1.51E+00	1.79E-02	1.79E-02	2.11E+00	9.03E-01	9.62E+00	1.51E+00	
		Patterson Lake North Arm - West Basin	1.96E-01	1.96E-01	1.20E+01	1.57E+00	1.47E+01	2.29E+00	5.81E-01	5.83E-01	2.25E+00	1.16E+00	1.01E+01	1.58E+00	
	Phytoplankton	Patterson Lake South Arm	2.92E-01	2.93E-01	1.28E+01	1.84E+00	1.50E+01	2.31E+00	1.74E-01	1.74E-01	2.20E+00	9.99E-01	9.65E+00	1.51E+00	
		Beet Lake	1.13E-01	1.13E-01	6.90E+00	1.26E+00	1.04E+01	1.63E+00	7.58E-02	7.60E-02	2.16E+00	9.44E-01	9.63E+00	1.51E+00	
		Lloyd Lake	2.68E-02	2.68E-02	2.60E+00	9.38E-01	9.67E+00	1.51E+00	2.32E-02	2.32E-02	2.11E+00	9.07E-01	9.63E+00	1.51E+00	
		Reference (Broach Lake)	3.42E-02	3.42E-02	4.52E+00	1.35E+00	6.88E+00	4.80E+01	3.42E-02	3.42E-02	4.52E+00	1.35E+00	6.88E+00	4.80E+01	
	Danthia	Patterson Lake North Arm - West Basin	3.08E-01	3.09E-01	2.51E+01	2.16E+00	9.42E+00	6.58E+01	1.11E+00	1.11E+00	4.83E+00	1.74E+00	7.20E+00	5.03E+01	
	Benthic Invertebrates	Patterson Lake South Arm	1.92E-01	1.92E-01	1.94E+01	1.84E+00	8.44E+00	5.89E+01	3.31E-01	3.32E-01	4.71E+00	1.49E+00	6.90E+00	4.82E+01	
	lilvertebrates	Beet Lake	9.24E-02	9.26E-02	1.24E+01	1.56E+00	7.11E+00	4.96E+01	1.45E-01	1.45E-01	4.62E+00	1.41E+00	6.89E+00	4.81E+01	
		Lloyd Lake	3.95E-02	3.96E-02	5.33E+00	1.37E+00	6.90E+00	4.81E+01	4.42E-02	4.42E-02	4.53E+00	1.35E+00	6.88E+00	4.80E+01	
	Zooplankton	Reference (Broach Lake)	3.56E-02	3.56E-02	4.54E+00	1.38E+00	1.17E+01	8.13E+01	3.56E-02	3.56E-02	4.54E+00	1.38E+00	1.17E+01	8.13E+01	
		Patterson Lake North Arm - West Basin	3.88E-01	3.89E-01	2.58E+01	2.41E+00	1.78E+01	1.23E+02	1.15E+00	1.16E+00	4.86E+00	1.79E+00	1.22E+01	8.51E+01	
		Patterson Lake South Arm	5.79E-01	5.81E-01	2.75E+01	2.83E+00	1.81E+01	1.25E+02	3.44E-01	3.46E-01	4.74E+00	1.53E+00	1.17E+01	8.16E+01	
		Beet Lake	1.08E-01	1.08E-01	8.79E+00	1.60E+00	1.19E+01	8.30E+01	1.50E-01	1.51E-01	4.65E+00	1.45E+00	1.17E+01	8.14E+01	
Aquatic Animals		Lloyd Lake	5.31E-02	5.32E-02	5.59E+00	1.44E+00	1.17E+01	8.17E+01	4.60E-02	4.60E-02	4.56E+00	1.39E+00	1.17E+01	8.13E+01	
Aquatic Aillinais		Reference (Broach Lake)	7.12E-03	7.12E-03	5.45E-02	6.65E-02	3.04E-01	7.73E-01	7.12E-03	7.12E-03	5.45E-02	6.65E-02	3.04E-01	7.73E-01	
		Patterson Lake North Arm - West Basin	7.76E-02	7.79E-02	3.10E-01	1.16E-01	4.64E-01	1.17E+00	2.30E-01	2.31E-01	5.83E-02	8.57E-02	3.18E-01	8.09E-01	
	Northern Pike	Patterson Lake South Arm	1.16E-01	1.16E-01	3.30E-01	1.36E-01	4.72E-01	1.18E+00	6.89E-02	6.91E-02	5.69E-02	7.35E-02	3.05E-01	7.75E-01	
		Beet Lake	4.49E-02	4.50E-02	1.78E-01	9.31E-02	3.28E-01	8.34E-01	3.01E-02	3.02E-02	5.58E-02	6.95E-02	3.04E-01	7.73E-01	
		Lloyd Lake	1.06E-02	1.06E-02	6.71E-02	6.90E-02	3.05E-01	7.76E-01	9.20E-03	9.20E-03	5.47E-02	6.67E-02	3.04E-01	7.73E-01	
		Reference (Broach Lake)	1.76E-03	1.76E-03			1.00E-01	6.94E-01	1.76E-03	1.76E-03	5.45E-02	•	1.00E-01	6.94E-01	
		Patterson Lake North Arm - West Basin	1.83E-02	1.84E-02	3.08E-01	1.12E-01	1.49E-01	1.03E+00	5.70E-02	5.73E-02	5.82E-02	8.51E-02	1.05E-01	7.26E-01	
	Whitefish	Patterson Lake South Arm	2.35E-02	2.36E-02	2.87E-01	1.22E-01	1.50E-01	1.03E+00	1.71E-02	1.71E-02	5.68E-02	7.30E-02	1.00E-01	6.96E-01	
		Beet Lake	9.28E-03	9.31E-03	1.60E-01	8.72E-02	1.07E-01	7.43E-01	7.45E-03	7.47E-03	5.57E-02	6.90E-02	1.00E-01	6.94E-01	
		Lloyd Lake	2.46E-03	2.46E-03	6.53E-02	6.80E-02	1.00E-01	6.97E-01	2.28E-03	2.28E-03	5.46E-02	6.63E-02	1.00E-01	6.94E-01	
		Reference (Broach Lake)	4.95E-01	4.95E-01	6.24E-02	1.08E+00	3.20E-01	4.33E-01	4.95E-01	4.95E-01	6.24E-02	1.08E+00	3.20E-01	4.33E-01	
		Patterson Lake North Arm - West Basin	2.27E+00	2.27E+00	1.07E-01	2.61E+00	1.20E+00	1.47E+00	6.74E-01	6.74E-01	8.29E-02	1.56E+00	3.20E-01	4.33E-01	
	Blueberries	Patterson Lake South Arm	8.37E-01	8.37E-01	7.10E-02	1.37E+00	4.90E-01	6.33E-01	5.29E-01	5.29E-01	6.64E-02	1.17E+00	3.20E-01	4.33E-01	
		Beet Lake	6.82E-01	6.82E-01	6.71E-02	1.24E+00	4.13E-01	5.43E-01	5.14E-01	5.14E-01	6.46E-02	1.13E+00	3.20E-01	4.33E-01	
Terrestrial Plants		Lloyd Lake	5.04E-01	5.04E-01	6.26E-02	1.08E+00	3.24E-01	4.38E-01	4.96E-01	4.96E-01	6.25E-02	1.08E+00	3.20E-01	4.33E-01	
		Reference (Broach Lake)	9.90E-01	9.90E-01	1.25E-01	2.15E+00	6.40E-01	8.67E-01	9.90E-01	9.90E-01	1.25E-01	2.15E+00	6.40E-01	8.67E-01	
		Patterson Lake North Arm - West Basin	4.40E+00	4.40E+00	2.77E+00	6.14E+00	3.33E+00	3.88E+00	1.35E+00	1.35E+00	1.66E-01	3.12E+00	6.40E-01	8.67E-01	
	Browse	Patterson Lake South Arm	1.65E+00	1.65E+00	6.34E-01	2.92E+00	1.16E+00	1.45E+00	1.06E+00	1.06E+00	1.33E-01	2.34E+00	6.40E-01	8.67E-01	
		Beet Lake	1.35E+00	1.35E+00	4.03E-01	2.57E+00	9.23E-01	1.18E+00	1.03E+00	1.03E+00	1.29E-01	2.25E+00	6.40E-01	8.67E-01	
		Lloyd Lake	1.01E+00	1.01E+00	1.38E-01	2.17E+00	6.53E-01	8.82E-01	9.92E-01	9.92E-01	1.25E-01	2.16E+00	6.40E-01	8.67E-01	

Table C.12: Maximum Concentrations of Radiological COPCs in Biota for the RFD Scenario - Project Lifespan and Far-Future

			Maximum Concentration RFD Case (Bq/kg fw)											
	Biota	Location	Project Lifespan Far Future											
			Uranium-	Uranium-	Thorium-	•	Lead-	Polonium-	Uranium-	Uranium-	1	1	Lead-	Polonium-
			238	234	230	226	210	210	238	234	230	226	210	210
		Reference (Broach Lake)	9.90E-01	9.90E-01	1.25E-01	2.15E+00	6.40E-01	8.67E-01	9.90E-01	9.90E-01	1.25E-01	2.15E+00	6.40E-01	8.67E-01
		Patterson Lake North Arm - West Basin	4.40E+00	4.40E+00	2.47E+01	2.80E+01	2.52E+01	2.24E+01	1.35E+00	1.35E+00	1.66E-01	3.12E+00	6.40E-01	8.67E-01
	Labrador Tea	Patterson Lake South Arm	1.65E+00	1.65E+00	4.85E+00	7.14E+00	5.36E+00	5.02E+00	1.06E+00	1.06E+00	1.33E-01	2.34E+00	6.40E-01	8.67E-01
		Beet Lake	1.35E+00	1.35E+00	2.71E+00	4.88E+00	3.22E+00	3.14E+00	1.03E+00	1.03E+00	1.29E-01	2.25E+00	6.40E-01	8.67E-01
		Lloyd Lake	1.01E+00	1.01E+00	2.48E-01	2.28E+00	7.63E-01	9.75E-01	9.92E-01	9.92E-01	1.25E-01	2.16E+00	6.40E-01	8.67E-01
		Reference (Broach Lake)	2.08E-01	2.08E-01	6.77E-01	1.33E+00	1.91E+02	1.45E+02	2.08E-01	2.08E-01	6.77E-01	1.33E+00	1.91E+02	1.45E+02
		Patterson Lake North Arm - West Basin	2.08E-01	2.08E-01	1.41E+02	1.42E+02	3.24E+02	1.72E+02	2.08E-01	2.08E-01	6.77E-01	1.33E+00	1.91E+02	1.45E+02
	Lichen	Patterson Lake South Arm	2.08E-01	2.08E-01	2.77E+01	2.83E+01	2.17E+02	1.50E+02	2.08E-01	2.08E-01	6.77E-01	1.33E+00	1.91E+02	1.45E+02
		Beet Lake	2.08E-01	2.08E-01	1.54E+01	1.61E+01	2.05E+02	1.48E+02	2.08E-01	2.08E-01	6.77E-01	1.33E+00	1.91E+02	1.45E+02
		Lloyd Lake	2.08E-01	2.08E-01	1.38E+00	2.04E+00	1.92E+02	1.45E+02	2.08E-01	2.08E-01	6.77E-01	1.33E+00	1.91E+02	1.45E+02
		Reference (Broach Lake)	2.70E-02	2.70E-02	1.77E-01	1.61E+00	1.14E+00	3.12E-02	2.70E-02	2.70E-02	1.77E-01	1.61E+00	1.14E+00	3.12E-02
	Townstrial	Patterson Lake North Arm - West Basin	1.20E-01	1.20E-01	2.85E+00	5.25E+00	3.93E+00	2.46E+00	3.68E-02	3.68E-02	2.35E-01	2.34E+00	1.14E+00	3.12E-02
	Terrestrial Invertebrate	Patterson Lake South Arm	4.49E-02	4.49E-02	6.91E-01	2.31E+00	1.68E+00	4.99E-01	2.89E-02	2.89E-02	1.88E-01	1.75E+00	1.14E+00	3.12E-02
	ilivertebrate	Beet Lake	3.68E-02	3.68E-02	4.58E-01	2.00E+00	1.43E+00	2.87E-01	2.80E-02	2.80E-02	1.83E-01	1.69E+00	1.14E+00	3.12E-02
		Lloyd Lake	2.75E-02	2.75E-02	1.90E-01	1.63E+00	1.15E+00	4.34E-02	2.71E-02	2.71E-02	1.77E-01	1.62E+00	1.14E+00	3.12E-02
	Beaver	Reference (Broach Lake)	1.81E-02	1.81E-02	4.78E-03	1.84E-01	6.65E-02	2.94E-01	1.81E-02	1.81E-02	4.78E-03	1.84E-01	6.65E-02	2.94E-01
		Patterson Lake North Arm - West Basin	8.08E-02	8.09E-02	4.56E-02	5.07E-01	1.74E-01	1.02E+00	2.78E-02	2.78E-02	5.66E-03	2.65E-01	6.84E-02	2.98E-01
		Patterson Lake South Arm	3.04E-02	3.04E-02	2.30E-02	2.49E-01	1.03E-01	4.38E-01	2.02E-02	2.02E-02	5.03E-03	2.01E-01	6.67E-02	2.94E-01
		Beet Lake	2.47E-02	2.47E-02	1.35E-02	2.19E-01	7.83E-02	3.68E-01	1.91E-02	1.91E-02	4.91E-03	1.93E-01	6.66E-02	2.94E-01
		Lloyd Lake	1.84E-02	1.84E-02	5.52E-03	1.86E-01	6.71E-02	2.98E-01	1.81E-02	1.81E-02	4.79E-03	1.85E-01	6.66E-02	2.94E-01
		Reference (Broach Lake)	1.35E-02	1.35E-02	2.77E-03	1.36E-01	2.87E-02	2.39E-01	1.35E-02	1.35E-02	2.77E-03	1.36E-01	2.87E-02	2.39E-01
	Black Bear	Patterson Lake North Arm - West Basin	3.96E-02	3.96E-02	4.88E-03	2.35E-01	5.31E-02	4.49E-01	1.71E-02	1.71E-02	3.28E-03	1.69E-01	2.89E-02	2.41E-01
	Diack Deal	Beet Lake	1.62E-02	1.62E-02	4.01E-03	1.51E-01	3.60E-02	2.69E-01	1.40E-02	1.40E-02	3.70E-03	1.45E-01	3.37E-02	2.49E-01
		Lloyd Lake	1.37E-02	1.37E-02	2.89E-03	1.37E-01	2.90E-02	2.41E-01	1.35E-02	1.35E-02	2.82E-03	1.36E-01	2.88E-02	2.39E-01
Terrestrial		Reference (Broach Lake)	5.46E-04	5.46E-04	1.52E-03	1.88E-02	2.07E-02	4.47E-01	5.46E-04	5.46E-04	1.52E-03	1.88E-02	2.07E-02	4.47E-01
Animals	Grey Wolf	Patterson Lake North-West Basin	1.06E-03	1.06E-03	2.10E-03	3.27E-02	2.32E-02	4.94E-01	6.56E-04	6.56E-04	1.64E-03	2.10E-02	2.08E-02	4.48E-01
		Lloyd Lake	5.56E-04	5.56E-04	1.55E-03	1.91E-02	2.08E-02	4.47E-01	5.49E-04	5.49E-04	1.52E-03	1.89E-02	2.07E-02	4.47E-01
		Reference (Broach Lake)	3.02E-04	3.02E-04	3.86E-03	1.50E-02	2.57E-02	1.20E+00	3.02E-04	3.02E-04	3.86E-03	1.50E-02	2.57E-02	1.20E+00
		Patterson Lake North Arm - West Basin	1.98E-03	1.98E-03	1.89E-02	2.43E-02	3.46E-02	1.64E+00	4.54E-03	4.56E-03	4.28E-03	1.98E-02	2.67E-02	1.25E+00
	Mink	Patterson Lake South Arm	9.94E-04	9.97E-04	1.45E-02	1.94E-02	3.10E-02	1.47E+00	1.47E-03	1.48E-03	4.03E-03	1.64E-02	2.57E-02	1.20E+00
		Beet Lake	5.69E-04	5.70E-04	9.51E-03	1.69E-02	2.64E-02	1.23E+00	7.38E-04	7.40E-04	3.95E-03	1.56E-02	2.57E-02	1.20E+00
		Lloyd Lake	3.25E-04	3.25E-04	4.44E-03	1.51E-02	2.57E-02	1.20E+00	3.42E-04	3.42E-04	3.87E-03	1.50E-02	2.57E-02	1.20E+00
		Reference (Broach Lake)	1.64E-02	1.64E-02	7.27E-03	1.78E-01	1.06E-01	3.65E-01	1.64E-02	1.64E-02	7.27E-03	1.78E-01	1.06E-01	3.65E-01
		Patterson Lake North Arm - West Basin	7.39E-02	7.39E-02	5.77E-02	4.76E-01	2.25E-01	1.06E+00	2.89E-02	2.89E-02	8.25E-03	2.55E-01	1.10E-01	3.73E-01
	Moose	Patterson Lake South Arm	7.33E-02	7.33E-02	5.68E-02	4.73E-01	2.24E-01	1.05E+00	2.87E-02	2.87E-02	7.99E-03	2.52E-01	1.10E-01	3.72E-01
		Beet Lake	2.26E-02	2.26E-02	2.16E-02	2.12E-01	1.20E-01	4.33E-01	1.77E-02	1.77E-02	7.45E-03	1.87E-01	1.06E-01	3.65E-01
		Lloyd Lake	1.67E-02	1.67E-02	8.59E-03	1.80E-01	1.07E-01	3.68E-01	1.65E-02	1.65E-02	7.28E-03	1.79E-01	1.06E-01	3.65E-01
		Reference (Broach Lake)	2.90E-02	2.90E-02	1.99E+00	9.96E-02	3.34E+00	3.65E-03	2.90E-02	2.90E-02	1.99E+00	9.96E-02	3.34E+00	3.65E-03
	Moose Organs	Patterson Lake North Arm - West Basin	1.31E-01	1.31E-01	1.58E+01	2.66E-01	7.08E+00	1.06E-02	5.11E-02	5.11E-02	2.26E+00	1.42E-01	3.47E+00	3.73E-03
		Patterson Lake South Arm	1.30E-01	1.30E-01	1.56E+01	2.64E-01	7.05E+00	1.05E-02	5.07E-02	5.08E-02	2.19E+00	1.41E-01	3.46E+00	3.72E-03

Table C.12: Maximum Concentrations of Radiological COPCs in Biota for the RFD Scenario - Project Lifespan and Far-Future

		Maximum Concentration RFD Case (Bq/kg fw)											
Biota	Location	Project Lifespan Far Future											
		Uranium-	Uranium-	Thorium-	Radium-	Lead-	Polonium-	Uranium-	Uranium-	Thorium-	Radium-	Lead-	Polonium-
		238	234	230	226	210	210	238	234	230	226	210	210
	Beet Lake	3.99E-02	3.99E-02	5.93E+00	1.18E-01	3.78E+00	4.33E-03	3.13E-02	3.13E-02	2.04E+00	1.04E-01	3.34E+00	3.65E-03
	Lloyd Lake	2.95E-02	2.95E-02	2.35E+00	1.01E-01	3.37E+00	3.68E-03	2.91E-02	2.91E-02	1.99E+00	9.99E-02	3.34E+00	3.65E-03
	Reference (Broach Lake)	9.58E-04	9.58E-04	1.29E-02	6.20E-02	1.63E-01	2.41E+00	9.58E-04	9.58E-04	1.29E-02	6.20E-02	1.63E-01	2.41E+00
	Patterson Lake North Arm - West Basin	8.65E-03	8.68E-03	7.23E-02	1.00E-01	2.28E-01	3.30E+00	3.10E-02	3.11E-02	1.38E-02	7.99E-02	1.70E-01	2.53E+00
Muskrat	Patterson Lake South Arm	5.39E-03	5.41E-03	5.58E-02	8.57E-02	2.14E-01	2.96E+00	9.27E-03	9.31E-03	1.35E-02	6.86E-02	1.63E-01	2.42E+00
	Beet Lake	2.59E-03	2.60E-03	3.57E-02	7.18E-02	1.70E-01	2.49E+00	4.05E-03	4.06E-03	1.32E-02	6.48E-02	1.63E-01	2.42E+00
	Lloyd Lake	1.11E-03	1.11E-03	1.53E-02	6.29E-02	1.63E-01	2.42E+00	1.24E-03	1.24E-03	1.30E-02	6.23E-02	1.63E-01	2.41E+00
	Reference (Broach Lake)	1.01E-03	1.01E-03	7.96E-04	2.29E-02	7.58E-03	2.12E-02	1.01E-03	1.01E-03	7.96E-04	2.29E-02	7.58E-03	2.12E-02
	Patterson Lake North Arm - West Basin	4.60E-03	4.60E-03	4.04E-03	6.23E-02	1.90E-02	1.12E-01	1.43E-03	1.43E-03	1.05E-03	3.31E-02	7.59E-03	2.13E-02
Red Fox	Patterson Lake South Arm	1.70E-03	1.70E-03	1.50E-03	3.05E-02	9.78E-03	3.87E-02	1.09E-03	1.09E-03	8.46E-04	2.48E-02	7.58E-03	2.12E-02
	Beet Lake	1.38E-03	1.38E-03	1.18E-03	2.70E-02	8.79E-03	3.08E-02	1.05E-03	1.05E-03	8.24E-04	2.39E-02	7.58E-03	2.12E-02
	Lloyd Lake	1.02E-03	1.02E-03	8.18E-04	2.31E-02	7.64E-03	2.17E-02	1.01E-03	1.01E-03	7.98E-04	2.29E-02	7.58E-03	2.12E-02
	Reference (Broach Lake)	1.69E-02	1.69E-02	2.73E-03	1.67E-01	2.82E-02	2.03E-01	1.69E-02	1.69E-02	2.73E-03	1.67E-01	2.82E-02	2.03E-01
	Patterson Lake North Arm - West Basin	7.54E-02	7.54E-02	2.79E-02	4.59E-01	1.09E-01	8.43E-01	2.31E-02	2.31E-02	3.62E-03	2.42E-01	2.82E-02	2.03E-01
Snowshoe	Patterson Lake South Arm	2.82E-02	2.82E-02	7.65E-03	2.23E-01	4.37E-02	3.26E-01	1.81E-02	1.81E-02	2.90E-03	1.81E-01	2.82E-02	2.03E-01
Hare	Beet Lake	2.31E-02	2.31E-02	5.42E-03	1.98E-01	3.67E-02	2.71E-01	1.76E-02	1.76E-02	2.82E-03	1.75E-01	2.82E-02	2.03E-01
	Lloyd Lake	1.72E-02	1.72E-02	2.86E-03	1.68E-01	2.86E-02	2.06E-01	1.70E-02	1.70E-02	2.74E-03	1.67E-01	2.82E-02	2.03E-01
	Reference (Broach Lake)	6.15E-03	6.15E-03	8.06E-04	5.98E-02	9.13E-03	7.20E-02	6.15E-03	6.15E-03	8.06E-04	5.98E-02	9.13E-03	7.20E-02
	Patterson Lake North Arm - West Basin	2.78E-02	2.78E-02	5.90E-03	1.55E-01	3.39E-02	2.72E-01	8.41E-03	8.41E-03	1.07E-03	8.66E-02	9.13E-03	7.21E-02
Southern Red-	Patterson Lake South Arm	1.03E-02	1.03E-02	1.82E-03	7.81E-02	1.39E-02	1.10E-01	6.59E-03	6.59E-03	8.56E-04	6.50E-02	9.13E-03	7.21E-02
Backed Vole	Beet Lake	8.43E-03	8.43E-03	1.36E-03	6.98E-02	1.17E-02	9.31E-02	6.39E-03	6.39E-03	8.33E-04	6.26E-02	9.13E-03	7.21E-02
	Lloyd Lake	6.26E-03	6.26E-03	8.33E-04	6.03E-02	9.25E-03	7.30E-02	6.16E-03	6.16E-03	8.07E-04	5.99E-02	9.13E-03	7.20E-02
	Reference (Broach Lake)	7.48E-03	7.48E-03	8.89E-03	1.16E-01	9.35E-01	4.81E+00	7.48E-03	7.48E-03	8.89E-03	1.16E-01	9.35E-01	4.81E+00
Woodland	Patterson Lake North Arm - West Basin	2.22E-02	2.22E-02	1.40E-01	9.70E-01	1.29E+00	5.47E+00	1.71E-02	1.72E-02	9.63E-03	1.38E-01	9.41E-01	4.83E+00
Caribou	Beet Lake	8.92E-03	8.93E-03	2.45E-02	1.97E-01	9.72E-01	4.89E+00	8.28E-03	8.28E-03	8.99E-03	1.18E-01	9.39E-01	4.83E+00
	Lloyd Lake	6.26E-03	6.26E-03	8.33E-04	6.03E-02	9.25E-03	7.30E-02	6.16E-03	6.16E-03	8.07E-04	5.99E-02	9.13E-03	7.20E-02
	Reference (Broach Lake)	4.04E-01	4.04E-01	2.33E-03	3.80E-02	3.34E-02	1.30E+00	4.04E-01	4.04E-01	2.33E-03	3.80E-02	3.34E-02	1.30E+00
	Patterson Lake North Arm - West Basin	1.80E+00	1.80E+00	1.39E-02	9.96E-02	1.01E-01	4.90E+00	5.51E-01	5.51E-01	3.10E-03	5.50E-02	3.34E-02	1.30E+00
Grouse	Patterson Lake South Arm	6.73E-01	6.73E-01	4.59E-03	4.98E-02	4.63E-02	1.99E+00	4.32E-01	4.32E-01	2.48E-03	4.13E-02	3.34E-02	1.30E+00
	Beet Lake	5.51E-01	5.51E-01	3.57E-03	4.45E-02	4.05E-02	1.68E+00	4.19E-01	4.19E-01	2.41E-03	3.98E-02	3.34E-02	1.30E+00
	Lloyd Lake	4.11E-01	4.11E-01	2.39E-03	3.83E-02	3.37E-02	1.32E+00	4.05E-01	4.05E-01	2.34E-03	3.81E-02	3.34E-02	1.30E+00
	Reference (Broach Lake)	5.66E-02	5.66E-02	3.38E-04	5.34E-03	4.74E-03	1.83E-01	5.66E-02	5.66E-02	3.38E-04	5.34E-03	4.74E-03	1.83E-01
	Patterson Lake North Arm - West Basin	2.52E-01	2.52E-01	2.44E-03	1.45E-02	1.50E-02	7.27E-01	7.78E-02	7.78E-02	4.47E-04	7.74E-03	4.74E-03	1.84E-01
Canada Goose	Patterson Lake South Arm	9.42E-02	9.42E-02	7.61E-04	7.10E-03	6.71E-03	2.88E-01	6.07E-02	6.07E-02	3.59E-04	5.80E-03	4.74E-03	1.83E-01
	Beet Lake	7.72E-02	7.72E-02	5.69E-04	6.30E-03	5.81E-03	2.41E-01	5.88E-02	5.88E-02	3.50E-04	5.59E-03	4.74E-03	1.83E-01
	Lloyd Lake	5.76E-02	5.76E-02	3.50E-04	5.39E-03	4.79E-03	1.86E-01	5.67E-02	5.67E-02	3.39E-04	5.35E-03	4.74E-03	1.83E-01
	Reference (Broach Lake)	2.36E-04	2.36E-04	1.83E-02	4.04E-02	8.48E-02	4.23E+00	2.36E-04	2.36E-04	1.83E-02	4.04E-02	8.48E-02	4.23E+00
Little Brown	Patterson Lake North Arm - West Basin	2.12E-03	2.13E-03	1.02E-01	6.47E-02	1.16E-01	5.79E+00	7.62E-03	7.65E-03	1.96E-02	5.20E-02	8.88E-02	4.42E+00
Myotis	Patterson Lake South Arm	1.32E-03	1.33E-03	7.85E-02	5.50E-02	1.04E-01	5.18E+00	2.28E-03	2.29E-03	1.91E-02	4.47E-02	8.51E-02	4.24E+00
	Beet Lake	6.38E-04	6.39E-04	5.03E-02	4.67E-02	8.76E-02	4.37E+00	9.96E-04	9.99E-04	1.87E-02	4.22E-02	8.49E-02	4.23E+00
	Door Lake	0.55L 0-7	0.55L 0-T	J.03L 0L	1.07 L 02	0.7 02 02	1.57 - 100	J.JUL U-	J.JJL U-T	1.07 L 02	1.222 02	0.15L 0L	

Table C.12: Maximum Concentrations of Radiological COPCs in Biota for the RFD Scenario - Project Lifespan and Far-Future

			Maximum Concentration RFD Case (Bq/kg fw)											
	Biota	Location	Project Lifespan								Far F	uture		
			Uranium-	Uranium-	Thorium-	Radium-	Lead-	Polonium-	Uranium-	Uranium-	Thorium-	Radium-	Lead-	Polonium-
			238	234	230	226	210	210	238	234	230	226	210	210
		Lloyd Lake	2.72E-04	2.72E-04	2.16E-02	4.10E-02	8.50E-02	4.23E+00	3.04E-04	3.05E-04	1.83E-02	4.05E-02	8.48E-02	4.23E+00
		Reference (Broach Lake)	2.29E-03	2.29E-03	1.55E-03	1.81E-03	1.48E-02	4.07E+00	2.29E-03	2.29E-03	1.55E-03	1.81E-03	1.48E-02	4.07E+00
		Patterson Lake North Arm - West Basin	2.37E-02	2.38E-02	8.63E-03	2.97E-03	2.08E-02	5.60E+00	7.43E-02	7.45E-02	1.66E-03	2.34E-03	1.55E-02	4.26E+00
	Loon	Patterson Lake South Arm	2.79E-02	2.80E-02	6.66E-03	2.69E-03	1.94E-02	5.11E+00	2.22E-02	2.23E-02	1.62E-03	2.01E-03	1.49E-02	4.08E+00
		Beet Lake	1.11E-02	1.11E-02	4.27E-03	2.15E-03	1.55E-02	4.22E+00	9.70E-03	9.73E-03	1.59E-03	1.90E-03	1.48E-02	4.07E+00
		Lloyd Lake	3.11E-03	3.11E-03	1.83E-03	1.84E-03	1.49E-02	4.08E+00	2.96E-03	2.97E-03	1.55E-03	1.82E-03	1.48E-02	4.07E+00
	Mallard	Reference (Broach Lake)	2.41E-02	2.41E-02	1.51E-02	1.96E-02	1.95E-01	3.49E+01	2.41E-02	2.41E-02	1.51E-02	1.96E-02	1.94E-01	3.49E+01
		Patterson Lake North Arm - West Basin	2.17E-01	2.18E-01	8.39E-02	3.16E-02	2.71E-01	4.77E+01	7.80E-01	7.83E-01	1.61E-02	2.53E-02	2.04E-01	3.65E+01
		Patterson Lake South Arm	1.35E-01	1.36E-01	6.47E-02	2.68E-02	2.51E-01	4.28E+01	2.33E-01	2.34E-01	1.57E-02	2.17E-02	1.95E-01	3.50E+01
		Beet Lake	6.52E-02	6.54E-02	4.15E-02	2.27E-02	2.03E-01	3.60E+01	1.02E-01	1.02E-01	1.54E-02	2.05E-02	1.95E-01	3.49E+01
		Lloyd Lake	2.79E-02	2.79E-02	1.78E-02	1.99E-02	1.95E-01	3.49E+01	3.11E-02	3.12E-02	1.51E-02	1.97E-02	1.95E-01	3.49E+01
		Reference (Broach Lake)	1.74E-01	1.74E-01	4.20E-02	5.45E-02	3.34E-01	9.54E+01	1.74E-01	1.74E-01	4.20E-02	5.45E-02	3.34E-01	9.54E+01
		Patterson Lake North Arm - West Basin	8.81E-01	8.82E-01	2.13E-01	9.51E-02	4.59E-01	1.31E+02	8.67E-01	8.70E-01	4.60E-02	7.32E-02	3.47E-01	9.98E+01
	Rusty	Patterson Lake South Arm	3.63E-01	3.64E-01	1.64E-01	7.04E-02	4.01E-01	1.17E+02	3.62E-01	3.63E-01	4.38E-02	5.97E-02	3.35E-01	9.57E+01
	Blackbird	Beet Lake	2.63E-01	2.63E-01	1.07E-01	6.19E-02	3.45E-01	9.86E+01	2.45E-01	2.45E-01	4.29E-02	5.69E-02	3.34E-01	9.54E+01
		Lloyd Lake	1.79E-01	1.79E-01	4.87E-02	5.51E-02	3.34E-01	9.56E+01	1.80E-01	1.80E-01	4.20E-02	5.47E-02	3.34E-01	9.54E+01

Table C13: Estimated Non-carcinogen Total Risk to Human Receptors – Project Lifespan – Application Case and Upper Bound Scenario

		Project Lifespan HQs											
	СОРС	Water (internal)	Soil (internal)	Sediment (internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC				
					Base	Case							
	Cobalt	8.58E-04	9.04E-07	1.01E-06	0.00E+00	1.15E-03	7.21E-04	1.97E-02	2.25E-02				
	Copper	2.91E-05	2.95E-08	4.12E-08	0.00E+00	8.94E-04	2.69E-04	3.89E-02	4.01E-02				
	Molybdenum	1.08E-04	1.06E-07	1.62E-07	0.00E+00	1.21E-06	1.46E-04	8.23E-02	8.25E-02				
	Uranium	2.46E-03	8.62E-06	2.24E-05	0.00E+00	3.43E-03	1.92E-02	7.29E-02	9.80E-02				
				Арј	olication Case -	Total Project	Risk						
Camp Worker	Cobalt	8.63E-04	9.07E-07	1.02E-06	0.00E+00	1.25E-03	7.29E-04	2.01E-02	2.29E-02				
(Patterson Lake North	Copper	2.91E-05	3.00E-08	4.12E-08	0.00E+00	9.36E-04	2.71E-04	3.91E-02	4.03E-02				
Arm - West Basin)	Molybdenum	1.09E-04	1.08E-07	1.62E-07	0.00E+00	1.79E-06	1.52E-04	8.24E-02	8.26E-02				
	Uranium	2.46E-03	2.77E-05	2.24E-05	0.00E+00	4.71E-03	4.06E-02	8.68E-02	1.35E-01				
	Upper Bound Scenario - Total Project Risk												
	Cobalt	8.63E-04	9.07E-07	1.02E-06	0.00E+00	1.25E-03	7.29E-04	2.01E-02	2.29E-02				
	Copper	2.91E-05	3.00E-08	4.12E-08	0.00E+00	9.37E-04	2.71E-04	3.91E-02	4.03E-02				
	Molybdenum	1.09E-04	1.08E-07	1.62E-07	0.00E+00	2.97E-06	1.52E-04	8.24E-02	8.27E-02				
	Uranium	2.47E-03	2.77E-05	2.25E-05	0.00E+00	6.96E-03	4.06E-02	8.74E-02	1.37E-01				
					Base	Case							
	Cobalt	8.58E-04	9.04E-07	1.01E-06	0.00E+00	2.30E-03	1.44E-03	2.44E-02	2.90E-02				
	Copper	2.91E-05	2.95E-08	4.12E-08	0.00E+00	1.79E-03	5.38E-04	4.12E-02	4.35E-02				
	Molybdenum	1.08E-04	1.06E-07	1.62E-07	0.00E+00	2.42E-06	2.92E-04	6.98E-02	7.02E-02				
Subsistence Harvester	Uranium	2.46E-03	8.62E-06	2.24E-05	0.00E+00	6.85E-03	3.84E-02	8.20E-02	1.30E-01				
(Patterson Lake South				Арј	olication Case -	Total Project			İ				
Arm)	Cobalt	9.32E-04	9.05E-07	1.09E-06	0.00E+00	2.49E-03	1.46E-03	2.51E-02	3.00E-02				
	Copper	3.05E-05	2.96E-08	4.29E-08	0.00E+00	1.87E-03	5.41E-04	4.15E-02	4.39E-02				
	Molybdenum	1.61E-04	1.07E-07	2.32E-07	0.00E+00	3.57E-06	3.03E-04	6.99E-02	7.04E-02				
	Uranium	3.39E-03	1.12E-05	2.92E-05	0.00E+00	9.42E-03	8.12E-02	1.10E-01	2.04E-01				
				Upper	Bound Scenar	io - Total Proje	ect Risk						



Table C13: Estimated Non-carcinogen Total Risk to Human Receptors – Project Lifespan – Application Case and Upper Bound Scenario

		Project Lifespan HQs										
	СОРС	Water (internal)	Soil (internal)	Sediment (internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC			
	Cobalt	9.35E-04	9.05E-07	1.10E-06	0.00E+00	2.50E-03	1.46E-03	2.51E-02	3.00E-02			
	Copper	3.05E-05	2.96E-08	4.30E-08	0.00E+00	1.87E-03	5.41E-04	4.15E-02	4.39E-02			
	Molybdenum	2.68E-04	1.07E-07	3.79E-07	0.00E+00	5.93E-06	3.03E-04	7.01E-02	7.06E-02			
	Uranium	5.02E-03	1.12E-05	4.14E-05	0.00E+00	1.39E-02	8.12E-02	1.11E-01	2.11E-01			
					Base	Case						
	Cobalt	9.58E-04	5.91E-05	6.62E-05	0.00E+00	2.18E-03	2.84E-03	4.78E-02	5.39E-02			
	Copper	3.25E-05	1.93E-06	2.69E-06	0.00E+00	1.69E-03	1.06E-03	9.70E-02	9.98E-02			
	Molybdenum	1.47E-04	8.46E-06	1.29E-05	0.00E+00	2.79E-06	7.01E-04	2.64E-01	2.64E-01			
	Uranium	2.75E-03	5.64E-04	1.47E-03	0.00E+00	6.49E-03	7.57E-02	1.70E-01	2.57E-01			
				Ар	olication Case	- Total Project	Risk					
Subsistence Harvester	Cobalt	1.04E-03	5.91E-05	7.14E-05	0.00E+00	2.36E-03	2.86E-03	4.85E-02	5.49E-02			
One-year-old (Patterson	Copper	3.40E-05	1.93E-06	2.81E-06	0.00E+00	1.77E-03	1.07E-03	9.73E-02	1.00E-01			
Lake South Arm)	Molybdenum	2.19E-04	8.49E-06	1.85E-05	0.00E+00	4.12E-06	7.20E-04	2.64E-01	2.65E-01			
	Uranium	3.79E-03	7.34E-04	1.91E-03	0.00E+00	8.93E-03	1.40E-01	1.91E-01	3.46E-01			
		Upper Bound Scenario - Total Project Risk										
	Cobalt	1.04E-03	5.91E-05	7.16E-05	0.00E+00	2.37E-03	2.86E-03	4.85E-02	5.49E-02			
	Copper	3.41E-05	1.93E-06	2.81E-06	0.00E+00	1.78E-03	1.07E-03	9.73E-02	1.00E-01			
	Molybdenum	3.65E-04	8.49E-06	3.02E-05	0.00E+00	6.84E-06	7.20E-04	2.64E-01	2.65E-01			
	Uranium	5.61E-03	7.34E-04	2.70E-03	0.00E+00	1.32E-02	1.40E-01	1.91E-01	3.54E-01			
				ı		Case		ı	T			
	Cobalt	8.58E-04	9.04E-07	1.01E-06	0.00E+00	2.30E-03	1.44E-03	2.44E-02	2.90E-02			
Subsistence Harvester	Copper	2.91E-05	2.95E-08	4.12E-08	0.00E+00	1.79E-03	5.38E-04	4.12E-02	4.35E-02			
(Beet Lake)	Molybdenum	1.08E-04	1.06E-07	1.62E-07	0.00E+00	2.42E-06	2.92E-04	6.98E-02	7.02E-02			
	Uranium	2.46E-03	8.62E-06	2.24E-05	0.00E+00	6.85E-03	3.84E-02	8.20E-02	1.30E-01			
					olication Case							
	Cobalt	8.97E-04	9.05E-07	1.06E-06	0.00E+00	2.40E-03	1.46E-03	2.47E-02	2.94E-02			



Table C13: Estimated Non-carcinogen Total Risk to Human Receptors – Project Lifespan – Application Case and Upper Bound Scenario

				1		espan HQs		1				
	СОРС	Water (internal)	Soil (internal)	Sediment (internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC			
	Copper	2.98E-05	2.96E-08	4.21E-08	0.00E+00	1.83E-03	5.40E-04	4.12E-02	4.36E-02			
	Molybdenum	1.35E-04	1.07E-07	1.98E-07	0.00E+00	3.01E-06	2.98E-04	6.98E-02	7.02E-02			
	Uranium	2.80E-03	1.02E-05	2.49E-05	0.00E+00	7.79E-03	6.36E-02	9.24E-02	1.67E-01			
				Upper	Bound Scenar	io - Total Proj	ect Risk					
	Cobalt	8.98E-04	9.05E-07	1.06E-06	0.00E+00	2.40E-03	1.46E-03	2.47E-02	2.94E-02			
	Copper	2.98E-05	2.96E-08	4.21E-08	0.00E+00	1.83E-03	5.40E-04	4.12E-02	4.36E-02			
	Molybdenum	1.91E-04	1.07E-07	2.74E-07	0.00E+00	4.23E-06	2.98E-04	6.98E-02	7.03E-02			
	Uranium	3.40E-03	1.02E-05	2.94E-05	0.00E+00	9.44E-03	6.36E-02	9.28E-02	1.69E-01			
		Base Case										
	Cobalt	9.58E-04	5.91E-05	6.62E-05	0.00E+00	2.18E-03	2.84E-03	4.78E-02	5.39E-02			
	Copper	3.25E-05	1.93E-06	2.69E-06	0.00E+00	1.69E-03	1.06E-03	9.70E-02	9.98E-02			
	Molybdenum	1.47E-04	8.46E-06	1.29E-05	0.00E+00	2.79E-06	7.01E-04	2.64E-01	2.64E-01			
	Uranium	2.75E-03	5.64E-04	1.47E-03	0.00E+00	6.49E-03	7.57E-02	1.70E-01	2.57E-01			
		Application Case - Total Project Risk										
Coloristan ex Hamasatan	Cobalt	1.00E-03	5.91E-05	6.89E-05	0.00E+00	2.27E-03	2.86E-03	4.80E-02	5.43E-02			
Subsistence Harvester One-year-old (Beet Lake)	Copper	3.33E-05	1.93E-06	2.75E-06	0.00E+00	1.73E-03	1.06E-03	9.71E-02	9.99E-02			
One-year-old (beet take)	Molybdenum	1.84E-04	8.47E-06	1.58E-05	0.00E+00	3.47E-06	7.11E-04	2.64E-01	2.64E-01			
	Uranium	3.13E-03	6.64E-04	1.63E-03	0.00E+00	7.39E-03	1.14E-01	1.78E-01	3.04E-01			
				Upper	Bound Scenar	io - Total Proj	ect Risk					
	Cobalt	1.00E-03	5.91E-05	6.91E-05	0.00E+00	2.28E-03	2.86E-03	4.81E-02	5.43E-02			
	Copper	3.33E-05	1.93E-06	2.75E-06	0.00E+00	1.74E-03	1.06E-03	9.71E-02	9.99E-02			
	Molybdenum	2.60E-04	8.47E-06	2.18E-05	0.00E+00	4.88E-06	7.11E-04	2.64E-01	2.65E-01			
	Uranium	3.80E-03	6.64E-04	1.92E-03	0.00E+00	8.95E-03	1.14E-01	1.78E-01	3.07E-01			
Culculator as Hamiston		Base Case										
Subsistence Harvester (Lloyd Lake)	Cobalt	8.58E-04	9.04E-07	1.01E-06	0.00E+00	2.30E-03	1.44E-03	2.44E-02	2.90E-02			
(Lioya Lake)	Copper	2.91E-05	2.95E-08	4.12E-08	0.00E+00	1.79E-03	5.38E-04	4.12E-02	4.35E-02			



Table C13: Estimated Non-carcinogen Total Risk to Human Receptors – Project Lifespan – Application Case and Upper Bound Scenario

		Project Lifespan HQs											
	СОРС	Water (internal)	Soil (internal)	Sediment (internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC				
	Molybdenum	1.08E-04	1.06E-07	1.62E-07	0.00E+00	2.42E-06	2.92E-04	6.98E-02	7.02E-02				
	Uranium	2.46E-03	8.62E-06	2.24E-05	0.00E+00	6.85E-03	3.84E-02	8.20E-02	1.30E-01				
				Арј	olication Case	- Total Project	Risk						
	Cobalt	8.62E-04	9.05E-07	1.02E-06	0.00E+00	2.31E-03	1.46E-03	2.45E-02	2.91E-02				
	Copper	2.92E-05	2.95E-08	4.13E-08	0.00E+00	1.79E-03	5.39E-04	4.12E-02	4.36E-02				
	Molybdenum	1.11E-04	1.07E-07	1.66E-07	0.00E+00	2.48E-06	2.98E-04	6.98E-02	7.02E-02				
	Uranium	2.49E-03	8.68E-06	2.27E-05	0.00E+00	6.94E-03	3.93E-02	8.24E-02	1.31E-01				
				Upper	Bound Scenar	io - Total Proj	ect Risk						
	Cobalt	8.62E-04	9.05E-07	1.02E-06	0.00E+00	2.31E-03	1.46E-03	2.45E-02	2.91E-02				
	Copper	2.92E-05	2.95E-08	4.13E-08	0.00E+00	1.79E-03	5.39E-04	4.12E-02	4.36E-02				
	Molybdenum	1.17E-04	1.07E-07	1.74E-07	0.00E+00	2.60E-06	2.98E-04	6.98E-02	7.02E-02				
	Uranium	2.55E-03	8.68E-06	2.31E-05	0.00E+00	7.09E-03	3.93E-02	8.24E-02	1.31E-01				
					Base	Case							
	Cobalt	9.58E-04	5.91E-05	6.62E-05	0.00E+00	2.18E-03	2.84E-03	4.78E-02	5.39E-02				
	Copper	3.25E-05	1.93E-06	2.69E-06	0.00E+00	1.69E-03	1.06E-03	9.70E-02	9.98E-02				
	Molybdenum	1.47E-04	8.46E-06	1.29E-05	0.00E+00	2.79E-06	7.01E-04	2.64E-01	2.64E-01				
	Uranium	2.75E-03	5.64E-04	1.47E-03	0.00E+00	6.49E-03	7.57E-02	1.70E-01	2.57E-01				
				Арј	olication Case	- Total Project	Risk						
Subsistence Harvester	Cobalt	9.62E-04	5.91E-05	6.65E-05	0.00E+00	2.19E-03	2.86E-03	4.79E-02	5.40E-02				
One-year-old (Lloyd Lake)	Copper	3.26E-05	1.93E-06	2.70E-06	0.00E+00	1.70E-03	1.06E-03	9.70E-02	9.98E-02				
	Molybdenum	1.51E-04	8.47E-06	1.32E-05	0.00E+00	2.86E-06	7.11E-04	2.64E-01	2.64E-01				
	Uranium	2.78E-03	5.67E-04	1.48E-03	0.00E+00	6.58E-03	7.71E-02	1.71E-01	2.59E-01				
				Upper	Bound Scenar	io - Total Proj	ect Risk						
	Cobalt	9.62E-04	5.91E-05	6.65E-05	0.00E+00	2.19E-03	2.86E-03	4.79E-02	5.40E-02				
	Copper	3.26E-05	1.93E-06	2.70E-06	0.00E+00	1.70E-03	1.06E-03	9.70E-02	9.98E-02				
	Molybdenum	1.59E-04	8.47E-06	1.38E-05	0.00E+00	3.01E-06	7.11E-04	2.64E-01	2.64E-01				



Table C13: Estimated Non-carcinogen Total Risk to Human Receptors – Project Lifespan – Application Case and Upper Bound Scenario

					Project Lif	fespan HQs			
	СОРС	Water (internal)	Soil (internal)	Sediment (internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
	Uranium	2.84E-03	5.67E-04	1.51E-03	0.00E+00	6.72E-03	7.71E-02	1.71E-01	2.59E-01
					Base	Case			
	Cobalt	8.58E-04	9.04E-07	1.01E-06	0.00E+00	7.27E-04	1.13E-03	1.88E-02	2.15E-02
	Copper	2.91E-05	2.95E-08	4.12E-08	0.00E+00	5.66E-04	4.22E-04	3.86E-02	3.96E-02
	Molybdenum	1.08E-04	1.06E-07	1.62E-07	0.00E+00	7.66E-07	2.29E-04	8.60E-02	8.63E-02
	Uranium	2.46E-03	8.62E-06	2.24E-05	0.00E+00	2.17E-03	3.01E-02	6.79E-02	1.03E-01
				Арј	olication Case	- Total Project	Risk		
Seasonal Resident	Cobalt	9.02E-04	9.05E-07	1.06E-06	0.00E+00	7.65E-04	1.13E-03	1.89E-02	2.17E-02
(Patterson Lake South	Copper	2.99E-05	2.95E-08	4.22E-08	0.00E+00	5.82E-04	4.23E-04	3.87E-02	3.97E-02
Arm)	Molybdenum	1.40E-04	1.07E-07	2.04E-07	0.00E+00	9.85E-07	2.32E-04	8.60E-02	8.64E-02
	Uranium	3.02E-03	1.02E-05	2.65E-05	0.00E+00	2.66E-03	4.35E-02	7.31E-02	1.22E-01
				Upper	Bound Scenar	io - Total Proj	ect Risk		
	Cobalt	9.04E-04	9.05E-07	1.06E-06	0.00E+00	7.66E-04	1.13E-03	1.89E-02	2.18E-02
	Copper	3.00E-05	2.95E-08	4.23E-08	0.00E+00	5.82E-04	4.23E-04	3.87E-02	3.97E-02
	Molybdenum	2.04E-04	1.07E-07	2.92E-07	0.00E+00	1.43E-06	2.32E-04	8.61E-02	8.65E-02
	Uranium	4.00E-03	1.02E-05	3.38E-05	0.00E+00	3.51E-03	4.35E-02	7.34E-02	1.24E-01
					Base	Case			
	Cobalt	9.58E-04	5.91E-05	6.62E-05	0.00E+00	6.89E-04	2.47E-03	4.13E-02	4.56E-02
	Copper	3.25E-05	1.93E-06	2.69E-06	0.00E+00	5.36E-04	9.22E-04	9.32E-02	9.47E-02
	Molybdenum	1.47E-04	8.46E-06	1.29E-05	0.00E+00	8.84E-07	6.10E-04	2.77E-01	2.78E-01
Seasonal Resident One- year-old (Patterson Lake	Uranium	2.75E-03	5.64E-04	1.47E-03	0.00E+00	2.06E-03	6.58E-02	1.60E-01	2.33E-01
South Arm)				Арј	olication Case	- Total Project	Risk		
Journ Amily	Cobalt	1.01E-03	5.91E-05	6.94E-05	0.00E+00	7.25E-04	2.48E-03	4.15E-02	4.59E-02
	Copper	3.34E-05	1.93E-06	2.76E-06	0.00E+00	5.52E-04	9.24E-04	9.32E-02	9.47E-02
	Molybdenum	1.90E-04	8.48E-06	1.62E-05	0.00E+00	1.14E-06	6.16E-04	2.77E-01	2.78E-01
	Uranium	3.37E-03	6.66E-04	1.73E-03	0.00E+00	2.52E-03	8.97E-02	1.64E-01	2.62E-01



Table C13: Estimated Non-carcinogen Total Risk to Human Receptors – Project Lifespan – Application Case and Upper Bound Scenario

				,	Project Lif	espan HQs			
	СОРС	Water (internal)	Soil (internal)	Sediment (internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
				Upper	Bound Scenar	io - Total Proj	ect Risk		
	Cobalt	1.01E-03	5.91E-05	6.95E-05	0.00E+00	7.26E-04	2.48E-03	4.15E-02	4.59E-02
	Copper	3.34E-05	1.93E-06	2.76E-06	0.00E+00	5.52E-04	9.24E-04	9.32E-02	9.47E-02
	Molybdenum	2.78E-04	8.48E-06	2.33E-05	0.00E+00	1.65E-06	6.16E-04	2.77E-01	2.78E-01
	Uranium	4.46E-03	6.66E-04	2.21E-03	0.00E+00	3.33E-03	8.97E-02	1.64E-01	2.65E-01
					Base	Case			
	Cobalt	8.58E-04	9.04E-07	1.01E-06	0.00E+00	7.27E-04	1.13E-03	1.88E-02	2.15E-02
	Copper	2.91E-05	2.95E-08	4.12E-08	0.00E+00	5.66E-04	4.22E-04	3.86E-02	3.96E-02
	Molybdenum	1.08E-04	1.06E-07	1.62E-07	0.00E+00	7.66E-07	2.29E-04	8.60E-02	8.63E-02
	Uranium	2.46E-03	8.62E-06	2.24E-05	0.00E+00	2.17E-03	3.01E-02	6.79E-02	1.03E-01
				Apı	olication Case -	Total Project	Risk		
Seasonal Resident (Lloyd	Cobalt	8.60E-04	9.05E-07	1.02E-06	0.00E+00	7.31E-04	1.13E-03	1.88E-02	2.15E-02
Lake)	Copper	2.91E-05	2.95E-08	4.13E-08	0.00E+00	7.53E-04	4.22E-04	3.86E-02	3.98E-02
L uncy	Molybdenum	1.10E-04	1.06E-07	1.64E-07	0.00E+00	7.86E-07	2.31E-04	8.60E-02	8.64E-02
	Uranium	2.48E-03	8.66E-06	2.26E-05	0.00E+00	2.73E-03	3.04E-02	6.80E-02	1.04E-01
					Bound Scenar				
	Cobalt	8.60E-04	9.05E-07	1.02E-06	0.00E+00	7.31E-04	1.13E-03	1.88E-02	2.15E-02
	Copper	2.91E-05	2.95E-08	4.13E-08	0.00E+00	7.53E-04	4.22E-04	3.86E-02	3.98E-02
	Molybdenum	1.14E-04	1.06E-07	1.69E-07	0.00E+00	8.26E-07	2.31E-04	8.60E-02	8.64E-02
	Uranium	2.51E-03	8.66E-06	2.28E-05	0.00E+00	2.79E-03	3.04E-02	6.80E-02	1.04E-01
			T	Ī		Case	I	Ī	
	Cobalt	9.58E-04	5.91E-05	6.62E-05	0.00E+00	6.89E-04	2.47E-03	4.13E-02	4.56E-02
Seasonal Resident One-	Copper	3.25E-05	1.93E-06	2.69E-06	0.00E+00	5.36E-04	9.22E-04	9.32E-02	9.47E-02
year-old (Lloyd Lake)	Molybdenum	1.47E-04	8.46E-06	1.29E-05	0.00E+00	8.84E-07	6.10E-04	2.77E-01	2.78E-01
	Uranium	2.75E-03	5.64E-04	1.47E-03	0.00E+00	2.06E-03	6.58E-02	1.60E-01	2.33E-01
				Apı	olication Case -	Total Project	Risk		



Table C13: Estimated Non-carcinogen Total Risk to Human Receptors – Project Lifespan – Application Case and Upper Bound Scenario

					Project Lif	espan HQs						
	СОРС	Water (internal)	Soil (internal)	Sediment (internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC			
	Cobalt	9.60E-04	5.91E-05	6.64E-05	0.00E+00	6.91E-04	2.48E-03	4.14E-02	4.56E-02			
	Copper	3.25E-05	1.93E-06	2.70E-06	0.00E+00	5.37E-04	9.23E-04	9.32E-02	9.47E-02			
	Molybdenum	1.50E-04	8.47E-06	1.31E-05	0.00E+00	8.97E-07	6.13E-04	2.77E-01	2.78E-01			
	Uranium	2.77E-03	5.66E-04	1.47E-03	0.00E+00	2.07E-03	6.63E-02	1.60E-01	2.34E-01			
				Upper	Bound Scenar	io - Total Proj	ect Risk					
	Cobalt	9.60E-04	5.91E-05	6.64E-05	0.00E+00	6.91E-04	2.48E-03	4.14E-02	4.56E-02			
	Copper	3.25E-05	1.93E-06	2.70E-06	0.00E+00	5.37E-04	9.23E-04	9.32E-02	9.47E-02			
	Molybdenum	1.54E-04	8.47E-06	1.35E-05	0.00E+00	9.25E-07	6.13E-04	2.77E-01	2.78E-01			
	Uranium	2.80E-03	5.66E-04	1.49E-03	0.00E+00	2.10E-03	6.63E-02	1.60E-01	2.34E-01			
					Base	Case						
	Cobalt	-	-	-	-	-	-	-	-			
	Copper	-	-	-	-	-	-	-	-			
	Molybdenum	-	-	-	-	-	-	-	-			
	Uranium	-	-	-	-	-	-	-	-			
				Apı	plication Case	- Total Project	Risk					
Permanent Resident	Cobalt	-	-	-	-	-	-	-	-			
(Patterson Lake North	Copper	-	-	-	-	-	-	-	-			
Arm - West Basin)	Molybdenum	-	-	-	-	-	-	-	-			
	Uranium	-	-	-	-	-	-	-	-			
				Upper	Bound Scenar	io - Total Proj	ect Risk					
	Cobalt	-	-	-	-	-	-	-	-			
	Copper	-	-	-	-	-	-	-	-			
	Molybdenum	-	-	-	-	-	-	-	-			
	Uranium	-	-	-	-	-	-	-	-			
		Base Case										
	Cobalt	-	-	-	-	-	-	-	-			



Table C13: Estimated Non-carcinogen Total Risk to Human Receptors – Project Lifespan – Application Case and Upper Bound Scenario

			Project Lifespan HQs										
	СОРС	Water (internal)	Soil (internal)	Sediment (internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC				
	Copper	-	-	-	-	-	-	-	-				
	Molybdenum	-	-	-	-	-	-	-	-				
	Uranium	-	-	-	-	-	-	-	-				
		Application Case - Total Project Risk											
	Cobalt	-	-	-	-	-	-	-	-				
Permanent Resident One-	Copper	-	-	-	-	-	-	-	-				
year-old (Patterson Lake	Molybdenum	-	-	-	-	-	-	-	-				
North Arm - West Basin)	Uranium	-	-	-	-	-	-	-	-				
				Upper	Bound Scenar	io - Total Proj	ect Risk						
	Cobalt	-	-	-	-	-	-	-	-				
	Copper	-	-	-	-	-	-	-	-				
	Molybdenum	-	-	-	-	-	-	-	-				
	Uranium	-	-	-	-	-	-	-	-				



Table C14: Estimated Non-carcinogen Total Risk to Human Receptors – Far Future – Application Case and Upper Bound Scenario

		Far Future HQs										
	СОРС	Water (internal)	Soil (internal)	Sediment (internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC			
					Base	Case						
	Cobalt	-	-	-	-	-	-	-	-			
	Copper	-	-	-	-	-	-	-	-			
	Molybdenum	-	-	-	-	-	-	-	-			
	Uranium	-	-	-	-	-	-	-	-			
				Appl	ication Case	- Total Projec	t Risk					
Camp Worker	Cobalt	-	-	-	-	-	-	-	-			
(Patterson Lake North	Copper	-	-	-	-	-	-	-	-			
Arm - West Basin)	Molybdenum	-	-	-	-	-	-	-	-			
	Uranium	-	-	-	-	-	-	-	-			
		Upper Bound Scenario - Total Project Risk										
	Cobalt	-	-	-	-	-	-	-	-			
	Copper	-	-	-	-	-	-	-	-			
	Molybdenum	-	-	-	-	-	-	-	-			
	Uranium	-	-	-	-	-	-	-	-			
					Base	Case						
	Cobalt	8.58E-04	9.04E-07	1.01E-06	0.00E+00	2.30E-03	1.44E-03	2.44E-02	2.90E-02			
	Copper	2.91E-05	2.95E-08	4.12E-08	0.00E+00	1.79E-03	5.38E-04	4.12E-02	4.35E-02			
	Molybdenum	1.08E-04	1.06E-07	1.62E-07	0.00E+00	2.42E-06	2.92E-04	6.98E-02	7.02E-02			
Subsistence Harvester (Patterson Lake South	Uranium	2.46E-03	8.62E-06	2.24E-05	0.00E+00	6.85E-03	3.84E-02	8.20E-02	1.30E-01			
(Patterson Lake South Arm)				Appl	ication Case	- Total Projec	t Risk					
74111)	Cobalt	1.23E-03	9.05E-07	1.46E-06	0.00E+00	3.30E-03	1.44E-03	2.73E-02	3.33E-02			
	Copper	4.27E-05	2.95E-08	6.04E-08	0.00E+00	2.62E-03	5.38E-04	4.19E-02	4.51E-02			
	Molybdenum	7.79E-04	1.06E-07	1.17E-06	0.00E+00	1.74E-05	2.92E-04	7.05E-02	7.16E-02			
	Uranium	5.52E-03	8.90E-06	5.04E-05	0.00E+00	1.54E-02	3.96E-02	8.62E-02	1.47E-01			

Table C14: Estimated Non-carcinogen Total Risk to Human Receptors – Far Future – Application Case and Upper Bound Scenario

					Far Fut	ure HQs			
	СОРС	Water (internal)	Soil (internal)	Sediment (internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
		(internal)	(internal)		Sound Scenar			Ailinas	201.0
	Cobalt	1.44E-03	9.05E-07	1.70E-06	0.00E+00	3.84E-03	1.44E-03	2.89E-02	3.56E-02
	Copper	4.95E-05	2.95E-08	7.01E-08	0.00E+00	3.04E-03	5.38E-04	4.23E-02	4.60E-02
	Molybdenum	2.68E-03	1.06E-07	4.01E-06	0.00E+00	5.98E-05	2.92E-04	7.27E-02	7.57E-02
	Uranium	5.53E-03	8.90E-06	5.04E-05	0.00E+00	1.54E-02	3.96E-02	8.62E-02	1.47E-01
					Base	Case			
	Cobalt	9.58E-04	5.91E-05	6.62E-05	0.00E+00	2.18E-03	2.84E-03	4.78E-02	5.39E-02
	Copper	3.25E-05	1.93E-06	2.69E-06	0.00E+00	1.69E-03	1.06E-03	9.70E-02	9.98E-02
	Molybdenum	1.47E-04	8.46E-06	1.29E-05	0.00E+00	2.79E-06	7.01E-04	2.64E-01	2.64E-01
	Uranium	2.75E-03	5.64E-04	1.47E-03	0.00E+00	6.49E-03	7.57E-02	1.70E-01	2.57E-01
				Appl	ication Case	- Total Projec	t Risk		
Subsistence Harvester	Cobalt	1.38E-03	5.91E-05	9.54E-05	0.00E+00	3.13E-03	2.84E-03	5.08E-02	5.83E-02
One-year-old (Patterson	Copper	4.76E-05	1.93E-06	3.95E-06	0.00E+00	2.48E-03	1.06E-03	9.77E-02	1.01E-01
Lake South Arm)	Molybdenum	1.06E-03	8.46E-06	9.27E-05	0.00E+00	2.00E-05	7.01E-04	2.64E-01	2.66E-01
	Uranium	6.17E-03	5.81E-04	3.29E-03	0.00E+00	1.46E-02	7.81E-02	1.73E-01	2.76E-01
					Sound Scenar		_		l
	Cobalt	1.60E-03	5.91E-05	1.11E-04	0.00E+00	3.64E-03	2.84E-03	5.25E-02	6.07E-02
	Copper	5.52E-05	1.93E-06	4.58E-06	0.00E+00	2.88E-03	1.06E-03	9.81E-02	1.02E-01
	Molybdenum	3.65E-03	8.46E-06	3.19E-04	0.00E+00	6.91E-05	7.01E-04	2.67E-01	2.72E-01
	Uranium	6.17E-03	5.81E-04	3.29E-03	0.00E+00	1.46E-02	7.81E-02	1.73E-01	2.76E-01
		0.505.04	0.045.05	4.045.00		Case	4.445.00	2.445.00	2.005.00
	Cobalt	8.58E-04	9.04E-07	1.01E-06	0.00E+00	2.30E-03	1.44E-03	2.44E-02	2.90E-02
Subsistence Harvester	Copper	2.91E-05	2.95E-08	4.12E-08	0.00E+00	1.79E-03	5.38E-04	4.12E-02	4.35E-02
(Beet Lake)	Molybdenum	1.08E-04	1.06E-07	1.62E-07	0.00E+00	2.42E-06	2.92E-04	6.98E-02	7.02E-02
	Uranium	2.46E-03	8.62E-06	2.24E-05	0.00E+00	6.85E-03	3.84E-02	8.20E-02	1.30E-01
				Appl	ication Case	- Total Projec	t Risk		

Table C14: Estimated Non-carcinogen Total Risk to Human Receptors – Far Future – Application Case and Upper Bound Scenario

					Far Fut	ure HQs					
	СОРС	Water (internal)	Soil (internal)	Sediment (internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC		
	Cobalt	1.06E-03	9.05E-07	1.25E-06	0.00E+00	2.83E-03	1.44E-03	2.54E-02	3.08E-02		
	Copper	3.62E-05	2.95E-08	5.13E-08	0.00E+00	2.22E-03	5.38E-04	4.15E-02	4.43E-02		
	Molybdenum	4.57E-04	1.06E-07	6.85E-07	0.00E+00	1.02E-05	2.92E-04	7.00E-02	7.08E-02		
	Uranium	3.60E-03	8.79E-06	3.28E-05	0.00E+00	1.00E-02	3.91E-02	8.34E-02	1.36E-01		
				Upper E	Sound Scenar	io - Total Pro	ject Risk				
	Cobalt	1.16E-03	9.05E-07	1.38E-06	0.00E+00	3.11E-03	1.44E-03	2.60E-02	3.17E-02		
	Copper	3.97E-05	2.95E-08	5.63E-08	0.00E+00	2.44E-03	5.38E-04	4.16E-02	4.46E-02		
	Molybdenum	1.45E-03	1.06E-07	2.17E-06	0.00E+00	3.23E-05	2.92E-04	7.08E-02	7.26E-02		
	Uranium	3.60E-03	8.79E-06	3.28E-05	0.00E+00	1.00E-02	3.91E-02	8.34E-02	1.36E-01		
			I	ı		Case		l			
	Cobalt	9.58E-04	5.91E-05	6.62E-05	0.00E+00	2.18E-03	2.84E-03	4.78E-02	5.39E-02		
	Copper	3.25E-05	1.93E-06	2.69E-06	0.00E+00	1.69E-03	1.06E-03	9.70E-02	9.98E-02		
	Molybdenum	1.47E-04	8.46E-06	1.29E-05	0.00E+00	2.79E-06	7.01E-04	2.64E-01	2.64E-01		
	Uranium	2.75E-03	5.64E-04	1.47E-03	0.00E+00	6.49E-03	7.57E-02	1.70E-01	2.57E-01		
			l			- Total Projec					
Subsistence Harvester	Cobalt	1.18E-03	5.91E-05	8.17E-05	0.00E+00	2.68E-03	2.84E-03	4.89E-02	5.57E-02		
One-year-old (Beet Lake)	Copper	4.04E-05	1.93E-06	3.35E-06	0.00E+00	2.11E-03	1.06E-03	9.73E-02	1.00E-01		
	Molybdenum	6.22E-04	8.46E-06	5.45E-05	0.00E+00	1.18E-05	7.01E-04	2.64E-01	2.65E-01		
	Uranium	4.02E-03	5.74E-04	2.14E-03	0.00E+00	9.50E-03	7.71E-02	1.71E-01	2.65E-01		
			I			io - Total Pro		T			
	Cobalt	1.30E-03	5.91E-05	8.99E-05	0.00E+00	2.95E-03	2.84E-03	4.94E-02	5.66E-02		
	Copper	4.44E-05	1.93E-06	3.68E-06	0.00E+00	2.31E-03	1.06E-03	9.74E-02	1.01E-01		
	Molybdenum	1.97E-03	8.46E-06	1.72E-04	0.00E+00	3.73E-05	7.01E-04	2.65E-01	2.68E-01		
	Uranium	4.02E-03	5.74E-04	2.15E-03	0.00E+00	9.51E-03	7.71E-02	1.71E-01	2.65E-01		
Subsistence Harvester		Base Case									
(Lloyd Lake)	Cobalt	8.58E-04	9.04E-07	1.01E-06	0.00E+00	2.30E-03	1.44E-03	2.44E-02	2.90E-02		

Table C14: Estimated Non-carcinogen Total Risk to Human Receptors – Far Future – Application Case and Upper Bound Scenario

		Far Future HQs								
					1					
	СОРС	Water (internal)	Soil (internal)	Sediment (internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC	
	Copper	2.91E-05	2.95E-08	4.12E-08	0.00E+00	1.79E-03	5.38E-04	4.12E-02	4.35E-02	
	Molybdenum	1.08E-04	1.06E-07	1.62E-07	0.00E+00	2.42E-06	2.92E-04	6.98E-02	7.02E-02	
	Uranium	2.46E-03	8.62E-06	2.24E-05	0.00E+00	6.85E-03	3.84E-02	8.20E-02	1.30E-01	
				Appl	ication Case	- Total Projec	t Risk			
	Cobalt	8.78E-04	9.05E-07	1.04E-06	0.00E+00	2.35E-03	1.44E-03	2.45E-02	2.92E-02	
	Copper	2.98E-05	2.95E-08	4.22E-08	0.00E+00	1.83E-03	5.38E-04	4.12E-02	4.36E-02	
	Molybdenum	1.44E-04	1.06E-07	2.15E-07	0.00E+00	3.21E-06	2.92E-04	6.98E-02	7.02E-02	
	Uranium	2.56E-03	8.63E-06	2.34E-05	0.00E+00	7.14E-03	3.84E-02	8.21E-02	1.30E-01	
				Upper E	Bound Scenar	io - Total Pro	ject Risk			
	Cobalt	8.89E-04	9.05E-07	1.05E-06	0.00E+00	2.38E-03	1.44E-03	2.46E-02	2.93E-02	
	Copper	3.02E-05	2.95E-08	4.28E-08	0.00E+00	1.85E-03	5.38E-04	4.12E-02	4.37E-02	
	Molybdenum	2.45E-04	1.06E-07	3.67E-07	0.00E+00	5.47E-06	2.92E-04	6.99E-02	7.04E-02	
	Uranium	2.56E-03	8.63E-06	2.34E-05	0.00E+00	7.14E-03	3.84E-02	8.21E-02	1.30E-01	
					Base	Case				
	Cobalt	9.58E-04	5.91E-05	6.62E-05	0.00E+00	2.18E-03	2.84E-03	4.78E-02	5.39E-02	
	Copper	3.25E-05	1.93E-06	2.69E-06	0.00E+00	1.69E-03	1.06E-03	9.70E-02	9.98E-02	
	Molybdenum	1.47E-04	8.46E-06	1.29E-05	0.00E+00	2.79E-06	7.01E-04	2.64E-01	2.64E-01	
	Uranium	2.75E-03	5.64E-04	1.47E-03	0.00E+00	6.49E-03	7.57E-02	1.70E-01	2.57E-01	
				Appl	ication Case	- Total Projec	t Risk			
Subsistence Harvester One-year-old (Lloyd Lake)	Cobalt	9.80E-04	5.91E-05	6.78E-05	0.00E+00	2.23E-03	2.84E-03	4.79E-02	5.41E-02	
One-year-old (Lloyd Lake)	Copper	3.33E-05	1.93E-06	2.76E-06	0.00E+00	1.74E-03	1.06E-03	9.70E-02	9.99E-02	
	Molybdenum	1.96E-04	8.46E-06	1.71E-05	0.00E+00	3.71E-06	7.01E-04	2.64E-01	2.64E-01	
	Uranium	2.86E-03	5.64E-04	1.53E-03	0.00E+00	6.77E-03	7.57E-02	1.70E-01	2.58E-01	
				Upper E	Bound Scenar	io - Total Pro	ject Risk			
	Cobalt	9.93E-04	5.91E-05	6.87E-05	0.00E+00	2.26E-03	2.84E-03	4.80E-02	5.42E-02	
	Copper	3.37E-05	1.93E-06	2.79E-06	0.00E+00	1.76E-03	1.06E-03	9.71E-02	9.99E-02	

Table C14: Estimated Non-carcinogen Total Risk to Human Receptors – Far Future – Application Case and Upper Bound Scenario

		Far Future HQs										
	СОРС	Water (internal)	Soil (internal)	Sediment (internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC			
	Molybdenum	3.33E-04	8.46E-06	2.92E-05	0.00E+00	6.31E-06	7.01E-04	2.64E-01	2.65E-01			
	Uranium	2.86E-03	5.64E-04	1.53E-03	0.00E+00	6.77E-03	7.57E-02	1.70E-01	2.58E-01			
					Base	Case						
	Cobalt	8.58E-04	9.04E-07	1.01E-06	0.00E+00	7.27E-04	1.13E-03	1.88E-02	2.15E-02			
	Copper	2.91E-05	2.95E-08	4.12E-08	0.00E+00	5.66E-04	4.22E-04	3.86E-02	3.96E-02			
	Molybdenum	1.08E-04	1.06E-07	1.62E-07	0.00E+00	7.66E-07	2.29E-04	8.60E-02	8.63E-02			
	Uranium	2.46E-03	8.62E-06	2.24E-05	0.00E+00	2.17E-03	3.01E-02	6.79E-02	1.03E-01			
			Application Case - Total Project Risk									
Seasonal Resident	Cobalt	1.08E-03	9.04E-07	1.28E-06	0.00E+00	9.19E-04	1.13E-03	1.95E-02	2.26E-02			
(Patterson Lake South	Copper	3.72E-05	2.95E-08	5.27E-08	0.00E+00	7.24E-04	4.22E-04	3.88E-02	4.00E-02			
Arm)	Molybdenum	5.11E-04	1.06E-07	7.64E-07	0.00E+00	3.61E-06	2.29E-04	8.62E-02	8.69E-02			
	Uranium	4.30E-03	8.79E-06	3.92E-05	0.00E+00	3.79E-03	3.07E-02	6.90E-02	1.08E-01			
				Upper E	ound Scenar	io - Total Pro	ject Risk					
	Cobalt	1.20E-03	9.04E-07	1.42E-06	0.00E+00	1.02E-03	1.13E-03	1.99E-02	2.32E-02			
	Copper	4.13E-05	2.95E-08	5.85E-08	0.00E+00	8.04E-04	4.22E-04	3.89E-02	4.02E-02			
	Molybdenum	1.65E-03	1.06E-07	2.47E-06	0.00E+00	1.17E-05	2.29E-04	8.67E-02	8.86E-02			
	Uranium	4.30E-03	8.79E-06	3.92E-05	0.00E+00	3.79E-03	3.07E-02	6.90E-02	1.08E-01			
					Base	Case						
	Cobalt	9.58E-04	5.91E-05	6.62E-05	0.00E+00	6.89E-04	2.47E-03	4.13E-02	4.56E-02			
	Copper	3.25E-05	1.93E-06	2.69E-06	0.00E+00	5.36E-04	9.22E-04	9.32E-02	9.47E-02			
Seasonal Resident One-	Molybdenum	1.47E-04	8.46E-06	1.29E-05	0.00E+00	8.84E-07	6.10E-04	2.77E-01	2.78E-01			
year-old (Patterson Lake	Uranium	2.75E-03	5.64E-04	1.47E-03	0.00E+00	2.06E-03	6.58E-02	1.60E-01	2.33E-01			
South Arm)				Appl	ication Case -	Total Projec	t Risk					
	Cobalt	1.21E-03	5.91E-05	8.37E-05	0.00E+00	8.71E-04	2.47E-03	4.21E-02	4.68E-02			
	Copper	4.16E-05	1.93E-06	3.45E-06	0.00E+00	6.86E-04	9.22E-04	9.33E-02	9.50E-02			
	Molybdenum	6.94E-04	8.46E-06	6.08E-05	0.00E+00	4.16E-06	6.10E-04	2.78E-01	2.79E-01			



Table C14: Estimated Non-carcinogen Total Risk to Human Receptors – Far Future – Application Case and Upper Bound Scenario

			Far Future HQs									
	СОРС	Water (internal)	Soil (internal)	Sediment (internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC			
	Uranium	4.80E-03	5.74E-04	2.56E-03	0.00E+00	3.60E-03	6.70E-02	1.61E-01	2.40E-01			
				Upper B	Sound Scenar	io - Total Pro	ject Risk					
	Cobalt	1.34E-03	5.91E-05	9.30E-05	0.00E+00	9.68E-04	2.47E-03	4.25E-02	4.74E-02			
	Copper	4.61E-05	1.93E-06	3.82E-06	0.00E+00	7.62E-04	9.22E-04	9.34E-02	9.52E-02			
	Molybdenum	2.25E-03	8.46E-06	1.97E-04	0.00E+00	1.35E-05	6.10E-04	2.78E-01	2.81E-01			
	Uranium	4.80E-03	5.74E-04	2.56E-03	0.00E+00	3.60E-03	6.70E-02	1.61E-01	2.40E-01			
					Base	Case						
	Cobalt	8.58E-04	9.04E-07	1.01E-06	0.00E+00	7.27E-04	1.13E-03	1.88E-02	2.15E-02			
	Copper	2.91E-05	2.95E-08	4.12E-08	0.00E+00	5.66E-04	4.22E-04	3.86E-02	3.96E-02			
	Molybdenum	1.08E-04	1.06E-07	1.62E-07	0.00E+00	7.66E-07	2.29E-04	8.60E-02	8.63E-02			
	Uranium	2.46E-03	8.62E-06	2.24E-05	0.00E+00	2.17E-03	3.01E-02	6.79E-02	1.03E-01			
				Appl	ication Case -	Total Projec	t Risk		•			
Seasonal Resident (Lloyd	Cobalt	8.70E-04	9.04E-07	1.03E-06	0.00E+00	7.45E-04	1.13E-03	1.88E-02	2.16E-02			
Lake)	Copper	2.95E-05	2.95E-08	4.18E-08	0.00E+00	7.70E-04	4.22E-04	3.86E-02	3.99E-02			
	Molybdenum	1.30E-04	1.06E-07	1.94E-07	0.00E+00	1.02E-06	2.29E-04	8.60E-02	8.64E-02			
	Uranium	2.52E-03	8.63E-06	2.30E-05	0.00E+00	2.81E-03	3.01E-02	6.80E-02	1.03E-01			
				Upper B	Sound Scenar	io - Total Pro	ject Risk		•			
	Cobalt	8.76E-04	9.04E-07	1.04E-06	0.00E+00	7.54E-04	1.13E-03	1.88E-02	2.16E-02			
	Copper	2.98E-05	2.95E-08	4.21E-08	0.00E+00	7.79E-04	4.22E-04	3.86E-02	3.99E-02			
	Molybdenum	1.90E-04	1.06E-07	2.85E-07	0.00E+00	1.73E-06	2.29E-04	8.60E-02	8.65E-02			
	Uranium	2.52E-03	8.63E-06	2.30E-05	0.00E+00	2.81E-03	3.01E-02	6.80E-02	1.03E-01			
			T			Case						
Seasonal Resident One-	Cobalt	9.58E-04	5.91E-05	6.62E-05	0.00E+00	6.89E-04	2.47E-03	4.13E-02	4.56E-02			
year-old (Lloyd Lake)	Copper	3.25E-05	1.93E-06	2.69E-06	0.00E+00	5.36E-04	9.22E-04	9.32E-02	9.47E-02			
Jour Gra (210) a 2anc)	Molybdenum	1.47E-04	8.46E-06	1.29E-05	0.00E+00	8.84E-07	6.10E-04	2.77E-01	2.78E-01			
	Uranium	2.75E-03	5.64E-04	1.47E-03	0.00E+00	2.06E-03	6.58E-02	1.60E-01	2.33E-01			



Table C14: Estimated Non-carcinogen Total Risk to Human Receptors – Far Future – Application Case and Upper Bound Scenario

					Far Fut	ure HQs			
	СОРС	Water (internal)	Soil (internal)	Sediment (internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
				Appl	ication Case	- Total Projec	t Risk		
	Cobalt	9.71E-04	5.91E-05	6.72E-05	0.00E+00	6.99E-04	2.47E-03	4.14E-02	4.56E-02
	Copper	3.30E-05	1.93E-06	2.73E-06	0.00E+00	5.44E-04	9.22E-04	9.32E-02	9.47E-02
	Molybdenum	1.76E-04	8.46E-06	1.54E-05	0.00E+00	1.06E-06	6.10E-04	2.77E-01	2.78E-01
	Uranium	2.82E-03	5.64E-04	1.50E-03	0.00E+00	2.11E-03	6.58E-02	1.60E-01	2.33E-01
				Upper E	Bound Scenar	io - Total Pro	oject Risk		
	Cobalt	9.79E-04	5.91E-05	6.77E-05	0.00E+00	7.05E-04	2.47E-03	4.14E-02	4.57E-02
	Copper	3.32E-05	1.93E-06	2.75E-06	0.00E+00	5.48E-04	9.22E-04	9.32E-02	9.47E-02
	Molybdenum	2.59E-04	8.46E-06	2.27E-05	0.00E+00	1.55E-06	6.10E-04	2.77E-01	2.78E-01
	Uranium	2.82E-03	5.64E-04	1.50E-03	0.00E+00	2.11E-03	6.58E-02	1.60E-01	2.33E-01
					Base	Case			
	Cobalt	8.58E-04	9.04E-07	1.01E-06	0.00E+00	2.30E-03	1.44E-03	2.44E-02	2.90E-02
	Copper	2.91E-05	2.95E-08	4.12E-08	0.00E+00	1.79E-03	5.38E-04	4.12E-02	4.35E-02
	Molybdenum	1.08E-04	1.06E-07	1.62E-07	0.00E+00	2.42E-06	2.92E-04	6.98E-02	7.02E-02
	Uranium	2.46E-03	8.62E-06	2.24E-05	0.00E+00	6.85E-03	3.84E-02	8.20E-02	1.30E-01
				Appl	ication Case	- Total Projec	t Risk		
Permanent Resident	Cobalt	2.10E-03	9.07E-07	2.48E-06	0.00E+00	5.62E-03	1.44E-03	3.08E-02	4.00E-02
(Patterson Lake North	Copper	7.53E-05	2.96E-08	1.07E-07	0.00E+00	4.62E-03	5.39E-04	4.30E-02	4.82E-02
Arm - West Basin)	Molybdenum	2.42E-03	1.06E-07	3.62E-06	0.00E+00	5.39E-05	2.92E-04	7.16E-02	7.43E-02
	Uranium	2.46E-02	1.25E-05	2.24E-04	0.00E+00	6.85E-02	5.55E-02	1.13E-01	2.62E-01
				Upper E	Bound Scenar	io - Total Pro	ject Risk		
	Cobalt	2.76E-03	9.07E-07	3.26E-06	0.00E+00	7.39E-03	1.44E-03	3.42E-02	4.58E-02
	Copper	9.85E-05	2.96E-08	1.40E-07	0.00E+00	6.05E-03	5.39E-04	4.39E-02	5.06E-02
	Molybdenum	8.98E-03	1.06E-07	1.34E-05	0.00E+00	2.00E-04	2.92E-04	7.67E-02	8.62E-02
	Uranium	2.46E-02	1.25E-05	2.25E-04	0.00E+00	6.86E-02	5.55E-02	1.13E-01	2.62E-01
					Base	Case			

Table C14: Estimated Non-carcinogen Total Risk to Human Receptors – Far Future – Application Case and Upper Bound Scenario

					Far Fut	ure HQs			
	СОРС	Water (internal)	Soil (internal)	Sediment (internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
	Cobalt	9.58E-04	5.91E-05	6.62E-05	0.00E+00	2.18E-03	2.84E-03	4.78E-02	5.39E-02
	Copper	3.25E-05	1.93E-06	2.69E-06	0.00E+00	1.69E-03	1.06E-03	9.70E-02	9.98E-02
	Molybdenum	1.47E-04	8.46E-06	1.29E-05	0.00E+00	2.79E-06	7.01E-04	2.64E-01	2.64E-01
	Uranium	2.75E-03	5.64E-04	1.47E-03	0.00E+00	6.49E-03	7.57E-02	1.70E-01	2.57E-01
				Appl	ication Case	- Total Projec	t Risk		
	Cobalt	2.34E-03	5.92E-05	1.62E-04	0.00E+00	5.33E-03	2.85E-03	5.43E-02	6.50E-02
Permanent Resident One-	Copper	8.41E-05	1.93E-06	6.97E-06	0.00E+00	4.38E-03	1.06E-03	9.87E-02	1.04E-01
year-old (Patterson Lake North Arm - West Basin)	Molybdenum	3.29E-03	8.46E-06	2.88E-04	0.00E+00	6.23E-05	7.01E-04	2.66E-01	2.70E-01
Hortin Aim West Bushing	Uranium	2.75E-02	8.15E-04	1.47E-02	0.00E+00	6.50E-02	1.09E-01	1.93E-01	4.10E-01
				Upper E	Sound Scenar	io - Total Pro	ject Risk		
	Cobalt	3.08E-03	5.92E-05	2.13E-04	0.00E+00	7.01E-03	2.85E-03	5.77E-02	7.09E-02
	Copper	1.10E-04	1.93E-06	9.12E-06	0.00E+00	5.73E-03	1.06E-03	9.95E-02	1.06E-01
	Molybdenum	1.22E-02	8.46E-06	1.07E-03	0.00E+00	2.31E-04	7.01E-04	2.72E-01	2.86E-01
	Uranium	2.75E-02	8.15E-04	1.47E-02	0.00E+00	6.50E-02	1.09E-01	1.93E-01	4.10E-01

Table C15: Estimated Non-carcinogen Total Risk to Human Receptors – Operations – Reasonably Foreseeable Development Case

	RFD Case HQs										
	СОРС	Water (internal)	Soil (internal)	Sediment (internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC		
					Base	Case					
	Cobalt	8.58E-04	9.04E-07	1.01E-06	0.00E+00	1.15E-03	7.21E-04	1.97E-02	2.25E-02		
	Copper	2.91E-05	2.95E-08	4.12E-08	0.00E+00	8.94E-04	2.69E-04	3.89E-02	4.01E-02		
	Molybdenum	1.08E-04	1.06E-07	1.62E-07	0.00E+00	1.21E-06	1.46E-04	8.23E-02	8.25E-02		
	Uranium	2.46E-03	8.62E-06	2.24E-05	0.00E+00	3.43E-03	1.92E-02	7.29E-02	9.80E-02		
				Pro	ject Lifespan -	Total Project	Risk				
Camp Worker	Cobalt	8.63E-04	9.06E-07	1.02E-06	0.00E+00	1.33E-03	7.29E-04	2.01E-02	2.30E-02		
(Patterson Lake North	Copper	2.91E-05	3.00E-08	4.12E-08	0.00E+00	9.80E-04	2.70E-04	3.91E-02	4.03E-02		
Arm - West Basin)	Molybdenum	1.09E-04	1.08E-07	1.62E-07	0.00E+00	2.19E-06	1.49E-04	8.24E-02	8.26E-02		
	Uranium	2.46E-03	2.80E-05	2.24E-05	0.00E+00	1.06E-02	4.34E-02	8.81E-02	1.45E-01		
				<u> </u>	ar Future - To	tal Project Ri	sk				
	Cobalt	-	_	_	_	_	-	-	_		
	Copper	_	_	-	_	-	-	-	_		
	Molybdenum	_	_	_	_	-	-	-	_		
	Uranium	-	-	-	-	-	-	-	-		
					Base	Case					
	Cobalt	8.58E-04	9.04E-07	1.01E-06	0.00E+00	2.30E-03	1.44E-03	2.44E-02	2.90E-02		
	Copper	2.91E-05	2.95E-08	4.12E-08	0.00E+00	1.79E-03	5.38E-04	4.12E-02	4.35E-02		
	Molybdenum	1.08E-04	1.06E-07	1.62E-07	0.00E+00	2.42E-06	2.92E-04	6.98E-02	7.02E-02		
Subsistence Harvester	Uranium	2.46E-03	8.62E-06	2.24E-05	0.00E+00	6.85E-03	3.84E-02	8.20E-02	1.30E-01		
(Patterson Lake South				Pro	ject Lifespan -	Total Project	t Risk				
Arm)	Cobalt	1.00E-03	9.05E-07	1.12E-06	0.00E+00	2.65E-03	1.46E-03	2.51E-02	3.02E-02		
	Copper	3.20E-05	2.95E-08	4.37E-08	0.00E+00	1.96E-03	5.39E-04	4.15E-02	4.40E-02		
	Molybdenum	2.04E-04	1.07E-07	2.54E-07	0.00E+00	4.38E-06	2.98E-04	6.99E-02	7.05E-02		
	Uranium	7.81E-03	1.16E-05	3.71E-05	0.00E+00	2.12E-02	8.67E-02	1.12E-01	2.28E-01		
					ar Future - To	tal Project Ri	sk				



Table C15: Estimated Non-carcinogen Total Risk to Human Receptors – Operations – Reasonably Foreseeable Development Case

					RFD Ca	se HQs			
	СОРС	Water (internal)	Soil (internal)	Sediment (internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
	Cobalt	1.23E-03	9.05E-07	1.46E-06	0.00E+00	3.30E-03	1.44E-03	2.73E-02	3.33E-02
	Copper	4.27E-05	2.95E-08	6.04E-08	0.00E+00	2.62E-03	5.38E-04	4.19E-02	4.51E-02
	Molybdenum	7.79E-04	1.06E-07	1.17E-06	0.00E+00	1.74E-05	2.92E-04	7.05E-02	7.16E-02
	Uranium	5.52E-03	8.93E-06	5.04E-05	0.00E+00	1.54E-02	3.98E-02	8.64E-02	1.47E-01
					Base	Case			
	Cobalt	9.58E-04	5.91E-05	6.62E-05	0.00E+00	2.18E-03	2.84E-03	4.78E-02	5.39E-02
	Copper	3.25E-05	1.93E-06	2.69E-06	0.00E+00	1.69E-03	1.06E-03	9.70E-02	9.98E-02
	Molybdenum	1.47E-04	8.46E-06	1.29E-05	0.00E+00	2.79E-06	7.01E-04	2.64E-01	2.64E-01
	Uranium	2.75E-03	5.64E-04	1.47E-03	0.00E+00	6.49E-03	7.57E-02	1.70E-01	2.57E-01
				Pro	ject Lifespan -	Total Project	Risk		
Subsistence Harvester	Cobalt	1.12E-03	5.91E-05	7.34E-05	0.00E+00	2.52E-03	2.86E-03	4.85E-02	5.51E-02
One-year-old (Patterson	Copper	3.57E-05	1.93E-06	2.86E-06	0.00E+00	1.86E-03	1.06E-03	9.73E-02	1.00E-01
Lake South Arm)	Molybdenum	2.78E-04	8.47E-06	2.02E-05	0.00E+00	5.06E-06	7.11E-04	2.64E-01	2.65E-01
	Uranium	8.72E-03	7.56E-04	2.43E-03	0.00E+00	2.01E-02	1.48E-01	1.92E-01	3.73E-01
					ar Future - To	tal Project Ri	sk		
	Cobalt	1.38E-03	5.91E-05	9.54E-05	0.00E+00	3.13E-03	2.84E-03	5.08E-02	5.83E-02
	Copper	4.76E-05	1.93E-06	3.95E-06	0.00E+00	2.48E-03	1.06E-03	9.77E-02	1.01E-01
	Molybdenum	1.06E-03	8.46E-06	9.27E-05	0.00E+00	2.00E-05	7.01E-04	2.64E-01	2.66E-01
	Uranium	6.17E-03	5.84E-04	3.29E-03	0.00E+00	1.46E-02	7.84E-02	1.74E-01	2.77E-01
					Base	Case			
	Cobalt	8.58E-04	9.04E-07	1.01E-06	0.00E+00	2.30E-03	1.44E-03	2.44E-02	2.90E-02
Subsistence Harvester	Copper	2.91E-05	2.95E-08	4.12E-08	0.00E+00	1.79E-03	5.38E-04	4.12E-02	4.35E-02
(Beet Lake)	Molybdenum	1.08E-04	1.06E-07	1.62E-07	0.00E+00	2.42E-06	2.92E-04	6.98E-02	7.02E-02
(beet Lake)	Uranium	2.46E-03	8.62E-06	2.24E-05	0.00E+00	6.85E-03	3.84E-02	8.20E-02	1.30E-01
				Pro	ject Lifespan -	Total Project	Risk		
	Cobalt	9.22E-04	9.05E-07	1.07E-06	0.00E+00	2.45E-03	1.46E-03	2.47E-02	2.96E-02

Table C15: Estimated Non-carcinogen Total Risk to Human Receptors – Operations – Reasonably Foreseeable Development Case

	RFD Case HQs											
	СОРС	Water (internal)	Soil (internal)	Sediment (internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC			
	Copper	3.03E-05	2.95E-08	4.25E-08	0.00E+00	1.86E-03	5.39E-04	4.12E-02	4.37E-02			
	Molybdenum	1.51E-04	1.07E-07	2.10E-07	0.00E+00	3.30E-06	2.98E-04	6.98E-02	7.03E-02			
	Uranium	4.32E-03	1.03E-05	2.79E-05	0.00E+00	1.18E-02	6.54E-02	9.34E-02	1.75E-01			
					ar Future - To	tal Project Ri	sk					
	Cobalt	1.06E-03	9.05E-07	1.25E-06	0.00E+00	2.83E-03	1.44E-03	2.54E-02	3.08E-02			
	Copper	3.62E-05	2.95E-08	5.13E-08	0.00E+00	2.22E-03	5.38E-04	4.15E-02	4.43E-02			
	Molybdenum	4.57E-04	1.06E-07	6.85E-07	0.00E+00	1.02E-05	2.92E-04	7.00E-02	7.08E-02			
	Uranium	3.60E-03	8.80E-06	3.28E-05	0.00E+00	1.00E-02	3.92E-02	8.35E-02	1.36E-01			
		Base Case										
	Cobalt	9.58E-04	5.91E-05	6.62E-05	0.00E+00	2.18E-03	2.84E-03	4.78E-02	5.39E-02			
	Copper	3.25E-05	1.93E-06	2.69E-06	0.00E+00	1.69E-03	1.06E-03	9.70E-02	9.98E-02			
	Molybdenum	1.47E-04	8.46E-06	1.29E-05	0.00E+00	2.79E-06	7.01E-04	2.64E-01	2.64E-01			
	Uranium	2.75E-03	5.64E-04	1.47E-03	0.00E+00	6.49E-03	7.57E-02	1.70E-01	2.57E-01			
	Project Lifespan - Total Project Risk											
Culturation on Hammandam	Cobalt	1.03E-03	5.91E-05	7.01E-05	0.00E+00	2.33E-03	2.86E-03	4.81E-02	5.45E-02			
Subsistence Harvester One-year-old (Beet Lake)	Copper	3.39E-05	1.93E-06	2.78E-06	0.00E+00	1.76E-03	1.06E-03	9.71E-02	9.99E-02			
One-year-old (beet Lake)	Molybdenum	2.06E-04	8.47E-06	1.67E-05	0.00E+00	3.80E-06	7.11E-04	2.64E-01	2.65E-01			
	Uranium	4.82E-03	6.71E-04	1.82E-03	0.00E+00	1.12E-02	1.16E-01	1.78E-01	3.13E-01			
				ı	ar Future - To	tal Project Ri	sk					
	Cobalt	1.18E-03	5.91E-05	8.17E-05	0.00E+00	2.68E-03	2.84E-03	4.89E-02	5.57E-02			
	Copper	4.04E-05	1.93E-06	3.35E-06	0.00E+00	2.11E-03	1.06E-03	9.73E-02	1.00E-01			
	Molybdenum	6.22E-04	8.46E-06	5.45E-05	0.00E+00	1.18E-05	7.01E-04	2.64E-01	2.65E-01			
	Uranium	4.02E-03	5.75E-04	2.14E-03	0.00E+00	9.50E-03	7.72E-02	1.71E-01	2.65E-01			
Culturation on Harmon					Base	Case						
Subsistence Harvester (Lloyd Lake)	Cobalt	8.58E-04	9.04E-07	1.01E-06	0.00E+00	2.30E-03	1.44E-03	2.44E-02	2.90E-02			
(Libya Lake)	Copper	2.91E-05	2.95E-08	4.12E-08	0.00E+00	1.79E-03	5.38E-04	4.12E-02	4.35E-02			



Table C15: Estimated Non-carcinogen Total Risk to Human Receptors – Operations – Reasonably Foreseeable Development Case

		RFD Case HQs									
	СОРС	Water (internal)	Soil (internal)	Sediment (internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC		
	Molybdenum	1.08E-04	1.06E-07	1.62E-07	0.00E+00	2.42E-06	2.92E-04	6.98E-02	7.02E-02		
	Uranium	2.46E-03	8.62E-06	2.24E-05	0.00E+00	6.85E-03	3.84E-02	8.20E-02	1.30E-01		
				Pro	ject Lifespan -	Total Project	t Risk				
	Cobalt	8.64E-04	9.05E-07	1.02E-06	0.00E+00	2.31E-03	1.46E-03	2.45E-02	2.91E-02		
	Copper	2.92E-05	2.95E-08	4.14E-08	0.00E+00	1.79E-03	5.39E-04	4.12E-02	4.36E-02		
	Molybdenum	1.13E-04	1.07E-07	1.67E-07	0.00E+00	2.51E-06	2.98E-04	6.98E-02	7.02E-02		
	Uranium	2.63E-03	8.64E-06	2.29E-05	0.00E+00	7.31E-03	3.87E-02	8.22E-02	1.31E-01		
					Far Future - To	tal Project Ri	sk				
	Cobalt	8.78E-04	9.05E-07	1.04E-06	0.00E+00	2.35E-03	1.44E-03	2.45E-02	2.92E-02		
	Copper	2.98E-05	2.95E-08	4.22E-08	0.00E+00	1.83E-03	5.38E-04	4.12E-02	4.36E-02		
	Molybdenum	1.44E-04	1.06E-07	2.15E-07	0.00E+00	3.21E-06	2.92E-04	6.98E-02	7.02E-02		
	Uranium	2.56E-03	8.63E-06	2.34E-05	0.00E+00	7.14E-03	3.84E-02	8.21E-02	1.30E-01		
					Base	Case					
	Cobalt	9.58E-04	5.91E-05	6.62E-05	0.00E+00	2.18E-03	2.84E-03	4.78E-02	5.39E-02		
	Copper	3.25E-05	1.93E-06	2.69E-06	0.00E+00	1.69E-03	1.06E-03	9.70E-02	9.98E-02		
	Molybdenum	1.47E-04	8.46E-06	1.29E-05	0.00E+00	2.79E-06	7.01E-04	2.64E-01	2.64E-01		
	Uranium	2.75E-03	5.64E-04	1.47E-03	0.00E+00	6.49E-03	7.57E-02	1.70E-01	2.57E-01		
				Pro	ject Lifespan -	Total Project	t Risk				
Subsistence Harvester	Cobalt	9.65E-04	5.91E-05	6.66E-05	0.00E+00	2.19E-03	2.86E-03	4.79E-02	5.40E-02		
One-year-old (Lloyd Lake)	Copper	3.26E-05	1.93E-06	2.70E-06	0.00E+00	1.70E-03	1.06E-03	9.70E-02	9.98E-02		
	Molybdenum	1.53E-04	8.47E-06	1.33E-05	0.00E+00	2.89E-06	7.11E-04	2.64E-01	2.64E-01		
	Uranium	2.94E-03	5.65E-04	1.50E-03	0.00E+00	6.93E-03	7.61E-02	1.70E-01	2.59E-01		
				ا	Far Future - To	tal Project Ri	sk				
	Cobalt	9.80E-04	5.91E-05	6.78E-05	0.00E+00	2.23E-03	2.84E-03	4.79E-02	5.41E-02		
	Copper	3.33E-05	1.93E-06	2.76E-06	0.00E+00	1.74E-03	1.06E-03	9.70E-02	9.99E-02		
	Molybdenum	1.96E-04	8.46E-06	1.71E-05	0.00E+00	3.71E-06	7.01E-04	2.64E-01	2.64E-01		



Table C15: Estimated Non-carcinogen Total Risk to Human Receptors – Operations – Reasonably Foreseeable Development Case

					RFD Ca	se HQs			
	СОРС	Water (internal)	Soil (internal)	Sediment (internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
	Uranium	2.86E-03	5.64E-04	1.53E-03	0.00E+00	6.77E-03	7.57E-02	1.70E-01	2.58E-01
				•	Base	Case			
	Cobalt	8.58E-04	9.04E-07	1.01E-06	0.00E+00	7.27E-04	1.13E-03	1.88E-02	2.15E-02
	Copper	2.91E-05	2.95E-08	4.12E-08	0.00E+00	5.66E-04	4.22E-04	3.86E-02	3.96E-02
	Molybdenum	1.08E-04	1.06E-07	1.62E-07	0.00E+00	7.66E-07	2.29E-04	8.60E-02	8.63E-02
	Uranium	2.46E-03	8.62E-06	2.24E-05	0.00E+00	2.17E-03	3.01E-02	6.79E-02	1.03E-01
				Pro	ject Lifespan -	Total Project	t Risk		
Seasonal Resident	Cobalt	9.45E-04	9.05E-07	1.08E-06	0.00E+00	7.96E-04	1.13E-03	1.89E-02	2.18E-02
(Patterson Lake South	Copper	3.08E-05	2.95E-08	4.27E-08	0.00E+00	5.99E-04	4.22E-04	3.87E-02	3.97E-02
Arm)	Molybdenum	1.66E-04	1.06E-07	2.17E-07	0.00E+00	1.14E-06	2.31E-04	8.60E-02	8.64E-02
	Uranium	5.67E-03	1.04E-05	3.13E-05	0.00E+00	4.90E-03	4.52E-02	7.36E-02	1.29E-01
				i	ar Future - To	tal Project Ri	sk		
	Cobalt	1.08E-03	9.04E-07	1.28E-06	0.00E+00	9.19E-04	1.13E-03	1.95E-02	2.26E-02
	Copper	3.72E-05	2.95E-08	5.27E-08	0.00E+00	7.24E-04	4.22E-04	3.88E-02	4.00E-02
	Molybdenum	5.11E-04	1.06E-07	7.64E-07	0.00E+00	3.61E-06	2.29E-04	8.62E-02	8.69E-02
	Uranium	4.30E-03	8.81E-06	3.92E-05	0.00E+00	3.79E-03	3.08E-02	6.90E-02	1.08E-01
					Base	Case			
	Cobalt	9.58E-04	5.91E-05	6.62E-05	0.00E+00	6.89E-04	2.47E-03	4.13E-02	4.56E-02
	Copper	3.25E-05	1.93E-06	2.69E-06	0.00E+00	5.36E-04	9.22E-04	9.32E-02	9.47E-02
Seasonal Resident One-	Molybdenum	1.47E-04	8.46E-06	1.29E-05	0.00E+00	8.84E-07	6.10E-04	2.77E-01	2.78E-01
year-old (Patterson Lake	Uranium	2.75E-03	5.64E-04	1.47E-03	0.00E+00	2.06E-03	6.58E-02	1.60E-01	2.33E-01
South Arm)				•	ject Lifespan -	•		-	
	Cobalt	1.06E-03	5.91E-05	7.06E-05	0.00E+00	7.54E-04	2.48E-03	4.15E-02	4.59E-02
	Copper	3.44E-05	1.93E-06	2.79E-06	0.00E+00	5.68E-04	9.23E-04	9.32E-02	9.48E-02
	Molybdenum	2.26E-04	8.47E-06	1.73E-05	0.00E+00	1.32E-06	6.13E-04	2.77E-01	2.78E-01
	Uranium	6.33E-03	6.79E-04	2.04E-03	0.00E+00	4.64E-03	9.28E-02	1.65E-01	2.71E-01

Table C15: Estimated Non-carcinogen Total Risk to Human Receptors – Operations – Reasonably Foreseeable Development Case

	RFD Case HQs											
	СОРС	Water (internal)	Soil (internal)	Sediment (internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC			
				F	ar Future - To	tal Project Ri	sk					
	Cobalt	1.21E-03	5.91E-05	8.37E-05	0.00E+00	8.71E-04	2.47E-03	4.21E-02	4.68E-02			
	Copper	4.16E-05	1.93E-06	3.45E-06	0.00E+00	6.86E-04	9.22E-04	9.33E-02	9.50E-02			
	Molybdenum	6.94E-04	8.46E-06	6.08E-05	0.00E+00	4.16E-06	6.10E-04	2.78E-01	2.79E-01			
	Uranium	4.80E-03	5.76E-04	2.56E-03	0.00E+00	3.60E-03	6.72E-02	1.61E-01	2.40E-01			
					Base	Case						
	Cobalt	8.58E-04	9.04E-07	1.01E-06	0.00E+00	7.27E-04	1.13E-03	1.88E-02	2.15E-02			
	Copper	2.91E-05	2.95E-08	4.12E-08	0.00E+00	5.66E-04	4.22E-04	3.86E-02	3.96E-02			
	Molybdenum	1.08E-04	1.06E-07	1.62E-07	0.00E+00	7.66E-07	2.29E-04	8.60E-02	8.63E-02			
	Uranium	2.46E-03	8.62E-06	2.24E-05	0.00E+00	2.17E-03	3.01E-02	6.79E-02	1.03E-01			
		Project Lifespan - Total Project Risk										
Seasonal Resident (Lloyd	Cobalt	8.62E-04	9.05E-07	1.02E-06	0.00E+00	7.33E-04	1.13E-03	1.88E-02	2.15E-02			
Lake)	Copper	2.92E-05	2.95E-08	4.13E-08	0.00E+00	7.54E-04	4.22E-04	3.86E-02	3.98E-02			
Lukey	Molybdenum	1.11E-04	1.06E-07	1.65E-07	0.00E+00	7.95E-07	2.31E-04	8.60E-02	8.64E-02			
	Uranium	2.56E-03	8.64E-06	2.27E-05	0.00E+00	2.88E-03	3.02E-02	6.80E-02	1.04E-01			
		_			ar Future - To	tal Project Ri	sk					
	Cobalt	8.70E-04	9.04E-07	1.03E-06	0.00E+00	7.45E-04	1.13E-03	1.88E-02	2.16E-02			
	Copper	2.95E-05	2.95E-08	4.18E-08	0.00E+00	7.70E-04	4.22E-04	3.86E-02	3.99E-02			
	Molybdenum	1.30E-04	1.06E-07	1.94E-07	0.00E+00	1.02E-06	2.29E-04	8.60E-02	8.64E-02			
	Uranium	2.52E-03	8.63E-06	2.30E-05	0.00E+00	2.81E-03	3.01E-02	6.80E-02	1.03E-01			
						Case						
	Cobalt	9.58E-04	5.91E-05	6.62E-05	0.00E+00	6.89E-04	2.47E-03	4.13E-02	4.56E-02			
Seasonal Resident One-	Copper	3.25E-05	1.93E-06	2.69E-06	0.00E+00	5.36E-04	9.22E-04	9.32E-02	9.47E-02			
year-old (Lloyd Lake)	Molybdenum	1.47E-04	8.46E-06	1.29E-05	0.00E+00	8.84E-07	6.10E-04	2.77E-01	2.78E-01			
	Uranium	2.75E-03	5.64E-04	1.47E-03	0.00E+00	2.06E-03	6.58E-02	1.60E-01	2.33E-01			
				Pro	ject Lifespan -	Total Projec	t Risk					



Table C15: Estimated Non-carcinogen Total Risk to Human Receptors – Operations – Reasonably Foreseeable Development Case

					RFD Ca	se HQs			
	СОРС	Water (internal)	Soil (internal)	Sediment (internal)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial Animals	Total by COPC
	Cobalt	9.62E-04	5.91E-05	6.65E-05	0.00E+00	6.92E-04	2.48E-03	4.14E-02	4.56E-02
	Copper	3.26E-05	1.93E-06	2.70E-06	0.00E+00	5.38E-04	9.23E-04	9.32E-02	9.47E-02
	Molybdenum	1.51E-04	8.47E-06	1.31E-05	0.00E+00	9.03E-07	6.13E-04	2.77E-01	2.78E-01
	Uranium	2.86E-03	5.64E-04	1.49E-03	0.00E+00	2.14E-03	6.60E-02	1.60E-01	2.33E-01
				F	ar Future - To	tal Project Ri	sk	,	
	Cobalt	9.71E-04	5.91E-05	6.72E-05	0.00E+00	6.99E-04	2.47E-03	4.14E-02	4.56E-02
	Copper	3.30E-05	1.93E-06	2.73E-06	0.00E+00	5.44E-04	9.22E-04	9.32E-02	9.47E-02
	Molybdenum	1.76E-04	8.46E-06	1.54E-05	0.00E+00	1.06E-06	6.10E-04	2.77E-01	2.78E-01
	Uranium	2.82E-03	5.64E-04	1.50E-03	0.00E+00	2.11E-03	6.58E-02	1.60E-01	2.33E-01
					Base Ca	se Adult			
	Cobalt	8.58E-04	9.04E-07	1.01E-06	0.00E+00	2.30E-03	1.44E-03	2.44E-02	2.90E-02
	Copper	2.91E-05	2.95E-08	4.12E-08	0.00E+00	1.79E-03	5.38E-04	4.12E-02	4.35E-02
	Molybdenum	1.08E-04	1.06E-07	1.62E-07	0.00E+00	2.42E-06	2.92E-04	6.98E-02	7.02E-02
	Uranium	2.46E-03	8.62E-06	2.24E-05	0.00E+00	6.85E-03	3.84E-02	8.20E-02	1.30E-01
					Base Case O	ne-year-old			
	Cobalt	9.58E-04	5.91E-05	6.62E-05	0.00E+00	2.18E-03	2.84E-03	4.78E-02	5.39E-02
Permanent Resident	Copper	3.25E-05	1.93E-06	2.69E-06	0.00E+00	1.69E-03	1.06E-03	9.70E-02	9.98E-02
(Patterson Lake North	Molybdenum	1.47E-04	8.46E-06	1.29E-05	0.00E+00	2.79E-06	7.01E-04	2.64E-01	2.64E-01
Arm - West Basin)	Uranium	2.75E-03	5.64E-04	1.47E-03	0.00E+00	6.49E-03	7.57E-02	1.70E-01	2.57E-01
				Far	future Adult -	Total Project	t Risk		
	Cobalt	2.10E-03	9.05E-07	2.48E-06	0.00E+00	5.62E-03	1.44E-03	3.08E-02	3.99E-02
	Copper	7.53E-05	2.96E-08	1.07E-07	0.00E+00	4.62E-03	5.39E-04	4.30E-02	4.82E-02
	Molybdenum	2.42E-03	1.06E-07	3.62E-06	0.00E+00	5.39E-05	2.92E-04	7.16E-02	7.43E-02
	Uranium	2.46E-02	1.25E-05	2.24E-04	0.00E+00	6.85E-02	5.57E-02	1.13E-01	2.62E-01
				Far futu	ire One-year-o	old - Total Pro	oject Risk		
	Cobalt	2.34E-03	5.92E-05	1.62E-04	0.00E+00	5.33E-03	2.84E-03	5.43E-02	6.50E-02

Table C15: Estimated Non-carcinogen Total Risk to Human Receptors – Operations – Reasonably Foreseeable Development Case

				RFD Ca	se HQs			
СОРС	Water	Soil	Sediment	Aquatic	Aquatic	Terrestrial	Terrestrial	Total by
COPC	(internal)	(internal)	(internal)	Plants	Animals	Plants	Animals	COPC
Copper	8.41E-05	1.93E-06	6.97E-06	0.00E+00	4.38E-03	1.06E-03	9.87E-02	1.04E-01
Molybdenum	3.29E-03	8.46E-06	2.88E-04	0.00E+00	6.23E-05	7.01E-04	2.66E-01	2.70E-01
Uranium	2.75E-02	8.17E-04	1.47E-02	0.00E+00	6.50E-02	1.10E-01	1.93E-01	4.10E-01



Table C16: Estimated Total Radiation Doses to Human Receptors – Operations – Application Case, Upper Bound Scenario

						Tota	l Dose by Pat	hway - Proje	ct Lifespan (r	nSv/yr)				
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
		•	•			•		Base Case				•		
	Uranium-238	0.00E+00	0.00E+00	6.08E-06	4.61E-11	7.42E-08	8.23E-04	5.54E-08	5.91E-07	0.00E+00	8.47E-06	1.65E-04	4.15E-04	1.42E-03
	Uranium-234	0.00E+00	0.00E+00	6.62E-06	8.04E-12	8.08E-08	4.15E-06	6.04E-08	2.14E-09	0.00E+00	9.22E-06	1.80E-04	4.52E-04	6.52E-04
	Thorium-230	0.00E+00	0.00E+00	7.25E-04	4.91E-10	2.27E-06	3.47E-05	7.89E-07	1.95E-08	0.00E+00	4.85E-04	9.73E-05	2.70E-03	4.05E-03
	Radium-226	0.00E+00	0.00E+00	5.89E-04	2.63E-07	2.71E-06	7.02E-02	3.26E-06	3.91E-04	0.00E+00	7.86E-04	2.24E-03	9.32E-03	8.35E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	5.91E-03	1.64E-03	2.93E-02	4.17E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	3.73E-02	3.86E-03	1.32E-01	1.75E-01
	Total by Pathway	0.00E+00	0.00E+00	5.00E-03	2.64E-07	2.73E-05	7.45E-02	1.28E-04	3.94E-04	0.00E+00	4.45E-02	8.18E-03	1.74E-01	0.31
							1	Application C	ase					
	Uranium-238	8.41E-04	2.31E-12	6.10E-06	4.63E-11	2.07E-07	2.29E-03	5.55E-08	5.92E-07	0.00E+00	1.96E-05	2.12E-04	4.77E-04	3.85E-03
	Uranium-234	1.01E-03	5.64E-12	6.64E-06	8.07E-12	2.25E-07	1.15E-05	6.04E-08	2.15E-09	0.00E+00	2.14E-05	2.31E-04	5.20E-04	1.81E-03
Camp Worker (Patterson Lake North	Thorium-230	4.06E-03	1.37E-11	7.26E-04	4.92E-10	2.96E-06	4.53E-05	7.90E-07	1.95E-08	0.00E+00	1.15E-03	6.07E-04	9.26E-03	1.58E-02
Arm - West Basin)	Radium-226	1.01E-03	2.62E-10	5.89E-04	2.63E-07	3.65E-06	9.44E-02	3.26E-06	3.91E-04	0.00E+00	8.86E-04	3.12E-03	1.01E-02	1.10E-01
	Lead-210	3.19E-04	4.12E-11	1.33E-03	1.17E-09	1.64E-05	3.80E-03	4.46E-05	2.53E-06	0.00E+00	6.00E-03	3.54E-03	3.57E-02	5.07E-02
	Polonium-210	9.56E-04	3.59E-13	2.35E-03	7.06E-12	9.90E-06	2.93E-07	7.90E-05	1.08E-08	0.00E+00	3.79E-02	6.90E-03	1.41E-01	1.89E-01
	Total by Pathway	8.20E-03	3.25E-10	5.01E-03	2.65E-07	3.33E-05	1.01E-01	1.28E-04	3.94E-04	0.00E+00	4.59E-02	1.46E-02	1.97E-01	0.37
							Up _l	oer Bound Sc	enario					
	Uranium-238	8.41E-04	2.31E-12	6.10E-06	4.63E-11	2.07E-07	2.29E-03	5.55E-08	5.92E-07	0.00E+00	3.90E-05	2.12E-04	4.82E-04	3.87E-03
	Uranium-234	1.01E-03	5.64E-12	6.64E-06	8.07E-12	2.25E-07	1.15E-05	6.04E-08	2.15E-09	0.00E+00	4.25E-05	2.31E-04	5.25E-04	1.83E-03
	Thorium-230	4.06E-03	1.37E-11	7.26E-04	4.92E-10	2.96E-06	4.53E-05	7.90E-07	1.95E-08	0.00E+00	1.16E-03	6.07E-04	9.31E-03	1.59E-02
	Radium-226	1.01E-03	2.62E-10	5.89E-04	2.63E-07	3.65E-06	9.44E-02	3.26E-06	3.91E-04	0.00E+00	8.90E-04	3.12E-03	1.01E-02	1.10E-01
	Lead-210	3.19E-04	4.12E-11	1.33E-03	1.17E-09	1.64E-05	3.80E-03	4.46E-05	2.53E-06	0.00E+00	6.36E-03	3.54E-03	4.51E-02	6.06E-02
	Polonium-210	9.56E-04	3.59E-13	2.35E-03	7.06E-12	9.90E-06	2.93E-07	7.90E-05	1.08E-08	0.00E+00	4.02E-02	6.90E-03	1.46E-01	1.96E-01
	Total by Pathway	8.20E-03	3.25E-10	5.01E-03	2.65E-07	3.33E-05	1.01E-01	1.28E-04	3.94E-04	0.00E+00	4.87E-02	1.46E-02	2.11E-01	0.39
				,	,			Base Case	,					
	Uranium-238	0.00E+00	0.00E+00	6.08E-06	4.61E-11	7.42E-08	8.23E-04	5.54E-08	5.91E-07	0.00E+00	1.69E-05	3.30E-04	5.22E-04	1.70E-03
	Uranium-234	0.00E+00	0.00E+00	6.62E-06	8.04E-12	8.08E-08	4.15E-06	6.04E-08	2.14E-09	0.00E+00	1.84E-05	3.60E-04	5.68E-04	9.57E-04
Subsistence Harvester (Patterson Lake South	Thorium-230	0.00E+00	0.00E+00	7.25E-04	4.91E-10	2.27E-06	3.47E-05	7.89E-07	1.95E-08	0.00E+00	9.70E-04	1.94E-04	4.54E-03	6.47E-03
Arm)	Radium-226	0.00E+00	0.00E+00	5.89E-04	2.63E-07	2.71E-06	7.02E-02	3.26E-06	3.91E-04	0.00E+00	1.57E-03	4.47E-03	1.03E-02	8.76E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	1.18E-02	3.28E-03	3.96E-02	5.95E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	7.46E-02	7.71E-03	2.23E-01	3.08E-01
	Total by Pathway	0.00E+00	0.00E+00	5.00E-03	2.64E-07	2.73E-05	7.45E-02	1.28E-04	3.94E-04	0.00E+00	8.90E-02	1.63E-02	2.78E-01	0.46

Table C16: Estimated Total Radiation Doses to Human Receptors – Operations – Application Case, Upper Bound Scenario

						Tota	l Dose by Pat	:hway - Proje	ct Lifespan (r	mSv/yr)				
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
			•					Application C	ase			•		•
	Uranium-238	1.30E-04	3.57E-13	1.42E-05	1.08E-10	9.47E-08	1.05E-03	1.14E-07	1.22E-06	0.00E+00	3.92E-05	4.24E-04	6.47E-04	2.31E-03
	Uranium-234	1.57E-04	8.74E-13	1.55E-05	1.88E-11	1.03E-07	5.30E-06	1.25E-07	4.42E-09	0.00E+00	4.28E-05	4.62E-04	7.04E-04	1.39E-03
	Thorium-230	6.29E-04	2.12E-12	1.72E-03	1.17E-09	2.38E-06	3.64E-05	1.81E-06	4.48E-08	0.00E+00	2.29E-03	1.21E-03	1.76E-02	2.35E-02
	Radium-226	1.57E-04	4.06E-11	6.66E-04	2.97E-07	2.86E-06	7.40E-02	3.60E-06	4.32E-04	0.00E+00	1.77E-03	6.23E-03	1.18E-02	9.50E-02
	Lead-210	4.94E-05	6.39E-12	1.35E-03	1.19E-09	1.51E-05	3.51E-03	4.52E-05	2.56E-06	0.00E+00	1.20E-02	7.08E-03	5.23E-02	7.63E-02
	Polonium-210	1.48E-04	5.57E-14	2.38E-03	7.16E-12	7.68E-06	2.28E-07	7.99E-05	1.10E-08	0.00E+00	7.57E-02	1.38E-02	2.41E-01	3.33E-01
	Total by Pathway	1.27E-03	5.04E-11	6.15E-03	3.00E-07	2.82E-05	7.86E-02	1.31E-04	4.35E-04	0.00E+00	9.18E-02	2.92E-02	3.24E-01	0.53
							Upj	oer Bound Sc	enario					
	Uranium-238	1.30E-04	3.57E-13	2.82E-05	2.14E-10	9.47E-08	1.05E-03	2.19E-07	2.34E-06	0.00E+00	7.79E-05	4.24E-04	6.57E-04	2.37E-03
	Uranium-234	1.57E-04	8.74E-13	3.08E-05	3.74E-11	1.03E-07	5.30E-06	2.39E-07	8.49E-09	0.00E+00	8.50E-05	4.62E-04	7.15E-04	1.46E-03
	Thorium-230	6.29E-04	2.12E-12	1.74E-03	1.18E-09	2.38E-06	3.64E-05	1.82E-06	4.51E-08	0.00E+00	2.31E-03	1.21E-03	1.77E-02	2.36E-02
	Radium-226	1.57E-04	4.06E-11	6.69E-04	2.99E-07	2.86E-06	7.40E-02	3.62E-06	4.33E-04	0.00E+00	1.78E-03	6.23E-03	1.18E-02	9.50E-02
	Lead-210	4.94E-05	6.39E-12	1.43E-03	1.26E-09	1.51E-05	3.51E-03	4.72E-05	2.67E-06	0.00E+00	1.27E-02	7.08E-03	7.12E-02	9.60E-02
	Polonium-210	1.48E-04	5.57E-14	2.53E-03	7.61E-12	7.68E-06	2.28E-07	8.35E-05	1.15E-08	0.00E+00	8.03E-02	1.38E-02	2.51E-01	3.48E-01
	Total by Pathway	1.27E-03	5.04E-11	6.43E-03	3.01E-07	2.82E-05	7.86E-02	1.37E-04	4.39E-04	0.00E+00	9.72E-02	2.92E-02	3.54E-01	0.57
						,		Base Case		,				
	Uranium-238	0.00E+00	0.00E+00	4.22E-06	1.14E-11	3.02E-06	1.10E-03	2.25E-06	7.69E-07	0.00E+00	9.99E-06	4.05E-04	5.67E-04	2.09E-03
	Uranium-234	0.00E+00	0.00E+00	4.58E-06	1.99E-12	3.27E-06	5.56E-06	2.44E-06	2.79E-09	0.00E+00	1.08E-05	4.39E-04	6.15E-04	1.08E-03
	Thorium-230	0.00E+00	0.00E+00	3.69E-04	1.22E-10	6.75E-05	3.95E-05	2.35E-05	2.53E-08	0.00E+00	4.19E-04	1.75E-04	2.77E-03	3.86E-03
	Radium-226	0.00E+00	0.00E+00	5.26E-04	6.50E-08	1.42E-04	9.40E-02	1.71E-04	5.10E-04	0.00E+00	1.19E-03	7.05E-03	1.67E-02	1.20E-01
	Lead-210	0.00E+00	0.00E+00	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	1.36E-02	7.86E-03	8.28E-02	1.15E-01
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	1.21E-01	2.60E-02	3.94E-01	5.56E-01
Subsistence Harvester One-Year-Old (Patterson	Total by Pathway	0.00E+00	0.00E+00	7.19E-03	6.54E-08	2.21E-03	9.97E-02	1.26E-02	5.14E-04	0.00E+00	1.36E-01	4.20E-02	4.98E-01	0.80
Lake South Arm)			1				1	Application C	ase		1	1		
	Uranium-238	1.30E-04	4.65E-13	9.85E-06	2.66E-11	3.85E-06	1.41E-03	4.64E-06	1.58E-06	0.00E+00	2.31E-05	5.21E-04	6.24E-04	2.72E-03
	Uranium-234	1.52E-04	1.14E-12	1.07E-05	4.65E-12	4.17E-06	7.09E-06	5.04E-06	5.75E-09	0.00E+00	2.51E-05	5.64E-04	6.76E-04	1.44E-03
	Thorium-230	4.84E-04	2.75E-12	8.76E-04	2.89E-10	7.07E-05	4.14E-05	5.39E-05	5.82E-08	0.00E+00	9.90E-04	7.31E-04	9.09E-03	1.23E-02
	Radium-226	1.52E-04	5.27E-11	5.95E-04	7.36E-08	1.49E-04	9.90E-02	1.88E-04	5.63E-04	0.00E+00	1.34E-03	9.05E-03	1.74E-02	1.28E-01
	Lead-210	5.11E-05	8.29E-12	1.83E-03	2.41E-10	1.20E-03	4.70E-03	3.59E-03	3.32E-06	0.00E+00	1.38E-02	1.41E-02	9.85E-02	1.38E-01
	Polonium-210	1.52E-04	7.25E-14	4.55E-03	1.77E-12	8.59E-04	2.97E-07	8.94E-03	1.42E-08	0.00E+00	1.23E-01	4.04E-02	4.13E-01	5.90E-01
	Total by Pathway	1.12E-03	6.55E-11	7.88E-03	7.41E-08	2.29E-03	1.05E-01	1.28E-02	5.68E-04	0.00E+00	1.39E-01	6.53E-02	5.39E-01	0.87

Table C16: Estimated Total Radiation Doses to Human Receptors – Operations – Application Case, Upper Bound Scenario

						Tota	l Dose by Pat	hway - Proje	ct Lifespan (ı	mSv/yr)				
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
							Up _l	per Bound Sc	enario					
	Uranium-238	1.30E-04	4.65E-13	1.96E-05	5.29E-11	3.85E-06	1.41E-03	8.90E-06	3.04E-06	0.00E+00	4.59E-05	5.21E-04	6.28E-04	2.77E-03
	Uranium-234	1.52E-04	1.14E-12	2.13E-05	9.27E-12	4.17E-06	7.09E-06	9.67E-06	1.10E-08	0.00E+00	4.99E-05	5.64E-04	6.81E-04	1.49E-03
	Thorium-230	4.84E-04	2.75E-12	8.83E-04	2.91E-10	7.07E-05	4.14E-05	5.43E-05	5.86E-08	0.00E+00	9.98E-04	7.31E-04	9.14E-03	1.24E-02
	Radium-226	1.52E-04	5.27E-11	5.98E-04	7.39E-08	1.49E-04	9.90E-02	1.89E-04	5.66E-04	0.00E+00	1.35E-03	9.05E-03	1.74E-02	1.28E-01
	Lead-210	5.11E-05	8.29E-12	1.94E-03	2.56E-10	1.20E-03	4.70E-03	3.76E-03	3.47E-06	0.00E+00	1.47E-02	1.41E-02	1.22E-01	1.63E-01
	Polonium-210	1.52E-04	7.25E-14	4.84E-03	1.88E-12	8.59E-04	2.97E-07	9.34E-03	1.49E-08	0.00E+00	1.30E-01	4.04E-02	4.24E-01	6.10E-01
	Total by Pathway	1.12E-03	6.55E-11	8.30E-03	7.46E-08	2.29E-03	1.05E-01	1.34E-02	5.72E-04	0.00E+00	1.47E-01	6.53E-02	5.74E-01	0.92
								Base Case						
	Uranium-238	0.00E+00	0.00E+00	6.08E-06	4.61E-11	7.42E-08	8.23E-04	5.54E-08	5.91E-07	0.00E+00	1.69E-05	3.30E-04	5.22E-04	1.70E-03
	Uranium-234	0.00E+00	0.00E+00	6.62E-06	8.04E-12	8.08E-08	4.15E-06	6.04E-08	2.14E-09	0.00E+00	1.84E-05	3.60E-04	5.68E-04	9.57E-04
	Thorium-230	0.00E+00	0.00E+00	7.25E-04	4.91E-10	2.27E-06	3.47E-05	7.89E-07	1.95E-08	0.00E+00	9.70E-04	1.94E-04	4.54E-03	6.47E-03
	Radium-226	0.00E+00	0.00E+00	5.89E-04	2.63E-07	2.71E-06	7.02E-02	3.26E-06	3.91E-04	0.00E+00	1.57E-03	4.47E-03	1.03E-02	8.76E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	1.18E-02	3.28E-03	3.96E-02	5.95E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	7.46E-02	7.71E-03	2.23E-01	3.08E-01
	Total by Pathway	0.00E+00	0.00E+00	5.00E-03	2.64E-07	2.73E-05	7.45E-02	1.28E-04	3.94E-04	0.00E+00	8.90E-02	1.63E-02	2.78E-01	0.46
							1	Application C	ase					
	Uranium-238	7.59E-05	2.08E-13	9.05E-06	6.87E-11	8.62E-08	9.56E-04	7.70E-08	8.21E-07	0.00E+00	2.51E-05	3.85E-04	5.71E-04	2.02E-03
	Uranium-234	9.16E-05	5.10E-13	9.87E-06	1.20E-11	9.39E-08	4.82E-06	8.39E-08	2.98E-09	0.00E+00	2.74E-05	4.19E-04	6.22E-04	1.17E-03
Subsistence Harvester	Thorium-230	3.67E-04	1.24E-12	1.25E-03	8.46E-10	2.33E-06	3.57E-05	1.33E-06	3.28E-08	0.00E+00	1.66E-03	7.87E-04	7.10E-03	1.12E-02
(Beet Lake)	Radium-226	9.16E-05	2.36E-11	6.21E-04	2.77E-07	2.80E-06	7.24E-02	3.41E-06	4.08E-04	0.00E+00	1.66E-03	5.50E-03	1.07E-02	9.14E-02
	Lead-210	2.88E-05	3.72E-12	1.33E-03	1.17E-09	1.50E-05	3.49E-03	4.47E-05	2.53E-06	0.00E+00	1.18E-02	5.49E-03	4.06E-02	6.28E-02
	Polonium-210	8.64E-05	3.25E-14	2.35E-03	7.07E-12	7.51E-06	2.23E-07	7.91E-05	1.08E-08	0.00E+00	7.47E-02	1.13E-02	2.28E-01	3.17E-01
	Total by Pathway	7.41E-04	2.94E-11	5.57E-03	2.79E-07	2.79E-05	7.69E-02	1.29E-04	4.12E-04	0.00E+00	8.99E-02	2.38E-02	2.88E-01	0.49
							Up _l	oer Bound Sc	enario					
	Uranium-238	7.59E-05	2.08E-13	1.42E-05	1.08E-10	8.62E-08	9.56E-04	1.15E-07	1.23E-06	0.00E+00	3.93E-05	3.85E-04	5.74E-04	2.05E-03
	Uranium-234	9.16E-05	5.10E-13	1.55E-05	1.88E-11	9.39E-08	4.82E-06	1.26E-07	4.47E-09	0.00E+00	4.29E-05	4.19E-04	6.25E-04	1.20E-03
	Thorium-230	3.67E-04	1.24E-12	1.26E-03	8.51E-10	2.33E-06	3.57E-05	1.33E-06	3.30E-08	0.00E+00	1.67E-03	7.87E-04	7.13E-03	1.13E-02
	Radium-226	9.16E-05	2.36E-11	6.23E-04	2.78E-07	2.80E-06	7.24E-02	3.41E-06	4.09E-04	0.00E+00	1.66E-03	5.50E-03	1.07E-02	9.14E-02
	Lead-210	2.88E-05	3.72E-12	1.34E-03	1.18E-09	1.50E-05	3.49E-03	4.50E-05	2.54E-06	0.00E+00	1.19E-02	5.49E-03	4.07E-02	6.31E-02
	Polonium-210	8.64E-05	3.25E-14	2.37E-03	7.12E-12	7.51E-06	2.23E-07	7.96E-05	1.09E-08	0.00E+00	7.53E-02	1.13E-02	2.29E-01	3.19E-01
	Total by Pathway	7.41E-04	2.94E-11	5.62E-03	2.80E-07	2.79E-05	7.69E-02	1.30E-04	4.13E-04	0.00E+00	9.07E-02	2.38E-02	2.89E-01	0.49

Table C16: Estimated Total Radiation Doses to Human Receptors – Operations – Application Case, Upper Bound Scenario

						Tota	l Dose by Pat	:hway - Proje	ct Lifespan (r	nSv/yr)				
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
								Base Case						
	Uranium-238	0.00E+00	0.00E+00	4.22E-06	1.14E-11	3.02E-06	1.10E-03	2.25E-06	7.69E-07	0.00E+00	9.99E-06	4.05E-04	5.67E-04	2.09E-03
	Uranium-234	0.00E+00	0.00E+00	4.58E-06	1.99E-12	3.27E-06	5.56E-06	2.44E-06	2.79E-09	0.00E+00	1.08E-05	4.39E-04	6.15E-04	1.08E-03
	Thorium-230	0.00E+00	0.00E+00	3.69E-04	1.22E-10	6.75E-05	3.95E-05	2.35E-05	2.53E-08	0.00E+00	4.19E-04	1.75E-04	2.77E-03	3.86E-03
	Radium-226	0.00E+00	0.00E+00	5.26E-04	6.50E-08	1.42E-04	9.40E-02	1.71E-04	5.10E-04	0.00E+00	1.19E-03	7.05E-03	1.67E-02	1.20E-01
	Lead-210	0.00E+00	0.00E+00	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	1.36E-02	7.86E-03	8.28E-02	1.15E-01
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	1.21E-01	2.60E-02	3.94E-01	5.56E-01
	Total by Pathway	0.00E+00	0.00E+00	7.19E-03	6.54E-08	2.21E-03	9.97E-02	1.26E-02	5.14E-04	0.00E+00	1.36E-01	4.20E-02	4.98E-01	0.80
							-	Application C	ase					
	Uranium-238	7.57E-05	2.71E-13	6.29E-06	1.70E-11	3.51E-06	1.28E-03	3.13E-06	1.07E-06	0.00E+00	1.48E-05	4.73E-04	5.89E-04	2.45E-03
	Uranium-234	8.86E-05	6.62E-13	6.83E-06	2.97E-12	3.80E-06	6.45E-06	3.39E-06	3.87E-09	0.00E+00	1.61E-05	5.12E-04	6.38E-04	1.28E-03
Subsistence Harvester	Thorium-230	2.82E-04	1.61E-12	6.35E-04	2.10E-10	6.94E-05	4.06E-05	3.95E-05	4.26E-08	0.00E+00	7.19E-04	4.99E-04	4.00E-03	6.28E-03
One-Year-Old (Beet Lake)	Radium-226	8.86E-05	3.07E-11	5.55E-04	6.87E-08	1.46E-04	9.69E-02	1.78E-04	5.32E-04	0.00E+00	1.26E-03	8.22E-03	1.69E-02	1.25E-01
	Lead-210	2.98E-05	4.83E-12	1.81E-03	2.38E-10	1.20E-03	4.67E-03	3.56E-03	3.29E-06	0.00E+00	1.37E-02	1.15E-02	8.39E-02	1.20E-01
	Polonium-210	8.86E-05	4.22E-14	4.50E-03	1.75E-12	8.40E-04	2.91E-07	8.84E-03	1.41E-08	0.00E+00	1.21E-01	3.44E-02	4.00E-01	5.70E-01
	Total by Pathway	6.53E-04	3.81E-11	7.51E-03	6.91E-08	2.26E-03	1.03E-01	1.26E-02	5.37E-04	0.00E+00	1.37E-01	5.56E-02	5.06E-01	0.83
							Upp	oer Bound Sc	enario					
	Uranium-238	7.57E-05	2.71E-13	9.88E-06	2.67E-11	3.51E-06	1.28E-03	4.69E-06	1.60E-06	0.00E+00	2.32E-05	4.73E-04	5.91E-04	2.46E-03
	Uranium-234	8.86E-05	6.62E-13	1.07E-05	4.67E-12	3.80E-06	6.45E-06	5.10E-06	5.81E-09	0.00E+00	2.52E-05	5.12E-04	6.40E-04	1.29E-03
	Thorium-230	2.82E-04	1.61E-12	6.39E-04	2.11E-10	6.94E-05	4.06E-05	3.97E-05	4.28E-08	0.00E+00	7.23E-04	4.99E-04	4.01E-03	6.31E-03
	Radium-226	8.86E-05	3.07E-11	5.56E-04	6.88E-08	1.46E-04	9.69E-02	1.78E-04	5.34E-04	0.00E+00	1.26E-03	8.22E-03	1.69E-02	1.25E-01
	Lead-210	2.98E-05	4.83E-12	1.82E-03	2.40E-10	1.20E-03	4.67E-03	3.58E-03	3.31E-06	0.00E+00	1.38E-02	1.15E-02	8.41E-02	1.21E-01
	Polonium-210	8.86E-05	4.22E-14	4.53E-03	1.77E-12	8.40E-04	2.91E-07	8.90E-03	1.42E-08	0.00E+00	1.22E-01	3.44E-02	4.01E-01	5.72E-01
	Total by Pathway	6.53E-04	3.81E-11	7.57E-03	6.93E-08	2.26E-03	1.03E-01	1.27E-02	5.39E-04	0.00E+00	1.38E-01	5.56E-02	5.08E-01	0.83
								Base Case						
	Uranium-238	0.00E+00	0.00E+00	6.08E-06	4.61E-11	7.42E-08	8.23E-04	5.54E-08	5.91E-07	0.00E+00	1.69E-05	3.30E-04	5.22E-04	1.70E-03
	Uranium-234	0.00E+00	0.00E+00	6.62E-06	8.04E-12	8.08E-08	4.15E-06	6.04E-08	2.14E-09	0.00E+00	1.84E-05	3.60E-04	5.68E-04	9.57E-04
Subsistence Harvester	Thorium-230	0.00E+00	0.00E+00	7.25E-04	4.91E-10	2.27E-06	3.47E-05	7.89E-07	1.95E-08	0.00E+00	9.70E-04	1.94E-04	4.54E-03	6.47E-03
(Lloyd Lake)	Radium-226	0.00E+00	0.00E+00	5.89E-04	2.63E-07	2.71E-06	7.02E-02	3.26E-06	3.91E-04	0.00E+00	1.57E-03	4.47E-03	1.03E-02	8.76E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	1.18E-02	3.28E-03	3.96E-02	5.95E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	7.46E-02	7.71E-03	2.23E-01	3.08E-01
	Total by Pathway	0.00E+00	0.00E+00	5.00E-03	2.64E-07	2.73E-05	7.45E-02	1.28E-04	3.94E-04	0.00E+00	8.90E-02	1.63E-02	2.78E-01	0.46

Table C16: Estimated Total Radiation Doses to Human Receptors – Operations – Application Case, Upper Bound Scenario

			· 			Tota	l Dose by Pat	thway - Proje	ct Lifespan (ı	mSv/yr)	· 	·	· •	
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
							1	Application C	Case					
	Uranium-238	2.54E-06	6.97E-15	6.35E-06	4.82E-11	7.46E-08	8.28E-04	5.74E-08	6.12E-07	0.00E+00	1.77E-05	3.32E-04	5.23E-04	1.71E-03
	Uranium-234	3.07E-06	1.70E-14	6.92E-06	8.40E-12	8.13E-08	4.17E-06	6.25E-08	2.22E-09	0.00E+00	1.92E-05	3.62E-04	5.70E-04	9.65E-04
	Thorium-230	1.23E-05	4.13E-14	7.79E-04	5.28E-10	2.27E-06	3.48E-05	8.44E-07	2.09E-08	0.00E+00	1.04E-03	2.14E-04	4.76E-03	6.84E-03
	Radium-226	3.07E-06	7.91E-13	5.92E-04	2.64E-07	2.71E-06	7.03E-02	3.27E-06	3.92E-04	0.00E+00	1.58E-03	4.50E-03	1.03E-02	8.77E-02
	Lead-210	9.63E-07	1.25E-13	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	1.18E-02	3.35E-03	3.96E-02	5.96E-02
	Polonium-210	2.89E-06	1.09E-15	2.35E-03	7.05E-12	7.28E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	7.46E-02	7.83E-03	2.23E-01	3.08E-01
	Total by Pathway	2.48E-05	9.82E-13	5.06E-03	2.66E-07	2.73E-05	7.46E-02	1.28E-04	3.96E-04	0.00E+00	8.91E-02	1.66E-02	2.79E-01	0.46
							Up _l	per Bound Sc	enario					
	Uranium-238	2.54E-06	6.97E-15	6.83E-06	5.18E-11	7.46E-08	8.28E-04	6.09E-08	6.49E-07	0.00E+00	1.90E-05	3.32E-04	5.24E-04	1.71E-03
	Uranium-234	3.07E-06	1.70E-14	7.44E-06	9.03E-12	8.13E-08	4.17E-06	6.63E-08	2.35E-09	0.00E+00	2.07E-05	3.62E-04	5.70E-04	9.68E-04
	Thorium-230	1.23E-05	4.13E-14	7.79E-04	5.28E-10	2.27E-06	3.48E-05	8.45E-07	2.09E-08	0.00E+00	1.04E-03	2.14E-04	4.76E-03	6.85E-03
	Radium-226	3.07E-06	7.91E-13	5.92E-04	2.64E-07	2.71E-06	7.03E-02	3.28E-06	3.93E-04	0.00E+00	1.58E-03	4.50E-03	1.03E-02	8.77E-02
	Lead-210	9.63E-07	1.25E-13	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	1.18E-02	3.35E-03	3.96E-02	5.96E-02
	Polonium-210	2.89E-06	1.09E-15	2.35E-03	7.06E-12	7.28E-06	2.16E-07	7.90E-05	1.08E-08	0.00E+00	7.46E-02	7.83E-03	2.23E-01	3.08E-01
	Total by Pathway	2.48E-05	9.82E-13	5.06E-03	2.66E-07	2.73E-05	7.46E-02	1.28E-04	3.96E-04	0.00E+00	8.91E-02	1.66E-02	2.79E-01	0.46
								Base Case						
	Uranium-238	0.00E+00	0.00E+00	4.22E-06	1.14E-11	3.02E-06	1.10E-03	2.25E-06	7.69E-07	0.00E+00	9.99E-06	4.05E-04	5.67E-04	2.09E-03
	Uranium-234	0.00E+00	0.00E+00	4.58E-06	1.99E-12	3.27E-06	5.56E-06	2.44E-06	2.79E-09	0.00E+00	1.08E-05	4.39E-04	6.15E-04	1.08E-03
	Thorium-230	0.00E+00	0.00E+00	3.69E-04	1.22E-10	6.75E-05	3.95E-05	2.35E-05	2.53E-08	0.00E+00	4.19E-04	1.75E-04	2.77E-03	3.86E-03
	Radium-226	0.00E+00	0.00E+00	5.26E-04	6.50E-08	1.42E-04	9.40E-02	1.71E-04	5.10E-04	0.00E+00	1.19E-03	7.05E-03	1.67E-02	1.20E-01
	Lead-210	0.00E+00	0.00E+00	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	1.36E-02	7.86E-03	8.28E-02	1.15E-01
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	1.21E-01	2.60E-02	3.94E-01	5.56E-01
Subsistence Harvester	Total by Pathway	0.00E+00	0.00E+00	7.19E-03	6.54E-08	2.21E-03	9.97E-02	1.26E-02	5.14E-04	0.00E+00	1.36E-01	4.20E-02	4.98E-01	0.80
One-Year-Old (Lloyd Lake)								Application C	Case					
,	Uranium-238	2.53E-06	9.06E-15	4.41E-06	1.19E-11	3.04E-06	1.11E-03	2.33E-06	7.96E-07	0.00E+00	1.04E-05	4.08E-04	5.68E-04	2.11E-03
	Uranium-234	2.96E-06	2.22E-14	4.78E-06	2.08E-12	3.29E-06	5.58E-06	2.53E-06	2.89E-09	0.00E+00	1.13E-05	4.42E-04	6.15E-04	1.09E-03
	Thorium-230	9.43E-06	5.37E-14	3.96E-04	1.31E-10	6.76E-05	3.95E-05	2.51E-05	2.71E-08	0.00E+00	4.50E-04	1.85E-04	2.88E-03	4.05E-03
	Radium-226	2.96E-06	1.03E-12	5.29E-04	6.54E-08	1.42E-04	9.41E-02	1.71E-04	5.12E-04	0.00E+00	1.20E-03	7.09E-03	1.67E-02	1.20E-01
	Lead-210	9.97E-07	1.62E-13	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	1.36E-02	7.98E-03	8.28E-02	1.16E-01
	Polonium-210	2.96E-06	1.41E-15	4.49E-03	1.75E-12	8.14E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	1.21E-01	2.63E-02	3.95E-01	5.56E-01
	Total by Pathway	2.18E-05	1.28E-12	7.22E-03	6.58E-08	2.22E-03	9.98E-02	1.26E-02	5.16E-04	0.00E+00	1.36E-01	4.24E-02	4.98E-01	0.80

Table C16: Estimated Total Radiation Doses to Human Receptors – Operations – Application Case, Upper Bound Scenario

						Tota	l Dose by Pat	thway - Proje	ect Lifespan (ı	mSv/yr)				
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
							Up _l	oer Bound Sc	enario					•
	Uranium-238	2.53E-06	9.06E-15	4.74E-06	1.28E-11	3.04E-06	1.11E-03	2.48E-06	8.44E-07	0.00E+00	1.12E-05	4.08E-04	5.68E-04	2.11E-03
	Uranium-234	2.96E-06	2.22E-14	5.14E-06	2.24E-12	3.29E-06	5.58E-06	2.68E-06	3.06E-09	0.00E+00	1.21E-05	4.42E-04	6.16E-04	1.09E-03
	Thorium-230	9.43E-06	5.37E-14	3.97E-04	1.31E-10	6.76E-05	3.95E-05	2.51E-05	2.71E-08	0.00E+00	4.50E-04	1.85E-04	2.88E-03	4.05E-03
	Radium-226	2.96E-06	1.03E-12	5.29E-04	6.54E-08	1.42E-04	9.41E-02	1.71E-04	5.12E-04	0.00E+00	1.20E-03	7.09E-03	1.67E-02	1.20E-01
	Lead-210	9.97E-07	1.62E-13	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	1.36E-02	7.98E-03	8.28E-02	1.16E-01
	Polonium-210	2.96E-06	1.41E-15	4.49E-03	1.75E-12	8.14E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	1.21E-01	2.63E-02	3.95E-01	5.56E-01
	Total by Pathway	2.18E-05	1.28E-12	7.23E-03	6.58E-08	2.22E-03	9.98E-02	1.26E-02	5.16E-04	0.00E+00	1.36E-01	4.24E-02	4.98E-01	0.80
								Base Case	:					
	Uranium-238	0.00E+00	0.00E+00	6.08E-06	4.61E-11	7.42E-08	8.23E-04	5.54E-08	5.91E-07	0.00E+00	5.36E-06	2.59E-04	3.62E-04	1.46E-03
	Uranium-234	0.00E+00	0.00E+00	6.62E-06	8.04E-12	8.08E-08	4.15E-06	6.04E-08	2.14E-09	0.00E+00	5.84E-06	2.82E-04	3.95E-04	6.94E-04
	Thorium-230	0.00E+00	0.00E+00	7.25E-04	4.91E-10	2.27E-06	3.47E-05	7.89E-07	1.95E-08	0.00E+00	3.07E-04	1.52E-04	2.32E-03	3.54E-03
	Radium-226	0.00E+00	0.00E+00	5.89E-04	2.63E-07	2.71E-06	7.02E-02	3.26E-06	3.91E-04	0.00E+00	4.98E-04	3.51E-03	8.84E-03	8.40E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	3.74E-03	2.57E-03	2.70E-02	3.82E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	2.36E-02	6.05E-03	1.23E-01	1.56E-01
	Total by Pathway	0.00E+00	0.00E+00	5.00E-03	2.64E-07	2.73E-05	7.45E-02	1.28E-04	3.94E-04	0.00E+00	2.82E-02	1.28E-02	1.62E-01	0.28
							1	Application C	Case					
	Uranium-238	7.82E-05	2.14E-13	1.09E-05	8.30E-11	8.65E-08	9.60E-04	9.07E-08	9.67E-07	0.00E+00	9.59E-06	3.04E-04	3.86E-04	1.75E-03
	Uranium-234	9.43E-05	5.25E-13	1.19E-05	1.45E-11	9.42E-08	4.84E-06	9.89E-08	3.51E-09	0.00E+00	1.05E-05	3.31E-04	4.20E-04	8.72E-04
Seasonal Resident (Patterson Lake South	Thorium-230	3.77E-04	1.27E-12	1.32E-03	8.97E-10	2.33E-06	3.57E-05	1.40E-06	3.47E-08	0.00E+00	5.59E-04	3.66E-04	5.37E-03	8.04E-03
Arm)	Radium-226	9.43E-05	2.43E-11	6.35E-04	2.83E-07	2.80E-06	7.25E-02	3.47E-06	4.15E-04	0.00E+00	5.36E-04	4.01E-03	9.13E-03	8.73E-02
•	Lead-210	2.97E-05	3.83E-12	1.34E-03	1.18E-09	1.50E-05	3.49E-03	4.49E-05	2.54E-06	0.00E+00	3.78E-03	3.55E-03	2.99E-02	4.22E-02
	Polonium-210	8.90E-05	3.34E-14	2.37E-03	7.12E-12	7.52E-06	2.23E-07	7.95E-05	1.09E-08	0.00E+00	2.38E-02	7.70E-03	1.27E-01	1.61E-01
	Total by Pathway	7.63E-04	3.02E-11	5.69E-03	2.85E-07	2.79E-05	7.69E-02	1.30E-04	4.19E-04	0.00E+00	2.87E-02	1.63E-02	1.72E-01	0.30
							Up _l	oer Bound Sc	enario					
	Uranium-238	7.82E-05	2.14E-13	1.94E-05	1.47E-10	8.65E-08	9.60E-04	1.54E-07	1.64E-06	0.00E+00	1.69E-05	3.04E-04	3.88E-04	1.77E-03
	Uranium-234	9.43E-05	5.25E-13	2.11E-05	2.57E-11	9.42E-08	4.84E-06	1.68E-07	5.95E-09	0.00E+00	1.85E-05	3.31E-04	4.23E-04	8.93E-04
	Thorium-230	3.77E-04	1.27E-12	1.33E-03	9.02E-10	2.33E-06	3.57E-05	1.41E-06	3.49E-08	0.00E+00	5.62E-04	3.66E-04	5.39E-03	8.07E-03
	Radium-226	9.43E-05	2.43E-11	6.37E-04	2.84E-07	2.80E-06	7.25E-02	3.47E-06	4.16E-04	0.00E+00	5.37E-04	4.01E-03	9.13E-03	8.73E-02
	Lead-210	2.97E-05	3.83E-12	1.39E-03	1.22E-09	1.50E-05	3.49E-03	4.62E-05	2.61E-06	0.00E+00	3.91E-03	3.55E-03	3.43E-02	4.67E-02
	Polonium-210	8.90E-05	3.34E-14	2.46E-03	7.38E-12	7.52E-06	2.23E-07	8.17E-05	1.12E-08	0.00E+00	2.47E-02	7.70E-03	1.30E-01	1.65E-01
	Total by Pathway	7.63E-04	3.02E-11	5.86E-03	2.87E-07	2.79E-05	7.69E-02	1.33E-04	4.21E-04	0.00E+00	2.98E-02	1.63E-02	1.79E-01	0.31

Table C16: Estimated Total Radiation Doses to Human Receptors – Operations – Application Case, Upper Bound Scenario

						Tota	l Dose by Pat	:hway - Proje	ct Lifespan (r	nSv/yr)				
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
								Base Case						
	Uranium-238	0.00E+00	0.00E+00	4.22E-06	1.14E-11	3.02E-06	1.10E-03	2.25E-06	7.69E-07	0.00E+00	3.16E-06	3.52E-04	4.97E-04	1.96E-03
	Uranium-234	0.00E+00	0.00E+00	4.58E-06	1.99E-12	3.27E-06	5.56E-06	2.44E-06	2.79E-09	0.00E+00	3.43E-06	3.82E-04	5.38E-04	9.40E-04
	Thorium-230	0.00E+00	0.00E+00	3.69E-04	1.22E-10	6.75E-05	3.95E-05	2.35E-05	2.53E-08	0.00E+00	1.33E-04	1.52E-04	1.68E-03	2.46E-03
	Radium-226	0.00E+00	0.00E+00	5.26E-04	6.50E-08	1.42E-04	9.40E-02	1.71E-04	5.10E-04	0.00E+00	3.78E-04	6.13E-03	1.65E-02	1.18E-01
	Lead-210	0.00E+00	0.00E+00	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	4.32E-03	6.84E-03	6.72E-02	8.95E-02
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	3.83E-02	2.26E-02	2.73E-01	3.49E-01
	Total by Pathway	0.00E+00	0.00E+00	7.19E-03	6.54E-08	2.21E-03	9.97E-02	1.26E-02	5.14E-04	0.00E+00	4.32E-02	3.65E-02	3.60E-01	0.56
							1	Application C	ase	,	,			
	Uranium-238	7.79E-05	2.79E-13	7.60E-06	2.05E-11	3.52E-06	1.28E-03	3.69E-06	1.26E-06	0.00E+00	5.66E-06	4.13E-04	5.08E-04	2.30E-03
	Uranium-234	9.12E-05	6.82E-13	8.24E-06	3.59E-12	3.81E-06	6.48E-06	4.00E-06	4.56E-09	0.00E+00	6.14E-06	4.47E-04	5.50E-04	1.12E-03
Seasonal Resident One- Year-Old (Patterson Lake	Thorium-230	2.90E-04	1.65E-12	6.73E-04	2.22E-10	6.94E-05	4.06E-05	4.17E-05	4.50E-08	0.00E+00	2.41E-04	2.70E-04	3.17E-03	4.80E-03
South Arm)	Radium-226	9.12E-05	3.16E-11	5.67E-04	7.02E-08	1.46E-04	9.70E-02	1.81E-04	5.42E-04	0.00E+00	4.06E-04	6.80E-03	1.66E-02	1.22E-01
,	Lead-210	3.07E-05	4.97E-12	1.82E-03	2.39E-10	1.20E-03	4.67E-03	3.58E-03	3.31E-06	0.00E+00	4.36E-03	8.67E-03	7.09E-02	9.52E-02
	Polonium-210	9.12E-05	4.35E-14	4.53E-03	1.76E-12	8.41E-04	2.91E-07	8.89E-03	1.42E-08	0.00E+00	3.87E-02	2.72E-02	2.77E-01	3.57E-01
	Total by Pathway	6.72E-04	3.93E-11	7.60E-03	7.06E-08	2.26E-03	1.03E-01	1.27E-02	5.46E-04	0.00E+00	4.37E-02	4.38E-02	3.69E-01	0.58
							Upp	er Bound Sc	enario					
	Uranium-238	7.79E-05	2.79E-13	1.35E-05	3.63E-11	3.52E-06	1.28E-03	6.24E-06	2.13E-06	0.00E+00	1.00E-05	4.13E-04	5.09E-04	2.32E-03
	Uranium-234	9.12E-05	6.82E-13	1.46E-05	6.36E-12	3.81E-06	6.48E-06	6.78E-06	7.74E-09	0.00E+00	1.09E-05	4.47E-04	5.51E-04	1.13E-03
	Thorium-230	2.90E-04	1.65E-12	6.77E-04	2.23E-10	6.94E-05	4.06E-05	4.20E-05	4.53E-08	0.00E+00	2.43E-04	2.70E-04	3.18E-03	4.81E-03
	Radium-226	9.12E-05	3.16E-11	5.69E-04	7.04E-08	1.46E-04	9.70E-02	1.82E-04	5.43E-04	0.00E+00	4.08E-04	6.80E-03	1.66E-02	1.22E-01
	Lead-210	3.07E-05	4.97E-12	1.89E-03	2.48E-10	1.20E-03	4.67E-03	3.67E-03	3.40E-06	0.00E+00	4.52E-03	8.67E-03	7.64E-02	1.01E-01
	Polonium-210	9.12E-05	4.35E-14	4.70E-03	1.83E-12	8.41E-04	2.91E-07	9.13E-03	1.46E-08	0.00E+00	4.01E-02	2.72E-02	2.80E-01	3.62E-01
	Total by Pathway	6.72E-04	3.93E-11	7.86E-03	7.09E-08	2.26E-03	1.03E-01	1.30E-02	5.49E-04	0.00E+00	4.53E-02	4.38E-02	3.77E-01	0.59
								Base Case						
	Uranium-238	0.00E+00	0.00E+00	6.08E-06	4.61E-11	7.42E-08	8.23E-04	5.54E-08	5.91E-07	0.00E+00	5.36E-06	2.59E-04	3.62E-04	1.46E-03
	Uranium-234	0.00E+00	0.00E+00	6.62E-06	8.04E-12	8.08E-08	4.15E-06	6.04E-08	2.14E-09	0.00E+00	5.84E-06	2.82E-04	3.95E-04	6.94E-04
Seasonal Resident (Lloyd	Thorium-230	0.00E+00	0.00E+00	7.25E-04	4.91E-10	2.27E-06	3.47E-05	7.89E-07	1.95E-08	0.00E+00	3.07E-04	1.52E-04	2.32E-03	3.54E-03
Lake)	Radium-226	0.00E+00	0.00E+00	5.89E-04	2.63E-07	2.71E-06	7.02E-02	3.26E-06	3.91E-04	0.00E+00	4.98E-04	3.51E-03	8.84E-03	8.40E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	3.74E-03	2.57E-03	2.70E-02	3.82E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	2.36E-02	6.05E-03	1.23E-01	1.56E-01
	Total by Pathway	0.00E+00	0.00E+00	5.00E-03	2.64E-07	2.73E-05	7.45E-02	1.28E-04	3.94E-04	0.00E+00	2.82E-02	1.28E-02	1.62E-01	0.28

Table C16: Estimated Total Radiation Doses to Human Receptors – Operations – Application Case, Upper Bound Scenario

						Tota	l Dose by Pat	thway - Proje	ct Lifespan (r	nSv/yr)				
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
		•						Application C	ase					
	Uranium-238	1.52E-06	4.18E-15	6.24E-06	4.74E-11	7.45E-08	8.26E-04	5.66E-08	6.04E-07	0.00E+00	6.95E-06	2.60E-04	3.63E-04	1.46E-03
	Uranium-234	1.84E-06	1.02E-14	6.80E-06	8.25E-12	8.11E-08	4.16E-06	6.16E-08	2.19E-09	0.00E+00	7.57E-06	2.83E-04	3.95E-04	6.99E-04
	Thorium-230	7.36E-06	2.48E-14	7.57E-04	5.13E-10	2.27E-06	3.48E-05	8.22E-07	2.03E-08	0.00E+00	3.30E-04	1.57E-04	2.37E-03	3.66E-03
	Radium-226	1.84E-06	4.75E-13	5.90E-04	2.64E-07	2.71E-06	7.02E-02	3.27E-06	3.92E-04	0.00E+00	5.01E-04	3.52E-03	8.84E-03	8.41E-02
	Lead-210	5.78E-07	7.47E-14	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	4.50E-03	2.59E-03	2.70E-02	3.89E-02
	Polonium-210	1.73E-06	6.52E-16	2.35E-03	7.05E-12	7.28E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	2.41E-02	6.08E-03	1.23E-01	1.56E-01
	Total by Pathway	1.49E-05	5.89E-13	5.04E-03	2.65E-07	2.73E-05	7.46E-02	1.28E-04	3.95E-04	0.00E+00	2.95E-02	1.29E-02	1.62E-01	0.28
							Up _l	per Bound Sc	enario					
	Uranium-238	1.52E-06	4.18E-15	6.53E-06	4.95E-11	7.45E-08	8.26E-04	5.87E-08	6.26E-07	0.00E+00	7.47E-06	2.60E-04	3.63E-04	1.47E-03
	Uranium-234	1.84E-06	1.02E-14	7.11E-06	8.63E-12	8.11E-08	4.16E-06	6.39E-08	2.27E-09	0.00E+00	8.14E-06	2.83E-04	3.95E-04	7.00E-04
	Thorium-230	7.36E-06	2.48E-14	7.57E-04	5.13E-10	2.27E-06	3.48E-05	8.22E-07	2.03E-08	0.00E+00	3.30E-04	1.57E-04	2.37E-03	3.66E-03
	Radium-226	1.84E-06	4.75E-13	5.91E-04	2.64E-07	2.71E-06	7.02E-02	3.27E-06	3.92E-04	0.00E+00	5.01E-04	3.52E-03	8.84E-03	8.41E-02
	Lead-210	5.78E-07	7.47E-14	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	4.50E-03	2.59E-03	2.70E-02	3.90E-02
	Polonium-210	1.73E-06	6.52E-16	2.35E-03	7.06E-12	7.28E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	2.42E-02	6.08E-03	1.23E-01	1.56E-01
	Total by Pathway	1.49E-05	5.89E-13	5.04E-03	2.65E-07	2.73E-05	7.46E-02	1.28E-04	3.95E-04	0.00E+00	2.95E-02	1.29E-02	1.62E-01	0.28
		•						Base Case						
	Uranium-238	0.00E+00	0.00E+00	4.22E-06	1.14E-11	3.02E-06	1.10E-03	2.25E-06	7.69E-07	0.00E+00	3.16E-06	3.52E-04	4.97E-04	1.96E-03
	Uranium-234	0.00E+00	0.00E+00	4.58E-06	1.99E-12	3.27E-06	5.56E-06	2.44E-06	2.79E-09	0.00E+00	3.43E-06	3.82E-04	5.38E-04	9.40E-04
	Thorium-230	0.00E+00	0.00E+00	3.69E-04	1.22E-10	6.75E-05	3.95E-05	2.35E-05	2.53E-08	0.00E+00	1.33E-04	1.52E-04	1.68E-03	2.46E-03
	Radium-226	0.00E+00	0.00E+00	5.26E-04	6.50E-08	1.42E-04	9.40E-02	1.71E-04	5.10E-04	0.00E+00	3.78E-04	6.13E-03	1.65E-02	1.18E-01
	Lead-210	0.00E+00	0.00E+00	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	4.32E-03	6.84E-03	6.72E-02	8.95E-02
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	3.83E-02	2.26E-02	2.73E-01	3.49E-01
Seasonal Resident One-	Total by Pathway	0.00E+00	0.00E+00	7.19E-03	6.54E-08	2.21E-03	9.97E-02	1.26E-02	5.14E-04	0.00E+00	4.32E-02	3.65E-02	3.60E-01	0.56
Year-Old (Lloyd Lake)								Application C	ase					
	Uranium-238	1.52E-06	5.44E-15	4.34E-06	1.17E-11	3.03E-06	1.11E-03	2.30E-06	7.85E-07	0.00E+00	3.25E-06	3.54E-04	4.97E-04	1.97E-03
	Uranium-234	1.78E-06	1.33E-14	4.70E-06	2.04E-12	3.28E-06	5.57E-06	2.49E-06	2.85E-09	0.00E+00	3.52E-06	3.83E-04	5.39E-04	9.43E-04
	Thorium-230	5.66E-06	3.22E-14	3.85E-04	1.27E-10	6.76E-05	3.95E-05	2.45E-05	2.64E-08	0.00E+00	1.39E-04	1.54E-04	1.70E-03	2.52E-03
	Radium-226	1.78E-06	6.17E-13	5.28E-04	6.53E-08	1.42E-04	9.40E-02	1.71E-04	5.11E-04	0.00E+00	3.79E-04	6.14E-03	1.65E-02	1.18E-01
	Lead-210	5.98E-07	9.70E-14	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	4.32E-03	6.87E-03	6.72E-02	8.96E-02
	Polonium-210	1.78E-06	8.48E-16	4.49E-03	1.75E-12	8.14E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	3.83E-02	2.27E-02	2.74E-01	3.49E-01
	Total by Pathway	1.31E-05	7.66E-13	7.21E-03	6.56E-08	2.22E-03	9.98E-02	1.26E-02	5.15E-04	0.00E+00	4.32E-02	3.66E-02	3.60E-01	0.56

Table C16: Estimated Total Radiation Doses to Human Receptors – Operations – Application Case, Upper Bound Scenario

						Tota	l Dose by Pat	thway - Proje	ct Lifespan (r	nSv/yr)				
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
							Upj	per Bound Sc	enario					
	Uranium-238	1.52E-06	5.44E-15	4.54E-06	1.22E-11	3.03E-06	1.11E-03	2.39E-06	8.14E-07	0.00E+00	3.40E-06	3.54E-04	4.97E-04	1.97E-03
	Uranium-234	1.78E-06	1.33E-14	4.92E-06	2.14E-12	3.28E-06	5.57E-06	2.59E-06	2.95E-09	0.00E+00	3.68E-06	3.83E-04	5.39E-04	9.44E-04
	Thorium-230	5.66E-06	3.22E-14	3.85E-04	1.27E-10	6.76E-05	3.95E-05	2.45E-05	2.64E-08	0.00E+00	1.39E-04	1.54E-04	1.70E-03	2.52E-03
	Radium-226	1.78E-06	6.17E-13	5.28E-04	6.53E-08	1.42E-04	9.40E-02	1.71E-04	5.11E-04	0.00E+00	3.79E-04	6.14E-03	1.65E-02	1.18E-01
	Lead-210	5.98E-07	9.70E-14	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	4.32E-03	6.87E-03	6.72E-02	8.96E-02
	Polonium-210	1.78E-06	8.48E-16	4.49E-03	1.75E-12	8.14E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	3.84E-02	2.27E-02	2.74E-01	3.49E-01
	Total by Pathway	1.31E-05	7.66E-13	7.21E-03	6.56E-08	2.22E-03	9.98E-02	1.26E-02	5.15E-04	0.00E+00	4.32E-02	3.66E-02	3.60E-01	0.56
								Base Case	· !					
	Uranium-238	-	-	-	-	-	-	-	-	-	-	-	-	-
	Uranium-234	-	-	-	-	-	-	-	-	-	-	-	-	-
	Thorium-230	-	-	-	-	-	-	-	-	-	-	-	-	-
	Radium-226	-	-	-	-	-	-	-	-	-	-	-	-	-
	Lead-210	-	-	-	-	-	-	-	-	-	-	-	-	-
	Polonium-210	-	-	-	-	-	-	-	-	-	-	-	-	-
	Total by Pathway	-	-	-	-	-	-	-	-	-	-	-	-	-
								Application C	ase					
	Uranium-238	-	-	-	-	-	-	-	-	-	-	-	-	-
	Uranium-234	-	-	-	-	-	-	-	-	-	-	-	-	-
Permanent Resident (Patterson Lake North	Thorium-230	-	-	-	-	-	-	-	-	-	-	-	-	-
Arm - West Basin)	Radium-226	-	-	-	-	-	-	-	-	-	-	-	-	-
,	Lead-210	-	-	-	-	-	-	-	-	-	-	-	-	-
	Polonium-210	-	-	-	-	-	-	-	-	-	-	-	-	-
	Total by Pathway	-	-	-	-	-	-	-	-	-	-	-	-	-
							Up _l	er Bound Sc	enario					
	Uranium-238	-	-	-	-	-	-	-	-	-	-	-	-	-
	Uranium-234	-	-	-	-	-	-	-	-	-	-	-	-	-
	Thorium-230	-	-	-	-	-	-	-	-	-	-	-	-	-
	Radium-226	-	-	-	-	-	-	-	-	-	-	-	-	-
	Lead-210	-	-	-	-	-	-	-	-	-	-	-	-	-
	Polonium-210	-	-	-	-	-	-	-	-	-	-	-	-	-
	Total by Pathway	-	-	-	-	-	-	-	-	-	-	-	-	-

Table C16: Estimated Total Radiation Doses to Human Receptors – Operations – Application Case, Upper Bound Scenario

						Tota	l Dose by Pat	thway - Proje	ct Lifespan (r	nSv/yr)				
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
								Base Case						
	Uranium-238	-	-	-	-	-	-	-	-	-	-	-	-	-
	Uranium-234	-	-	-	-	-	-	-	-	-	-	-	-	-
	Thorium-230	-	-	-	-	-	-	-	-	-	-	-	-	-
	Radium-226	-	-	-	-	-	-	-	-	-	-	-	-	-
	Lead-210	-	-	-	-	-	-	-	-	-	-	-	-	-
	Polonium-210	-	-	-	-	-	-	-	-	-	-	-	-	-
	Total by Pathway	-	-	-	-	-	-	-	-	-	-	-	-	-
								Application C	lase					
	Uranium-238	-	-	-	-	-	-	-	-	-	-	-	-	-
	Uranium-234	-	-	-	-	-	-	-	-	-	-	-	-	-
Permanent Resident One- Year-Old (Patterson Lake	Thorium-230	-	-	-	-	-	-	-	-	-	-	-	-	-
North Arm - West Basin)	Radium-226	-	-	-	-	-	-	-	-	-	-	-	-	-
,	Lead-210	-	-	-	-	-	-	-	-	-	-	-	-	-
	Polonium-210	-	-	-	-	-	-	-	-	-	-	-	-	-
	Total by Pathway	-	-	-	-	-	-	-	-	-	-	-	-	-
							Up _l	per Bound Sc	enario					
	Uranium-238	-	-	-	-	-	-	-	-	-	-	-	-	-
	Uranium-234	-	-	-	-	-	-	-	-	-	-	-	-	-
	Thorium-230	-	-	-	-	-	-	-	-	-	-	-	-	-
	Radium-226	-	-	-	-	-	-	-	-	-	-	-	-	-
	Lead-210	-	-	-	-	-	-	-	-	-	-	-	-	-
	Polonium-210	-	-	-	-	-	-	-	-	-	-	-	-	-
	Total by Pathway	-	-	-	-	-	-	-	-	-	-	-	-	-

Table C17: Estimated Total Radiation Doses to Human Receptors – Far-Future – Application Case, Upper Bound Scenario

						To	otal Dose by	Pathway - Fa	r-Future (mS	v/yr)				
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
								Base Case						
	Uranium-238	-	-		-	-	-	-	-	-	-	-	ı	-
	Uranium-234	-	-	-	-	-	-	-	-	-	-	-	-	-
	Thorium-230	-	-	-	-	-	-	-	-	-	-	-	ı	-
	Radium-226	-	-	-	-	-	-	-	-	-	-	-	-	-
	Lead-210	-	-	-	-	-	-	-	-	-	-	-	-	-
	Polonium-210	-	-	-	-	-	-	-	-	-	-	-	ı	-
	Total by Pathway	-	-	-	-	-	-	-	-	-	-	-	-	-
							1	Application C	ase					
	Uranium-238	-	-	-	-	-	-	-	-	-	-	-	-	-
	Uranium-234	-	-	-	-	-	-	-	-	-	-	-	-	-
Camp Worker (Patterson Lake North	Thorium-230	-	-	-	-	-	-	-	-	-	-	-	ı	-
Arm - West Basin)	Radium-226	-	-		-	-	-	-	-	-	-	-	ı	-
,	Lead-210	-	-	•	-	-	-	-	-	-	-	-	•	-
	Polonium-210	-	-	-	-	-	-	-	-	-	-	-	•	-
	Total by Pathway	-	-	-	-	-	-	-	-	-	-	-	-	-
							Upp	oer Bound Sc	enario					
	Uranium-238	-	-	-	-	-	-	-	-	-	-	-	ı	-
	Uranium-234	-	-	-	-	-	-	-	-	-	-	-	ı	-
	Thorium-230	-	-	-	-	-	-	-	-	-	-	-	-	-
	Radium-226	-	-	-	-	-	-	-	-	-	-	-	-	-
	Lead-210	-	-	-	-	-	-	-	-	-	-	-	ı	-
	Polonium-210	-	-	-	-	-	-	-	-	-	-	-	-	-
	Total by Pathway	-	-	-	-	-	-	-	-	-	-	-	-	-
								Base Case						
	Uranium-238	0.00E+00	0.00E+00	6.08E-06	4.61E-11	7.42E-08	8.23E-04	5.54E-08	5.91E-07	0.00E+00	1.69E-05	3.30E-04	5.22E-04	1.70E-03
	Uranium-234	0.00E+00	0.00E+00	6.62E-06	8.04E-12	8.08E-08	4.15E-06	6.04E-08	2.14E-09	0.00E+00	1.84E-05	3.60E-04	5.68E-04	9.57E-04
Subsistence Harvester (Patterson Lake South	Thorium-230	0.00E+00	0.00E+00	7.25E-04	4.91E-10	2.27E-06	3.47E-05	7.89E-07	1.95E-08	0.00E+00	9.70E-04	1.94E-04	4.54E-03	6.47E-03
(Patterson Lake South Arm)	Radium-226	0.00E+00	0.00E+00	5.89E-04	2.63E-07	2.71E-06	7.02E-02	3.26E-06	3.91E-04	0.00E+00	1.57E-03	4.47E-03	1.03E-02	8.76E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	1.18E-02	3.28E-03	3.96E-02	5.95E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	7.46E-02	7.71E-03	2.23E-01	3.07E-01
	Total by Pathway	0.00E+00	0.00E+00	5.00E-03	2.64E-07	2.73E-05	7.45E-02	1.28E-04	3.94E-04	0.00E+00	8.90E-02	1.63E-02	2.78E-01	0.46

Table C17: Estimated Total Radiation Doses to Human Receptors – Far-Future – Application Case, Upper Bound Scenario

						To	otal Dose by	Pathway - Fa	r-Future (mS	v/yr)				
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
								Application C	Case					
	Uranium-238	0.00E+00	0.00E+00	3.25E-05	2.46E-10	7.64E-08	8.47E-04	2.96E-07	3.16E-06	0.00E+00	9.04E-05	3.40E-04	5.56E-04	1.87E-03
	Uranium-234	0.00E+00	0.00E+00	3.55E-05	4.31E-11	8.32E-08	4.27E-06	3.23E-07	1.15E-08	0.00E+00	9.88E-05	3.70E-04	6.06E-04	1.12E-03
	Thorium-230	0.00E+00	0.00E+00	7.40E-04	5.01E-10	2.33E-06	3.57E-05	8.05E-07	1.99E-08	0.00E+00	9.90E-04	2.00E-04	4.73E-03	6.70E-03
	Radium-226	0.00E+00	0.00E+00	6.20E-04	2.77E-07	2.81E-06	7.27E-02	3.43E-06	4.12E-04	0.00E+00	1.66E-03	4.63E-03	1.07E-02	9.08E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.47E-05	2.53E-06	0.00E+00	1.18E-02	3.28E-03	4.00E-02	5.99E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.06E-12	7.27E-06	2.16E-07	7.90E-05	1.08E-08	0.00E+00	7.47E-02	7.71E-03	2.23E-01	3.08E-01
	Total by Pathway	0.00E+00	0.00E+00	5.11E-03	2.79E-07	2.75E-05	7.71E-02	1.29E-04	4.17E-04	0.00E+00	8.94E-02	1.65E-02	2.80E-01	0.47
							Up _l	per Bound Sc	enario					
	Uranium-238	0.00E+00	0.00E+00	3.25E-05	2.47E-10	7.64E-08	8.47E-04	2.96E-07	3.16E-06	0.00E+00	9.05E-05	3.40E-04	5.56E-04	1.87E-03
	Uranium-234	0.00E+00	0.00E+00	3.55E-05	4.31E-11	8.32E-08	4.27E-06	3.24E-07	1.15E-08	0.00E+00	9.88E-05	3.70E-04	6.06E-04	1.12E-03
	Thorium-230	0.00E+00	0.00E+00	7.40E-04	5.01E-10	2.33E-06	3.57E-05	8.05E-07	1.99E-08	0.00E+00	9.90E-04	2.00E-04	4.73E-03	6.70E-03
	Radium-226	0.00E+00	0.00E+00	6.20E-04	2.77E-07	2.81E-06	7.27E-02	3.44E-06	4.12E-04	0.00E+00	1.66E-03	4.63E-03	1.07E-02	9.08E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.47E-05	2.53E-06	0.00E+00	1.18E-02	3.28E-03	4.00E-02	5.99E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.06E-12	7.27E-06	2.16E-07	7.90E-05	1.08E-08	0.00E+00	7.47E-02	7.71E-03	2.23E-01	3.08E-01
	Total by Pathway	0.00E+00	0.00E+00	5.11E-03	2.79E-07	2.75E-05	7.71E-02	1.29E-04	4.17E-04	0.00E+00	8.94E-02	1.65E-02	2.80E-01	0.47
								Base Case	:					
	Uranium-238	0.00E+00	0.00E+00	4.22E-06	1.14E-11	3.02E-06	1.10E-03	2.25E-06	7.69E-07	0.00E+00	9.99E-06	4.05E-04	5.67E-04	2.09E-03
	Uranium-234	0.00E+00	0.00E+00	4.58E-06	1.99E-12	3.27E-06	5.56E-06	2.44E-06	2.79E-09	0.00E+00	1.08E-05	4.39E-04	6.15E-04	1.08E-03
	Thorium-230	0.00E+00	0.00E+00	3.69E-04	1.22E-10	6.75E-05	3.95E-05	2.35E-05	2.53E-08	0.00E+00	4.19E-04	1.75E-04	2.77E-03	3.86E-03
	Radium-226	0.00E+00	0.00E+00	5.26E-04	6.50E-08	1.42E-04	9.40E-02	1.71E-04	5.10E-04	0.00E+00	1.19E-03	7.05E-03	1.67E-02	1.20E-01
	Lead-210	0.00E+00	0.00E+00	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	1.36E-02	7.86E-03	8.28E-02	1.15E-01
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	1.21E-01	2.60E-02	3.94E-01	5.56E-01
Subsistence Harvester One-Year-Old (Patterson	Total by Pathway	0.00E+00	0.00E+00	7.19E-03	6.54E-08	2.21E-03	9.97E-02	1.26E-02	5.14E-04	0.00E+00	1.36E-01	4.20E-02	4.98E-01	0.80
Lake South Arm)								Application C	Case					
,	Uranium-238	0.00E+00	0.00E+00	2.26E-05	6.09E-11	3.11E-06	1.13E-03	1.20E-05	4.11E-06	0.00E+00	5.33E-05	4.17E-04	5.83E-04	2.23E-03
	Uranium-234	0.00E+00	0.00E+00	2.45E-05	1.07E-11	3.37E-06	5.72E-06	1.31E-05	1.49E-08	0.00E+00	5.80E-05	4.52E-04	6.32E-04	1.19E-03
	Thorium-230	0.00E+00	0.00E+00	3.76E-04	1.24E-10	6.93E-05	4.06E-05	2.40E-05	2.59E-08	0.00E+00	4.28E-04	1.79E-04	2.86E-03	3.98E-03
	Radium-226	0.00E+00	0.00E+00	5.54E-04	6.85E-08	1.47E-04	9.73E-02	1.80E-04	5.37E-04	0.00E+00	1.26E-03	7.31E-03	1.69E-02	1.24E-01
	Lead-210	0.00E+00	0.00E+00	1.81E-03	2.38E-10	1.19E-03	4.63E-03	3.55E-03	3.29E-06	0.00E+00	1.37E-02	7.86E-03	8.33E-02	1.16E-01
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.84E-03	1.41E-08	0.00E+00	1.21E-01	2.60E-02	3.95E-01	5.56E-01
	Total by Pathway	0.00E+00	0.00E+00	7.28E-03	6.89E-08	2.22E-03	1.03E-01	1.26E-02	5.44E-04	0.00E+00	1.37E-01	4.22E-02	4.99E-01	0.80

Table C17: Estimated Total Radiation Doses to Human Receptors – Far-Future – Application Case, Upper Bound Scenario

						To	otal Dose by	Pathway - Fa	r-Future (mS	v/yr)				
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
							Up _l	per Bound Sc	enario					
	Uranium-238	0.00E+00	0.00E+00	2.26E-05	6.09E-11	3.11E-06	1.13E-03	1.20E-05	4.11E-06	0.00E+00	5.34E-05	4.17E-04	5.83E-04	2.23E-03
	Uranium-234	0.00E+00	0.00E+00	2.45E-05	1.07E-11	3.37E-06	5.72E-06	1.31E-05	1.49E-08	0.00E+00	5.80E-05	4.52E-04	6.32E-04	1.19E-03
	Thorium-230	0.00E+00	0.00E+00	3.76E-04	1.24E-10	6.93E-05	4.06E-05	2.40E-05	2.59E-08	0.00E+00	4.28E-04	1.79E-04	2.86E-03	3.98E-03
	Radium-226	0.00E+00	0.00E+00	5.54E-04	6.85E-08	1.47E-04	9.73E-02	1.80E-04	5.37E-04	0.00E+00	1.26E-03	7.31E-03	1.69E-02	1.24E-01
	Lead-210	0.00E+00	0.00E+00	1.81E-03	2.38E-10	1.19E-03	4.63E-03	3.55E-03	3.29E-06	0.00E+00	1.37E-02	7.86E-03	8.33E-02	1.16E-01
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.84E-03	1.41E-08	0.00E+00	1.21E-01	2.60E-02	3.95E-01	5.56E-01
	Total by Pathway	0.00E+00	0.00E+00	7.28E-03	6.90E-08	2.22E-03	1.03E-01	1.26E-02	5.45E-04	0.00E+00	1.37E-01	4.22E-02	4.99E-01	0.80
								Base Case	:					_
	Uranium-238	0.00E+00	0.00E+00	6.08E-06	4.61E-11	7.42E-08	8.23E-04	5.54E-08	5.91E-07	0.00E+00	1.69E-05	3.30E-04	5.22E-04	1.70E-03
	Uranium-234	0.00E+00	0.00E+00	6.62E-06	8.04E-12	8.08E-08	4.15E-06	6.04E-08	2.14E-09	0.00E+00	1.84E-05	3.60E-04	5.68E-04	9.57E-04
	Thorium-230	0.00E+00	0.00E+00	7.25E-04	4.91E-10	2.27E-06	3.47E-05	7.89E-07	1.95E-08	0.00E+00	9.70E-04	1.94E-04	4.54E-03	6.47E-03
	Radium-226	0.00E+00	0.00E+00	5.89E-04	2.63E-07	2.71E-06	7.02E-02	3.26E-06	3.91E-04	0.00E+00	1.57E-03	4.47E-03	1.03E-02	8.76E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	1.18E-02	3.28E-03	3.96E-02	5.95E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	7.46E-02	7.71E-03	2.23E-01	3.07E-01
	Total by Pathway	0.00E+00	0.00E+00	5.00E-03	2.64E-07	2.73E-05	7.45E-02	1.28E-04	3.94E-04	0.00E+00	8.90E-02	1.63E-02	2.78E-01	0.46
								Application C	ase					
	Uranium-238	0.00E+00	0.00E+00	1.59E-05	1.21E-10	7.55E-08	8.37E-04	1.45E-07	1.55E-06	0.00E+00	4.42E-05	3.36E-04	5.34E-04	1.77E-03
	Uranium-234	0.00E+00	0.00E+00	1.73E-05	2.11E-11	8.22E-08	4.22E-06	1.58E-07	5.62E-09	0.00E+00	4.83E-05	3.66E-04	5.81E-04	1.02E-03
Subsistence Harvester	Thorium-230	0.00E+00	0.00E+00	7.33E-04	4.97E-10	2.30E-06	3.53E-05	7.97E-07	1.97E-08	0.00E+00	9.81E-04	1.97E-04	4.59E-03	6.54E-03
(Beet Lake)	Radium-226	0.00E+00	0.00E+00	6.02E-04	2.69E-07	2.77E-06	7.17E-02	3.34E-06	4.00E-04	0.00E+00	1.61E-03	4.56E-03	1.04E-02	8.93E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	1.18E-02	3.28E-03	3.96E-02	5.95E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	7.46E-02	7.71E-03	2.23E-01	3.08E-01
	Total by Pathway	0.00E+00	0.00E+00	5.04E-03	2.71E-07	2.74E-05	7.60E-02	1.28E-04	4.04E-04	0.00E+00	8.91E-02	1.65E-02	2.78E-01	0.47
							Up _l	per Bound Sc	enario					
	Uranium-238	0.00E+00	0.00E+00	1.59E-05	1.21E-10	7.55E-08	8.37E-04	1.45E-07	1.55E-06	0.00E+00	4.43E-05	3.36E-04	5.34E-04	1.77E-03
	Uranium-234	0.00E+00	0.00E+00	1.74E-05	2.11E-11	8.22E-08	4.22E-06	1.58E-07	5.62E-09	0.00E+00	4.83E-05	3.66E-04	5.81E-04	1.02E-03
	Thorium-230	0.00E+00	0.00E+00	7.33E-04	4.97E-10	2.30E-06	3.53E-05	7.97E-07	1.97E-08	0.00E+00	9.81E-04	1.97E-04	4.59E-03	6.54E-03
	Radium-226	0.00E+00	0.00E+00	6.02E-04	2.69E-07	2.77E-06	7.17E-02	3.34E-06	4.00E-04	0.00E+00	1.61E-03	4.56E-03	1.04E-02	8.93E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	1.18E-02	3.28E-03	3.96E-02	5.95E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	7.46E-02	7.71E-03	2.23E-01	3.08E-01
	Total by Pathway	0.00E+00	0.00E+00	5.04E-03	2.71E-07	2.74E-05	7.60E-02	1.28E-04	4.04E-04	0.00E+00	8.91E-02	1.65E-02	2.78E-01	0.47

Table C17: Estimated Total Radiation Doses to Human Receptors – Far-Future – Application Case, Upper Bound Scenario

						To	otal Dose by	Pathway - Fa	r-Future (mS	v/yr)				
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
								Base Case						
	Uranium-238	0.00E+00	0.00E+00	4.22E-06	1.14E-11	3.02E-06	1.10E-03	2.25E-06	7.69E-07	0.00E+00	9.99E-06	4.05E-04	5.67E-04	2.09E-03
	Uranium-234	0.00E+00	0.00E+00	4.58E-06	1.99E-12	3.27E-06	5.56E-06	2.44E-06	2.79E-09	0.00E+00	1.08E-05	4.39E-04	6.15E-04	1.08E-03
	Thorium-230	0.00E+00	0.00E+00	3.69E-04	1.22E-10	6.75E-05	3.95E-05	2.35E-05	2.53E-08	0.00E+00	4.19E-04	1.75E-04	2.77E-03	3.86E-03
	Radium-226	0.00E+00	0.00E+00	5.26E-04	6.50E-08	1.42E-04	9.40E-02	1.71E-04	5.10E-04	0.00E+00	1.19E-03	7.05E-03	1.67E-02	1.20E-01
	Lead-210	0.00E+00	0.00E+00	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	1.36E-02	7.86E-03	8.28E-02	1.15E-01
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	1.21E-01	2.60E-02	3.94E-01	5.56E-01
	Total by Pathway	0.00E+00	0.00E+00	7.19E-03	6.54E-08	2.21E-03	9.97E-02	1.26E-02	5.14E-04	0.00E+00	1.36E-01	4.20E-02	4.98E-01	0.80
								Application C	ase					
	Uranium-238	0.00E+00	0.00E+00	1.10E-05	2.98E-11	3.07E-06	1.12E-03	5.89E-06	2.01E-06	0.00E+00	2.61E-05	4.12E-04	5.73E-04	2.15E-03
	Uranium-234	0.00E+00	0.00E+00	1.20E-05	5.21E-12	3.33E-06	5.65E-06	6.40E-06	7.30E-09	0.00E+00	2.83E-05	4.47E-04	6.20E-04	1.12E-03
Subsistence Harvester	Thorium-230	0.00E+00	0.00E+00	3.73E-04	1.23E-10	6.86E-05	4.01E-05	2.37E-05	2.56E-08	0.00E+00	4.24E-04	1.77E-04	2.79E-03	3.90E-03
One-Year-Old (Beet Lake)	Radium-226	0.00E+00	0.00E+00	5.38E-04	6.65E-08	1.45E-04	9.59E-02	1.74E-04	5.22E-04	0.00E+00	1.22E-03	7.20E-03	1.68E-02	1.23E-01
	Lead-210	0.00E+00	0.00E+00	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	1.36E-02	7.86E-03	8.28E-02	1.15E-01
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	1.21E-01	2.60E-02	3.94E-01	5.56E-01
	Total by Pathway	0.00E+00	0.00E+00	7.23E-03	6.69E-08	2.22E-03	1.02E-01	1.26E-02	5.27E-04	0.00E+00	1.36E-01	4.21E-02	4.98E-01	0.80
							Upp	oer Bound Sc	enario					
	Uranium-238	0.00E+00	0.00E+00	1.10E-05	2.98E-11	3.07E-06	1.12E-03	5.89E-06	2.01E-06	0.00E+00	2.61E-05	4.12E-04	5.73E-04	2.15E-03
	Uranium-234	0.00E+00	0.00E+00	1.20E-05	5.22E-12	3.33E-06	5.65E-06	6.40E-06	7.31E-09	0.00E+00	2.84E-05	4.47E-04	6.20E-04	1.12E-03
	Thorium-230	0.00E+00	0.00E+00	3.73E-04	1.23E-10	6.86E-05	4.01E-05	2.37E-05	2.56E-08	0.00E+00	4.24E-04	1.77E-04	2.79E-03	3.90E-03
	Radium-226	0.00E+00	0.00E+00	5.38E-04	6.65E-08	1.45E-04	9.59E-02	1.74E-04	5.22E-04	0.00E+00	1.22E-03	7.20E-03	1.68E-02	1.23E-01
	Lead-210	0.00E+00	0.00E+00	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	1.36E-02	7.86E-03	8.28E-02	1.15E-01
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	1.21E-01	2.60E-02	3.94E-01	5.56E-01
	Total by Pathway	0.00E+00	0.00E+00	7.23E-03	6.69E-08	2.22E-03	1.02E-01	1.26E-02	5.27E-04	0.00E+00	1.36E-01	4.21E-02	4.98E-01	0.80
								Base Case				•		
	Uranium-238	0.00E+00	0.00E+00	6.08E-06	4.61E-11	7.42E-08	8.23E-04	5.54E-08	5.91E-07	0.00E+00	1.69E-05	3.30E-04	5.22E-04	1.70E-03
	Uranium-234	0.00E+00	0.00E+00	6.62E-06	8.04E-12	8.08E-08	4.15E-06	6.04E-08	2.14E-09	0.00E+00	1.84E-05	3.60E-04	5.68E-04	9.57E-04
Subsistence Harvester	Thorium-230	0.00E+00	0.00E+00	7.25E-04	4.91E-10	2.27E-06	3.47E-05	7.89E-07	1.95E-08	0.00E+00	9.70E-04	1.94E-04	4.54E-03	6.47E-03
(Lloyd Lake)	Radium-226	0.00E+00	0.00E+00	5.89E-04	2.63E-07	2.71E-06	7.02E-02	3.26E-06	3.91E-04	0.00E+00	1.57E-03	4.47E-03	1.03E-02	8.76E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	1.18E-02	3.28E-03	3.96E-02	5.95E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	7.46E-02	7.71E-03	2.23E-01	3.07E-01
	Total by Pathway	0.00E+00	0.00E+00	5.00E-03	2.64E-07	2.73E-05	7.45E-02	1.28E-04	3.94E-04	0.00E+00	8.90E-02	1.63E-02	2.78E-01	0.46

Table C17: Estimated Total Radiation Doses to Human Receptors – Far-Future – Application Case, Upper Bound Scenario

						To	otal Dose by	Pathway - Fa	r-Future (mS	v/yr)				
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
							-	Application C	ase					
	Uranium-238	0.00E+00	0.00E+00	6.97E-06	5.29E-11	7.43E-08	8.24E-04	6.35E-08	6.78E-07	0.00E+00	1.94E-05	3.31E-04	5.22E-04	1.70E-03
	Uranium-234	0.00E+00	0.00E+00	7.59E-06	9.22E-12	8.09E-08	4.15E-06	6.92E-08	2.46E-09	0.00E+00	2.11E-05	3.60E-04	5.69E-04	9.62E-04
	Thorium-230	0.00E+00	0.00E+00	7.25E-04	4.92E-10	2.27E-06	3.48E-05	7.89E-07	1.95E-08	0.00E+00	9.71E-04	1.95E-04	4.54E-03	6.47E-03
	Radium-226	0.00E+00	0.00E+00	5.90E-04	2.63E-07	2.71E-06	7.03E-02	3.27E-06	3.92E-04	0.00E+00	1.57E-03	4.47E-03	1.03E-02	8.76E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	1.18E-02	3.28E-03	3.96E-02	5.95E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	7.46E-02	7.71E-03	2.23E-01	3.07E-01
	Total by Pathway	0.00E+00	0.00E+00	5.00E-03	2.65E-07	2.73E-05	7.46E-02	1.28E-04	3.95E-04	0.00E+00	8.90E-02	1.63E-02	2.78E-01	0.46
							Upp	per Bound Sc	enario					
	Uranium-238	0.00E+00	0.00E+00	6.97E-06	5.29E-11	7.43E-08	8.24E-04	6.35E-08	6.78E-07	0.00E+00	1.94E-05	3.31E-04	5.22E-04	1.70E-03
	Uranium-234	0.00E+00	0.00E+00	7.59E-06	9.22E-12	8.09E-08	4.15E-06	6.92E-08	2.46E-09	0.00E+00	2.11E-05	3.60E-04	5.69E-04	9.62E-04
	Thorium-230	0.00E+00	0.00E+00	7.25E-04	4.92E-10	2.27E-06	3.48E-05	7.89E-07	1.95E-08	0.00E+00	9.71E-04	1.95E-04	4.54E-03	6.47E-03
	Radium-226	0.00E+00	0.00E+00	5.90E-04	2.63E-07	2.71E-06	7.03E-02	3.27E-06	3.92E-04	0.00E+00	1.57E-03	4.47E-03	1.03E-02	8.76E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	1.18E-02	3.28E-03	3.96E-02	5.95E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	7.46E-02	7.71E-03	2.23E-01	3.07E-01
	Total by Pathway	0.00E+00	0.00E+00	5.00E-03	2.65E-07	2.73E-05	7.46E-02	1.28E-04	3.95E-04	0.00E+00	8.90E-02	1.63E-02	2.78E-01	0.46
								Base Case						
	Uranium-238	0.00E+00	0.00E+00	4.22E-06	1.14E-11	3.02E-06	1.10E-03	2.25E-06	7.69E-07	0.00E+00	9.99E-06	4.05E-04	5.67E-04	2.09E-03
	Uranium-234	0.00E+00	0.00E+00	4.58E-06	1.99E-12	3.27E-06	5.56E-06	2.44E-06	2.79E-09	0.00E+00	1.08E-05	4.39E-04	6.15E-04	1.08E-03
	Thorium-230	0.00E+00	0.00E+00	3.69E-04	1.22E-10	6.75E-05	3.95E-05	2.35E-05	2.53E-08	0.00E+00	4.19E-04	1.75E-04	2.77E-03	3.86E-03
	Radium-226	0.00E+00	0.00E+00	5.26E-04	6.50E-08	1.42E-04	9.40E-02	1.71E-04	5.10E-04	0.00E+00	1.19E-03	7.05E-03	1.67E-02	1.20E-01
	Lead-210	0.00E+00	0.00E+00	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	1.36E-02	7.86E-03	8.28E-02	1.15E-01
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	1.21E-01	2.60E-02	3.94E-01	5.56E-01
Subsistence Harvester One-Year-Old (Lloyd	Total by Pathway	0.00E+00	0.00E+00	7.19E-03	6.54E-08	2.21E-03	9.97E-02	1.26E-02	5.14E-04	0.00E+00	1.36E-01	4.20E-02	4.98E-01	0.80
Lake)								Application C	ase					
	Uranium-238	0.00E+00	0.00E+00	4.84E-06	1.31E-11	3.02E-06	1.10E-03	2.58E-06	8.81E-07	0.00E+00	1.14E-05	4.06E-04	5.68E-04	2.10E-03
	Uranium-234	0.00E+00	0.00E+00	5.25E-06	2.28E-12	3.27E-06	5.56E-06	2.80E-06	3.20E-09	0.00E+00	1.24E-05	4.39E-04	6.15E-04	1.08E-03
	Thorium-230	0.00E+00	0.00E+00	3.69E-04	1.22E-10	6.76E-05	3.95E-05	2.35E-05	2.54E-08	0.00E+00	4.19E-04	1.75E-04	2.77E-03	3.87E-03
	Radium-226	0.00E+00	0.00E+00	5.27E-04	6.52E-08	1.42E-04	9.40E-02	1.71E-04	5.11E-04	0.00E+00	1.19E-03	7.06E-03	1.67E-02	1.20E-01
	Lead-210	0.00E+00	0.00E+00	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	1.36E-02	7.86E-03	8.28E-02	1.15E-01
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	1.21E-01	2.60E-02	3.94E-01	5.56E-01
	Total by Pathway	0.00E+00	0.00E+00	7.20E-03	6.56E-08	2.22E-03	9.98E-02	1.26E-02	5.15E-04	0.00E+00	1.36E-01	4.20E-02	4.98E-01	0.80

Table C17: Estimated Total Radiation Doses to Human Receptors – Far-Future – Application Case, Upper Bound Scenario

						To	otal Dose by	Pathway - Fa	r-Future (mS	v/yr)				
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
							Upp	oer Bound Sc	enario					
	Uranium-238	0.00E+00	0.00E+00	4.84E-06	1.31E-11	3.02E-06	1.10E-03	2.58E-06	8.81E-07	0.00E+00	1.14E-05	4.06E-04	5.68E-04	2.10E-03
	Uranium-234	0.00E+00	0.00E+00	5.25E-06	2.28E-12	3.27E-06	5.56E-06	2.80E-06	3.20E-09	0.00E+00	1.24E-05	4.39E-04	6.15E-04	1.08E-03
	Thorium-230	0.00E+00	0.00E+00	3.69E-04	1.22E-10	6.76E-05	3.95E-05	2.35E-05	2.54E-08	0.00E+00	4.19E-04	1.75E-04	2.77E-03	3.87E-03
	Radium-226	0.00E+00	0.00E+00	5.27E-04	6.52E-08	1.42E-04	9.40E-02	1.71E-04	5.11E-04	0.00E+00	1.19E-03	7.06E-03	1.67E-02	1.20E-01
	Lead-210	0.00E+00	0.00E+00	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	1.36E-02	7.86E-03	8.28E-02	1.15E-01
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	1.21E-01	2.60E-02	3.94E-01	5.56E-01
	Total by Pathway	0.00E+00	0.00E+00	7.20E-03	6.56E-08	2.22E-03	9.98E-02	1.26E-02	5.15E-04	0.00E+00	1.36E-01	4.20E-02	4.98E-01	0.80
							,	Base Case			,			
	Uranium-238	0.00E+00	0.00E+00	6.08E-06	4.61E-11	7.42E-08	8.23E-04	5.54E-08	5.91E-07	0.00E+00	5.36E-06	2.59E-04	3.62E-04	1.46E-03
	Uranium-234	0.00E+00	0.00E+00	6.62E-06	8.04E-12	8.08E-08	4.15E-06	6.04E-08	2.14E-09	0.00E+00	5.84E-06	2.82E-04	3.95E-04	6.94E-04
	Thorium-230	0.00E+00	0.00E+00	7.25E-04	4.91E-10	2.27E-06	3.47E-05	7.89E-07	1.95E-08	0.00E+00	3.07E-04	1.52E-04	2.32E-03	3.54E-03
	Radium-226	0.00E+00	0.00E+00	5.89E-04	2.63E-07	2.71E-06	7.02E-02	3.26E-06	3.91E-04	0.00E+00	4.98E-04	3.51E-03	8.84E-03	8.40E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	3.74E-03	2.57E-03	2.70E-02	3.82E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	2.36E-02	6.05E-03	1.23E-01	1.55E-01
	Total by Pathway	0.00E+00	0.00E+00	5.00E-03	2.64E-07	2.73E-05	7.45E-02	1.28E-04	3.94E-04	0.00E+00	2.82E-02	1.28E-02	1.62E-01	0.28
							1	Application C	ase					
	Uranium-238	0.00E+00	0.00E+00	2.19E-05	1.66E-10	7.55E-08	8.38E-04	2.00E-07	2.13E-06	0.00E+00	1.93E-05	2.64E-04	3.71E-04	1.52E-03
	Uranium-234	0.00E+00	0.00E+00	2.39E-05	2.91E-11	8.23E-08	4.22E-06	2.18E-07	7.75E-09	0.00E+00	2.11E-05	2.87E-04	4.04E-04	7.40E-04
Seasonal Resident (Patterson Lake South	Thorium-230	0.00E+00	0.00E+00	7.34E-04	4.97E-10	2.30E-06	3.53E-05	7.99E-07	1.97E-08	0.00E+00	3.11E-04	1.55E-04	2.36E-03	3.60E-03
Arm)	Radium-226	0.00E+00	0.00E+00	6.07E-04	2.71E-07	2.77E-06	7.17E-02	3.37E-06	4.03E-04	0.00E+00	5.14E-04	3.58E-03	8.91E-03	8.57E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.53E-06	0.00E+00	3.75E-03	2.57E-03	2.71E-02	3.83E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.06E-12	7.27E-06	2.16E-07	7.90E-05	1.08E-08	0.00E+00	2.37E-02	6.05E-03	1.23E-01	1.56E-01
	Total by Pathway	0.00E+00	0.00E+00	5.06E-03	2.73E-07	2.74E-05	7.61E-02	1.28E-04	4.08E-04	0.00E+00	2.83E-02	1.29E-02	1.63E-01	0.29
							Upp	er Bound Sc	enario					
	Uranium-238	0.00E+00	0.00E+00	2.19E-05	1.66E-10	7.55E-08	8.38E-04	2.00E-07	2.13E-06	0.00E+00	1.93E-05	2.64E-04	3.71E-04	1.52E-03
	Uranium-234	0.00E+00	0.00E+00	2.39E-05	2.91E-11	8.23E-08	4.22E-06	2.18E-07	7.75E-09	0.00E+00	2.11E-05	2.87E-04	4.04E-04	7.40E-04
	Thorium-230	0.00E+00	0.00E+00	7.34E-04	4.97E-10	2.30E-06	3.53E-05	7.99E-07	1.97E-08	0.00E+00	3.11E-04	1.55E-04	2.36E-03	3.60E-03
	Radium-226	0.00E+00	0.00E+00	6.07E-04	2.71E-07	2.77E-06	7.17E-02	3.37E-06	4.03E-04	0.00E+00	5.14E-04	3.58E-03	8.91E-03	8.57E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.53E-06	0.00E+00	3.75E-03	2.57E-03	2.71E-02	3.83E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.06E-12	7.27E-06	2.16E-07	7.90E-05	1.08E-08	0.00E+00	2.37E-02	6.05E-03	1.23E-01	1.56E-01
	Total by Pathway	0.00E+00	0.00E+00	5.06E-03	2.73E-07	2.74E-05	7.61E-02	1.28E-04	4.08E-04	0.00E+00	2.83E-02	1.29E-02	1.63E-01	0.29

Table C17: Estimated Total Radiation Doses to Human Receptors – Far-Future – Application Case, Upper Bound Scenario

						To	otal Dose by	Pathway - Fa	r-Future (mS	v/yr)				
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
		•				•		Base Case				•		
	Uranium-238	0.00E+00	0.00E+00	4.22E-06	1.14E-11	3.02E-06	1.10E-03	2.25E-06	7.69E-07	0.00E+00	3.16E-06	3.52E-04	4.97E-04	1.96E-03
	Uranium-234	0.00E+00	0.00E+00	4.58E-06	1.99E-12	3.27E-06	5.56E-06	2.44E-06	2.79E-09	0.00E+00	3.43E-06	3.82E-04	5.38E-04	9.40E-04
	Thorium-230	0.00E+00	0.00E+00	3.69E-04	1.22E-10	6.75E-05	3.95E-05	2.35E-05	2.53E-08	0.00E+00	1.33E-04	1.52E-04	1.68E-03	2.46E-03
	Radium-226	0.00E+00	0.00E+00	5.26E-04	6.50E-08	1.42E-04	9.40E-02	1.71E-04	5.10E-04	0.00E+00	3.78E-04	6.13E-03	1.65E-02	1.18E-01
	Lead-210	0.00E+00	0.00E+00	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	4.32E-03	6.84E-03	6.72E-02	8.95E-02
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	3.83E-02	2.26E-02	2.73E-01	3.49E-01
	Total by Pathway	0.00E+00	0.00E+00	7.19E-03	6.54E-08	2.21E-03	9.97E-02	1.26E-02	5.14E-04	0.00E+00	4.32E-02	3.65E-02	3.60E-01	0.56
								Application C	ase					
	Uranium-238	0.00E+00	0.00E+00	1.52E-05	4.11E-11	3.07E-06	1.12E-03	8.12E-06	2.77E-06	0.00E+00	1.14E-05	3.59E-04	5.01E-04	2.02E-03
	Uranium-234	0.00E+00	0.00E+00	1.65E-05	7.19E-12	3.33E-06	5.65E-06	8.83E-06	1.01E-08	0.00E+00	1.24E-05	3.88E-04	5.43E-04	9.78E-04
Seasonal Resident One- Year-Old (Patterson Lake	Thorium-230	0.00E+00	0.00E+00	3.73E-04	1.23E-10	6.86E-05	4.01E-05	2.38E-05	2.57E-08	0.00E+00	1.34E-04	1.54E-04	1.70E-03	2.49E-03
South Arm)	Radium-226	0.00E+00	0.00E+00	5.43E-04	6.71E-08	1.45E-04	9.60E-02	1.76E-04	5.26E-04	0.00E+00	3.90E-04	6.26E-03	1.65E-02	1.21E-01
	Lead-210	0.00E+00	0.00E+00	1.81E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	4.32E-03	6.84E-03	6.73E-02	8.96E-02
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	3.84E-02	2.26E-02	2.74E-01	3.49E-01
	Total by Pathway	0.00E+00	0.00E+00	7.24E-03	6.75E-08	2.22E-03	1.02E-01	1.26E-02	5.32E-04	0.00E+00	4.32E-02	3.66E-02	3.60E-01	0.56
		1	T			Ī	Up _l	per Bound Sc	enario			Ī		
	Uranium-238	0.00E+00	0.00E+00	1.52E-05	4.11E-11	3.07E-06	1.12E-03	8.13E-06	2.77E-06	0.00E+00	1.14E-05	3.59E-04	5.01E-04	2.02E-03
	Uranium-234	0.00E+00	0.00E+00	1.66E-05	7.20E-12	3.33E-06	5.65E-06	8.83E-06	1.01E-08	0.00E+00	1.24E-05	3.88E-04	5.43E-04	9.78E-04
	Thorium-230	0.00E+00	0.00E+00	3.73E-04	1.23E-10	6.86E-05	4.01E-05	2.38E-05	2.57E-08	0.00E+00	1.34E-04	1.54E-04	1.70E-03	2.49E-03
	Radium-226	0.00E+00	0.00E+00	5.43E-04	6.71E-08	1.45E-04	9.60E-02	1.76E-04	5.26E-04	0.00E+00	3.90E-04	6.26E-03	1.65E-02	1.21E-01
	Lead-210	0.00E+00	0.00E+00	1.81E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	4.32E-03	6.84E-03	6.73E-02	8.96E-02
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	3.84E-02	2.26E-02	2.74E-01	3.49E-01
	Total by Pathway	0.00E+00	0.00E+00	7.24E-03	6.75E-08	2.22E-03	1.02E-01	1.26E-02	5.32E-04	0.00E+00	4.32E-02	3.66E-02	3.60E-01	0.56
		1				I		Base Case				I		
	Uranium-238	0.00E+00	0.00E+00	6.08E-06	4.61E-11	7.42E-08	8.23E-04	5.54E-08	5.91E-07	0.00E+00	5.36E-06	2.59E-04	3.62E-04	1.46E-03
	Uranium-234	0.00E+00	0.00E+00	6.62E-06	8.04E-12	8.08E-08	4.15E-06	6.04E-08	2.14E-09	0.00E+00	5.84E-06	2.82E-04	3.95E-04	6.94E-04
Seasonal Resident (Lloyd	Thorium-230	0.00E+00	0.00E+00	7.25E-04	4.91E-10	2.27E-06	3.47E-05	7.89E-07	1.95E-08	0.00E+00	3.07E-04	1.52E-04	2.32E-03	3.54E-03
Lake)	Radium-226	0.00E+00	0.00E+00	5.89E-04	2.63E-07	2.71E-06	7.02E-02	3.26E-06	3.91E-04	0.00E+00	4.98E-04	3.51E-03	8.84E-03	8.40E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	3.74E-03	2.57E-03	2.70E-02	3.82E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	2.36E-02	6.05E-03	1.23E-01	1.55E-01
	Total by Pathway	0.00E+00	0.00E+00	5.00E-03	2.64E-07	2.73E-05	7.45E-02	1.28E-04	3.94E-04	0.00E+00	2.82E-02	1.28E-02	1.62E-01	0.28

Table C17: Estimated Total Radiation Doses to Human Receptors – Far-Future – Application Case, Upper Bound Scenario

						To	otal Dose by	Pathway - Fa	r-Future (mS	v/yr)				
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
							1	Application C	ase					
	Uranium-238	0.00E+00	0.00E+00	6.61E-06	5.02E-11	7.43E-08	8.24E-04	6.03E-08	6.43E-07	0.00E+00	7.63E-06	2.59E-04	3.63E-04	1.46E-03
	Uranium-234	0.00E+00	0.00E+00	7.20E-06	8.74E-12	8.09E-08	4.15E-06	6.57E-08	2.33E-09	0.00E+00	8.31E-06	2.82E-04	3.95E-04	6.97E-04
	Thorium-230	0.00E+00	0.00E+00	7.25E-04	4.91E-10	2.27E-06	3.48E-05	7.89E-07	1.95E-08	0.00E+00	3.08E-04	1.53E-04	2.32E-03	3.54E-03
	Radium-226	0.00E+00	0.00E+00	5.89E-04	2.63E-07	2.71E-06	7.02E-02	3.27E-06	3.91E-04	0.00E+00	5.00E-04	3.51E-03	8.84E-03	8.41E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	4.50E-03	2.57E-03	2.70E-02	3.89E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	2.41E-02	6.05E-03	1.23E-01	1.56E-01
	Total by Pathway	0.00E+00	0.00E+00	5.00E-03	2.65E-07	2.73E-05	7.46E-02	1.28E-04	3.95E-04	0.00E+00	2.95E-02	1.28E-02	1.62E-01	0.28
							Up _l	oer Bound Sc	enario					
	Uranium-238	0.00E+00	0.00E+00	6.61E-06	5.02E-11	7.43E-08	8.24E-04	6.03E-08	6.43E-07	0.00E+00	7.63E-06	2.59E-04	3.63E-04	1.46E-03
	Uranium-234	0.00E+00	0.00E+00	7.20E-06	8.75E-12	8.09E-08	4.15E-06	6.57E-08	2.33E-09	0.00E+00	8.31E-06	2.82E-04	3.95E-04	6.97E-04
	Thorium-230	0.00E+00	0.00E+00	7.25E-04	4.91E-10	2.27E-06	3.48E-05	7.89E-07	1.95E-08	0.00E+00	3.08E-04	1.53E-04	2.32E-03	3.54E-03
	Radium-226	0.00E+00	0.00E+00	5.89E-04	2.63E-07	2.71E-06	7.02E-02	3.27E-06	3.91E-04	0.00E+00	5.00E-04	3.51E-03	8.84E-03	8.41E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	4.50E-03	2.57E-03	2.70E-02	3.89E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	2.41E-02	6.05E-03	1.23E-01	1.56E-01
	Total by Pathway	0.00E+00	0.00E+00	5.00E-03	2.65E-07	2.73E-05	7.46E-02	1.28E-04	3.95E-04	0.00E+00	2.95E-02	1.28E-02	1.62E-01	0.28
								Base Case						
	Uranium-238	0.00E+00	0.00E+00	4.22E-06	1.14E-11	3.02E-06	1.10E-03	2.25E-06	7.69E-07	0.00E+00	3.16E-06	3.52E-04	4.97E-04	1.96E-03
	Uranium-234	0.00E+00	0.00E+00	4.58E-06	1.99E-12	3.27E-06	5.56E-06	2.44E-06	2.79E-09	0.00E+00	3.43E-06	3.82E-04	5.38E-04	9.40E-04
	Thorium-230	0.00E+00	0.00E+00	3.69E-04	1.22E-10	6.75E-05	3.95E-05	2.35E-05	2.53E-08	0.00E+00	1.33E-04	1.52E-04	1.68E-03	2.46E-03
	Radium-226	0.00E+00	0.00E+00	5.26E-04	6.50E-08	1.42E-04	9.40E-02	1.71E-04	5.10E-04	0.00E+00	3.78E-04	6.13E-03	1.65E-02	1.18E-01
	Lead-210	0.00E+00	0.00E+00	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	4.32E-03	6.84E-03	6.72E-02	8.95E-02
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	3.83E-02	2.26E-02	2.73E-01	3.49E-01
Seasonal Resident One-	Total by Pathway	0.00E+00	0.00E+00	7.19E-03	6.54E-08	2.21E-03	9.97E-02	1.26E-02	5.14E-04	0.00E+00	4.32E-02	3.65E-02	3.60E-01	0.56
Year-Old (Lloyd Lake)								Application C	ase					
	Uranium-238	0.00E+00	0.00E+00	4.59E-06	1.24E-11	3.02E-06	1.10E-03	2.45E-06	8.36E-07	0.00E+00	3.44E-06	3.53E-04	4.97E-04	1.97E-03
	Uranium-234	0.00E+00	0.00E+00	4.98E-06	2.17E-12	3.27E-06	5.56E-06	2.66E-06	3.03E-09	0.00E+00	3.73E-06	3.82E-04	5.39E-04	9.41E-04
	Thorium-230	0.00E+00	0.00E+00	3.69E-04	1.22E-10	6.75E-05	3.95E-05	2.35E-05	2.54E-08	0.00E+00	1.33E-04	1.52E-04	1.68E-03	2.46E-03
	Radium-226	0.00E+00	0.00E+00	5.27E-04	6.51E-08	1.42E-04	9.40E-02	1.71E-04	5.11E-04	0.00E+00	3.78E-04	6.13E-03	1.65E-02	1.18E-01
	Lead-210	0.00E+00	0.00E+00	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	4.32E-03	6.84E-03	6.72E-02	8.95E-02
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	3.83E-02	2.26E-02	2.73E-01	3.49E-01
	Total by Pathway	0.00E+00	0.00E+00	7.20E-03	6.55E-08	2.21E-03	9.98E-02	1.26E-02	5.15E-04	0.00E+00	4.32E-02	3.65E-02	3.60E-01	0.56

Table C17: Estimated Total Radiation Doses to Human Receptors – Far-Future – Application Case, Upper Bound Scenario

						To	otal Dose by	Pathway - Fa	r-Future (mS	v/yr)				
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
			•		•		Upj	per Bound Sc	enario					
	Uranium-238	0.00E+00	0.00E+00	4.59E-06	1.24E-11	3.02E-06	1.10E-03	2.45E-06	8.36E-07	0.00E+00	3.44E-06	3.53E-04	4.97E-04	1.97E-03
	Uranium-234	0.00E+00	0.00E+00	4.98E-06	2.17E-12	3.27E-06	5.56E-06	2.66E-06	3.03E-09	0.00E+00	3.73E-06	3.82E-04	5.39E-04	9.41E-04
	Thorium-230	0.00E+00	0.00E+00	3.69E-04	1.22E-10	6.75E-05	3.95E-05	2.35E-05	2.54E-08	0.00E+00	1.33E-04	1.52E-04	1.68E-03	2.46E-03
	Radium-226	0.00E+00	0.00E+00	5.27E-04	6.51E-08	1.42E-04	9.40E-02	1.71E-04	5.11E-04	0.00E+00	3.78E-04	6.13E-03	1.65E-02	1.18E-01
	Lead-210	0.00E+00	0.00E+00	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	4.32E-03	6.84E-03	6.72E-02	8.95E-02
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	3.83E-02	2.26E-02	2.73E-01	3.49E-01
	Total by Pathway	0.00E+00	0.00E+00	7.20E-03	6.55E-08	2.21E-03	9.98E-02	1.26E-02	5.15E-04	0.00E+00	4.32E-02	3.65E-02	3.60E-01	0.56
								Base Case						
	Uranium-238	0.00E+00	0.00E+00	6.08E-06	4.61E-11	7.42E-08	8.23E-04	5.54E-08	5.91E-07	0.00E+00	1.69E-05	3.30E-04	5.22E-04	1.70E-03
	Uranium-234	0.00E+00	0.00E+00	6.62E-06	8.04E-12	8.08E-08	4.15E-06	6.04E-08	2.14E-09	0.00E+00	1.84E-05	3.60E-04	5.68E-04	9.57E-04
	Thorium-230	0.00E+00	0.00E+00	7.25E-04	4.91E-10	2.27E-06	3.47E-05	7.89E-07	1.95E-08	0.00E+00	9.70E-04	1.94E-04	4.54E-03	6.47E-03
	Radium-226	0.00E+00	0.00E+00	5.89E-04	2.63E-07	2.71E-06	7.02E-02	3.26E-06	3.91E-04	0.00E+00	1.57E-03	4.47E-03	1.03E-02	8.76E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	1.18E-02	3.28E-03	3.96E-02	5.95E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	7.46E-02	7.71E-03	2.23E-01	3.07E-01
	Total by Pathway	0.00E+00	0.00E+00	5.00E-03	2.64E-07	2.73E-05	7.45E-02	1.28E-04	3.94E-04	0.00E+00	8.90E-02	1.63E-02	2.78E-01	0.46
							1	Application C	ase					
	Uranium-238	0.00E+00	0.00E+00	1.97E-04	1.49E-09	1.01E-07	1.12E-03	1.79E-06	1.91E-05	0.00E+00	5.48E-04	4.48E-04	7.60E-04	3.09E-03
	Uranium-234	0.00E+00	0.00E+00	2.15E-04	2.61E-10	1.10E-07	5.63E-06	1.96E-06	6.97E-08	0.00E+00	5.99E-04	4.88E-04	8.28E-04	2.14E-03
Permanent Resident (Patterson Lake North	Thorium-230	0.00E+00	0.00E+00	7.75E-04	5.25E-10	3.00E-06	4.60E-05	8.43E-07	2.08E-08	0.00E+00	1.04E-03	2.57E-04	5.07E-03	7.19E-03
Arm - West Basin)	Radium-226	0.00E+00	0.00E+00	7.59E-04	3.39E-07	3.91E-06	1.01E-01	4.20E-06	5.04E-04	0.00E+00	2.03E-03	6.44E-03	1.22E-02	1.23E-01
ŕ	Lead-210	0.00E+00	0.00E+00	1.39E-03	1.22E-09	1.49E-05	3.46E-03	4.67E-05	2.64E-06	0.00E+00	1.24E-02	3.28E-03	4.05E-02	6.10E-02
	Polonium-210	0.00E+00	0.00E+00	2.46E-03	7.38E-12	7.28E-06	2.16E-07	8.26E-05	1.13E-08	0.00E+00	7.80E-02	7.72E-03	2.30E-01	3.18E-01
	Total by Pathway	0.00E+00	0.00E+00	5.79E-03	3.42E-07	2.93E-05	1.06E-01	1.38E-04	5.26E-04	0.00E+00	9.46E-02	1.86E-02	2.89E-01	0.51
							Upp	per Bound Sc	enario				,	
	Uranium-238	0.00E+00	0.00E+00	1.97E-04	1.49E-09	1.01E-07	1.12E-03	1.80E-06	1.92E-05	0.00E+00	5.48E-04	4.48E-04	7.60E-04	3.09E-03
	Uranium-234	0.00E+00	0.00E+00	2.15E-04	2.61E-10	1.10E-07	5.63E-06	1.96E-06	6.97E-08	0.00E+00	6.00E-04	4.88E-04	8.28E-04	2.14E-03
	Thorium-230	0.00E+00	0.00E+00	7.75E-04	5.25E-10	3.00E-06	4.60E-05	8.43E-07	2.08E-08	0.00E+00	1.04E-03	2.57E-04	5.07E-03	7.19E-03
	Radium-226	0.00E+00	0.00E+00	7.59E-04	3.39E-07	3.91E-06	1.01E-01	4.21E-06	5.04E-04	0.00E+00	2.03E-03	6.44E-03	1.22E-02	1.23E-01
	Lead-210	0.00E+00	0.00E+00	1.39E-03	1.22E-09	1.49E-05	3.46E-03	4.67E-05	2.64E-06	0.00E+00	1.24E-02	3.28E-03	4.05E-02	6.10E-02
	Polonium-210	0.00E+00	0.00E+00	2.46E-03	7.38E-12	7.28E-06	2.16E-07	8.26E-05	1.13E-08	0.00E+00	7.80E-02	7.72E-03	2.30E-01	3.18E-01
	Total by Pathway	0.00E+00	0.00E+00	5.79E-03	3.42E-07	2.93E-05	1.06E-01	1.38E-04	5.26E-04	0.00E+00	9.46E-02	1.86E-02	2.89E-01	0.51

Table C17: Estimated Total Radiation Doses to Human Receptors – Far-Future – Application Case, Upper Bound Scenario

						To	otal Dose by	Pathway - Fa	r-Future (mS	v/yr)				
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
								Base Case						
	Uranium-238	0.00E+00	0.00E+00	4.22E-06	1.14E-11	3.02E-06	1.10E-03	2.25E-06	7.69E-07	0.00E+00	9.99E-06	4.05E-04	5.67E-04	2.09E-03
	Uranium-234	0.00E+00	0.00E+00	4.58E-06	1.99E-12	3.27E-06	5.56E-06	2.44E-06	2.79E-09	0.00E+00	1.08E-05	4.39E-04	6.15E-04	1.08E-03
	Thorium-230	0.00E+00	0.00E+00	3.69E-04	1.22E-10	6.75E-05	3.95E-05	2.35E-05	2.53E-08	0.00E+00	4.19E-04	1.75E-04	2.77E-03	3.86E-03
	Radium-226	0.00E+00	0.00E+00	5.26E-04	6.50E-08	1.42E-04	9.40E-02	1.71E-04	5.10E-04	0.00E+00	1.19E-03	7.05E-03	1.67E-02	1.20E-01
	Lead-210	0.00E+00	0.00E+00	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	1.36E-02	7.86E-03	8.28E-02	1.15E-01
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	1.21E-01	2.60E-02	3.94E-01	5.56E-01
	Total by Pathway	0.00E+00	0.00E+00	7.19E-03	6.54E-08	2.21E-03	9.97E-02	1.26E-02	5.14E-04	0.00E+00	1.36E-01	4.20E-02	4.98E-01	0.80
							1	Application C	ase					
	Uranium-238	0.00E+00	0.00E+00	1.37E-04	3.69E-10	4.09E-06	1.49E-03	7.30E-05	2.49E-05	0.00E+00	3.23E-04	5.50E-04	6.75E-04	3.28E-03
	Uranium-234	0.00E+00	0.00E+00	1.49E-04	6.47E-11	4.44E-06	7.53E-06	7.94E-05	9.06E-08	0.00E+00	3.52E-04	5.96E-04	7.32E-04	1.92E-03
Permanent Resident One- Year-Old (Patterson Lake	Thorium-230	0.00E+00	0.00E+00	3.94E-04	1.30E-10	8.94E-05	5.23E-05	2.51E-05	2.71E-08	0.00E+00	4.48E-04	2.31E-04	3.02E-03	4.26E-03
North Arm - West Basin)	Radium-226	0.00E+00	0.00E+00	6.78E-04	8.39E-08	2.04E-04	1.35E-01	2.20E-04	6.57E-04	0.00E+00	1.54E-03	1.02E-02	1.75E-02	1.66E-01
	Lead-210	0.00E+00	0.00E+00	1.89E-03	2.48E-10	1.19E-03	4.63E-03	3.71E-03	3.43E-06	0.00E+00	1.43E-02	7.86E-03	8.39E-02	1.17E-01
	Polonium-210	0.00E+00	0.00E+00	4.69E-03	1.83E-12	8.14E-04	2.82E-07	9.23E-03	1.47E-08	0.00E+00	1.27E-01	2.60E-02	4.04E-01	5.71E-01
	Total by Pathway	0.00E+00	0.00E+00	7.94E-03	8.47E-08	2.30E-03	1.42E-01	1.33E-02	6.86E-04	0.00E+00	1.44E-01	4.54E-02	5.09E-01	0.86
							Upp	oer Bound Sc	enario			1		
	Uranium-238	0.00E+00	0.00E+00	1.37E-04	3.69E-10	4.09E-06	1.49E-03	7.30E-05	2.49E-05	0.00E+00	3.24E-04	5.50E-04	6.75E-04	3.28E-03
	Uranium-234	0.00E+00	0.00E+00	1.49E-04	6.47E-11	4.44E-06	7.53E-06	7.94E-05	9.06E-08	0.00E+00	3.52E-04	5.96E-04	7.32E-04	1.92E-03
	Thorium-230	0.00E+00	0.00E+00	3.94E-04	1.30E-10	8.94E-05	5.23E-05	2.51E-05	2.71E-08	0.00E+00	4.48E-04	2.31E-04	3.02E-03	4.26E-03
	Radium-226	0.00E+00	0.00E+00	6.79E-04	8.39E-08	2.04E-04	1.35E-01	2.20E-04	6.58E-04	0.00E+00	1.54E-03	1.02E-02	1.75E-02	1.66E-01
	Lead-210	0.00E+00	0.00E+00	1.89E-03	2.48E-10	1.19E-03	4.63E-03	3.71E-03	3.43E-06	0.00E+00	1.43E-02	7.86E-03	8.39E-02	1.17E-01
	Polonium-210	0.00E+00	0.00E+00	4.69E-03	1.83E-12	8.14E-04	2.82E-07	9.24E-03	1.47E-08	0.00E+00	1.27E-01	2.60E-02	4.04E-01	5.71E-01
	Total by Pathway	0.00E+00	0.00E+00	7.94E-03	8.47E-08	2.30E-03	1.42E-01	1.33E-02	6.86E-04	0.00E+00	1.44E-01	4.54E-02	5.09E-01	0.86

Table C18: Estimated Total Radiation Doses to Human Receptors – Reasonably Foreseeable Development Case

			Cam	p Worker (P	atterson Lak	e North Arm	- West Basin)	Incremental	Dose by Pat	hway RFD Ca	se (mSv/yr)			
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
								Base Case						
	Uranium-238	0.00E+00	0.00E+00	6.08E-06	4.61E-11	7.42E-08	8.23E-04	5.54E-08	5.91E-07	0.00E+00	8.47E-06	1.65E-04	4.15E-04	1.42E-03
	Uranium-234	0.00E+00	0.00E+00	6.62E-06	8.04E-12	8.08E-08	4.15E-06	6.04E-08	2.14E-09	0.00E+00	9.22E-06	1.80E-04	4.52E-04	6.52E-04
	Thorium-230	0.00E+00	0.00E+00	7.25E-04	4.91E-10	2.27E-06	3.47E-05	7.89E-07	1.95E-08	0.00E+00	4.85E-04	9.73E-05	2.70E-03	4.05E-03
	Radium-226	0.00E+00	0.00E+00	5.89E-04	2.63E-07	2.71E-06	7.02E-02	3.26E-06	3.91E-04	0.00E+00	7.86E-04	2.24E-03	9.32E-03	8.35E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	5.91E-03	1.64E-03	2.93E-02	4.17E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	3.73E-02	3.86E-03	1.32E-01	1.75E-01
	Total by Pathway	0.00E+00	0.00E+00	5.00E-03	2.64E-07	2.73E-05	7.45E-02	1.28E-04	3.94E-04	0.00E+00	4.45E-02	8.18E-03	1.74E-01	0.31
								RFD Case						
	Uranium-238	8.63E-04	2.37E-12	6.10E-06	4.63E-11	2.10E-07	2.33E-03	5.55E-08	5.92E-07	0.00E+00	7.07E-05	2.22E-04	4.87E-04	3.98E-03
	Uranium-234	1.04E-03	5.79E-12	6.64E-06	8.07E-12	2.29E-07	1.17E-05	6.04E-08	2.15E-09	0.00E+00	7.72E-05	2.41E-04	5.31E-04	1.91E-03
Camp Worker (Patterson Lake North	Thorium-230	4.17E-03	1.40E-11	7.26E-04	4.92E-10	2.98E-06	4.56E-05	7.90E-07	1.95E-08	0.00E+00	1.62E-03	7.09E-04	9.33E-03	1.66E-02
Arm - West Basin)	Radium-226	1.04E-03	2.69E-10	5.89E-04	2.63E-07	3.67E-06	9.50E-02	3.26E-06	3.91E-04	0.00E+00	1.16E-03	3.30E-03	1.01E-02	1.12E-01
	Lead-210	3.27E-04	4.23E-11	1.33E-03	1.17E-09	1.64E-05	3.81E-03	4.46E-05	2.53E-06	0.00E+00	7.50E-03	3.93E-03	3.58E-02	5.28E-02
	Polonium-210	9.82E-04	3.69E-13	2.35E-03	7.06E-12	9.97E-06	2.95E-07	7.90E-05	1.08E-08	0.00E+00	4.68E-02	7.51E-03	1.48E-01	2.05E-01
	Total by Pathway	8.42E-03	3.34E-10	5.01E-03	2.65E-07	3.35E-05	1.01E-01	1.28E-04	3.94E-04	0.00E+00	5.72E-02	1.59E-02	2.04E-01	0.39
					,		,	Far-Future	•	,	,			
	Uranium-238	-	-	-	-	-	-	-	-	-	-	-	-	-
	Uranium-234	-	-	-	-	-	-	-	-	-	-	-	-	-
	Thorium-230	-	-	-	-	-	-	-	-	-	-	-	-	-
	Radium-226	-	-	-	-	-	-	-	-	-	-	-	-	-
	Lead-210	-	-	-	-	-	-	-	-	-	-	-	-	-
	Polonium-210	-	-	-	-	-	-	-	-	-	-	-	-	-
	Total by Pathway	-	-	-	-	-	-	-	-	-	-	-	-	-
								Base Case						
	Uranium-238	0.00E+00	0.00E+00	6.08E-06	4.61E-11	7.42E-08	8.23E-04	5.54E-08	5.91E-07	0.00E+00	1.69E-05	3.30E-04	5.22E-04	1.70E-03
Colodor	Uranium-234	0.00E+00	0.00E+00	6.62E-06	8.04E-12	8.08E-08	4.15E-06	6.04E-08	2.14E-09	0.00E+00	1.84E-05	3.60E-04	5.68E-04	9.57E-04
Subsistence Harvester (Patterson Lake South	Thorium-230	0.00E+00	0.00E+00	7.25E-04	4.91E-10	2.27E-06	3.47E-05	7.89E-07	1.95E-08	0.00E+00	9.70E-04	1.94E-04	4.54E-03	6.47E-03
Arm)	Radium-226	0.00E+00	0.00E+00	5.89E-04	2.63E-07	2.71E-06	7.02E-02	3.26E-06	3.91E-04	0.00E+00	1.57E-03	4.47E-03	1.03E-02	8.76E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	1.18E-02	3.28E-03	3.96E-02	5.95E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	7.46E-02	7.71E-03	2.23E-01	3.08E-01
	Total by Pathway	0.00E+00	0.00E+00	5.00E-03	2.64E-07	2.73E-05	7.45E-02	1.28E-04	3.94E-04	0.00E+00	8.90E-02	1.63E-02	2.78E-01	0.46

Table C18: Estimated Total Radiation Doses to Human Receptors – Reasonably Foreseeable Development Case

			Can	np Worker (P	atterson Lake	North Arm	- West Basin)	Incremental	Dose by Pat	hway RFD Ca	se (mSv/yr)			
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
								RFD Case						
	Uranium-238	1.57E-04	4.30E-13	5.25E-05	3.98E-10	9.89E-08	1.10E-03	1.83E-07	1.95E-06	0.00E+00	1.41E-04	4.43E-04	6.66E-04	2.56E-03
	Uranium-234	1.89E-04	1.05E-12	5.73E-05	6.96E-11	1.08E-07	5.53E-06	2.00E-07	7.10E-09	0.00E+00	1.54E-04	4.82E-04	7.26E-04	1.61E-03
	Thorium-230	7.56E-04	2.55E-12	2.56E-03	1.73E-09	2.40E-06	3.67E-05	2.08E-06	5.15E-08	0.00E+00	3.23E-03	1.42E-03	1.78E-02	2.58E-02
	Radium-226	1.89E-04	4.88E-11	8.95E-04	4.00E-07	2.89E-06	7.47E-02	3.85E-06	4.62E-04	0.00E+00	2.31E-03	6.59E-03	1.19E-02	9.71E-02
	Lead-210	5.94E-05	7.68E-12	1.69E-03	1.49E-09	1.52E-05	3.52E-03	4.97E-05	2.81E-06	0.00E+00	1.50E-02	7.85E-03	5.26E-02	8.07E-02
	Polonium-210	1.78E-04	6.70E-14	2.97E-03	8.93E-12	7.76E-06	2.30E-07	8.78E-05	1.21E-08	0.00E+00	9.35E-02	1.50E-02	2.55E-01	3.67E-01
	Total by Pathway	1.53E-03	6.05E-11	8.23E-03	4.03E-07	2.84E-05	7.94E-02	1.44E-04	4.67E-04	0.00E+00	1.14E-01	3.18E-02	3.39E-01	0.57
								Far-Future						
	Uranium-238	0.00E+00	0.00E+00	3.25E-05	2.46E-10	7.68E-08	8.52E-04	2.96E-07	3.16E-06	0.00E+00	9.04E-05	3.42E-04	5.58E-04	1.88E-03
	Uranium-234	0.00E+00	0.00E+00	3.55E-05	4.31E-11	8.37E-08	4.30E-06	3.23E-07	1.15E-08	0.00E+00	9.88E-05	3.72E-04	6.08E-04	1.12E-03
	Thorium-230	0.00E+00	0.00E+00	7.40E-04	5.01E-10	2.34E-06	3.58E-05	8.05E-07	1.99E-08	0.00E+00	9.90E-04	2.01E-04	4.73E-03	6.70E-03
	Radium-226	0.00E+00	0.00E+00	6.20E-04	2.77E-07	2.83E-06	7.32E-02	3.43E-06	4.12E-04	0.00E+00	1.66E-03	4.66E-03	1.07E-02	9.13E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.47E-05	2.53E-06	0.00E+00	1.18E-02	3.28E-03	4.00E-02	5.99E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.06E-12	7.27E-06	2.16E-07	7.90E-05	1.08E-08	0.00E+00	7.47E-02	7.71E-03	2.23E-01	3.08E-01
	Total by Pathway	0.00E+00	0.00E+00	5.11E-03	2.79E-07	2.75E-05	7.76E-02	1.29E-04	4.17E-04	0.00E+00	8.94E-02	1.66E-02	2.80E-01	0.47
								Base Case						_
	Uranium-238	0.00E+00	0.00E+00	4.22E-06	1.14E-11	3.02E-06	1.10E-03	2.25E-06	7.69E-07	0.00E+00	9.99E-06	4.05E-04	5.67E-04	2.09E-03
	Uranium-234	0.00E+00	0.00E+00	4.58E-06	1.99E-12	3.27E-06	5.56E-06	2.44E-06	2.79E-09	0.00E+00	1.08E-05	4.39E-04	6.15E-04	1.08E-03
	Thorium-230	0.00E+00	0.00E+00	3.69E-04	1.22E-10	6.75E-05	3.95E-05	2.35E-05	2.53E-08	0.00E+00	4.19E-04	1.75E-04	2.77E-03	3.86E-03
	Radium-226	0.00E+00	0.00E+00	5.26E-04	6.50E-08	1.42E-04	9.40E-02	1.71E-04	5.10E-04	0.00E+00	1.19E-03	7.05E-03	1.67E-02	1.20E-01
	Lead-210	0.00E+00	0.00E+00	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	1.36E-02	7.86E-03	8.28E-02	1.15E-01
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	1.21E-01	2.60E-02	3.94E-01	5.56E-01
Subsistence Harvester One-Year-Old (Patterson	Total by Pathway	0.00E+00	0.00E+00	7.19E-03	6.54E-08	2.21E-03	9.97E-02	1.26E-02	5.14E-04	0.00E+00	1.36E-01	4.20E-02	4.98E-01	0.80
Lake South Arm)						,		RFD Case		,				
	Uranium-238	1.56E-04	5.59E-13	3.65E-05	9.84E-11	4.02E-06	1.47E-03	7.45E-06	2.54E-06	0.00E+00	8.33E-05	5.44E-04	6.33E-04	2.93E-03
	Uranium-234	1.83E-04	1.37E-12	3.96E-05	1.72E-11	4.36E-06	7.40E-06	8.09E-06	9.23E-09	0.00E+00	9.06E-05	5.90E-04	6.85E-04	1.61E-03
	Thorium-230	5.81E-04	3.31E-12	1.30E-03	4.29E-10	7.14E-05	4.18E-05	6.21E-05	6.70E-08	0.00E+00	1.40E-03	8.44E-04	9.16E-03	1.35E-02
	Radium-226	1.83E-04	6.34E-11	8.00E-04	9.90E-08	1.51E-04	1.00E-01	2.01E-04	6.02E-04	0.00E+00	1.75E-03	9.46E-03	1.74E-02	1.31E-01
	Lead-210	6.14E-05	9.96E-12	2.30E-03	3.03E-10	1.21E-03	4.71E-03	3.95E-03	3.65E-06	0.00E+00	1.73E-02	1.53E-02	9.88E-02	1.44E-01
	Polonium-210	1.83E-04	8.71E-14	5.68E-03	2.21E-12	8.68E-04	3.01E-07	9.82E-03	1.57E-08	0.00E+00	1.52E-01	4.33E-02	4.31E-01	6.42E-01
	Total by Pathway	1.35E-03	7.87E-11	1.02E-02	9.98E-08	2.31E-03	1.06E-01	1.41E-02	6.09E-04	0.00E+00	1.72E-01	7.00E-02	5.58E-01	0.93

Table C18: Estimated Total Radiation Doses to Human Receptors – Reasonably Foreseeable Development Case

			Can	np Worker (P	atterson Lake	e North Arm	- West Basin	Incremental	Dose by Pat	hway RFD Ca	se (mSv/yr)			
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
								Far-Future						
	Uranium-238	0.00E+00	0.00E+00	2.26E-05	6.09E-11	3.12E-06	1.14E-03	1.20E-05	4.11E-06	0.00E+00	5.33E-05	4.20E-04	5.84E-04	2.24E-03
	Uranium-234	0.00E+00	0.00E+00	2.45E-05	1.07E-11	3.38E-06	5.75E-06	1.31E-05	1.49E-08	0.00E+00	5.80E-05	4.55E-04	6.33E-04	1.19E-03
	Thorium-230	0.00E+00	0.00E+00	3.76E-04	1.24E-10	6.97E-05	4.08E-05	2.40E-05	2.59E-08	0.00E+00	4.28E-04	1.80E-04	2.86E-03	3.98E-03
	Radium-226	0.00E+00	0.00E+00	5.54E-04	6.85E-08	1.48E-04	9.80E-02	1.80E-04	5.37E-04	0.00E+00	1.26E-03	7.36E-03	1.69E-02	1.25E-01
	Lead-210	0.00E+00	0.00E+00	1.81E-03	2.38E-10	1.19E-03	4.63E-03	3.55E-03	3.29E-06	0.00E+00	1.37E-02	7.86E-03	8.33E-02	1.16E-01
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.84E-03	1.41E-08	0.00E+00	1.21E-01	2.60E-02	3.95E-01	5.56E-01
	Total by Pathway	0.00E+00	0.00E+00	7.28E-03	6.89E-08	2.22E-03	1.04E-01	1.26E-02	5.44E-04	0.00E+00	1.37E-01	4.23E-02	4.99E-01	0.80
								Base Case						
	Uranium-238	0.00E+00	0.00E+00	6.08E-06	4.61E-11	7.42E-08	8.23E-04	5.54E-08	5.91E-07	0.00E+00	1.69E-05	3.30E-04	5.22E-04	1.70E-03
	Uranium-234	0.00E+00	0.00E+00	6.62E-06	8.04E-12	8.08E-08	4.15E-06	6.04E-08	2.14E-09	0.00E+00	1.84E-05	3.60E-04	5.68E-04	9.57E-04
	Thorium-230	0.00E+00	0.00E+00	7.25E-04	4.91E-10	2.27E-06	3.47E-05	7.89E-07	1.95E-08	0.00E+00	9.70E-04	1.94E-04	4.54E-03	6.47E-03
	Radium-226	0.00E+00	0.00E+00	5.89E-04	2.63E-07	2.71E-06	7.02E-02	3.26E-06	3.91E-04	0.00E+00	1.57E-03	4.47E-03	1.03E-02	8.76E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	1.18E-02	3.28E-03	3.96E-02	5.95E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	7.46E-02	7.71E-03	2.23E-01	3.08E-01
	Total by Pathway	0.00E+00	0.00E+00	5.00E-03	2.64E-07	2.73E-05	7.45E-02	1.28E-04	3.94E-04	0.00E+00	8.90E-02	1.63E-02	2.78E-01	0.46
								RFD Case						
	Uranium-238	8.56E-05	2.35E-13	2.22E-05	1.68E-10	8.77E-08	9.73E-04	1.03E-07	1.09E-06	0.00E+00	6.01E-05	3.92E-04	5.79E-04	2.11E-03
	Uranium-234	1.03E-04	5.75E-13	2.42E-05	2.94E-11	9.55E-08	4.90E-06	1.12E-07	3.97E-09	0.00E+00	6.56E-05	4.27E-04	6.30E-04	1.26E-03
Subsistence Harvester	Thorium-230	4.13E-04	1.39E-12	1.55E-03	1.05E-09	2.34E-06	3.58E-05	1.48E-06	3.65E-08	0.00E+00	1.99E-03	8.63E-04	8.41E-03	1.33E-02
(Beet Lake)	Radium-226	1.03E-04	2.67E-11	7.07E-04	3.15E-07	2.81E-06	7.27E-02	3.52E-06	4.22E-04	0.00E+00	1.86E-03	5.63E-03	1.07E-02	9.21E-02
	Lead-210	3.25E-05	4.20E-12	1.38E-03	1.22E-09	1.51E-05	3.49E-03	4.53E-05	2.57E-06	0.00E+00	1.23E-02	5.77E-03	4.13E-02	6.43E-02
	Polonium-210	9.74E-05	3.66E-14	2.44E-03	7.33E-12	7.54E-06	2.23E-07	8.02E-05	1.10E-08	0.00E+00	7.74E-02	1.17E-02	2.30E-01	3.22E-01
	Total by Pathway	8.35E-04	3.31E-11	6.12E-03	3.18E-07	2.79E-05	7.72E-02	1.31E-04	4.25E-04	0.00E+00	9.36E-02	2.48E-02	2.92E-01	0.50
								Far-Future	•					
	Uranium-238	0.00E+00	0.00E+00	1.59E-05	1.21E-10	7.57E-08	8.39E-04	1.45E-07	1.55E-06	0.00E+00	4.42E-05	3.37E-04	5.34E-04	1.77E-03
	Uranium-234	0.00E+00	0.00E+00	1.73E-05	2.11E-11	8.24E-08	4.23E-06	1.58E-07	5.62E-09	0.00E+00	4.83E-05	3.67E-04	5.82E-04	1.02E-03
	Thorium-230	0.00E+00	0.00E+00	7.33E-04	4.97E-10	2.31E-06	3.53E-05	7.97E-07	1.97E-08	0.00E+00	9.81E-04	1.98E-04	4.59E-03	6.54E-03
	Radium-226	0.00E+00	0.00E+00	6.02E-04	2.69E-07	2.78E-06	7.19E-02	3.34E-06	4.00E-04	0.00E+00	1.61E-03	4.58E-03	1.04E-02	8.95E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	1.18E-02	3.28E-03	3.96E-02	5.95E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	7.46E-02	7.71E-03	2.23E-01	3.08E-01
	Total by Pathway	0.00E+00	0.00E+00	5.04E-03	2.71E-07	2.74E-05	7.62E-02	1.28E-04	4.04E-04	0.00E+00	8.91E-02	1.65E-02	2.78E-01	0.47

Table C18: Estimated Total Radiation Doses to Human Receptors – Reasonably Foreseeable Development Case

			Can	np Worker (P	atterson Lak	e North Arm	- West Basin) Incremental	Dose by Pat	hway RFD Ca	se (mSv/yr)			
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
		!			-			Base Case	1		<u> </u>		-	
	Uranium-238	0.00E+00	0.00E+00	4.22E-06	1.14E-11	3.02E-06	1.10E-03	2.25E-06	7.69E-07	0.00E+00	9.99E-06	4.05E-04	5.67E-04	2.09E-03
	Uranium-234	0.00E+00	0.00E+00	4.58E-06	1.99E-12	3.27E-06	5.56E-06	2.44E-06	2.79E-09	0.00E+00	1.08E-05	4.39E-04	6.15E-04	1.08E-03
	Thorium-230	0.00E+00	0.00E+00	3.69E-04	1.22E-10	6.75E-05	3.95E-05	2.35E-05	2.53E-08	0.00E+00	4.19E-04	1.75E-04	2.77E-03	3.86E-03
	Radium-226	0.00E+00	0.00E+00	5.26E-04	6.50E-08	1.42E-04	9.40E-02	1.71E-04	5.10E-04	0.00E+00	1.19E-03	7.05E-03	1.67E-02	1.20E-01
	Lead-210	0.00E+00	0.00E+00	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	1.36E-02	7.86E-03	8.28E-02	1.15E-01
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	1.21E-01	2.60E-02	3.94E-01	5.56E-01
	Total by Pathway	0.00E+00	0.00E+00	7.19E-03	6.54E-08	2.21E-03	9.97E-02	1.26E-02	5.14E-04	0.00E+00	1.36E-01	4.20E-02	4.98E-01	0.80
								RFD Case						
	Uranium-238	8.53E-05	3.05E-13	1.54E-05	4.16E-11	3.57E-06	1.30E-03	4.17E-06	1.42E-06	0.00E+00	3.54E-05	4.81E-04	5.93E-04	2.52E-03
	Uranium-234	9.99E-05	7.47E-13	1.68E-05	7.29E-12	3.86E-06	6.56E-06	4.53E-06	5.17E-09	0.00E+00	3.85E-05	5.21E-04	6.42E-04	1.33E-03
Subsistence Harvester	Thorium-230	3.18E-04	1.81E-12	7.88E-04	2.60E-10	6.96E-05	4.07E-05	4.40E-05	4.75E-08	0.00E+00	8.59E-04	5.40E-04	4.63E-03	7.29E-03
One-Year-Old (Beet Lake)	Radium-226	9.99E-05	3.46E-11	6.31E-04	7.81E-08	1.47E-04	9.73E-02	1.84E-04	5.50E-04	0.00E+00	1.41E-03	8.37E-03	1.69E-02	1.26E-01
	Lead-210	3.36E-05	5.45E-12	1.88E-03	2.47E-10	1.20E-03	4.67E-03	3.61E-03	3.33E-06	0.00E+00	1.42E-02	1.19E-02	8.48E-02	1.22E-01
	Polonium-210	9.99E-05	4.76E-14	4.66E-03	1.82E-12	8.43E-04	2.92E-07	8.97E-03	1.43E-08	0.00E+00	1.26E-01	3.55E-02	4.03E-01	5.78E-01
	Total by Pathway	7.36E-04	4.30E-11	7.99E-03	7.86E-08	2.26E-03	1.03E-01	1.28E-02	5.55E-04	0.00E+00	1.42E-01	5.73E-02	5.10E-01	0.84
								Far-Future						
	Uranium-238	0.00E+00	0.00E+00	1.10E-05	2.98E-11	3.08E-06	1.12E-03	5.89E-06	2.01E-06	0.00E+00	2.61E-05	4.13E-04	5.73E-04	2.16E-03
	Uranium-234	0.00E+00	0.00E+00	1.20E-05	5.21E-12	3.33E-06	5.66E-06	6.40E-06	7.30E-09	0.00E+00	2.83E-05	4.48E-04	6.21E-04	1.12E-03
	Thorium-230	0.00E+00	0.00E+00	3.73E-04	1.23E-10	6.87E-05	4.02E-05	2.37E-05	2.56E-08	0.00E+00	4.24E-04	1.78E-04	2.79E-03	3.90E-03
	Radium-226	0.00E+00	0.00E+00	5.38E-04	6.65E-08	1.45E-04	9.62E-02	1.74E-04	5.22E-04	0.00E+00	1.22E-03	7.22E-03	1.68E-02	1.23E-01
	Lead-210	0.00E+00	0.00E+00	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	1.36E-02	7.86E-03	8.28E-02	1.15E-01
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	1.21E-01	2.60E-02	3.94E-01	5.56E-01
	Total by Pathway	0.00E+00	0.00E+00	7.23E-03	6.69E-08	2.22E-03	1.02E-01	1.26E-02	5.27E-04	0.00E+00	1.36E-01	4.21E-02	4.98E-01	0.80
								Base Case						
	Uranium-238	0.00E+00	0.00E+00	6.08E-06	4.61E-11	7.42E-08	8.23E-04	5.54E-08	5.91E-07	0.00E+00	1.69E-05	3.30E-04	5.22E-04	1.70E-03
	Uranium-234	0.00E+00	0.00E+00	6.62E-06	8.04E-12	8.08E-08	4.15E-06	6.04E-08	2.14E-09	0.00E+00	1.84E-05	3.60E-04	5.68E-04	9.57E-04
Subsistence Harvester	Thorium-230	0.00E+00	0.00E+00	7.25E-04	4.91E-10	2.27E-06	3.47E-05	7.89E-07	1.95E-08	0.00E+00	9.70E-04	1.94E-04	4.54E-03	6.47E-03
(Lloyd Lake)	Radium-226	0.00E+00	0.00E+00	5.89E-04	2.63E-07	2.71E-06	7.02E-02	3.26E-06	3.91E-04	0.00E+00	1.57E-03	4.47E-03	1.03E-02	8.76E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	1.18E-02	3.28E-03	3.96E-02	5.95E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	7.46E-02	7.71E-03	2.23E-01	3.08E-01
	Total by Pathway	0.00E+00	0.00E+00	5.00E-03	2.64E-07	2.73E-05	7.45E-02	1.28E-04	3.94E-04	0.00E+00	8.90E-02	1.63E-02	2.78E-01	0.46

Table C18: Estimated Total Radiation Doses to Human Receptors – Reasonably Foreseeable Development Case

			Can	np Worker (P	atterson Lake	North Arm	- West Basin)	Incremental	Dose by Pat	hway RFD Ca	se (mSv/yr)			
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
								RFD Case						1
	Uranium-238	4.08E-06	1.12E-14	7.58E-06	5.75E-11	7.49E-08	8.31E-04	5.98E-08	6.37E-07	0.00E+00	2.09E-05	3.33E-04	5.25E-04	1.72E-03
	Uranium-234	4.92E-06	2.74E-14	8.26E-06	1.00E-11	8.15E-08	4.19E-06	6.51E-08	2.31E-09	0.00E+00	2.28E-05	3.63E-04	5.71E-04	9.75E-04
	Thorium-230	1.97E-05	6.64E-14	8.08E-04	5.48E-10	2.27E-06	3.48E-05	8.60E-07	2.13E-08	0.00E+00	1.07E-03	2.26E-04	4.90E-03	7.06E-03
	Radium-226	4.92E-06	1.27E-12	6.00E-04	2.68E-07	2.72E-06	7.03E-02	3.29E-06	3.94E-04	0.00E+00	1.60E-03	4.53E-03	1.03E-02	8.78E-02
	Lead-210	1.55E-06	2.00E-13	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.53E-06	0.00E+00	1.18E-02	3.40E-03	3.97E-02	5.97E-02
	Polonium-210	4.64E-06	1.75E-15	2.35E-03	7.07E-12	7.29E-06	2.16E-07	7.90E-05	1.08E-08	0.00E+00	7.47E-02	7.90E-03	2.23E-01	3.08E-01
	Total by Pathway	3.98E-05	1.58E-12	5.11E-03	2.70E-07	2.73E-05	7.47E-02	1.28E-04	3.97E-04	0.00E+00	8.93E-02	1.67E-02	2.79E-01	0.47
								Far-Future						
	Uranium-238	0.00E+00	0.00E+00	6.97E-06	5.29E-11	7.43E-08	8.24E-04	6.35E-08	6.78E-07	0.00E+00	1.94E-05	3.31E-04	5.23E-04	1.70E-03
	Uranium-234	0.00E+00	0.00E+00	7.59E-06	9.22E-12	8.09E-08	4.15E-06	6.92E-08	2.46E-09	0.00E+00	2.11E-05	3.60E-04	5.69E-04	9.62E-04
	Thorium-230	0.00E+00	0.00E+00	7.25E-04	4.92E-10	2.27E-06	3.48E-05	7.89E-07	1.95E-08	0.00E+00	9.71E-04	1.95E-04	4.54E-03	6.47E-03
	Radium-226	0.00E+00	0.00E+00	5.90E-04	2.63E-07	2.71E-06	7.03E-02	3.27E-06	3.92E-04	0.00E+00	1.57E-03	4.48E-03	1.03E-02	8.77E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	1.18E-02	3.28E-03	3.96E-02	5.95E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	7.46E-02	7.71E-03	2.23E-01	3.07E-01
	Total by Pathway	0.00E+00	0.00E+00	5.00E-03	2.65E-07	2.73E-05	7.46E-02	1.28E-04	3.95E-04	0.00E+00	8.90E-02	1.64E-02	2.78E-01	0.46
								Base Case						
	Uranium-238	0.00E+00	0.00E+00	4.22E-06	1.14E-11	3.02E-06	1.10E-03	2.25E-06	7.69E-07	0.00E+00	9.99E-06	4.05E-04	5.67E-04	2.09E-03
	Uranium-234	0.00E+00	0.00E+00	4.58E-06	1.99E-12	3.27E-06	5.56E-06	2.44E-06	2.79E-09	0.00E+00	1.08E-05	4.39E-04	6.15E-04	1.08E-03
	Thorium-230	0.00E+00	0.00E+00	3.69E-04	1.22E-10	6.75E-05	3.95E-05	2.35E-05	2.53E-08	0.00E+00	4.19E-04	1.75E-04	2.77E-03	3.86E-03
	Radium-226	0.00E+00	0.00E+00	5.26E-04	6.50E-08	1.42E-04	9.40E-02	1.71E-04	5.10E-04	0.00E+00	1.19E-03	7.05E-03	1.67E-02	1.20E-01
	Lead-210	0.00E+00	0.00E+00	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	1.36E-02	7.86E-03	8.28E-02	1.15E-01
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	1.21E-01	2.60E-02	3.94E-01	5.56E-01
Subsistence Harvester	Total by Pathway	0.00E+00	0.00E+00	7.19E-03	6.54E-08	2.21E-03	9.97E-02	1.26E-02	5.14E-04	0.00E+00	1.36E-01	4.20E-02	4.98E-01	0.80
One-Year-Old (Lloyd Lake)								RFD Case						
,	Uranium-238	4.07E-06	1.46E-14	5.27E-06	1.42E-11	3.05E-06	1.11E-03	2.43E-06	8.29E-07	0.00E+00	1.24E-05	4.09E-04	5.69E-04	2.12E-03
	Uranium-234	4.76E-06	3.56E-14	5.71E-06	2.48E-12	3.30E-06	5.60E-06	2.63E-06	3.00E-09	0.00E+00	1.34E-05	4.43E-04	6.16E-04	1.09E-03
	Thorium-230	1.51E-05	8.63E-14	4.11E-04	1.36E-10	6.76E-05	3.96E-05	2.56E-05	2.76E-08	0.00E+00	4.64E-04	1.92E-04	2.94E-03	4.16E-03
	Radium-226	4.76E-06	1.65E-12	5.36E-04	6.63E-08	1.42E-04	9.41E-02	1.72E-04	5.14E-04	0.00E+00	1.21E-03	7.11E-03	1.67E-02	1.21E-01
	Lead-210	1.60E-06	2.60E-13	1.81E-03	2.38E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	1.37E-02	8.06E-03	8.29E-02	1.16E-01
	Polonium-210	4.76E-06	2.27E-15	4.50E-03	1.75E-12	8.15E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	1.21E-01	2.65E-02	3.95E-01	5.57E-01
	Total by Pathway	3.51E-05	2.05E-12	7.26E-03	6.67E-08	2.22E-03	9.99E-02	1.26E-02	5.18E-04	0.00E+00	1.37E-01	4.27E-02	4.99E-01	0.80

Table C18: Estimated Total Radiation Doses to Human Receptors – Reasonably Foreseeable Development Case

			Can	np Worker (P	atterson Lake	e North Arm	- West Basin)	Incremental	Dose by Pat	hway RFD Ca	se (mSv/yr)			
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
								Far-Future						
	Uranium-238	0.00E+00	0.00E+00	4.84E-06	1.31E-11	3.02E-06	1.10E-03	2.58E-06	8.81E-07	0.00E+00	1.14E-05	4.06E-04	5.68E-04	2.10E-03
	Uranium-234	0.00E+00	0.00E+00	5.25E-06	2.28E-12	3.27E-06	5.56E-06	2.80E-06	3.20E-09	0.00E+00	1.24E-05	4.40E-04	6.15E-04	1.08E-03
	Thorium-230	0.00E+00	0.00E+00	3.69E-04	1.22E-10	6.76E-05	3.95E-05	2.35E-05	2.54E-08	0.00E+00	4.19E-04	1.75E-04	2.77E-03	3.87E-03
	Radium-226	0.00E+00	0.00E+00	5.27E-04	6.52E-08	1.42E-04	9.41E-02	1.71E-04	5.11E-04	0.00E+00	1.19E-03	7.06E-03	1.67E-02	1.20E-01
	Lead-210	0.00E+00	0.00E+00	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	1.36E-02	7.86E-03	8.28E-02	1.15E-01
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	1.21E-01	2.60E-02	3.94E-01	5.56E-01
	Total by Pathway	0.00E+00	0.00E+00	7.20E-03	6.56E-08	2.22E-03	9.98E-02	1.26E-02	5.15E-04	0.00E+00	1.36E-01	4.20E-02	4.98E-01	0.80
		•						Base Case						
	Uranium-238	0.00E+00	0.00E+00	6.08E-06	4.61E-11	7.42E-08	8.23E-04	5.54E-08	5.91E-07	0.00E+00	5.36E-06	2.59E-04	3.62E-04	1.46E-03
	Uranium-234	0.00E+00	0.00E+00	6.62E-06	8.04E-12	8.08E-08	4.15E-06	6.04E-08	2.14E-09	0.00E+00	5.84E-06	2.82E-04	3.95E-04	6.94E-04
	Thorium-230	0.00E+00	0.00E+00	7.25E-04	4.91E-10	2.27E-06	3.47E-05	7.89E-07	1.95E-08	0.00E+00	3.07E-04	1.52E-04	2.32E-03	3.54E-03
	Radium-226	0.00E+00	0.00E+00	5.89E-04	2.63E-07	2.71E-06	7.02E-02	3.26E-06	3.91E-04	0.00E+00	4.98E-04	3.51E-03	8.84E-03	8.40E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	3.74E-03	2.57E-03	2.70E-02	3.82E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	2.36E-02	6.05E-03	1.23E-01	1.56E-01
	Total by Pathway	0.00E+00	0.00E+00	5.00E-03	2.64E-07	2.73E-05	7.45E-02	1.28E-04	3.94E-04	0.00E+00	2.82E-02	1.28E-02	1.62E-01	0.28
								RFD Case						
	Uranium-238	9.39E-05	2.58E-13	3.39E-05	2.57E-10	8.90E-08	9.87E-04	1.32E-07	1.41E-06	0.00E+00	2.90E-05	3.13E-04	3.90E-04	1.85E-03
	Uranium-234	1.13E-04	6.30E-13	3.70E-05	4.50E-11	9.69E-08	4.98E-06	1.44E-07	5.12E-09	0.00E+00	3.17E-05	3.40E-04	4.24E-04	9.52E-04
Seasonal Resident	Thorium-230	4.54E-04	1.53E-12	1.82E-03	1.24E-09	2.35E-06	3.59E-05	1.57E-06	3.87E-08	0.00E+00	7.37E-04	4.09E-04	5.40E-03	8.87E-03
(Patterson Lake South Arm)	Radium-226	1.13E-04	2.93E-11	7.73E-04	3.45E-07	2.82E-06	7.29E-02	3.62E-06	4.33E-04	0.00E+00	6.39E-04	4.11E-03	9.15E-03	8.81E-02
,	Lead-210	3.56E-05	4.61E-12	1.55E-03	1.36E-09	1.51E-05	3.50E-03	4.76E-05	2.70E-06	0.00E+00	4.35E-03	3.75E-03	3.00E-02	4.32E-02
	Polonium-210	1.07E-04	4.02E-14	2.72E-03	8.18E-12	7.57E-06	2.24E-07	8.43E-05	1.16E-08	0.00E+00	2.72E-02	8.03E-03	1.31E-01	1.69E-01
	Total by Pathway	9.17E-04	3.63E-11	6.94E-03	3.48E-07	2.80E-05	7.74E-02	1.37E-04	4.38E-04	0.00E+00	3.30E-02	1.69E-02	1.77E-01	0.31
								Far-Future						
	Uranium-238	0.00E+00	0.00E+00	2.19E-05	1.66E-10	7.58E-08	8.41E-04	2.00E-07	2.13E-06	0.00E+00	1.93E-05	2.65E-04	3.71E-04	1.52E-03
	Uranium-234	0.00E+00	0.00E+00	2.39E-05	2.91E-11	8.25E-08	4.24E-06	2.18E-07	7.75E-09	0.00E+00	2.11E-05	2.88E-04	4.04E-04	7.42E-04
	Thorium-230	0.00E+00	0.00E+00	7.34E-04	4.97E-10	2.31E-06	3.54E-05	7.99E-07	1.97E-08	0.00E+00	3.11E-04	1.55E-04	2.36E-03	3.60E-03
	Radium-226	0.00E+00	0.00E+00	6.07E-04	2.71E-07	2.78E-06	7.20E-02	3.37E-06	4.03E-04	0.00E+00	5.14E-04	3.60E-03	8.92E-03	8.61E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.53E-06	0.00E+00	3.75E-03	2.57E-03	2.71E-02	3.83E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.06E-12	7.27E-06	2.16E-07	7.90E-05	1.08E-08	0.00E+00	2.37E-02	6.05E-03	1.23E-01	1.56E-01
	Total by Pathway	0.00E+00	0.00E+00	5.06E-03	2.73E-07	2.74E-05	7.64E-02	1.28E-04	4.08E-04	0.00E+00	2.83E-02	1.29E-02	1.63E-01	0.29

Table C18: Estimated Total Radiation Doses to Human Receptors – Reasonably Foreseeable Development Case

			Can	np Worker (P	atterson Lake	e North Arm	- West Basin)	Incremental	Dose by Patl	nway RFD Ca	se (mSv/yr)			
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
								Base Case				-		
	Uranium-238	0.00E+00	0.00E+00	4.22E-06	1.14E-11	3.02E-06	1.10E-03	2.25E-06	7.69E-07	0.00E+00	3.16E-06	3.52E-04	4.97E-04	1.96E-03
	Uranium-234	0.00E+00	0.00E+00	4.58E-06	1.99E-12	3.27E-06	5.56E-06	2.44E-06	2.79E-09	0.00E+00	3.43E-06	3.82E-04	5.38E-04	9.40E-04
	Thorium-230	0.00E+00	0.00E+00	3.69E-04	1.22E-10	6.75E-05	3.95E-05	2.35E-05	2.53E-08	0.00E+00	1.33E-04	1.52E-04	1.68E-03	2.46E-03
	Radium-226	0.00E+00	0.00E+00	5.26E-04	6.50E-08	1.42E-04	9.40E-02	1.71E-04	5.10E-04	0.00E+00	3.78E-04	6.13E-03	1.65E-02	1.18E-01
	Lead-210	0.00E+00	0.00E+00	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	4.32E-03	6.84E-03	6.72E-02	8.95E-02
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	3.83E-02	2.26E-02	2.73E-01	3.49E-01
	Total by Pathway	0.00E+00	0.00E+00	7.19E-03	6.54E-08	2.21E-03	9.97E-02	1.26E-02	5.14E-04	0.00E+00	4.32E-02	3.65E-02	3.60E-01	0.56
			T			ı	T	RFD Case						
	Uranium-238	9.37E-05	3.35E-13	2.36E-05	6.36E-11	3.62E-06	1.32E-03	5.37E-06	1.83E-06	0.00E+00	1.71E-05	4.25E-04	5.09E-04	2.40E-03
6 10 11 10	Uranium-234	1.10E-04	8.20E-13	2.56E-05	1.11E-11	3.92E-06	6.66E-06	5.83E-06	6.65E-09	0.00E+00	1.86E-05	4.61E-04	5.52E-04	1.18E-03
Seasonal Resident One- Year-Old (Patterson Lake	Thorium-230	3.49E-04	1.99E-12	9.28E-04	3.06E-10	6.98E-05	4.09E-05	4.66E-05	5.03E-08	0.00E+00	3.18E-04	2.94E-04	3.18E-03	5.23E-03
South Arm)	Radium-226	1.10E-04	3.80E-11	6.90E-04	8.54E-08	1.47E-04	9.76E-02	1.89E-04	5.65E-04	0.00E+00	4.84E-04	6.93E-03	1.66E-02	1.23E-01
	Lead-210	3.69E-05	5.98E-12	2.10E-03	2.77E-10	1.20E-03	4.68E-03	3.79E-03	3.50E-06	0.00E+00	5.02E-03	9.04E-03	7.09E-02	9.68E-02
	Polonium-210	1.10E-04	5.23E-14	5.20E-03	2.03E-12	8.46E-04	2.93E-07	9.43E-03	1.50E-08	0.00E+00	4.42E-02	2.81E-02	2.82E-01	3.70E-01
	Total by Pathway	8.08E-04	4.72E-11	8.97E-03	8.60E-08	2.27E-03	1.04E-01	1.35E-02	5.71E-04	0.00E+00	5.00E-02	4.52E-02	3.74E-01	0.60
			ı			ı	ı	Far-Future				1		
	Uranium-238	0.00E+00	0.00E+00	1.52E-05	4.11E-11	3.08E-06	1.12E-03	8.12E-06	2.77E-06	0.00E+00	1.14E-05	3.60E-04	5.01E-04	2.03E-03
	Uranium-234	0.00E+00	0.00E+00	1.65E-05	7.19E-12	3.34E-06	5.67E-06	8.83E-06	1.01E-08	0.00E+00	1.24E-05	3.90E-04	5.43E-04	9.79E-04
	Thorium-230	0.00E+00	0.00E+00	3.73E-04	1.23E-10	6.88E-05	4.03E-05	2.38E-05	2.57E-08	0.00E+00	1.34E-04	1.55E-04	1.70E-03	2.50E-03
	Radium-226	0.00E+00	0.00E+00	5.43E-04	6.71E-08	1.45E-04	9.64E-02	1.76E-04	5.26E-04	0.00E+00	3.90E-04	6.29E-03	1.65E-02	1.21E-01
	Lead-210	0.00E+00	0.00E+00	1.81E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	4.32E-03	6.84E-03	6.73E-02	8.96E-02
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	3.84E-02	2.26E-02	2.74E-01	3.49E-01
	Total by Pathway	0.00E+00	0.00E+00	7.24E-03	6.75E-08	2.22E-03	1.02E-01	1.26E-02	5.32E-04	0.00E+00	4.32E-02	3.67E-02	3.60E-01	0.56
						ı		Base Case				1		
	Uranium-238	0.00E+00	0.00E+00	6.08E-06	4.61E-11	7.42E-08	8.23E-04	5.54E-08	5.91E-07	0.00E+00	5.36E-06	2.59E-04	3.62E-04	1.46E-03
	Uranium-234	0.00E+00	0.00E+00	6.62E-06	8.04E-12	8.08E-08	4.15E-06	6.04E-08	2.14E-09	0.00E+00	5.84E-06	2.82E-04	3.95E-04	6.94E-04
Seasonal Resident (Lloyd	Thorium-230	0.00E+00	0.00E+00	7.25E-04	4.91E-10	2.27E-06	3.47E-05	7.89E-07	1.95E-08	0.00E+00	3.07E-04	1.52E-04	2.32E-03	3.54E-03
Lake)	Radium-226	0.00E+00	0.00E+00	5.89E-04	2.63E-07	2.71E-06	7.02E-02	3.26E-06	3.91E-04	0.00E+00	4.98E-04	3.51E-03	8.84E-03	8.40E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	3.74E-03	2.57E-03	2.70E-02	3.82E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	2.36E-02	6.05E-03	1.23E-01	1.56E-01
	Total by Pathway	0.00E+00	0.00E+00	5.00E-03	2.64E-07	2.73E-05	7.45E-02	1.28E-04	3.94E-04	0.00E+00	2.82E-02	1.28E-02	1.62E-01	0.28

Table C18: Estimated Total Radiation Doses to Human Receptors – Reasonably Foreseeable Development Case

			Can	np Worker (P	atterson Lak	North Arm	- West Basin)	Incrementa	Dose by Pat	hway RFD Ca	se (mSv/yr)			
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
								RFD Case						
	Uranium-238	2.45E-06	6.72E-15	6.98E-06	5.30E-11	7.46E-08	8.28E-04	5.80E-08	6.19E-07	0.00E+00	8.27E-06	2.61E-04	3.63E-04	1.47E-03
	Uranium-234	2.95E-06	1.64E-14	7.60E-06	9.23E-12	8.13E-08	4.17E-06	6.32E-08	2.24E-09	0.00E+00	9.01E-06	2.84E-04	3.95E-04	7.03E-04
	Thorium-230	1.18E-05	3.98E-14	7.75E-04	5.25E-10	2.27E-06	3.48E-05	8.31E-07	2.05E-08	0.00E+00	3.41E-04	1.59E-04	2.40E-03	3.72E-03
	Radium-226	2.95E-06	7.62E-13	5.95E-04	2.66E-07	2.71E-06	7.03E-02	3.28E-06	3.93E-04	0.00E+00	5.08E-04	3.52E-03	8.84E-03	8.41E-02
	Lead-210	9.29E-07	1.20E-13	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.53E-06	0.00E+00	4.51E-03	2.60E-03	2.70E-02	3.90E-02
	Polonium-210	2.79E-06	1.05E-15	2.35E-03	7.06E-12	7.28E-06	2.16E-07	7.90E-05	1.08E-08	0.00E+00	2.42E-02	6.10E-03	1.23E-01	1.56E-01
	Total by Pathway	2.39E-05	9.47E-13	5.06E-03	2.68E-07	2.73E-05	7.46E-02	1.28E-04	3.96E-04	0.00E+00	2.96E-02	1.29E-02	1.62E-01	0.29
								Far-Future	9					
	Uranium-238	0.00E+00	0.00E+00	6.61E-06	5.02E-11	7.43E-08	8.24E-04	6.03E-08	6.43E-07	0.00E+00	7.63E-06	2.59E-04	3.63E-04	1.46E-03
	Uranium-234	0.00E+00	0.00E+00	7.20E-06	8.74E-12	8.09E-08	4.15E-06	6.57E-08	2.33E-09	0.00E+00	8.31E-06	2.82E-04	3.95E-04	6.97E-04
	Thorium-230	0.00E+00	0.00E+00	7.25E-04	4.91E-10	2.27E-06	3.48E-05	7.89E-07	1.95E-08	0.00E+00	3.08E-04	1.53E-04	2.32E-03	3.54E-03
	Radium-226	0.00E+00	0.00E+00	5.89E-04	2.63E-07	2.71E-06	7.03E-02	3.27E-06	3.91E-04	0.00E+00	5.00E-04	3.51E-03	8.84E-03	8.41E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	4.50E-03	2.57E-03	2.70E-02	3.89E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	2.41E-02	6.05E-03	1.23E-01	1.56E-01
	Total by Pathway	0.00E+00	0.00E+00	5.00E-03	2.65E-07	2.73E-05	7.46E-02	1.28E-04	3.95E-04	0.00E+00	2.95E-02	1.28E-02	1.62E-01	0.28
								Base Case						
	Uranium-238	0.00E+00	0.00E+00	4.22E-06	1.14E-11	3.02E-06	1.10E-03	2.25E-06	7.69E-07	0.00E+00	3.16E-06	3.52E-04	4.97E-04	1.96E-03
	Uranium-234	0.00E+00	0.00E+00	4.58E-06	1.99E-12	3.27E-06	5.56E-06	2.44E-06	2.79E-09	0.00E+00	3.43E-06	3.82E-04	5.38E-04	9.40E-04
	Thorium-230	0.00E+00	0.00E+00	3.69E-04	1.22E-10	6.75E-05	3.95E-05	2.35E-05	2.53E-08	0.00E+00	1.33E-04	1.52E-04	1.68E-03	2.46E-03
	Radium-226	0.00E+00	0.00E+00	5.26E-04	6.50E-08	1.42E-04	9.40E-02	1.71E-04	5.10E-04	0.00E+00	3.78E-04	6.13E-03	1.65E-02	1.18E-01
	Lead-210	0.00E+00	0.00E+00	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	4.32E-03	6.84E-03	6.72E-02	8.95E-02
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	3.83E-02	2.26E-02	2.73E-01	3.49E-01
Seasonal Resident One-	Total by Pathway	0.00E+00	0.00E+00	7.19E-03	6.54E-08	2.21E-03	9.97E-02	1.26E-02	5.14E-04	0.00E+00	4.32E-02	3.65E-02	3.60E-01	0.56
Year-Old (Lloyd Lake)								RFD Case		,			,	
	Uranium-238	2.44E-06	8.73E-15	4.85E-06	1.31E-11	3.03E-06	1.11E-03	2.36E-06	8.05E-07	0.00E+00	3.61E-06	3.54E-04	4.97E-04	1.98E-03
	Uranium-234	2.86E-06	2.14E-14	5.26E-06	2.29E-12	3.29E-06	5.58E-06	2.56E-06	2.92E-09	0.00E+00	3.92E-06	3.84E-04	5.39E-04	9.46E-04
	Thorium-230	9.09E-06	5.18E-14	3.94E-04	1.30E-10	6.76E-05	3.95E-05	2.47E-05	2.67E-08	0.00E+00	1.41E-04	1.56E-04	1.72E-03	2.55E-03
	Radium-226	2.86E-06	9.91E-13	5.32E-04	6.58E-08	1.42E-04	9.41E-02	1.71E-04	5.12E-04	0.00E+00	3.82E-04	6.15E-03	1.65E-02	1.18E-01
	Lead-210	9.61E-07	1.56E-13	1.81E-03	2.38E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	4.32E-03	6.89E-03	6.72E-02	8.96E-02
	Polonium-210	2.86E-06	1.36E-15	4.49E-03	1.75E-12	8.14E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	3.84E-02	2.28E-02	2.74E-01	3.49E-01
	Total by Pathway	2.11E-05	1.23E-12	7.23E-03	6.62E-08	2.22E-03	9.98E-02	1.26E-02	5.16E-04	0.00E+00	4.32E-02	3.67E-02	3.60E-01	0.56

Table C18: Estimated Total Radiation Doses to Human Receptors – Reasonably Foreseeable Development Case

			Can	np Worker (P	atterson Lake	e North Arm	- West Basin)	Incremental	Dose by Pat	hway RFD Ca	se (mSv/yr)			
	Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
								Far-Future				1		
	Uranium-238	0.00E+00	0.00E+00	4.59E-06	1.24E-11	3.02E-06	1.10E-03	2.45E-06	8.36E-07	0.00E+00	3.44E-06	3.53E-04	4.97E-04	1.97E-03
	Uranium-234	0.00E+00	0.00E+00	4.98E-06	2.17E-12	3.27E-06	5.56E-06	2.66E-06	3.03E-09	0.00E+00	3.73E-06	3.82E-04	5.39E-04	9.41E-04
	Thorium-230	0.00E+00	0.00E+00	3.69E-04	1.22E-10	6.76E-05	3.95E-05	2.35E-05	2.54E-08	0.00E+00	1.33E-04	1.52E-04	1.68E-03	2.46E-03
	Radium-226	0.00E+00	0.00E+00	5.27E-04	6.51E-08	1.42E-04	9.40E-02	1.71E-04	5.11E-04	0.00E+00	3.78E-04	6.13E-03	1.65E-02	1.18E-01
	Lead-210	0.00E+00	0.00E+00	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	4.32E-03	6.84E-03	6.72E-02	8.95E-02
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	3.83E-02	2.26E-02	2.73E-01	3.49E-01
	Total by Pathway	0.00E+00	0.00E+00	7.20E-03	6.55E-08	2.22E-03	9.98E-02	1.26E-02	5.15E-04	0.00E+00	4.32E-02	3.65E-02	3.60E-01	0.56
		•						Base Case Ad	lult					
	Uranium-238	0.00E+00	0.00E+00	6.08E-06	4.61E-11	7.42E-08	8.23E-04	5.54E-08	5.91E-07	0.00E+00	1.69E-05	3.30E-04	5.22E-04	1.70E-03
	Uranium-234	0.00E+00	0.00E+00	6.62E-06	8.04E-12	8.08E-08	4.15E-06	6.04E-08	2.14E-09	0.00E+00	1.84E-05	3.60E-04	5.68E-04	9.57E-04
	Thorium-230	0.00E+00	0.00E+00	7.25E-04	4.91E-10	2.27E-06	3.47E-05	7.89E-07	1.95E-08	0.00E+00	9.70E-04	1.94E-04	4.54E-03	6.47E-03
	Radium-226	0.00E+00	0.00E+00	5.89E-04	2.63E-07	2.71E-06	7.02E-02	3.26E-06	3.91E-04	0.00E+00	1.57E-03	4.47E-03	1.03E-02	8.76E-02
	Lead-210	0.00E+00	0.00E+00	1.33E-03	1.17E-09	1.49E-05	3.46E-03	4.46E-05	2.52E-06	0.00E+00	1.18E-02	3.28E-03	3.96E-02	5.95E-02
	Polonium-210	0.00E+00	0.00E+00	2.35E-03	7.05E-12	7.27E-06	2.16E-07	7.89E-05	1.08E-08	0.00E+00	7.46E-02	7.71E-03	2.23E-01	3.07E-01
	Total by Pathway	0.00E+00	0.00E+00	5.00E-03	2.64E-07	2.73E-05	7.45E-02	1.28E-04	3.94E-04	0.00E+00	8.90E-02	1.63E-02	2.78E-01	0.46
		•					Base	Case One-Yo	ear-Old					
	Uranium-238	0.00E+00	0.00E+00	4.22E-06	1.14E-11	3.02E-06	1.10E-03	2.25E-06	7.69E-07	0.00E+00	9.99E-06	4.05E-04	5.67E-04	2.09E-03
	Uranium-234	0.00E+00	0.00E+00	4.58E-06	1.99E-12	3.27E-06	5.56E-06	2.44E-06	2.79E-09	0.00E+00	1.08E-05	4.39E-04	6.15E-04	1.08E-03
Permanent Resident (Patterson Lake North	Thorium-230	0.00E+00	0.00E+00	3.69E-04	1.22E-10	6.75E-05	3.95E-05	2.35E-05	2.53E-08	0.00E+00	4.19E-04	1.75E-04	2.77E-03	3.86E-03
Arm - West Basin)	Radium-226	0.00E+00	0.00E+00	5.26E-04	6.50E-08	1.42E-04	9.40E-02	1.71E-04	5.10E-04	0.00E+00	1.19E-03	7.05E-03	1.67E-02	1.20E-01
	Lead-210	0.00E+00	0.00E+00	1.80E-03	2.37E-10	1.19E-03	4.63E-03	3.55E-03	3.28E-06	0.00E+00	1.36E-02	7.86E-03	8.28E-02	1.15E-01
	Polonium-210	0.00E+00	0.00E+00	4.49E-03	1.75E-12	8.13E-04	2.82E-07	8.83E-03	1.41E-08	0.00E+00	1.21E-01	2.60E-02	3.94E-01	5.56E-01
	Total by Pathway	0.00E+00	0.00E+00	7.19E-03	6.54E-08	2.21E-03	9.97E-02	1.26E-02	5.14E-04	0.00E+00	1.36E-01	4.20E-02	4.98E-01	0.80
							Ī	ar-Future Ac	dult					
	Uranium-238	0.00E+00	0.00E+00	1.97E-04	1.49E-09	1.01E-07	1.12E-03	1.79E-06	1.91E-05	0.00E+00	5.48E-04	4.50E-04	7.62E-04	3.10E-03
	Uranium-234	0.00E+00	0.00E+00	2.15E-04	2.61E-10	1.10E-07	5.66E-06	1.96E-06	6.97E-08	0.00E+00	5.99E-04	4.90E-04	8.30E-04	2.14E-03
	Thorium-230	0.00E+00	0.00E+00	7.75E-04	5.25E-10	3.01E-06	4.62E-05	8.43E-07	2.08E-08	0.00E+00	1.04E-03	2.58E-04	5.08E-03	7.20E-03
	Radium-226	0.00E+00	0.00E+00	7.59E-04	3.39E-07	3.93E-06	1.02E-01	4.20E-06	5.04E-04	0.00E+00	2.03E-03	6.48E-03	1.22E-02	1.24E-01
	Lead-210	0.00E+00	0.00E+00	1.39E-03	1.22E-09	1.49E-05	3.46E-03	4.67E-05	2.64E-06	0.00E+00	1.24E-02	3.28E-03	4.05E-02	6.10E-02
	Polonium-210	0.00E+00	0.00E+00	2.46E-03	7.38E-12	7.28E-06	2.16E-07	8.26E-05	1.13E-08	0.00E+00	7.80E-02	7.72E-03	2.30E-01	3.18E-01
	Total by Pathway	0.00E+00	0.00E+00	5.79E-03	3.42E-07	2.93E-05	1.06E-01	1.38E-04	5.26E-04	0.00E+00	9.46E-02	1.87E-02	2.89E-01	0.52

Table C18: Estimated Total Radiation Doses to Human Receptors – Reasonably Foreseeable Development Case

		Can	ıp Worker (P	atterson Lake	North Arm	- West Basin)	Incremental	Dose by Pat	hway RFD Ca	se (mSv/yr)			
Radionuclide	Air (Internal)	Air (External)	Water (Internal)	Water (External)	Soil (Internal)	Soil (External)	Sediment (Internal)	Sediment (External)	Aquatic Plants	Aquatic Animals	Terrestrial Plants	Terrestrial animals	Total by Radionuclide
						Far-I	uture One-Y	ear-Old					'
Uranium-238	0.00E+00	0.00E+00	1.37E-04	3.69E-10	4.11E-06	1.50E-03	7.30E-05	2.49E-05	0.00E+00	3.23E-04	5.52E-04	6.76E-04	3.29E-03
Uranium-234	0.00E+00	0.00E+00	1.49E-04	6.47E-11	4.46E-06	7.57E-06	7.94E-05	9.06E-08	0.00E+00	3.52E-04	5.98E-04	7.32E-04	1.92E-03
Thorium-230	0.00E+00	0.00E+00	3.94E-04	1.30E-10	8.98E-05	5.25E-05	2.51E-05	2.71E-08	0.00E+00	4.48E-04	2.32E-04	3.03E-03	4.27E-03
Radium-226	0.00E+00	0.00E+00	6.78E-04	8.39E-08	2.05E-04	1.36E-01	2.20E-04	6.57E-04	0.00E+00	1.54E-03	1.02E-02	1.75E-02	1.67E-01
Lead-210	0.00E+00	0.00E+00	1.89E-03	2.48E-10	1.19E-03	4.63E-03	3.71E-03	3.43E-06	0.00E+00	1.43E-02	7.86E-03	8.39E-02	1.17E-01
Polonium-210	0.00E+00	0.00E+00	4.69E-03	1.83E-12	8.14E-04	2.82E-07	9.23E-03	1.47E-08	0.00E+00	1.27E-01	2.60E-02	4.04E-01	5.71E-01
Total by Pathway	0.00E+00	0.00E+00	7.94E-03	8.47E-08	2.30E-03	1.42E-01	1.33E-02	6.86E-04	0.00E+00	1.44E-01	4.55E-02	5.09E-01	0.87

Table C.19: Maximum Radiation Dose to Ecological Receptors for the Application Case and Upper Bound Scenario - Project Lifespan

	Benthic Invertebrates Total Dose Operations (mGy/d)												
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose						
		Α	pplication Case										
Reference (Broach Lake)	4.73E-09	6.18E-09	1.80E-08	8.07E-05	3.79E-05	1.72E-09	1.19E-04						
Patterson Lake North Arm - West Basin	4.26E-08	5.58E-08	1.00E-07	1.30E-04	5.19E-05	2.35E-09	1.82E-04						
Patterson Lake South Arm	1.47E-08	1.93E-08	6.47E-08	9.76E-05	3.89E-05	1.76E-09	1.37E-04						
Beet Lake	8.41E-09	1.10E-08	4.26E-08	8.79E-05	3.81E-05	1.72E-09	1.26E-04						
Lloyd Lake	5.06E-09	6.61E-09	2.06E-08	8.14E-05	3.79E-05	1.72E-09	1.19E-04						
		Uppe	er Bound Scenari	0									
Reference (Broach Lake)	4.73E-09	6.18E-09	1.80E-08	8.07E-05	3.79E-05	1.72E-09	1.19E-04						
Patterson Lake North Arm - West Basin	1.10E-07	1.44E-07	1.01E-07	1.32E-04	1.04E-04	4.71E-09	2.36E-04						
Patterson Lake South Arm	3.26E-08	4.28E-08	6.54E-08	9.84E-05	4.23E-05	1.92E-09	1.41E-04						
Beet Lake	1.50E-08	1.96E-08	4.29E-08	8.82E-05	3.85E-05	1.75E-09	1.27E-04						
Lloyd Lake	5.66E-09	7.39E-09	2.06E-08	8.15E-05	3.80E-05	1.72E-09	1.19E-04						

	Zooplankton Total Dose Operations (mGy/d)												
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose						
		A	pplication Case										
Reference (Broach Lake)	2.05E-05	2.39E-05	2.94E-03	4.57E-03	2.80E-05	6.05E-02	6.81E-02						
Patterson Lake North Arm - West Basin	2.23E-04	2.62E-04	1.67E-02	7.95E-03	4.28E-05	9.18E-02	1.17E-01						
Patterson Lake South Arm	7.51E-05	8.78E-05	1.11E-02	5.76E-03	2.88E-05	6.23E-02	7.93E-02						
Beet Lake	4.06E-05	4.74E-05	7.21E-03	5.08E-03	2.81E-05	6.07E-02	7.31E-02						
Lloyd Lake	2.23E-05	2.61E-05	3.38E-03	4.62E-03	2.80E-05	6.05E-02	6.86E-02						
		Uppe	er Bound Scenari	0									
Reference (Broach Lake)	2.05E-05	2.39E-05	2.94E-03	4.57E-03	2.80E-05	6.05E-02	6.81E-02						
Patterson Lake North Arm - West Basin	6.09E-04	7.13E-04	1.69E-02	8.10E-03	1.07E-04	2.34E-01	2.61E-01						
Patterson Lake South Arm	1.70E-04	1.99E-04	1.12E-02	5.82E-03	3.23E-05	7.00E-02	8.74E-02						
Beet Lake	7.54E-05	8.82E-05	7.26E-03	5.10E-03	2.85E-05	6.17E-02	7.43E-02						
Lloyd Lake	2.55E-05	2.98E-05	3.39E-03	4.62E-03	2.80E-05	6.06E-02	6.87E-02						

	Northern Pike Total Dose Operations (mGy/d)												
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose						
		Α	pplication Case										
Reference (Broach Lake)	4.13E-06	4.70E-06	3.54E-05	2.22E-04	1.79E-06	5.64E-04	8.33E-04						
Patterson Lake North Arm - West Basin	4.50E-05	5.14E-05	2.02E-04	3.87E-04	2.74E-06	8.56E-04	1.54E-03						
Patterson Lake South Arm	1.51E-05	1.72E-05	1.33E-04	2.80E-04	1.85E-06	5.81E-04	1.03E-03						
Beet Lake	8.17E-06	9.31E-06	8.67E-05	2.47E-04	1.80E-06	5.67E-04	9.20E-04						
Lloyd Lake	4.50E-06	5.12E-06	4.07E-05	2.25E-04	1.79E-06	5.64E-04	8.41E-04						
		Upp	er Bound Scenari	0									
Reference (Broach Lake)	4.13E-06	4.70E-06	3.54E-05	2.22E-04	1.79E-06	5.64E-04	8.33E-04						
Patterson Lake North Arm - West Basin	1.23E-04	1.40E-04	2.03E-04	3.94E-04	6.88E-06	2.18E-03	3.05E-03						
Patterson Lake South Arm	3.42E-05	3.90E-05	1.34E-04	2.83E-04	2.07E-06	6.53E-04	1.15E-03						
Beet Lake	1.52E-05	1.73E-05	8.74E-05	2.48E-04	1.83E-06	5.76E-04	9.46E-04						
Lloyd Lake	5.14E-06	5.86E-06	4.08E-05	2.25E-04	1.80E-06	5.65E-04	8.43E-04						

		Whitefish Tota	al Dose Operation	ns (mGy/d)			
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose
		A	pplication Case				
Reference (Broach Lake)	7.69E-07	8.97E-07	2.65E-05	1.86E-04	8.29E-07	4.31E-05	2.58E-04
Patterson Lake North Arm - West Basin	8.38E-06	9.82E-06	1.51E-04	3.21E-04	1.22E-06	6.55E-05	5.57E-04
Patterson Lake South Arm	2.81E-06	3.29E-06	9.95E-05	2.33E-04	8.53E-07	4.44E-05	3.84E-04
Beet Lake	1.52E-06	1.78E-06	6.49E-05	2.06E-04	8.33E-07	4.33E-05	3.18E-04
Lloyd Lake	8.37E-07	9.77E-07	3.05E-05	1.88E-04	8.30E-07	4.31E-05	2.64E-04
		Uppe	er Bound Scenari	o			
Reference (Broach Lake)	7.69E-07	8.97E-07	2.65E-05	1.86E-04	8.29E-07	4.31E-05	2.58E-04
Patterson Lake North Arm - West Basin	2.28E-05	2.67E-05	1.52E-04	3.27E-04	2.82E-06	1.67E-04	6.99E-04
Patterson Lake South Arm	6.37E-06	7.45E-06	1.00E-04	2.36E-04	9.46E-07	4.99E-05	4.01E-04
Beet Lake	2.83E-06	3.31E-06	6.54E-05	2.07E-04	8.45E-07	4.40E-05	3.23E-04
Lloyd Lake	9.58E-07	1.12E-06	3.05E-05	1.88E-04	8.30E-07	4.32E-05	2.64E-04

Table C.19: Maximum Radiation Dose to Ecological Receptors for the Application Case and Upper Bound Scenario - Project Lifespan

	Macrophytes Total Dose Operations (mGy/d)												
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose						
		A	pplication Case										
Reference (Broach Lake)	1.04E-05	1.18E-05	1.37E-03	2.99E-03	5.06E-05	1.10E-03	5.53E-03						
Patterson Lake North Arm - West Basin	1.13E-04	1.30E-04	7.79E-03	5.20E-03	7.73E-05	1.67E-03	1.50E-02						
Patterson Lake South Arm	3.81E-05	4.35E-05	5.14E-03	3.77E-03	5.22E-05	1.13E-03	1.02E-02						
Beet Lake	2.06E-05	2.35E-05	3.35E-03	3.32E-03	5.08E-05	1.11E-03	7.88E-03						
Lloyd Lake	1.13E-05	1.29E-05	1.57E-03	3.02E-03	5.06E-05	1.10E-03	5.77E-03						
		Uppe	er Bound Scenari	o									
Reference (Broach Lake)	1.04E-05	1.18E-05	1.37E-03	2.99E-03	5.06E-05	1.10E-03	5.53E-03						
Patterson Lake North Arm - West Basin	3.09E-04	3.53E-04	7.87E-03	5.30E-03	1.94E-04	4.26E-03	1.83E-02						
Patterson Lake South Arm	8.62E-05	9.84E-05	5.19E-03	3.81E-03	5.85E-05	1.27E-03	1.05E-02						
Beet Lake	3.83E-05	4.37E-05	3.38E-03	3.34E-03	5.17E-05	1.12E-03	7.97E-03						
Lloyd Lake	1.30E-05	1.48E-05	1.58E-03	3.02E-03	5.07E-05	1.10E-03	5.78E-03						

	Phytoplankton Total Dose Operations (mGy/d)												
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose						
		Α	pplication Case										
Reference (Broach Lake)	1.04E-05	1.18E-05	1.36E-03	3.00E-03	1.93E-05	1.11E-03	5.51E-03						
Patterson Lake North Arm - West Basin	1.13E-04	1.29E-04	7.75E-03	5.21E-03	2.96E-05	1.68E-03	1.49E-02						
Patterson Lake South Arm	3.81E-05	4.34E-05	5.12E-03	3.78E-03	1.99E-05	1.14E-03	1.01E-02						
Beet Lake	2.06E-05	2.34E-05	3.34E-03	3.33E-03	1.94E-05	1.11E-03	7.84E-03						
Lloyd Lake	1.13E-05	1.29E-05	1.57E-03	3.03E-03	1.94E-05	1.11E-03	5.75E-03						
		Upp	er Bound Scenari	0									
Reference (Broach Lake)	1.04E-05	1.18E-05	1.36E-03	3.00E-03	1.93E-05	1.11E-03	5.51E-03						
Patterson Lake North Arm - West Basin	3.09E-04	3.52E-04	7.83E-03	5.31E-03	7.42E-05	4.28E-03	1.82E-02						
Patterson Lake South Arm	8.61E-05	9.82E-05	5.17E-03	3.82E-03	2.24E-05	1.28E-03	1.05E-02						
Beet Lake	3.82E-05	4.36E-05	3.36E-03	3.35E-03	1.97E-05	1.13E-03	7.94E-03						
Lloyd Lake	1.30E-05	1.47E-05	1.57E-03	3.03E-03	1.94E-05	1.11E-03	5.75E-03						

	Terrestrial Invertebrate Total Dose Operations (mGy/d)										
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
	Application Case										
Reference (Broach Lake)	1.57E-05	1.78E-05	1.15E-04	5.73E-03	6.72E-06	2.28E-05	5.91E-03				
Patterson Lake North Arm - West Basin	6.88E-05	7.83E-05	1.82E-03	1.78E-02	2.16E-05	1.77E-03	2.16E-02				
Patterson Lake South Arm	2.43E-05	2.77E-05	3.93E-04	7.71E-03	9.14E-06	3.07E-04	8.47E-03				
Beet Lake	2.07E-05	2.36E-05	2.77E-04	6.88E-03	8.13E-06	1.88E-04	7.40E-03				
Lloyd Lake	1.58E-05	1.80E-05	1.20E-04	5.77E-03	6.76E-06	2.83E-05	5.96E-03				
		Uppe	er Bound Scenari	0							
Reference (Broach Lake)	1.57E-05	1.78E-05	1.15E-04	5.73E-03	6.72E-06	2.28E-05	5.91E-03				
Patterson Lake North Arm - West Basin	6.88E-05	7.83E-05	1.82E-03	1.78E-02	2.16E-05	1.77E-03	2.16E-02				
Patterson Lake South Arm	2.43E-05	2.77E-05	3.93E-04	7.71E-03	9.14E-06	3.07E-04	8.47E-03				
Beet Lake	2.07E-05	2.36E-05	2.77E-04	6.88E-03	8.13E-06	1.88E-04	7.40E-03				
Lloyd Lake	1.58E-05	1.80E-05	1.20E-04	5.77E-03	6.76E-06	2.83E-05	5.96E-03				

Browse Total Dose Operations (mGy/d)										
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose			
Application Case										
Patterson Lake North Arm - West Basin	2.53E-03	2.88E-03	1.80E-03	2.15E-02	2.10E-04	2.79E-03	3.18E-02			
Patterson Lake South Arm	8.97E-04	1.02E-03	3.80E-04	1.02E-02	1.73E-04	9.85E-04	1.36E-02			
		Uppe	er Bound Scenari	0						
Patterson Lake North Arm - West Basin	2.53E-03	2.88E-03	1.80E-03	2.15E-02	2.10E-04	2.79E-03	3.18E-02			
Patterson Lake South Arm	8.97E-04	1.02E-03	3.80E-04	1.02E-02	1.73E-04	9.85E-04	1.36E-02			

Table C.19: Maximum Radiation Dose to Ecological Receptors for the Application Case and Upper Bound Scenario - Project Lifespan

Labrador Tea Total Dose Operations (mGy/d)											
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
	Application Case										
Patterson Lake North Arm - West Basin	2.53E-03	2.88E-03	1.58E-02	9.27E-02	3.26E-04	1.61E-02	1.30E-01				
Patterson Lake South Arm	8.97E-04	1.02E-03	2.66E-03	2.18E-02	1.92E-04	3.16E-03	2.97E-02				
		Uppe	er Bound Scenari	0							
Patterson Lake North Arm - West Basin	2.53E-03	2.88E-03	1.58E-02	9.27E-02	3.26E-04	1.61E-02	1.30E-01				
Patterson Lake South Arm	8.97E-04	1.02E-03	2.66E-03	2.18E-02	1.92E-04	3.16E-03	2.97E-02				

Lichen Total Dose Operations (mGy/d)										
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose			
Application Case										
Patterson Lake North Arm - West Basin	1.21E-04	1.37E-04	8.96E-02	4.62E-01	1.40E-03	1.26E-01	6.79E-01			
Patterson Lake South Arm	1.21E-04	1.37E-04	1.50E-02	7.91E-02	9.26E-04	1.09E-01	2.05E-01			
		Uppe	er Bound Scenari	o						
Patterson Lake North Arm - West Basin	1.21E-04	1.37E-04	8.96E-02	4.62E-01	1.40E-03	1.26E-01	6.79E-01			
Patterson Lake South Arm	1.21E-04	1.37E-04	1.50E-02	7.91E-02	9.26E-04	1.09E-01	2.05E-01			

Blueberries Total Dose Operations (mGy/d)										
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose			
Application Case										
Patterson Lake North Arm - West Basin	1.31E-03	1.49E-03	6.96E-05	9.15E-03	8.42E-06	1.06E-03	1.31E-02			
Patterson Lake South Arm	4.53E-04	5.16E-04	4.55E-05	4.74E-03	3.88E-06	4.38E-04	6.20E-03			
		Uppe	er Bound Scenari	0						
Patterson Lake North Arm - West Basin	1.31E-03	1.49E-03	6.96E-05	9.15E-03	8.42E-06	1.06E-03	1.31E-02			
Patterson Lake South Arm	4.53E-04	5.16E-04	4.55E-05	4.74E-03	3.88E-06	4.38E-04	6.20E-03			

	Beaver Total Dose Operations (mGy/d)										
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
Application Case											
Reference (Broach Lake)	1.05E-05	1.19E-05	3.11E-06	6.36E-04	4.75E-07	2.15E-04	8.77E-04				
Patterson Lake North Arm - West Basin	4.63E-05	5.27E-05	2.94E-05	1.72E-03	1.13E-06	7.41E-04	2.59E-03				
Patterson Lake South Arm	1.64E-05	1.86E-05	1.08E-05	8.19E-04	5.66E-07	2.98E-04	1.16E-03				
Beet Lake	1.39E-05	1.58E-05	7.32E-06	7.41E-04	5.24E-07	2.62E-04	1.04E-03				
Lloyd Lake	1.06E-05	1.21E-05	3.42E-06	6.40E-04	4.77E-07	2.16E-04	8.83E-04				
		Uppe	er Bound Scenari	0							
Reference (Broach Lake)	1.05E-05	1.19E-05	3.11E-06	6.36E-04	4.75E-07	2.15E-04	8.77E-04				
Patterson Lake North Arm - West Basin	4.73E-05	5.38E-05	2.95E-05	1.72E-03	1.76E-06	8.55E-04	2.70E-03				
Patterson Lake South Arm	1.66E-05	1.89E-05	1.09E-05	8.19E-04	6.02E-07	3.05E-04	1.17E-03				
Beet Lake	1.40E-05	1.59E-05	7.36E-06	7.41E-04	5.28E-07	2.63E-04	1.04E-03				
Lloyd Lake	1.06E-05	1.21E-05	3.43E-06	6.40E-04	4.77E-07	2.16E-04	8.83E-04				

Black Bear Total Dose Operations (mGy/d)											
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
	Application Case										
Reference (Broach Lake)	7.81E-06	8.89E-06	1.88E-06	8.19E-04	8.32E-07	1.74E-04	1.01E-03				
Patterson Lake North Arm - West Basin	2.23E-05	2.54E-05	3.20E-06	1.26E-03	1.04E-06	3.23E-04	1.64E-03				
Beet Lake	9.24E-06	1.05E-05	2.69E-06	1.08E-03	1.23E-06	1.94E-04	1.30E-03				
Lloyd Lake	7.90E-06	8.99E-06	1.94E-06	8.23E-04	8.34E-07	1.76E-04	1.02E-03				
		Uppe	er Bound Scenari	0							
Reference (Broach Lake)	7.81E-06	8.89E-06	1.88E-06	8.19E-04	8.32E-07	1.74E-04	1.01E-03				
Patterson Lake North Arm - West Basin	2.26E-05	2.58E-05	3.21E-06	1.26E-03	1.06E-06	3.82E-04	1.70E-03				
Beet Lake	9.27E-06	1.06E-05	2.70E-06	1.08E-03	1.23E-06	1.95E-04	1.30E-03				
Lloyd Lake	7.91E-06	9.00E-06	1.94E-06	8.23E-04	8.34E-07	1.76E-04	1.02E-03				

Table C.19: Maximum Radiation Dose to Ecological Receptors for the Application Case and Upper Bound Scenario - Project Lifespan

Grey Wolf Total Dose Operations (mGy/d)											
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
	Application Case										
Reference (Broach Lake)	3.26E-07	3.77E-07	1.17E-06	7.63E-04	1.92E-06	3.26E-04	1.09E-03				
Patterson Lake North Arm - West Basin	6.21E-07	7.18E-07	1.56E-06	9.19E-04	2.02E-06	3.59E-04	1.28E-03				
Lloyd Lake	3.29E-07	3.81E-07	1.19E-06	7.65E-04	1.92E-06	3.26E-04	1.10E-03				
		Uppe	er Bound Scenari	io							
Reference (Broach Lake)	3.26E-07	3.77E-07	1.17E-06	7.63E-04	1.92E-06	3.26E-04	1.09E-03				
Patterson Lake North Arm - West Basin	6.39E-07	7.38E-07	1.56E-06	9.19E-04	2.02E-06	3.71E-04	1.29E-03				
Lloyd Lake	3.30E-07	3.82E-07	1.19E-06	7.65E-04	1.92E-06	3.26E-04	1.10E-03				

		Mink Total I	Dose Operations	(mGy/d)						
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose			
Application Case										
Reference (Broach Lake)	1.76E-07	2.00E-07	2.51E-06	6.44E-05	2.38E-07	8.73E-04	9.40E-04			
Patterson Lake North Arm - West Basin	1.15E-06	1.32E-06	1.23E-05	1.04E-04	3.23E-07	1.20E-03	1.32E-03			
Patterson Lake South Arm	3.99E-07	4.55E-07	7.97E-06	7.64E-05	2.44E-07	8.95E-04	9.80E-04			
Beet Lake	2.69E-07	3.07E-07	5.38E-06	6.97E-05	2.39E-07	8.76E-04	9.52E-04			
Lloyd Lake	1.82E-07	2.07E-07	2.80E-06	6.48E-05	2.38E-07	8.73E-04	9.41E-04			
		Uppe	er Bound Scenari	0						
Reference (Broach Lake)	1.76E-07	2.00E-07	2.51E-06	6.44E-05	2.38E-07	8.73E-04	9.40E-04			
Patterson Lake North Arm - West Basin	2.29E-06	2.62E-06	1.25E-05	1.06E-04	6.21E-07	2.39E-03	2.51E-03			
Patterson Lake South Arm	7.00E-07	7.99E-07	8.05E-06	7.68E-05	2.64E-07	9.75E-04	1.06E-03			
Beet Lake	3.80E-07	4.33E-07	5.42E-06	6.99E-05	2.42E-07	8.87E-04	9.63E-04			
Lloyd Lake	1.92E-07	2.19E-07	2.81E-06	6.48E-05	2.38E-07	8.73E-04	9.42E-04			

	Moose Total Dose Operations (mGy/d)										
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
	Application Case										
Reference (Broach Lake)	9.50E-06	1.08E-05	4.80E-06	9.61E-04	1.30E-06	2.66E-04	1.25E-03				
Patterson Lake North Arm - West Basin	4.24E-05	4.82E-05	3.75E-05	2.18E-03	2.14E-06	7.71E-04	3.09E-03				
Patterson Lake South Arm	4.20E-05	4.78E-05	3.69E-05	1.97E-03	2.03E-06	7.67E-04	2.87E-03				
Beet Lake	1.26E-05	1.44E-05	1.14E-05	1.08E-03	1.36E-06	3.10E-04	1.43E-03				
Lloyd Lake	9.61E-06	1.09E-05	5.37E-06	9.66E-04	1.30E-06	2.68E-04	1.26E-03				
		Uppe	er Bound Scenari	0							
Reference (Broach Lake)	9.50E-06	1.08E-05	4.80E-06	9.61E-04	1.30E-06	2.66E-04	1.25E-03				
Patterson Lake North Arm - West Basin	4.44E-05	5.05E-05	3.77E-05	2.19E-03	3.22E-06	1.01E-03	3.33E-03				
Patterson Lake South Arm	4.40E-05	5.01E-05	3.71E-05	1.98E-03	3.10E-06	1.00E-03	3.11E-03				
Beet Lake	1.28E-05	1.46E-05	1.14E-05	1.08E-03	1.36E-06	3.11E-04	1.43E-03				
Lloyd Lake	9.63E-06	1.10E-05	5.38E-06	9.66E-04	1.30E-06	2.68E-04	1.26E-03				

		Muskrat Tota	l Dose Operation	s (mGy/d)							
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
Application Case											
Reference (Broach Lake)	5.56E-07	6.32E-07	8.41E-06	2.23E-04	1.03E-06	1.76E-03	2.00E-03				
Patterson Lake North Arm - West Basin	5.07E-06	5.79E-06	4.71E-05	3.63E-04	1.46E-06	2.42E-03	2.84E-03				
Patterson Lake South Arm	1.75E-06	2.00E-06	3.06E-05	2.72E-04	1.06E-06	1.81E-03	2.11E-03				
Beet Lake	9.95E-07	1.13E-06	2.01E-05	2.44E-04	1.04E-06	1.77E-03	2.04E-03				
Lloyd Lake	5.96E-07	6.78E-07	9.61E-06	2.25E-04	1.03E-06	1.76E-03	2.00E-03				
		Uppe	er Bound Scenari	0							
Reference (Broach Lake)	5.56E-07	6.32E-07	8.41E-06	2.23E-04	1.03E-06	1.76E-03	2.00E-03				
Patterson Lake North Arm - West Basin	1.31E-05	1.50E-05	4.76E-05	3.70E-04	3.15E-06	4.88E-03	5.33E-03				
Patterson Lake South Arm	3.88E-06	4.43E-06	3.09E-05	2.74E-04	1.16E-06	1.97E-03	2.28E-03				
Beet Lake	1.78E-06	2.03E-06	2.02E-05	2.45E-04	1.05E-06	1.79E-03	2.06E-03				
Lloyd Lake	6.67E-07	7.59E-07	9.63E-06	2.25E-04	1.03E-06	1.76E-03	2.00E-03				

Table C.19: Maximum Radiation Dose to Ecological Receptors for the Application Case and Upper Bound Scenario - Project Lifespan

		Red Fox Tota	Dose Operation	s (mGy/d)			
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose
		Α	pplication Case				
Reference (Broach Lake)	5.92E-07	6.79E-07	6.84E-07	7.09E-04	1.59E-06	1.55E-05	7.28E-04
Patterson Lake North Arm - West Basin	2.67E-06	3.06E-06	2.86E-06	1.25E-03	1.95E-06	8.07E-05	1.34E-03
Patterson Lake South Arm	9.33E-07	1.07E-06	1.08E-06	7.97E-04	1.65E-06	2.61E-05	8.28E-04
Beet Lake	7.90E-07	9.06E-07	9.11E-07	7.60E-04	1.62E-06	2.17E-05	7.86E-04
Lloyd Lake	5.99E-07	6.87E-07	6.94E-07	7.10E-04	1.59E-06	1.57E-05	7.30E-04
		Uppe	er Bound Scenari	0			
Reference (Broach Lake)	5.92E-07	6.79E-07	6.84E-07	7.09E-04	1.59E-06	1.55E-05	7.28E-04
Patterson Lake North Arm - West Basin	2.70E-06	3.09E-06	2.86E-06	1.25E-03	1.95E-06	8.14E-05	1.34E-03
Patterson Lake South Arm	9.38E-07	1.08E-06	1.08E-06	7.97E-04	1.65E-06	2.62E-05	8.28E-04
Beet Lake	7.92E-07	9.08E-07	9.12E-07	7.60E-04	1.62E-06	2.17E-05	7.86E-04
Lloyd Lake	5.99E-07	6.87E-07	6.94E-07	7.10E-04	1.59E-06	1.57E-05	7.30E-04

	Snowshoe Hare Total Dose Operations (mGy/d)										
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
	Application Case										
Reference (Broach Lake)	9.84E-06	1.12E-05	1.96E-06	1.26E-03	1.96E-06	1.48E-04	1.44E-03				
Patterson Lake North Arm - West Basin	4.32E-05	4.92E-05	1.82E-05	2.69E-03	2.76E-06	6.07E-04	3.41E-03				
Patterson Lake South Arm	1.53E-05	1.74E-05	4.64E-06	1.50E-03	2.09E-06	2.23E-04	1.76E-03				
Beet Lake	1.30E-05	1.48E-05	3.52E-06	1.40E-03	2.04E-06	1.92E-04	1.62E-03				
Lloyd Lake	9.95E-06	1.13E-05	2.01E-06	1.27E-03	1.97E-06	1.50E-04	1.44E-03				
		Upp	er Bound Scenari	0							
Reference (Broach Lake)	9.84E-06	1.12E-05	1.96E-06	1.26E-03	1.96E-06	1.48E-04	1.44E-03				
Patterson Lake North Arm - West Basin	4.32E-05	4.92E-05	1.82E-05	2.69E-03	2.76E-06	6.07E-04	3.41E-03				
Patterson Lake South Arm	1.53E-05	1.74E-05	4.64E-06	1.50E-03	2.09E-06	2.23E-04	1.76E-03				
Beet Lake	1.30E-05	1.48E-05	3.52E-06	1.40E-03	2.04E-06	1.92E-04	1.62E-03				
Lloyd Lake	9.95E-06	1.13E-05	2.01E-06	1.27E-03	1.97E-06	1.50E-04	1.44E-03				

	South	ern Red-Backed \	Vole Total Dose (Operations (mGy	//d)					
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose			
Application Case										
Reference (Broach Lake)	3.58E-06	4.08E-06	7.10E-07	9.01E-04	1.85E-06	5.26E-05	9.64E-04			
Patterson Lake North Arm - West Basin	1.59E-05	1.82E-05	4.07E-06	1.67E-03	2.33E-06	1.96E-04	1.91E-03			
Patterson Lake South Arm	5.59E-06	6.37E-06	1.28E-06	1.03E-03	1.93E-06	7.59E-05	1.12E-03			
Beet Lake	4.75E-06	5.42E-06	1.04E-06	9.75E-04	1.90E-06	6.62E-05	1.05E-03			
Lloyd Lake	3.62E-06	4.12E-06	7.22E-07	9.04E-04	1.85E-06	5.31E-05	9.67E-04			
		Uppe	er Bound Scenari	0						
Reference (Broach Lake)	3.58E-06	4.08E-06	7.10E-07	9.01E-04	1.85E-06	5.26E-05	9.64E-04			
Patterson Lake North Arm - West Basin	1.59E-05	1.82E-05	4.07E-06	1.67E-03	2.33E-06	1.96E-04	1.91E-03			
Patterson Lake South Arm	5.60E-06	6.38E-06	1.28E-06	1.03E-03	1.93E-06	7.60E-05	1.12E-03			
Beet Lake	4.75E-06	5.42E-06	1.04E-06	9.75E-04	1.90E-06	6.62E-05	1.05E-03			
Lloyd Lake	3.62E-06	4.12E-06	7.22E-07	9.04E-04	1.85E-06	5.31E-05	9.67E-04			

WoodLand Caribou Total Dose Operations (mGy/d)									
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose		
		A	pplication Case						
Reference (Broach Lake)	4.34E-06	4.94E-06	5.85E-06	7.54E-04	6.27E-06	3.51E-03	4.28E-03		
Patterson Lake North Arm - West Basin	1.23E-05	1.40E-05	8.67E-05	3.59E-03	8.33E-06	3.97E-03	7.69E-03		
Beet Lake	5.01E-06	5.70E-06	1.42E-05	1.00E-03	6.46E-06	3.56E-03	4.59E-03		
Lloyd Lake	4.40E-06	5.01E-06	6.78E-06	7.72E-04	6.28E-06	3.51E-03	4.31E-03		
		Uppe	er Bound Scenari	0					
Reference (Broach Lake)	4.34E-06	4.94E-06	5.85E-06	7.54E-04	6.27E-06	3.51E-03	4.28E-03		
Patterson Lake North Arm - West Basin	1.47E-05	1.67E-05	8.69E-05	3.60E-03	8.87E-06	4.16E-03	7.88E-03		
Beet Lake	5.20E-06	5.92E-06	1.43E-05	1.00E-03	6.46E-06	3.56E-03	4.60E-03		
Lloyd Lake	4.44E-06	5.05E-06	6.78E-06	7.72E-04	6.28E-06	3.51E-03	4.31E-03		

Table C.19: Maximum Radiation Dose to Ecological Receptors for the Application Case and Upper Bound Scenario - Project Lifespan

	Grouse Total Dose Operations (mGy/d)											
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose					
	Application Case											
Reference (Broach Lake)	2.34E-04	2.67E-04	1.70E-06	7.73E-04	1.88E-06	9.49E-04	2.23E-03					
Patterson Lake North Arm - West Basin	1.03E-03	1.17E-03	9.22E-06	1.40E-03	2.58E-06	3.53E-03	7.15E-03					
Patterson Lake South Arm	3.64E-04	4.14E-04	2.94E-06	8.75E-04	1.99E-06	1.37E-03	3.03E-03					
Beet Lake	3.10E-04	3.53E-04	2.42E-06	8.32E-04	1.94E-06	1.19E-03	2.69E-03					
Lloyd Lake	2.37E-04	2.69E-04	1.72E-06	7.75E-04	1.88E-06	9.57E-04	2.24E-03					
		Uppe	er Bound Scenari	0								
Reference (Broach Lake)	2.34E-04	2.67E-04	1.70E-06	7.73E-04	1.88E-06	9.49E-04	2.23E-03					
Patterson Lake North Arm - West Basin	1.03E-03	1.17E-03	9.22E-06	1.40E-03	2.58E-06	3.54E-03	7.15E-03					
Patterson Lake South Arm	3.64E-04	4.15E-04	2.94E-06	8.75E-04	1.99E-06	1.37E-03	3.03E-03					
Beet Lake	3.10E-04	3.53E-04	2.42E-06	8.32E-04	1.94E-06	1.19E-03	2.69E-03					
Lloyd Lake	2.37E-04	2.69E-04	1.72E-06	7.75E-04	1.88E-06	9.57E-04	2.24E-03					

		Canada Goose To	otal Dose Operat	ions (mGy/d)							
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
Application Case											
Reference (Broach Lake)	3.28E-05	3.74E-05	4.00E-07	6.64E-04	1.71E-06	1.34E-04	8.70E-04				
Patterson Lake North Arm - West Basin	1.44E-04	1.64E-04	1.85E-06	1.12E-03	2.09E-06	5.24E-04	1.95E-03				
Patterson Lake South Arm	5.10E-05	5.80E-05	6.46E-07	7.37E-04	1.77E-06	1.97E-04	1.05E-03				
Beet Lake	4.34E-05	4.94E-05	5.42E-07	7.07E-04	1.74E-06	1.71E-04	9.73E-04				
Lloyd Lake	3.32E-05	3.78E-05	4.05E-07	6.65E-04	1.71E-06	1.35E-04	8.73E-04				
		Upp	er Bound Scenari	0							
Reference (Broach Lake)	3.28E-05	3.74E-05	4.00E-07	6.64E-04	1.71E-06	1.34E-04	8.70E-04				
Patterson Lake North Arm - West Basin	1.44E-04	1.64E-04	1.85E-06	1.12E-03	2.09E-06	5.26E-04	1.95E-03				
Patterson Lake South Arm	5.11E-05	5.81E-05	6.46E-07	7.37E-04	1.77E-06	1.98E-04	1.05E-03				
Beet Lake	4.34E-05	4.94E-05	5.43E-07	7.07E-04	1.74E-06	1.71E-04	9.73E-04				
Lloyd Lake	3.32E-05	3.78E-05	4.05E-07	6.65E-04	1.71E-06	1.35E-04	8.73E-04				

	Little Brown Myotis Total Dose Operations (mGy/d)										
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
	Application Case										
Reference (Broach Lake)	1.37E-07	1.55E-07	1.19E-05	1.36E-04	4.92E-07	3.09E-03	3.24E-03				
Patterson Lake North Arm - West Basin	1.24E-06	1.41E-06	6.61E-05	2.19E-04	6.74E-07	4.22E-03	4.51E-03				
Patterson Lake South Arm	4.27E-07	4.88E-07	4.27E-05	1.65E-04	5.04E-07	3.16E-03	3.37E-03				
Beet Lake	2.44E-07	2.78E-07	2.81E-05	1.48E-04	4.94E-07	3.10E-03	3.27E-03				
Lloyd Lake	1.46E-07	1.67E-07	1.36E-05	1.37E-04	4.92E-07	3.09E-03	3.24E-03				
		Uppe	er Bound Scenari	0							
Reference (Broach Lake)	1.37E-07	1.55E-07	1.19E-05	1.36E-04	4.92E-07	3.09E-03	3.24E-03				
Patterson Lake North Arm - West Basin	3.18E-06	3.63E-06	6.68E-05	2.22E-04	1.35E-06	8.46E-03	8.76E-03				
Patterson Lake South Arm	9.45E-07	1.08E-06	4.32E-05	1.66E-04	5.49E-07	3.45E-03	3.66E-03				
Beet Lake	4.33E-07	4.94E-07	2.83E-05	1.49E-04	5.00E-07	3.14E-03	3.31E-03				
Lloyd Lake	1.64E-07	1.86E-07	1.36E-05	1.37E-04	4.92E-07	3.09E-03	3.24E-03				

	Loon Total Dose Operations (mGy/d)									
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose			
Application Case										
Reference (Broach Lake)	1.33E-06	1.51E-06	1.01E-06	6.06E-06	8.76E-08	2.97E-03	2.98E-03			
Patterson Lake North Arm - West Basin	1.37E-05	1.57E-05	5.65E-06	9.94E-06	1.23E-07	4.09E-03	4.14E-03			
Patterson Lake South Arm	4.59E-06	5.24E-06	3.63E-06	7.38E-06	8.99E-08	3.05E-03	3.07E-03			
Beet Lake	2.53E-06	2.89E-06	2.39E-06	6.62E-06	8.79E-08	2.98E-03	3.00E-03			
Lloyd Lake	1.44E-06	1.64E-06	1.15E-06	6.11E-06	8.76E-08	2.97E-03	2.98E-03			
		Uppe	er Bound Scenari	0						
Reference (Broach Lake)	1.33E-06	1.51E-06	1.01E-06	6.06E-06	8.76E-08	2.97E-03	2.98E-03			
Patterson Lake North Arm - West Basin	3.61E-05	4.12E-05	5.71E-06	1.01E-05	2.57E-07	8.35E-03	8.45E-03			
Patterson Lake South Arm	1.03E-05	1.17E-05	3.67E-06	7.45E-06	9.84E-08	3.32E-03	3.36E-03			
Beet Lake	4.61E-06	5.27E-06	2.41E-06	6.65E-06	8.91E-08	3.02E-03	3.04E-03			
Lloyd Lake	1.63E-06	1.86E-06	1.15E-06	6.12E-06	8.76E-08	2.97E-03	2.98E-03			

Table C.19: Maximum Radiation Dose to Ecological Receptors for the Application Case and Upper Bound Scenario - Project Lifespan

Mallard Total Dose Operations (mGy/d)									
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose		
		Α	pplication Case						
Reference (Broach Lake)	1.40E-05	1.59E-05	9.79E-06	7.84E-05	1.23E-06	2.55E-02	2.56E-02		
Patterson Lake North Arm - West Basin	1.27E-04	1.45E-04	5.46E-05	1.27E-04	1.73E-06	3.49E-02	3.53E-02		
Patterson Lake South Arm	4.40E-05	5.02E-05	3.54E-05	9.52E-05	1.26E-06	2.61E-02	2.63E-02		
Beet Lake	2.50E-05	2.85E-05	2.33E-05	8.56E-05	1.23E-06	2.55E-02	2.57E-02		
Lloyd Lake	1.50E-05	1.71E-05	1.12E-05	7.91E-05	1.23E-06	2.55E-02	2.56E-02		
		Uppe	er Bound Scenari	0					
Reference (Broach Lake)	1.40E-05	1.59E-05	9.79E-06	7.84E-05	1.23E-06	2.55E-02	2.56E-02		
Patterson Lake North Arm - West Basin	3.29E-04	3.76E-04	5.51E-05	1.29E-04	3.66E-06	7.01E-02	7.10E-02		
Patterson Lake South Arm	9.75E-05	1.11E-04	3.57E-05	9.60E-05	1.38E-06	2.84E-02	2.88E-02		
Beet Lake	4.46E-05	5.09E-05	2.34E-05	8.59E-05	1.25E-06	2.59E-02	2.61E-02		
Lloyd Lake	1.68E-05	1.91E-05	1.12E-05	7.92E-05	1.23E-06	2.55E-02	2.56E-02		

	Rusty Blackbird Total Dose Operations (mGy/d)										
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
	Application Case										
Reference (Broach Lake)	1.01E-04	1.15E-04	2.74E-05	8.28E-04	3.65E-06	6.96E-02	7.07E-02				
Patterson Lake North Arm - West Basin	5.07E-04	5.77E-04	1.39E-04	1.39E-03	4.70E-06	9.59E-02	9.85E-02				
Patterson Lake South Arm	1.77E-04	2.01E-04	9.02E-05	9.31E-04	3.76E-06	7.15E-02	7.29E-02				
Beet Lake	1.40E-04	1.59E-04	6.05E-05	8.84E-04	3.70E-06	7.00E-02	7.12E-02				
Lloyd Lake	1.03E-04	1.17E-04	3.08E-05	8.30E-04	3.65E-06	6.97E-02	7.07E-02				
		Uppe	er Bound Scenari	0							
Reference (Broach Lake)	1.01E-04	1.15E-04	2.74E-05	8.28E-04	3.65E-06	6.96E-02	7.07E-02				
Patterson Lake North Arm - West Basin	6.81E-04	7.76E-04	1.41E-04	1.39E-03	6.99E-06	1.91E-01	1.94E-01				
Patterson Lake South Arm	2.23E-04	2.54E-04	9.10E-05	9.32E-04	3.92E-06	7.78E-02	7.93E-02				
Beet Lake	1.57E-04	1.78E-04	6.09E-05	8.85E-04	3.72E-06	7.08E-02	7.21E-02				
Lloyd Lake	1.04E-04	1.19E-04	3.09E-05	8.30E-04	3.65E-06	6.97E-02	7.08E-02				

Table C.20: Maximum Radiation Dose to Ecological Receptors for the Application Case and Upper Bound Scenario - Far-Future

	Ве	nthic Invertebra	tes Total Dose F	ar Future (mGy,	/d)						
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
Application Case											
Reference (Broach Lake)	4.73E-09	6.18E-09	1.80E-08	8.07E-05	3.79E-05	1.72E-09	1.19E-04				
Patterson Lake North Arm - West Basin	1.53E-07	2.01E-07	1.93E-08	1.04E-04	3.97E-05	1.80E-09	1.44E-04				
Patterson Lake South Arm	4.58E-08	6.00E-08	1.88E-08	8.93E-05	3.80E-05	1.72E-09	1.28E-04				
Beet Lake	2.00E-08	2.62E-08	1.84E-08	8.44E-05	3.79E-05	1.72E-09	1.22E-04				
Lloyd Lake	6.11E-09	7.99E-09	1.81E-08	8.11E-05	3.79E-05	1.72E-09	1.19E-04				
		Upp	er Bound Scena	rio							
Reference (Broach Lake)	4.73E-09	6.18E-09	1.80E-08	8.07E-05	3.79E-05	1.72E-09	1.19E-04				
Patterson Lake North Arm - West Basin	1.53E-07	2.01E-07	1.93E-08	1.04E-04	3.97E-05	1.80E-09	1.44E-04				
Patterson Lake South Arm	4.58E-08	6.01E-08	1.88E-08	8.94E-05	3.80E-05	1.72E-09	1.28E-04				
Beet Lake	2.00E-08	2.62E-08	1.84E-08	8.45E-05	3.79E-05	1.72E-09	1.22E-04				
Lloyd Lake	6.11E-09	7.99E-09	1.81E-08	8.11E-05	3.79E-05	1.72E-09	1.19E-04				

	Zooplankton Total Dose Far Future (mGy/d)										
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
		A	Application Case								
Reference (Broach Lake)	2.05E-05	2.39E-05	2.94E-03	4.57E-03	2.80E-05	6.05E-02	6.81E-02				
Patterson Lake North Arm - West Basin	6.63E-04	7.77E-04	3.15E-03	5.89E-03	2.93E-05	6.33E-02	7.38E-02				
Patterson Lake South Arm	1.98E-04	2.32E-04	3.07E-03	5.05E-03	2.81E-05	6.07E-02	6.93E-02				
Beet Lake	8.66E-05	1.01E-04	3.01E-03	4.78E-03	2.80E-05	6.05E-02	6.85E-02				
Lloyd Lake	2.65E-05	3.09E-05	2.95E-03	4.59E-03	2.80E-05	6.05E-02	6.81E-02				
		Upp	er Bound Scena	rio							
Reference (Broach Lake)	2.05E-05	2.39E-05	2.94E-03	4.57E-03	2.80E-05	6.05E-02	6.81E-02				
Patterson Lake North Arm - West Basin	6.64E-04	7.78E-04	3.15E-03	5.89E-03	2.93E-05	6.33E-02	7.38E-02				
Patterson Lake South Arm	1.99E-04	2.33E-04	3.07E-03	5.06E-03	2.81E-05	6.07E-02	6.93E-02				
Beet Lake	8.67E-05	1.01E-04	3.01E-03	4.78E-03	2.80E-05	6.05E-02	6.85E-02				
Lloyd Lake	2.65E-05	3.09E-05	2.95E-03	4.59E-03	2.80E-05	6.05E-02	6.81E-02				

		Northern Pike	Total Dose Far F	uture (mGy/d)							
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
	Application Case										
Reference (Broach Lake)	4.13E-06	4.70E-06	3.54E-05	2.22E-04	1.79E-06	5.64E-04	8.33E-04				
Patterson Lake North Arm - West Basin	1.34E-04	1.53E-04	3.79E-05	2.87E-04	1.88E-06	5.90E-04	1.20E-03				
Patterson Lake South Arm	4.00E-05	4.56E-05	3.69E-05	2.46E-04	1.80E-06	5.66E-04	9.36E-04				
Beet Lake	1.74E-05	1.99E-05	3.62E-05	2.32E-04	1.79E-06	5.65E-04	8.72E-04				
Lloyd Lake	5.33E-06	6.07E-06	3.55E-05	2.23E-04	1.79E-06	5.64E-04	8.36E-04				
		Upp	er Bound Scena	rio							
Reference (Broach Lake)	4.13E-06	4.70E-06	3.54E-05	2.22E-04	1.79E-06	5.64E-04	8.33E-04				
Patterson Lake North Arm - West Basin	1.34E-04	1.53E-04	3.79E-05	2.87E-04	1.88E-06	5.90E-04	1.20E-03				
Patterson Lake South Arm	4.00E-05	4.57E-05	3.69E-05	2.46E-04	1.80E-06	5.66E-04	9.36E-04				
Beet Lake	1.75E-05	1.99E-05	3.62E-05	2.32E-04	1.79E-06	5.65E-04	8.72E-04				
Lloyd Lake	5.34E-06	6.08E-06	3.55E-05	2.23E-04	1.79E-06	5.64E-04	8.36E-04				

		Whitefish To	tal Dose Far Futi	ure (mGy/d)			
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose
		A	Application Case				
Reference (Broach Lake)	7.69E-07	8.97E-07	2.65E-05	1.86E-04	8.29E-07	4.31E-05	2.58E-04
Patterson Lake North Arm - West Basin	2.49E-05	2.91E-05	2.83E-05	2.39E-04	8.68E-07	4.51E-05	3.68E-04
Patterson Lake South Arm	7.44E-06	8.71E-06	2.76E-05	2.05E-04	8.32E-07	4.33E-05	2.93E-04
Beet Lake	3.25E-06	3.80E-06	2.71E-05	1.94E-04	8.30E-07	4.32E-05	2.72E-04
Lloyd Lake	9.93E-07	1.16E-06	2.66E-05	1.86E-04	8.29E-07	4.31E-05	2.59E-04
		Upp	er Bound Scena	rio			
Reference (Broach Lake)	7.69E-07	8.97E-07	2.65E-05	1.86E-04	8.29E-07	4.31E-05	2.58E-04
Patterson Lake North Arm - West Basin	2.49E-05	2.92E-05	2.83E-05	2.39E-04	8.68E-07	4.51E-05	3.68E-04
Patterson Lake South Arm	7.45E-06	8.72E-06	2.76E-05	2.05E-04	8.32E-07	4.33E-05	2.93E-04
Beet Lake	3.25E-06	3.81E-06	2.71E-05	1.94E-04	8.30E-07	4.32E-05	2.72E-04
Lloyd Lake	9.94E-07	1.16E-06	2.66E-05	1.86E-04	8.29E-07	4.31E-05	2.59E-04

Table C.20: Maximum Radiation Dose to Ecological Receptors for the Application Case and Upper Bound Scenario - Far-Future

		Macrophytes T	otal Dose Far Fu	ıture (mGy/d)							
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
	Application Case										
Reference (Broach Lake)	1.04E-05	1.18E-05	1.37E-03	2.99E-03	5.06E-05	1.10E-03	5.53E-03				
Patterson Lake North Arm - West Basin	3.37E-04	3.85E-04	1.47E-03	3.85E-03	5.30E-05	1.15E-03	7.25E-03				
Patterson Lake South Arm	1.01E-04	1.15E-04	1.43E-03	3.31E-03	5.08E-05	1.10E-03	6.11E-03				
Beet Lake	4.40E-05	5.02E-05	1.40E-03	3.13E-03	5.07E-05	1.10E-03	5.77E-03				
Lloyd Lake	1.34E-05	1.53E-05	1.37E-03	3.00E-03	5.06E-05	1.10E-03	5.56E-03				
		Upp	er Bound Scena	rio							
Reference (Broach Lake)	1.04E-05	1.18E-05	1.37E-03	2.99E-03	5.06E-05	1.10E-03	5.53E-03				
Patterson Lake North Arm - West Basin	3.37E-04	3.85E-04	1.47E-03	3.86E-03	5.30E-05	1.15E-03	7.25E-03				
Patterson Lake South Arm	1.01E-04	1.15E-04	1.43E-03	3.31E-03	5.08E-05	1.10E-03	6.11E-03				
Beet Lake	4.40E-05	5.02E-05	1.40E-03	3.13E-03	5.07E-05	1.10E-03	5.77E-03				
Lloyd Lake	1.34E-05	1.53E-05	1.37E-03	3.00E-03	5.06E-05	1.10E-03	5.56E-03				

		Phytoplankton '	Total Dose Far F	uture (mGy/d)							
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
	Application Case										
Reference (Broach Lake)	1.04E-05	1.18E-05	1.36E-03	3.00E-03	1.93E-05	1.11E-03	5.51E-03				
Patterson Lake North Arm - West Basin	3.37E-04	3.84E-04	1.46E-03	3.86E-03	2.02E-05	1.16E-03	7.22E-03				
Patterson Lake South Arm	1.01E-04	1.15E-04	1.42E-03	3.32E-03	1.94E-05	1.11E-03	6.08E-03				
Beet Lake	4.39E-05	5.01E-05	1.39E-03	3.13E-03	1.94E-05	1.11E-03	5.75E-03				
Lloyd Lake	1.34E-05	1.53E-05	1.37E-03	3.01E-03	1.93E-05	1.11E-03	5.53E-03				
		Upp	er Bound Scena	rio							
Reference (Broach Lake)	1.04E-05	1.18E-05	1.36E-03	3.00E-03	1.93E-05	1.11E-03	5.51E-03				
Patterson Lake North Arm - West Basin	3.37E-04	3.84E-04	1.46E-03	3.87E-03	2.03E-05	1.16E-03	7.22E-03				
Patterson Lake South Arm	1.01E-04	1.15E-04	1.42E-03	3.32E-03	1.94E-05	1.11E-03	6.08E-03				
Beet Lake	4.40E-05	5.01E-05	1.39E-03	3.13E-03	1.94E-05	1.11E-03	5.75E-03				
Lloyd Lake	1.34E-05	1.53E-05	1.37E-03	3.01E-03	1.93E-05	1.11E-03	5.53E-03				

	Ter	restrial Invertebi	rate Total Dose I	Far Future (mGy	//d)						
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
	Application Case										
Reference (Broach Lake)	1.57E-05	1.78E-05	1.15E-04	5.73E-03	6.72E-06	2.28E-05	5.91E-03				
Patterson Lake North Arm - West Basin	2.13E-05	2.42E-05	1.52E-04	8.26E-03	6.72E-06	2.28E-05	8.49E-03				
Patterson Lake South Arm	1.66E-05	1.89E-05	1.21E-04	6.15E-03	6.72E-06	2.28E-05	6.33E-03				
Beet Lake	1.62E-05	1.84E-05	1.19E-04	5.97E-03	6.72E-06	2.28E-05	6.16E-03				
Lloyd Lake	1.57E-05	1.79E-05	1.15E-04	5.74E-03	6.72E-06	2.28E-05	5.92E-03				
		Upp	er Bound Scena	rio							
Reference (Broach Lake)	1.57E-05	1.78E-05	1.15E-04	5.73E-03	6.72E-06	2.28E-05	5.91E-03				
Patterson Lake North Arm - West Basin	2.13E-05	2.42E-05	1.52E-04	8.26E-03	6.72E-06	2.28E-05	8.49E-03				
Patterson Lake South Arm	1.66E-05	1.89E-05	1.21E-04	6.15E-03	6.72E-06	2.28E-05	6.33E-03				
Beet Lake	1.62E-05	1.84E-05	1.19E-04	5.97E-03	6.72E-06	2.28E-05	6.16E-03				
Lloyd Lake	1.57E-05	1.79E-05	1.15E-04	5.74E-03	6.72E-06	2.28E-05	5.92E-03				

	Browse Total Dose Far Future (mGy/d)										
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
	Application Case										
Patterson Lake North Arm - West Basin	7.83E-04	8.91E-04	1.36E-04	1.15E-02	1.65E-04	6.33E-04	1.41E-02				
Patterson Lake South Arm	6.11E-04	6.96E-04	1.08E-04	8.55E-03	1.65E-04	6.33E-04	1.08E-02				
		Upp	er Bound Scena	rio							
Patterson Lake North Arm - West Basin	7.83E-04	8.91E-04	1.36E-04	1.15E-02	1.65E-04	6.33E-04	1.41E-02				
Patterson Lake South Arm	6.11E-04	6.96E-04	1.08E-04	8.55E-03	1.65E-04	6.33E-04	1.08E-02				

Table C.20: Maximum Radiation Dose to Ecological Receptors for the Application Case and Upper Bound Scenario - Far-Future

Labrador Tea Total Dose Far Future (mGy/d)											
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
	Application Case										
Patterson Lake North Arm - West Basin	7.83E-04	8.91E-04	1.36E-04	1.15E-02	1.65E-04	6.33E-04	1.41E-02				
Patterson Lake South Arm	6.11E-04	6.96E-04	1.08E-04	8.55E-03	1.65E-04	6.33E-04	1.08E-02				
		Upp	er Bound Scena	rio							
Patterson Lake North Arm - West Basin	7.83E-04	8.91E-04	1.36E-04	1.15E-02	1.65E-04	6.33E-04	1.41E-02				
Patterson Lake South Arm	6.11E-04	6.96E-04	1.08E-04	8.55E-03	1.65E-04	6.33E-04	1.08E-02				

	Lichen Total Dose Operations (mGy/d)										
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
	Application Case										
Patterson Lake North Arm - West Basin	1.21E-04	1.37E-04	4.38E-04	4.44E-03	8.33E-04	1.06E-01	1.12E-01				
Patterson Lake South Arm	1.21E-04	1.37E-04	4.38E-04	4.44E-03	8.33E-04	1.06E-01	1.12E-01				
		Upp	er Bound Scena	rio							
Patterson Lake North Arm - West Basin	1.21E-04	1.37E-04	4.38E-04	4.44E-03	8.33E-04	1.06E-01	1.12E-01				
Patterson Lake South Arm	1.21E-04	1.37E-04	4.38E-04	4.44E-03	8.33E-04	1.06E-01	1.12E-01				

	Blueberries Total Dose Far Future (mGy/d)									
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose			
		A	Application Case							
Patterson Lake North Arm - West Basin	3.89E-04	4.43E-04	5.39E-05	5.59E-03	3.00E-06	3.16E-04	6.80E-03			
Patterson Lake South Arm	3.04E-04	3.46E-04	4.29E-05	4.16E-03	3.00E-06	3.16E-04	5.17E-03			
		Upp	er Bound Scena	rio						
Patterson Lake North Arm - West Basin	3.89E-04	4.43E-04	5.39E-05	5.59E-03	3.00E-06	3.16E-04	6.80E-03			
Patterson Lake South Arm	3.04E-04	3.46E-04	4.29E-05	4.16E-03	3.00E-06	3.16E-04	5.17E-03			

		Beaver Tota	al Dose Far Futui	re (mGy/d)							
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
Application Case											
Reference (Broach Lake)	1.05E-05	1.19E-05	3.11E-06	6.36E-04	4.75E-07	2.15E-04	8.77E-04				
Patterson Lake North Arm - West Basin	1.61E-05	1.84E-05	3.68E-06	9.09E-04	4.90E-07	2.18E-04	1.17E-03				
Patterson Lake South Arm	1.16E-05	1.32E-05	3.26E-06	6.84E-04	4.76E-07	2.15E-04	9.27E-04				
Beet Lake	1.10E-05	1.26E-05	3.19E-06	6.63E-04	4.75E-07	2.15E-04	9.05E-04				
Lloyd Lake	1.05E-05	1.20E-05	3.11E-06	6.37E-04	4.75E-07	2.15E-04	8.78E-04				
		Upp	er Bound Scena	rio							
Reference (Broach Lake)	1.05E-05	1.19E-05	3.11E-06	6.36E-04	4.75E-07	2.15E-04	8.77E-04				
Patterson Lake North Arm - West Basin	1.61E-05	1.84E-05	3.68E-06	9.09E-04	4.90E-07	2.18E-04	1.17E-03				
Patterson Lake South Arm	1.16E-05	1.32E-05	3.26E-06	6.84E-04	4.76E-07	2.15E-04	9.27E-04				
Beet Lake	1.10E-05	1.26E-05	3.19E-06	6.63E-04	4.75E-07	2.15E-04	9.05E-04				
Lloyd Lake	1.05E-05	1.20E-05	3.11E-06	6.37E-04	4.75E-07	2.15E-04	8.78E-04				

	Black Bear Total Dose Far Future (mGy/d)									
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose			
		P	Application Case							
Reference (Broach Lake)	7.81E-06	8.89E-06	1.88E-06	8.19E-04	8.32E-07	1.74E-04	1.01E-03			
Patterson Lake North Arm - West Basin	9.86E-06	1.12E-05	2.21E-06	1.01E-03	8.33E-07	1.76E-04	1.21E-03			
Beet Lake	8.12E-06	9.24E-06	2.52E-06	1.05E-03	1.21E-06	1.82E-04	1.26E-03			
Lloyd Lake	7.83E-06	8.91E-06	1.91E-06	8.21E-04	8.33E-07	1.75E-04	1.02E-03			
		Upp	er Bound Scena	rio						
Reference (Broach Lake)	7.81E-06	8.89E-06	1.88E-06	8.19E-04	8.32E-07	1.74E-04	1.01E-03			
Patterson Lake North Arm - West Basin	9.86E-06	1.12E-05	2.21E-06	1.01E-03	8.33E-07	1.76E-04	1.21E-03			
Beet Lake	8.12E-06	9.24E-06	2.52E-06	1.05E-03	1.21E-06	1.82E-04	1.26E-03			
Lloyd Lake	7.83E-06	8.91E-06	1.91E-06	8.21E-04	8.33E-07	1.75E-04	1.02E-03			

Table C.20: Maximum Radiation Dose to Ecological Receptors for the Application Case and Upper Bound Scenario - Far-Future

	Grey Wolf Total Dose Far Future (mGy/d)										
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
		,	Application Case								
Reference (Broach Lake)	3.26E-07	3.77E-07	1.17E-06	7.63E-04	1.92E-06	3.26E-04	1.09E-03				
Patterson Lake North Arm - West Basin	3.90E-07	4.50E-07	1.26E-06	8.45E-04	1.92E-06	3.27E-04	1.18E-03				
Lloyd Lake	3.27E-07	3.79E-07	1.18E-06	7.64E-04	1.92E-06	3.26E-04	1.09E-03				
		Upp	er Bound Scena	rio							
Reference (Broach Lake)	3.26E-07	3.77E-07	1.17E-06	7.63E-04	1.92E-06	3.26E-04	1.09E-03				
Patterson Lake North Arm - West Basin	3.90E-07	4.50E-07	1.26E-06	8.45E-04	1.92E-06	3.27E-04	1.18E-03				
Lloyd Lake	3.27E-07	3.79E-07	1.18E-06	7.64E-04	1.92E-06	3.26E-04	1.09E-03				

		Mink Total	Dose Far Future	e (mGy/d)							
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
Application Case											
Reference (Broach Lake)	1.76E-07	2.00E-07	2.51E-06	6.44E-05	2.38E-07	8.73E-04	9.40E-04				
Patterson Lake North Arm - West Basin	2.66E-06	3.04E-06	2.79E-06	8.52E-05	2.48E-07	9.13E-04	1.01E-03				
Patterson Lake South Arm	8.59E-07	9.81E-07	2.62E-06	7.07E-05	2.38E-07	8.75E-04	9.51E-04				
Beet Lake	4.31E-07	4.92E-07	2.57E-06	6.72E-05	2.38E-07	8.73E-04	9.44E-04				
Lloyd Lake	1.98E-07	2.26E-07	2.51E-06	6.46E-05	2.38E-07	8.73E-04	9.40E-04				
		Upp	er Bound Scena	rio							
Reference (Broach Lake)	1.76E-07	2.00E-07	2.51E-06	6.44E-05	2.38E-07	8.73E-04	9.40E-04				
Patterson Lake North Arm - West Basin	2.66E-06	3.04E-06	2.79E-06	8.52E-05	2.48E-07	9.13E-04	1.01E-03				
Patterson Lake South Arm	8.60E-07	9.82E-07	2.62E-06	7.07E-05	2.38E-07	8.75E-04	9.51E-04				
Beet Lake	4.31E-07	4.92E-07	2.57E-06	6.72E-05	2.38E-07	8.73E-04	9.44E-04				
Lloyd Lake	1.98E-07	2.26E-07	2.51E-06	6.46E-05	2.38E-07	8.73E-04	9.40E-04				

	Moose Total Dose Far Future (mGy/d)										
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
	Application Case										
Reference (Broach Lake)	9.50E-06	1.08E-05	4.80E-06	9.61E-04	1.30E-06	2.66E-04	1.25E-03				
Patterson Lake North Arm - West Basin	1.69E-05	1.92E-05	5.47E-06	1.37E-03	1.32E-06	2.73E-04	1.69E-03				
Patterson Lake South Arm	1.67E-05	1.91E-05	5.28E-06	1.23E-03	1.32E-06	2.72E-04	1.55E-03				
Beet Lake	1.02E-05	1.16E-05	4.92E-06	1.00E-03	1.30E-06	2.67E-04	1.30E-03				
Lloyd Lake	9.55E-06	1.09E-05	4.81E-06	9.63E-04	1.30E-06	2.66E-04	1.26E-03				
		Upp	er Bound Scena	rio							
Reference (Broach Lake)	9.50E-06	1.08E-05	4.80E-06	9.61E-04	1.30E-06	2.66E-04	1.25E-03				
Patterson Lake North Arm - West Basin	1.69E-05	1.92E-05	5.47E-06	1.37E-03	1.32E-06	2.73E-04	1.69E-03				
Patterson Lake South Arm	1.68E-05	1.91E-05	5.28E-06	1.23E-03	1.32E-06	2.72E-04	1.55E-03				
Beet Lake	1.02E-05	1.16E-05	4.92E-06	1.00E-03	1.30E-06	2.67E-04	1.30E-03				
Lloyd Lake	9.55E-06	1.09E-05	4.81E-06	9.63E-04	1.30E-06	2.66E-04	1.26E-03				

	Muskrat Total Dose Far Future (mGy/d)										
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
Application Case											
Reference (Broach Lake)	5.56E-07	6.32E-07	8.41E-06	2.23E-04	1.03E-06	1.76E-03	2.00E-03				
Patterson Lake North Arm - West Basin	1.80E-05	2.06E-05	8.99E-06	2.88E-04	1.08E-06	1.84E-03	2.18E-03				
Patterson Lake South Arm	5.38E-06	6.14E-06	8.77E-06	2.47E-04	1.04E-06	1.77E-03	2.04E-03				
Beet Lake	2.35E-06	2.68E-06	8.60E-06	2.33E-04	1.03E-06	1.76E-03	2.01E-03				
Lloyd Lake	7.18E-07	8.18E-07	8.43E-06	2.24E-04	1.03E-06	1.76E-03	2.00E-03				
		Upp	er Bound Scena	rio							
Reference (Broach Lake)	5.56E-07	6.32E-07	8.41E-06	2.23E-04	1.03E-06	1.76E-03	2.00E-03				
Patterson Lake North Arm - West Basin	1.80E-05	2.06E-05	8.99E-06	2.88E-04	1.08E-06	1.84E-03	2.18E-03				
Patterson Lake South Arm	5.38E-06	6.15E-06	8.77E-06	2.47E-04	1.04E-06	1.77E-03	2.04E-03				
Beet Lake	2.35E-06	2.68E-06	8.60E-06	2.33E-04	1.03E-06	1.76E-03	2.01E-03				
Lloyd Lake	7.18E-07	8.18E-07	8.43E-06	2.24E-04	1.03E-06	1.76E-03	2.00E-03				

Table C.20: Maximum Radiation Dose to Ecological Receptors for the Application Case and Upper Bound Scenario - Far-Future

	Red Fox Total Dose Far Future (mGy/d)										
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
	Application Case										
Reference (Broach Lake)	5.92E-07	6.79E-07	6.84E-07	7.09E-04	1.59E-06	1.55E-05	7.28E-04				
Patterson Lake North Arm - West Basin	8.41E-07	9.64E-07	9.00E-07	1.02E-03	1.59E-06	1.55E-05	1.04E-03				
Patterson Lake South Arm	6.37E-07	7.30E-07	7.20E-07	7.59E-04	1.59E-06	1.55E-05	7.79E-04				
Beet Lake	6.16E-07	7.06E-07	7.05E-07	7.38E-04	1.59E-06	1.55E-05	7.57E-04				
Lloyd Lake	5.93E-07	6.80E-07	6.85E-07	7.10E-04	1.59E-06	1.55E-05	7.29E-04				
		Upp	er Bound Scena	rio							
Reference (Broach Lake)	5.92E-07	6.79E-07	6.84E-07	7.09E-04	1.59E-06	1.55E-05	7.28E-04				
Patterson Lake North Arm - West Basin	8.41E-07	9.64E-07	9.00E-07	1.02E-03	1.59E-06	1.55E-05	1.04E-03				
Patterson Lake South Arm	6.37E-07	7.30E-07	7.20E-07	7.59E-04	1.59E-06	1.55E-05	7.79E-04				
Beet Lake	6.16E-07	7.06E-07	7.05E-07	7.38E-04	1.59E-06	1.55E-05	7.57E-04				
Lloyd Lake	5.93E-07	6.80E-07	6.85E-07	7.10E-04	1.59E-06	1.55E-05	7.29E-04				

	Snowshoe Hare Total Dose Far Future (mGy/d)										
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
	Application Case										
Reference (Broach Lake)	9.84E-06	1.12E-05	1.96E-06	1.26E-03	1.96E-06	1.48E-04	1.44E-03				
Patterson Lake North Arm - West Basin	1.34E-05	1.52E-05	2.59E-06	1.82E-03	1.96E-06	1.48E-04	2.00E-03				
Patterson Lake South Arm	1.04E-05	1.19E-05	2.06E-06	1.35E-03	1.96E-06	1.48E-04	1.53E-03				
Beet Lake	1.02E-05	1.16E-05	2.02E-06	1.32E-03	1.96E-06	1.48E-04	1.49E-03				
Lloyd Lake	9.85E-06	1.12E-05	1.96E-06	1.26E-03	1.96E-06	1.48E-04	1.44E-03				
		Upp	er Bound Scena	rio							
Reference (Broach Lake)	9.84E-06	1.12E-05	1.96E-06	1.26E-03	1.96E-06	1.48E-04	1.44E-03				
Patterson Lake North Arm - West Basin	1.34E-05	1.52E-05	2.59E-06	1.82E-03	1.96E-06	1.48E-04	2.00E-03				
Patterson Lake South Arm	1.04E-05	1.19E-05	2.06E-06	1.35E-03	1.96E-06	1.48E-04	1.53E-03				
Beet Lake	1.02E-05	1.16E-05	2.02E-06	1.32E-03	1.96E-06	1.48E-04	1.49E-03				
Lloyd Lake	9.85E-06	1.12E-05	1.96E-06	1.26E-03	1.96E-06	1.48E-04	1.44E-03				

	Southern Red-Backed Vole Total Dose Far Future (mGy/d)										
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
Application Case											
Reference (Broach Lake)	3.58E-06	4.08E-06	7.10E-07	9.01E-04	1.85E-06	5.26E-05	9.64E-04				
Patterson Lake North Arm - West Basin	4.87E-06	5.55E-06	9.37E-07	1.30E-03	1.85E-06	5.26E-05	1.36E-03				
Patterson Lake South Arm	3.79E-06	4.32E-06	7.47E-07	9.66E-04	1.85E-06	5.26E-05	1.03E-03				
Beet Lake	3.70E-06	4.22E-06	7.31E-07	9.39E-04	1.85E-06	5.26E-05	1.00E-03				
Lloyd Lake	3.58E-06	4.08E-06	7.10E-07	9.03E-04	1.85E-06	5.26E-05	9.65E-04				
		Upp	er Bound Scena	rio							
Reference (Broach Lake)	3.58E-06	4.08E-06	7.10E-07	9.01E-04	1.85E-06	5.26E-05	9.64E-04				
Patterson Lake North Arm - West Basin	4.87E-06	5.55E-06	9.37E-07	1.30E-03	1.85E-06	5.26E-05	1.36E-03				
Patterson Lake South Arm	3.79E-06	4.32E-06	7.47E-07	9.66E-04	1.85E-06	5.26E-05	1.03E-03				
Beet Lake	3.70E-06	4.22E-06	7.31E-07	9.39E-04	1.85E-06	5.26E-05	1.00E-03				
Lloyd Lake	3.58E-06	4.08E-06	7.10E-07	9.03E-04	1.85E-06	5.26E-05	9.65E-04				

	WoodLand Caribou Total Dose Far Future (mGy/d)									
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose			
		F	Application Case							
Reference (Broach Lake)	4.34E-06	4.94E-06	5.85E-06	7.54E-04	6.27E-06	3.51E-03	4.28E-03			
Patterson Lake North Arm - West Basin	1.01E-05	1.16E-05	6.35E-06	9.11E-04	6.31E-06	3.53E-03	4.47E-03			
Beet Lake	4.82E-06	5.48E-06	5.92E-06	7.67E-04	6.29E-06	3.52E-03	4.31E-03			
Lloyd Lake	4.43E-06	5.04E-06	5.86E-06	7.55E-04	6.27E-06	3.51E-03	4.29E-03			
		Upp	er Bound Scena	rio						
Reference (Broach Lake)	4.34E-06	4.94E-06	5.85E-06	7.54E-04	6.27E-06	3.51E-03	4.28E-03			
Patterson Lake North Arm - West Basin	1.01E-05	1.16E-05	6.35E-06	9.11E-04	6.31E-06	3.53E-03	4.47E-03			
Beet Lake	4.82E-06	5.48E-06	5.92E-06	7.67E-04	6.29E-06	3.52E-03	4.31E-03			
Lloyd Lake	4.43E-06	5.04E-06	5.86E-06	7.55E-04	6.27E-06	3.51E-03	4.29E-03			

Table C.20: Maximum Radiation Dose to Ecological Receptors for the Application Case and Upper Bound Scenario - Far-Future

	Grouse Total Dose Far Future (mGy/d)										
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
Application Case											
Reference (Broach Lake)	2.34E-04	2.67E-04	1.70E-06	7.73E-04	1.88E-06	9.49E-04	2.23E-03				
Patterson Lake North Arm - West Basin	3.18E-04	3.62E-04	2.24E-06	1.11E-03	1.88E-06	9.49E-04	2.75E-03				
Patterson Lake South Arm	2.48E-04	2.82E-04	1.79E-06	8.28E-04	1.88E-06	9.49E-04	2.31E-03				
Beet Lake	2.42E-04	2.76E-04	1.75E-06	8.05E-04	1.88E-06	9.49E-04	2.28E-03				
Lloyd Lake	2.35E-04	2.67E-04	1.70E-06	7.74E-04	1.88E-06	9.49E-04	2.23E-03				
		Upp	er Bound Scena	rio							
Reference (Broach Lake)	2.34E-04	2.67E-04	1.70E-06	7.73E-04	1.88E-06	9.49E-04	2.23E-03				
Patterson Lake North Arm - West Basin	3.18E-04	3.62E-04	2.24E-06	1.11E-03	1.88E-06	9.49E-04	2.75E-03				
Patterson Lake South Arm	2.48E-04	2.82E-04	1.79E-06	8.28E-04	1.88E-06	9.49E-04	2.31E-03				
Beet Lake	2.42E-04	2.76E-04	1.75E-06	8.05E-04	1.88E-06	9.49E-04	2.28E-03				
Lloyd Lake	2.35E-04	2.67E-04	1.70E-06	7.74E-04	1.88E-06	9.49E-04	2.23E-03				

	Canada Goose Total Dose Far Future (mGy/d)										
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
	Application Case										
Reference (Broach Lake)	3.28E-05	3.74E-05	4.00E-07	6.64E-04	1.71E-06	1.34E-04	8.70E-04				
Patterson Lake North Arm - West Basin	4.50E-05	5.12E-05	5.28E-07	9.56E-04	1.71E-06	1.34E-04	1.19E-03				
Patterson Lake South Arm	3.49E-05	3.97E-05	4.21E-07	7.11E-04	1.71E-06	1.34E-04	9.22E-04				
Beet Lake	3.40E-05	3.87E-05	4.12E-07	6.91E-04	1.71E-06	1.34E-04	9.00E-04				
Lloyd Lake	3.29E-05	3.74E-05	4.00E-07	6.64E-04	1.71E-06	1.34E-04	8.71E-04				
		Upp	er Bound Scena	rio							
Reference (Broach Lake)	3.28E-05	3.74E-05	4.00E-07	6.64E-04	1.71E-06	1.34E-04	8.70E-04				
Patterson Lake North Arm - West Basin	4.50E-05	5.12E-05	5.28E-07	9.56E-04	1.71E-06	1.34E-04	1.19E-03				
Patterson Lake South Arm	3.49E-05	3.97E-05	4.21E-07	7.11E-04	1.71E-06	1.34E-04	9.22E-04				
Beet Lake	3.40E-05	3.87E-05	4.12E-07	6.91E-04	1.71E-06	1.34E-04	9.00E-04				
Lloyd Lake	3.29E-05	3.74E-05	4.00E-07	6.64E-04	1.71E-06	1.34E-04	8.71E-04				

	Little Brown Myotis Total Dose Far Future (mGy/d)										
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
Application Case											
Reference (Broach Lake)	1.37E-07	1.55E-07	1.19E-05	1.36E-04	4.92E-07	3.09E-03	3.23E-03				
Patterson Lake North Arm - West Basin	4.42E-06	5.05E-06	1.27E-05	1.76E-04	5.15E-07	3.23E-03	3.43E-03				
Patterson Lake South Arm	1.32E-06	1.51E-06	1.24E-05	1.51E-04	4.93E-07	3.10E-03	3.26E-03				
Beet Lake	5.78E-07	6.59E-07	1.22E-05	1.42E-04	4.92E-07	3.09E-03	3.24E-03				
Lloyd Lake	1.77E-07	2.01E-07	1.19E-05	1.37E-04	4.92E-07	3.09E-03	3.24E-03				
		Upp	er Bound Scena	rio							
Reference (Broach Lake)	1.37E-07	1.55E-07	1.19E-05	1.36E-04	4.92E-07	3.09E-03	3.23E-03				
Patterson Lake North Arm - West Basin	4.43E-06	5.06E-06	1.27E-05	1.76E-04	5.15E-07	3.23E-03	3.43E-03				
Patterson Lake South Arm	1.32E-06	1.51E-06	1.24E-05	1.51E-04	4.94E-07	3.10E-03	3.26E-03				
Beet Lake	5.78E-07	6.60E-07	1.22E-05	1.42E-04	4.92E-07	3.09E-03	3.24E-03				
Lloyd Lake	1.77E-07	2.01E-07	1.19E-05	1.37E-04	4.92E-07	3.09E-03	3.24E-03				

		Loon Total	Dose Far Future	e (mGy/d)						
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose			
Application Case										
Reference (Broach Lake)	1.33E-06	1.51E-06	1.01E-06	6.06E-06	8.76E-08	2.97E-03	2.98E-03			
Patterson Lake North Arm - West Basin	4.31E-05	4.92E-05	1.08E-06	7.81E-06	9.16E-08	3.11E-03	3.21E-03			
Patterson Lake South Arm	1.29E-05	1.47E-05	1.05E-06	6.70E-06	8.78E-08	2.98E-03	3.02E-03			
Beet Lake	5.62E-06	6.42E-06	1.03E-06	6.34E-06	8.76E-08	2.97E-03	2.99E-03			
Lloyd Lake	1.72E-06	1.96E-06	1.01E-06	6.09E-06	8.76E-08	2.97E-03	2.98E-03			
		Upp	er Bound Scena	rio						
Reference (Broach Lake)	1.33E-06	1.51E-06	1.01E-06	6.06E-06	8.76E-08	2.97E-03	2.98E-03			
Patterson Lake North Arm - West Basin	4.31E-05	4.92E-05	1.08E-06	7.82E-06	9.16E-08	3.11E-03	3.21E-03			
Patterson Lake South Arm	1.29E-05	1.47E-05	1.05E-06	6.71E-06	8.78E-08	2.98E-03	3.02E-03			
Beet Lake	5.63E-06	6.42E-06	1.03E-06	6.34E-06	8.76E-08	2.97E-03	2.99E-03			
Lloyd Lake	1.72E-06	1.96E-06	1.01E-06	6.09E-06	8.76E-08	2.97E-03	2.98E-03			

Table C.20: Maximum Radiation Dose to Ecological Receptors for the Application Case and Upper Bound Scenario - Far-Future

	Mallard Total Dose Far Future (mGy/d)										
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
Application Case											
Reference (Broach Lake)	1.40E-05	1.59E-05	9.79E-06	7.84E-05	1.23E-06	2.55E-02	2.56E-02				
Patterson Lake North Arm - West Basin	4.53E-04	5.17E-04	1.05E-05	1.01E-04	1.29E-06	2.66E-02	2.77E-02				
Patterson Lake South Arm	1.35E-04	1.55E-04	1.02E-05	8.68E-05	1.23E-06	2.55E-02	2.59E-02				
Beet Lake	5.91E-05	6.75E-05	1.00E-05	8.20E-05	1.23E-06	2.55E-02	2.57E-02				
Lloyd Lake	1.81E-05	2.06E-05	9.82E-06	7.88E-05	1.23E-06	2.55E-02	2.56E-02				
		Upp	er Bound Scena	rio							
Reference (Broach Lake)	1.40E-05	1.59E-05	9.79E-06	7.84E-05	1.23E-06	2.55E-02	2.56E-02				
Patterson Lake North Arm - West Basin	4.53E-04	5.17E-04	1.05E-05	1.01E-04	1.29E-06	2.66E-02	2.77E-02				
Patterson Lake South Arm	1.35E-04	1.55E-04	1.02E-05	8.68E-05	1.23E-06	2.55E-02	2.59E-02				
Beet Lake	5.91E-05	6.75E-05	1.00E-05	8.20E-05	1.23E-06	2.55E-02	2.57E-02				
Lloyd Lake	1.81E-05	2.06E-05	9.82E-06	7.88E-05	1.23E-06	2.55E-02	2.56E-02				

	Rusty Blackbird Total Dose Far Future (mGy/d)										
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
	Application Case										
Reference (Broach Lake)	1.01E-04	1.15E-04	2.74E-05	8.28E-04	3.65E-06	6.96E-02	7.07E-02				
Patterson Lake North Arm - West Basin	5.15E-04	5.88E-04	3.02E-05	1.18E-03	3.73E-06	7.28E-02	7.52E-02				
Patterson Lake South Arm	2.12E-04	2.42E-04	2.87E-05	8.91E-04	3.65E-06	6.99E-02	7.12E-02				
Beet Lake	1.43E-04	1.63E-04	2.81E-05	8.63E-04	3.65E-06	6.97E-02	7.09E-02				
Lloyd Lake	1.04E-04	1.19E-04	2.75E-05	8.29E-04	3.65E-06	6.96E-02	7.07E-02				
		Upp	er Bound Scena	rio							
Reference (Broach Lake)	1.01E-04	1.15E-04	2.74E-05	8.28E-04	3.65E-06	6.96E-02	7.07E-02				
Patterson Lake North Arm - West Basin	5.16E-04	5.89E-04	3.02E-05	1.18E-03	3.73E-06	7.29E-02	7.52E-02				
Patterson Lake South Arm	2.12E-04	2.42E-04	2.87E-05	8.91E-04	3.65E-06	6.99E-02	7.12E-02				
Beet Lake	1.43E-04	1.63E-04	2.81E-05	8.63E-04	3.65E-06	6.97E-02	7.09E-02				
Lloyd Lake	1.04E-04	1.19E-04	2.75E-05	8.29E-04	3.65E-06	6.96E-02	7.07E-02				

Table C.21: Maximum Radiation Dose to Ecological Receptors for the RFD Scenario - Project Lifespan and Far-Future

		Benthic Inverteb	rates Total Dose	RFD Case (mGy	/d)						
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
	Application Case										
Reference (Broach Lake) 4.73E-09 6.18E-09 1.80E-08 8.07E-05 3.79E-05 1.72E-09 1.19E-04											
Patterson Lake North Arm - West Basin	4.26E-08	5.58E-08	1.00E-07	1.30E-04	5.19E-05	2.35E-09	1.82E-04				
Patterson Lake South Arm	2.65E-08	3.47E-08	7.73E-08	1.10E-04	4.65E-05	2.11E-09	1.57E-04				
Beet Lake	1.28E-08	1.67E-08	4.96E-08	9.35E-05	3.92E-05	1.78E-09	1.33E-04				
Lloyd Lake	5.47E-09	7.14E-09	2.13E-08	8.20E-05	3.80E-05	1.72E-09	1.20E-04				
		Up	oper Bound Scen	ario							
Reference (Broach Lake)	4.73E-09	6.18E-09	1.80E-08	8.07E-05	3.79E-05	1.72E-09	1.19E-04				
Patterson Lake North Arm - West Basin	1.53E-07	2.01E-07	1.93E-08	1.04E-04	3.97E-05	1.80E-09	1.44E-04				
Patterson Lake South Arm	4.58E-08	6.00E-08	1.88E-08	8.93E-05	3.80E-05	1.72E-09	1.28E-04				
Beet Lake	2.00E-08	2.62E-08	1.84E-08	8.44E-05	3.79E-05	1.72E-09	1.22E-04				
Lloyd Lake	6.11E-09	7.99E-09	1.81E-08	8.11E-05	3.79E-05	1.72E-09	1.19E-04				

		Zooplanktor	1 Total Dose RFD	Case (mGy/d)						
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose			
			Application Cas	e						
Reference (Broach Lake) 2.05E-05 2.39E-05 2.94E-03 4.57E-03 2.80E-05 6.05E-02 6.81E-02										
Patterson Lake North Arm - West Basin	2.23E-04	2.62E-04	1.67E-02	7.95E-03	4.28E-05	9.18E-02	1.17E-01			
Patterson Lake South Arm	3.33E-04	3.90E-04	1.78E-02	9.33E-03	4.35E-05	9.27E-02	1.21E-01			
Beet Lake	1.29E-04	1.51E-04	9.64E-03	6.40E-03	3.02E-05	6.52E-02	8.16E-02			
Lloyd Lake	3.06E-05	3.58E-05	3.63E-03	4.75E-03	2.81E-05	6.08E-02	6.92E-02			
		Up	per Bound Scen	ario						
Reference (Broach Lake)	2.05E-05	2.39E-05	2.94E-03	4.57E-03	2.80E-05	6.05E-02	6.81E-02			
Patterson Lake North Arm - West Basin	6.63E-04	7.77E-04	3.15E-03	5.89E-03	2.93E-05	6.33E-02	7.38E-02			
Patterson Lake South Arm	1.98E-04	2.32E-04	3.07E-03	5.05E-03	2.81E-05	6.07E-02	6.93E-02			
Beet Lake	8.66E-05	1.01E-04	3.01E-03	4.78E-03	2.80E-05	6.05E-02	6.85E-02			
Lloyd Lake	2.65E-05	3.09E-05	2.95E-03	4.59E-03	2.80E-05	6.05E-02	6.81E-02			

		Northern Pik	e Total Dose RFD	Case (mGy/d)							
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
			Application Cas	e							
Reference (Broach Lake) 4.13E-06 4.70E-06 3.54E-05 2.22E-04 1.79E-06 5.64E-04 8.33E-04											
Patterson Lake North Arm - West Basin	4.50E-05	5.14E-05	2.02E-04	3.87E-04	2.74E-06	8.56E-04	1.54E-03				
Patterson Lake South Arm	6.71E-05	7.67E-05	2.15E-04	4.54E-04	2.79E-06	8.65E-04	1.68E-03				
Beet Lake	2.60E-05	2.97E-05	1.16E-04	3.11E-04	1.94E-06	6.08E-04	1.09E-03				
Lloyd Lake	6.16E-06	7.02E-06	4.36E-05	2.31E-04	1.80E-06	5.67E-04	8.56E-04				
		Up	pper Bound Scen	ario							
Reference (Broach Lake)	4.13E-06	4.70E-06	3.54E-05	2.22E-04	1.79E-06	5.64E-04	8.33E-04				
Patterson Lake North Arm - West Basin	1.34E-04	1.53E-04	3.79E-05	2.87E-04	1.88E-06	5.90E-04	1.20E-03				
Patterson Lake South Arm	4.00E-05	4.56E-05	3.69E-05	2.46E-04	1.80E-06	5.66E-04	9.36E-04				
Beet Lake	1.74E-05	1.99E-05	3.62E-05	2.32E-04	1.79E-06	5.65E-04	8.72E-04				
Lloyd Lake	5.33E-06	6.07E-06	3.55E-05	2.23E-04	1.79E-06	5.64E-04	8.36E-04				

		Whitefish 1	Total Dose RFD C	ase (mGy/d)							
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
	Application Case										
Reference (Broach Lake) 7.69E-07 8.97E-07 2.65E-05 1.86E-04 8.29E-07 4.31E-05 2.58E-04											
Patterson Lake North Arm - West Basin	8.38E-06	9.82E-06	1.51E-04	3.21E-04	1.22E-06	6.55E-05	5.57E-04				
Patterson Lake South Arm	1.25E-05	1.46E-05	1.60E-04	3.69E-04	1.18E-06	6.61E-05	6.24E-04				
Beet Lake	4.84E-06	5.67E-06	8.68E-05	2.57E-04	8.81E-07	4.65E-05	4.01E-04				
Lloyd Lake	1.15E-06	1.34E-06	3.26E-05	1.92E-04	8.32E-07	4.33E-05	2.72E-04				
		U _l	oper Bound Scen	ario							
Reference (Broach Lake)	7.69E-07	8.97E-07	2.65E-05	1.86E-04	8.29E-07	4.31E-05	2.58E-04				
Patterson Lake North Arm - West Basin	2.49E-05	2.91E-05	2.83E-05	2.39E-04	8.68E-07	4.51E-05	3.68E-04				
Patterson Lake South Arm	7.44E-06	8.71E-06	2.76E-05	2.05E-04	8.32E-07	4.33E-05	2.93E-04				
Beet Lake	3.25E-06	3.80E-06	2.71E-05	1.94E-04	8.30E-07	4.32E-05	2.72E-04				
Lloyd Lake	9.93E-07	1.16E-06	2.66E-05	1.86E-04	8.29E-07	4.31E-05	2.59E-04				

Table C.21: Maximum Radiation Dose to Ecological Receptors for the RFD Scenario - Project Lifespan and Far-Future

		Macrophytes	s Total Dose RFD	Case (mGy/d)							
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
Application Case											
Reference (Broach Lake) 1.04E-05 1.18E-05 1.37E-03 2.99E-03 5.06E-05 1.10E-03 5.53E-03											
Patterson Lake North Arm - West Basin	1.13E-04	1.30E-04	7.79E-03	5.20E-03	7.73E-05	1.67E-03	1.50E-02				
Patterson Lake South Arm	1.69E-04	1.93E-04	8.30E-03	6.10E-03	7.85E-05	1.69E-03	1.65E-02				
Beet Lake	6.56E-05	7.49E-05	4.49E-03	4.19E-03	5.47E-05	1.19E-03	1.01E-02				
Lloyd Lake	1.55E-05	1.77E-05	1.69E-03	3.10E-03	5.09E-05	1.11E-03	5.98E-03				
		U _l	pper Bound Scen	ario							
Reference (Broach Lake)	1.04E-05	1.18E-05	1.37E-03	2.99E-03	5.06E-05	1.10E-03	5.53E-03				
Patterson Lake North Arm - West Basin	3.37E-04	3.85E-04	1.47E-03	3.85E-03	5.30E-05	1.15E-03	7.25E-03				
Patterson Lake South Arm	1.01E-04	1.15E-04	1.43E-03	3.31E-03	5.08E-05	1.10E-03	6.11E-03				
Beet Lake	4.40E-05	5.02E-05	1.40E-03	3.13E-03	5.07E-05	1.10E-03	5.77E-03				
Lloyd Lake	1.34E-05	1.53E-05	1.37E-03	3.00E-03	5.06E-05	1.10E-03	5.56E-03				

		Phytoplankto	n Total Dose RFI	O Case (mGy/d)							
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
			Application Case	e							
Reference (Broach Lake) 1.04E-05 1.18E-05 1.36E-03 3.00E-03 1.93E-05 1.11E-03 5.51E-03											
Patterson Lake North Arm - West Basin	1.13E-04	1.29E-04	7.75E-03	5.21E-03	2.96E-05	1.68E-03	1.49E-02				
Patterson Lake South Arm	1.69E-04	1.93E-04	8.26E-03	6.12E-03	3.01E-05	1.69E-03	1.65E-02				
Beet Lake	6.55E-05	7.47E-05	4.46E-03	4.20E-03	2.09E-05	1.19E-03	1.00E-02				
Lloyd Lake	1.55E-05	1.77E-05	1.68E-03	3.11E-03	1.94E-05	1.11E-03	5.95E-03				
		U _l	pper Bound Scen	ario							
Reference (Broach Lake)	1.04E-05	1.18E-05	1.36E-03	3.00E-03	1.93E-05	1.11E-03	5.51E-03				
Patterson Lake North Arm - West Basin	3.37E-04	3.84E-04	1.46E-03	3.86E-03	2.02E-05	1.16E-03	7.22E-03				
Patterson Lake South Arm	1.01E-04	1.15E-04	1.42E-03	3.32E-03	1.94E-05	1.11E-03	6.08E-03				
Beet Lake	4.39E-05	5.01E-05	1.39E-03	3.13E-03	1.94E-05	1.11E-03	5.75E-03				
Lloyd Lake	1.34E-05	1.53E-05	1.37E-03	3.01E-03	1.93E-05	1.11E-03	5.53E-03				

	T	errestrial Inverte	ebrate Total Dose	RFD Case (mG	y/d)					
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose			
			Application Cas	e						
Reference (Broach Lake) 1.57E-05 1.78E-05 1.15E-04 5.73E-03 6.72E-06 2.28E-05 5.91E-03										
Patterson Lake North Arm - West Basin	6.97E-05	7.93E-05	1.85E-03	1.81E-02	2.19E-05	1.80E-03	2.19E-02			
Patterson Lake South Arm	2.61E-05	2.97E-05	4.50E-04	8.11E-03	9.63E-06	3.64E-04	8.99E-03			
Beet Lake	2.14E-05	2.43E-05	2.98E-04	7.03E-03	8.31E-06	2.10E-04	7.59E-03			
Lloyd Lake	1.59E-05	1.81E-05	1.24E-04	5.80E-03	6.79E-06	3.17E-05	5.99E-03			
		U _l	per Bound Scen	ario						
Reference (Broach Lake)	1.57E-05	1.78E-05	1.15E-04	5.73E-03	6.72E-06	2.28E-05	5.91E-03			
Patterson Lake North Arm - West Basin	2.14E-05	2.43E-05	1.53E-04	8.31E-03	6.72E-06	2.28E-05	8.54E-03			
Patterson Lake South Arm	1.68E-05	1.91E-05	1.22E-04	6.23E-03	6.72E-06	2.28E-05	6.42E-03			
Beet Lake	1.63E-05	1.85E-05	1.19E-04	6.01E-03	6.72E-06	2.28E-05	6.19E-03			
Lloyd Lake	1.57E-05	1.79E-05	1.15E-04	5.75E-03	6.72E-06	2.28E-05	5.93E-03			

	Browse Total Dose RFD Case (mGy/d)										
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
			Application Case	e							
Patterson Lake North Arm - West Basin 2.57E-03 2.92E-03 1.83E-03 2.18E-02 2.11E-04 2.83E-03 3.22E-02											
Patterson Lake South Arm	9.61E-04	1.09E-03	4.36E-04	1.06E-02	1.74E-04	1.06E-03	1.44E-02				
		U _l	pper Bound Scen	ario							
Patterson Lake North Arm - West Basin 7.87E-04 8.96E-04 1.37E-04 1.16E-02 1.65E-04 6.33E-04 1.42E-02											
Patterson Lake South Arm	6.18E-04	7.03E-04	1.10E-04	8.67E-03	1.65E-04	6.33E-04	1.09E-02				

Table C.21: Maximum Radiation Dose to Ecological Receptors for the RFD Scenario - Project Lifespan and Far-Future

	Labrador Tea Total Dose RFD Case (mGy/d)									
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose			
			Application Cas	e						
Patterson Lake North Arm - West Basin 2.57E-03 2.92E-03 1.61E-02 9.42E-02 3.29E-04 1.64E-02 1.32E-01										
Patterson Lake South Arm	9.61E-04	1.09E-03	3.18E-03	2.46E-02	1.97E-04	3.67E-03	3.37E-02			
		Up	pper Bound Scen	ario						
Patterson Lake North Arm - West Basin 7.87E-04 8.96E-04 1.37E-04 1.16E-02 1.65E-04 6.33E-04 1.42E-02										
Patterson Lake South Arm	6.18E-04	7.03E-04	1.10E-04	8.67E-03	1.65E-04	6.33E-04	1.09E-02			

	Lichen Total Dose Operations (mGy/d)									
Location	cation Uranium-238 Uranium-234 Thorium-230 Radium-226 Lead-210 Polonium-210 Total Dos									
			Application Cas	e						
Patterson Lake North Arm - West Basin 1.21E-04 1.37E-04 9.12E-02 4.70E-01 1.41E-03 1.26E-01 6.90E-01										
Patterson Lake South Arm	1.21E-04	1.37E-04	1.79E-02	9.42E-02	9.45E-04	1.10E-01	2.23E-01			
		Up	oper Bound Scen	ario						
Patterson Lake North Arm - West Basin 1.21E-04 1.37E-04 4.38E-04 4.44E-03 8.33E-04 1.06E-01 1.12E-01										
Patterson Lake South Arm	1.21E-04	1.37E-04	4.38E-04	4.44E-03	8.33E-04	1.06E-01	1.12E-01			

	Blueberries Total Dose RFD Case (mGy/d)									
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose			
			Application Cas	e						
Patterson Lake North Arm - West Basin	1.33E-03	1.51E-03	7.01E-05	9.25E-03	8.51E-06	1.08E-03	1.32E-02			
Patterson Lake South Arm	4.87E-04	5.54E-04	4.64E-05	4.91E-03	4.06E-06	4.63E-04	6.47E-03			
		Up	pper Bound Scen	ario						
Patterson Lake North Arm - West Basin	3.91E-04	4.45E-04	5.42E-05	5.62E-03	3.00E-06	3.16E-04	6.83E-03			
Patterson Lake South Arm	3.07E-04	3.50E-04	4.33E-05	4.22E-03	3.00E-06	3.16E-04	5.23E-03			

		Beaver To	otal Dose RFD Ca	se (mGy/d)							
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose				
Application Case											
Reference (Broach Lake)	1.05E-05	1.19E-05	3.11E-06	6.36E-04	4.75E-07	2.15E-04	8.77E-04				
Patterson Lake North Arm - West Basin	4.69E-05	5.34E-05	2.97E-05	1.74E-03	1.14E-06	7.50E-04	2.62E-03				
Patterson Lake South Arm	1.79E-05	2.04E-05	1.52E-05	8.78E-04	7.12E-07	3.35E-04	1.27E-03				
Beet Lake	1.44E-05	1.64E-05	8.93E-06	7.63E-04	5.49E-07	2.72E-04	1.07E-03				
Lloyd Lake	1.07E-05	1.21E-05	3.60E-06	6.43E-04	4.79E-07	2.17E-04	8.87E-04				
		Uį	oper Bound Scen	ario							
Reference (Broach Lake)	1.05E-05	1.19E-05	3.11E-06	6.36E-04	4.75E-07	2.15E-04	8.77E-04				
Patterson Lake North Arm - West Basin	1.62E-05	1.84E-05	3.69E-06	9.14E-04	4.90E-07	2.18E-04	1.17E-03				
Patterson Lake South Arm	1.17E-05	1.34E-05	3.27E-06	6.92E-04	4.76E-07	2.15E-04	9.36E-04				
Beet Lake	1.11E-05	1.26E-05	3.20E-06	6.66E-04	4.75E-07	2.15E-04	9.08E-04				
Lloyd Lake	1.05E-05	1.20E-05	3.12E-06	6.38E-04	4.75E-07	2.15E-04	8.79E-04				

Black Bear Total Dose RFD Case (mGy/d)								
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose	
			Application Cas	e				
Reference (Broach Lake)	7.81E-06	8.89E-06	1.88E-06	8.19E-04	8.32E-07	1.74E-04	1.01E-03	
Patterson Lake North Arm - West Basin	2.32E-05	2.64E-05	3.30E-06	1.29E-03	1.05E-06	3.42E-04	1.68E-03	
Beet Lake	9.48E-06	1.08E-05	2.78E-06	1.09E-03	1.23E-06	1.98E-04	1.31E-03	
Lloyd Lake	7.97E-06	9.07E-06	1.96E-06	8.25E-04	8.35E-07	1.76E-04	1.02E-03	
		U _l	oper Bound Scen	ario				
Reference (Broach Lake)	7.81E-06	8.89E-06	1.88E-06	8.19E-04	8.32E-07	1.74E-04	1.01E-03	
Patterson Lake North Arm - West Basin	9.93E-06	1.13E-05	2.22E-06	1.02E-03	8.33E-07	1.76E-04	1.22E-03	
Beet Lake	8.14E-06	9.26E-06	2.53E-06	1.06E-03	1.21E-06	1.82E-04	1.26E-03	
Lloyd Lake	7.84E-06	8.92E-06	1.91E-06	8.22E-04	8.33E-07	1.75E-04	1.02E-03	

Table C.21: Maximum Radiation Dose to Ecological Receptors for the RFD Scenario - Project Lifespan and Far-Future

Grey Wolf Total Dose RFD Case (mGy/d)										
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose			
	Application Case									
Reference (Broach Lake)	3.26E-07	3.77E-07	1.17E-06	7.63E-04	1.92E-06	3.26E-04	1.09E-03			
Patterson Lake North Arm - West Basin	6.35E-07	7.34E-07	1.58E-06	9.25E-04	2.02E-06	3.60E-04	1.29E-03			
Lloyd Lake	3.32E-07	3.85E-07	1.19E-06	7.66E-04	1.92E-06	3.27E-04	1.10E-03			
Upper Bound Scenario										
Reference (Broach Lake)	3.26E-07	3.77E-07	1.17E-06	7.63E-04	1.92E-06	3.26E-04	1.09E-03			
Patterson Lake North Arm - West Basin	3.91E-07	4.52E-07	1.27E-06	8.48E-04	1.92E-06	3.27E-04	1.18E-03			
Lloyd Lake	3.27E-07	3.79E-07	1.18E-06	7.65E-04	1.92E-06	3.26E-04	1.09E-03			

Mink Total Dose RFD Case (mGy/d)									
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose		
Application Case									
Reference (Broach Lake)	1.76E-07	2.00E-07	2.51E-06	6.44E-05	2.38E-07	8.73E-04	9.40E-04		
Patterson Lake North Arm - West Basin	1.16E-06	1.32E-06	1.23E-05	1.04E-04	3.23E-07	1.20E-03	1.32E-03		
Patterson Lake South Arm	6.81E-07	7.77E-07	9.46E-06	8.47E-05	2.90E-07	1.07E-03	1.17E-03		
Beet Lake	3.67E-07	4.18E-07	6.20E-06	7.33E-05	2.46E-07	9.02E-04	9.82E-04		
Lloyd Lake	1.91E-07	2.18E-07	2.89E-06	6.52E-05	2.38E-07	8.74E-04	9.43E-04		
		Up	oper Bound Scen	ario					
Reference (Broach Lake)	1.76E-07	2.00E-07	2.51E-06	6.44E-05	2.38E-07	8.73E-04	9.40E-04		
Patterson Lake North Arm - West Basin	2.66E-06	3.04E-06	2.79E-06	8.53E-05	2.48E-07	9.13E-04	1.01E-03		
Patterson Lake South Arm	8.60E-07	9.83E-07	2.62E-06	7.09E-05	2.38E-07	8.75E-04	9.51E-04		
Beet Lake	4.31E-07	4.92E-07	2.57E-06	6.73E-05	2.38E-07	8.73E-04	9.44E-04		
Lloyd Lake	1.99E-07	2.26E-07	2.51E-06	6.46E-05	2.38E-07	8.73E-04	9.40E-04		

Moose Total Dose RFD Case (mGy/d)									
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose		
Application Case									
Reference (Broach Lake)	9.50E-06	1.08E-05	4.80E-06	9.61E-04	1.30E-06	2.66E-04	1.25E-03		
Patterson Lake North Arm - West Basin	4.29E-05	4.89E-05	3.78E-05	2.21E-03	2.15E-06	7.80E-04	3.12E-03		
Patterson Lake South Arm	4.26E-05	4.85E-05	3.72E-05	2.00E-03	2.04E-06	7.75E-04	2.90E-03		
Beet Lake	1.33E-05	1.51E-05	1.44E-05	1.11E-03	1.40E-06	3.21E-04	1.48E-03		
Lloyd Lake	9.70E-06	1.10E-05	5.68E-06	9.70E-04	1.30E-06	2.69E-04	1.27E-03		
		U _l	pper Bound Scen	ario					
Reference (Broach Lake)	9.50E-06	1.08E-05	4.80E-06	9.61E-04	1.30E-06	2.66E-04	1.25E-03		
Patterson Lake North Arm - West Basin	1.69E-05	1.93E-05	5.48E-06	1.38E-03	1.32E-06	2.73E-04	1.70E-03		
Patterson Lake South Arm	1.68E-05	1.91E-05	5.29E-06	1.24E-03	1.32E-06	2.72E-04	1.56E-03		
Beet Lake	1.03E-05	1.17E-05	4.92E-06	1.01E-03	1.30E-06	2.67E-04	1.30E-03		
Lloyd Lake	9.56E-06	1.09E-05	4.81E-06	9.64E-04	1.30E-06	2.66E-04	1.26E-03		

Muskrat Total Dose RFD Case (mGy/d)									
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose		
Application Case									
Reference (Broach Lake)	5.56E-07	6.32E-07	8.41E-06	2.23E-04	1.03E-06	1.76E-03	2.00E-03		
Patterson Lake North Arm - West Basin	5.07E-06	5.79E-06	4.71E-05	3.63E-04	1.46E-06	2.42E-03	2.84E-03		
Patterson Lake South Arm	3.45E-06	3.94E-06	4.03E-05	3.31E-04	1.36E-06	2.17E-03	2.55E-03		
Beet Lake	1.62E-06	1.84E-06	2.44E-05	2.68E-04	1.08E-06	1.82E-03	2.12E-03		
Lloyd Lake	6.53E-07	7.44E-07	1.00E-05	2.27E-04	1.03E-06	1.77E-03	2.01E-03		
		Up	per Bound Scen	ario					
Reference (Broach Lake)	0.00E+00	3.32E-03	6.32E-07	8.39E-06	1.76E-03	2.23E-04	5.31E-03		
Patterson Lake North Arm - West Basin	0.00E+00	3.76E-03	2.05E-05	9.01E-06	1.84E-03	2.87E-04	5.92E-03		
Patterson Lake South Arm	0.00E+00	3.40E-03	6.14E-06	8.76E-06	1.77E-03	2.46E-04	5.43E-03		
Beet Lake	0.00E+00	3.35E-03	2.68E-06	8.59E-06	1.76E-03	2.33E-04	5.36E-03		
Lloyd Lake	0.00E+00	3.32E-03	8.18E-07	8.41E-06	1.76E-03	2.24E-04	5.31E-03		

Table C.21: Maximum Radiation Dose to Ecological Receptors for the RFD Scenario - Project Lifespan and Far-Future

		Red Fox T	otal Dose RFD Ca	ase (mGy/d)					
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose		
Application Case									
Reference (Broach Lake)	5.92E-07	6.79E-07	6.84E-07	7.09E-04	1.59E-06	1.55E-05	7.28E-04		
Patterson Lake North Arm - West Basin	2.71E-06	3.11E-06	2.89E-06	1.26E-03	1.95E-06	8.19E-05	1.36E-03		
Patterson Lake South Arm	1.02E-06	1.17E-06	1.19E-06	8.16E-04	1.66E-06	2.84E-05	8.49E-04		
Beet Lake	8.20E-07	9.40E-07	9.53E-07	7.67E-04	1.63E-06	2.25E-05	7.94E-04		
Lloyd Lake	6.03E-07	6.92E-07	6.99E-07	7.11E-04	1.59E-06	1.58E-05	7.31E-04		
		Up	oper Bound Scen	ario					
Reference (Broach Lake)	5.92E-07	6.79E-07	6.84E-07	7.09E-04	1.59E-06	1.55E-05	7.28E-04		
Patterson Lake North Arm - West Basin	8.45E-07	9.69E-07	9.04E-07	1.03E-03	1.59E-06	1.55E-05	1.05E-03		
Patterson Lake South Arm	6.44E-07	7.38E-07	7.27E-07	7.70E-04	1.59E-06	1.55E-05	7.89E-04		
Beet Lake	6.18E-07	7.09E-07	7.08E-07	7.42E-04	1.59E-06	1.55E-05	7.61E-04		
Lloyd Lake	5.93E-07	6.81E-07	6.85E-07	7.10E-04	1.59E-06	1.55E-05	7.29E-04		

		Snowshoe Ha	re Total Dose RF	D Case (mGy/d)	ı.					
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose			
	Application Case									
Reference (Broach Lake)	9.84E-06	1.12E-05	1.96E-06	1.26E-03	1.96E-06	1.48E-04	1.44E-03			
Patterson Lake North Arm - West Basin	4.38E-05	4.98E-05	1.85E-05	2.72E-03	2.78E-06	6.15E-04	3.45E-03			
Patterson Lake South Arm	1.64E-05	1.87E-05	5.21E-06	1.54E-03	2.12E-06	2.38E-04	1.82E-03			
Beet Lake	1.34E-05	1.53E-05	3.73E-06	1.42E-03	2.05E-06	1.98E-04	1.65E-03			
Lloyd Lake	1.00E-05	1.14E-05	2.05E-06	1.27E-03	1.97E-06	1.51E-04	1.45E-03			
		U _l	per Bound Scen	ario						
Reference (Broach Lake)	9.84E-06	1.12E-05	1.96E-06	1.26E-03	1.96E-06	1.48E-04	1.44E-03			
Patterson Lake North Arm - West Basin	1.34E-05	1.53E-05	2.60E-06	1.83E-03	1.96E-06	1.48E-04	2.01E-03			
Patterson Lake South Arm	1.05E-05	1.20E-05	2.08E-06	1.37E-03	1.96E-06	1.48E-04	1.55E-03			
Beet Lake	1.02E-05	1.16E-05	2.03E-06	1.32E-03	1.96E-06	1.48E-04	1.50E-03			
Lloyd Lake	9.86E-06	1.12E-05	1.96E-06	1.27E-03	1.96E-06	1.48E-04	1.44E-03			

	So	uthern Red-Back	ed Vole Total Do	se RFD Case (m	Gy/d)		
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose
			Application Cas	e			
Reference (Broach Lake)	3.58E-06	4.08E-06	7.10E-07	9.01E-04	1.85E-06	5.26E-05	9.64E-04
Patterson Lake North Arm - West Basin	1.62E-05	1.84E-05	4.13E-06	1.69E-03	2.34E-06	1.98E-04	1.93E-03
Patterson Lake South Arm	6.01E-06	6.85E-06	1.41E-06	1.05E-03	1.95E-06	8.07E-05	1.15E-03
Beet Lake	4.90E-06	5.59E-06	1.09E-06	9.84E-04	1.90E-06	6.79E-05	1.07E-03
Lloyd Lake	3.64E-06	4.15E-06	7.29E-07	9.05E-04	1.86E-06	5.33E-05	9.69E-04
		Up	pper Bound Scen	ario			
Reference (Broach Lake)	3.58E-06	4.08E-06	7.10E-07	9.01E-04	1.85E-06	5.26E-05	9.64E-04
Patterson Lake North Arm - West Basin	4.89E-06	5.58E-06	9.41E-07	1.31E-03	1.85E-06	5.26E-05	1.37E-03
Patterson Lake South Arm	3.83E-06	4.37E-06	7.54E-07	9.79E-04	1.85E-06	5.26E-05	1.04E-03
Beet Lake	3.72E-06	4.24E-06	7.34E-07	9.44E-04	1.85E-06	5.26E-05	1.01E-03
Lloyd Lake	3.59E-06	4.09E-06	7.11E-07	9.03E-04	1.85E-06	5.26E-05	9.66E-04

		WoodLand Cari	bou Total Dose F	RFD Case (mGy/	d)		
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose
			Application Case	e			
Reference (Broach Lake)	4.34E-06	4.94E-06	5.85E-06	7.54E-04	6.27E-06	3.51E-03	4.28E-03
Patterson Lake North Arm - West Basin	1.31E-05	1.49E-05	9.13E-05	3.75E-03	8.51E-06	4.01E-03	7.89E-03
Beet Lake	5.27E-06	6.00E-06	1.63E-05	1.04E-03	6.50E-06	3.57E-03	4.65E-03
Lloyd Lake	4.46E-06	5.08E-06	7.29E-06	7.84E-04	6.29E-06	3.51E-03	4.32E-03
		Up	pper Bound Scen	ario			
Reference (Broach Lake)	4.34E-06	4.94E-06	5.85E-06	7.54E-04	6.27E-06	3.51E-03	4.28E-03
Patterson Lake North Arm - West Basin	1.02E-05	1.16E-05	6.36E-06	9.17E-04	6.31E-06	3.53E-03	4.48E-03
Beet Lake	4.82E-06	5.49E-06	5.92E-06	7.69E-04	6.29E-06	3.52E-03	4.32E-03
Lloyd Lake	4.43E-06	5.04E-06	5.87E-06	7.56E-04	6.27E-06	3.51E-03	4.29E-03

Table C.21: Maximum Radiation Dose to Ecological Receptors for the RFD Scenario - Project Lifespan and Far-Future

		Grouse To	otal Dose RFD Ca	se (mGy/d)			
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose
			Application Cas	e			
Reference (Broach Lake)	2.34E-04	2.67E-04	1.70E-06	7.73E-04	1.88E-06	9.49E-04	2.23E-03
Patterson Lake North Arm - West Basin	1.05E-03	1.19E-03	9.35E-06	1.41E-03	2.60E-06	3.58E-03	7.24E-03
Patterson Lake South Arm	3.91E-04	4.45E-04	3.20E-06	8.95E-04	2.02E-06	1.46E-03	3.19E-03
Beet Lake	3.20E-04	3.64E-04	2.51E-06	8.40E-04	1.95E-06	1.23E-03	2.75E-03
Lloyd Lake	2.38E-04	2.71E-04	1.74E-06	7.76E-04	1.88E-06	9.62E-04	2.25E-03
		Up	oper Bound Scen	ario			
Reference (Broach Lake)	2.34E-04	2.67E-04	1.70E-06	7.73E-04	1.88E-06	9.49E-04	2.23E-03
Patterson Lake North Arm - West Basin	3.20E-04	3.64E-04	2.25E-06	1.12E-03	1.88E-06	9.49E-04	2.76E-03
Patterson Lake South Arm	2.51E-04	2.85E-04	1.80E-06	8.39E-04	1.88E-06	9.49E-04	2.33E-03
Beet Lake	2.43E-04	2.77E-04	1.76E-06	8.09E-04	1.88E-06	9.49E-04	2.28E-03
Lloyd Lake	2.35E-04	2.67E-04	1.70E-06	7.74E-04	1.88E-06	9.49E-04	2.23E-03

		Canada Goos	e Total Dose RFD	Case (mGy/d)			
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose
			Application Cas	e			
Reference (Broach Lake)	3.28E-05	3.74E-05	4.00E-07	6.64E-04	1.71E-06	1.34E-04	8.70E-04
Patterson Lake North Arm - West Basin	1.46E-04	1.66E-04	1.87E-06	1.12E-03	2.09E-06	5.31E-04	1.97E-03
Patterson Lake South Arm	5.48E-05	6.24E-05	7.05E-07	7.52E-04	1.78E-06	2.11E-04	1.08E-03
Beet Lake	4.48E-05	5.10E-05	5.64E-07	7.12E-04	1.75E-06	1.76E-04	9.86E-04
Lloyd Lake	3.34E-05	3.80E-05	4.08E-07	6.66E-04	1.71E-06	1.36E-04	8.75E-04
		U _l	per Bound Scen	ario			
Reference (Broach Lake)	3.28E-05	3.74E-05	4.00E-07	6.64E-04	1.71E-06	1.34E-04	8.70E-04
Patterson Lake North Arm - West Basin	4.52E-05	5.14E-05	5.30E-07	9.61E-04	1.71E-06	1.34E-04	1.19E-03
Patterson Lake South Arm	3.52E-05	4.01E-05	4.25E-07	7.21E-04	1.71E-06	1.34E-04	9.32E-04
Beet Lake	3.41E-05	3.88E-05	4.14E-07	6.95E-04	1.71E-06	1.34E-04	9.04E-04
Lloyd Lake	3.29E-05	3.74E-05	4.00E-07	6.65E-04	1.71E-06	1.34E-04	8.71E-04

		Little Brown My	otis Total Dose F	RFD Case (mGy/	d)		
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose
			Application Cas	e			
Reference (Broach Lake)	1.37E-07	1.55E-07	1.19E-05	1.36E-04	4.92E-07	3.09E-03	3.24E-03
Patterson Lake North Arm - West Basin	1.24E-06	1.41E-06	6.61E-05	2.19E-04	6.74E-07	4.22E-03	4.51E-03
Patterson Lake South Arm	7.73E-07	8.82E-07	5.10E-05	1.86E-04	6.04E-07	3.78E-03	4.02E-03
Beet Lake	3.72E-07	4.24E-07	3.27E-05	1.58E-04	5.08E-07	3.19E-03	3.38E-03
Lloyd Lake	1.58E-07	1.80E-07	1.40E-05	1.38E-04	4.93E-07	3.09E-03	3.24E-03
		Up	pper Bound Scen	ario			
Reference (Broach Lake)	1.37E-07	1.55E-07	1.19E-05	1.36E-04	4.92E-07	3.09E-03	3.23E-03
Patterson Lake North Arm - West Basin	4.42E-06	5.05E-06	1.27E-05	1.76E-04	5.15E-07	3.23E-03	3.43E-03
Patterson Lake South Arm	1.32E-06	1.51E-06	1.24E-05	1.51E-04	4.93E-07	3.10E-03	3.26E-03
Beet Lake	5.78E-07	6.59E-07	1.22E-05	1.42E-04	4.92E-07	3.09E-03	3.24E-03
Lloyd Lake	1.77E-07	2.01E-07	1.19E-05	1.37E-04	4.92E-07	3.09E-03	3.24E-03

		Loon To	tal Dose RFD Cas	e (mGy/d)			
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose
			Application Cas	e			
Reference (Broach Lake)	1.33E-06	1.51E-06	1.01E-06	6.06E-06	8.76E-08	2.97E-03	2.98E-03
Patterson Lake North Arm - West Basin	1.37E-05	1.57E-05	5.65E-06	9.94E-06	1.23E-07	4.09E-03	4.14E-03
Patterson Lake South Arm	1.62E-05	1.85E-05	4.35E-06	8.98E-06	1.15E-07	3.73E-03	3.78E-03
Beet Lake	6.44E-06	7.36E-06	2.78E-06	7.18E-06	9.15E-08	3.08E-03	3.10E-03
Lloyd Lake	1.80E-06	2.05E-06	1.19E-06	6.17E-06	8.78E-08	2.98E-03	2.99E-03
		Uį	pper Bound Scen	ario			
Reference (Broach Lake)	1.33E-06	1.51E-06	1.01E-06	6.06E-06	8.76E-08	2.97E-03	2.98E-03
Patterson Lake North Arm - West Basin	4.31E-05	4.92E-05	1.08E-06	7.81E-06	9.16E-08	3.11E-03	3.21E-03
Patterson Lake South Arm	1.29E-05	1.47E-05	1.05E-06	6.70E-06	8.78E-08	2.98E-03	3.02E-03
Beet Lake	5.62E-06	6.42E-06	1.03E-06	6.34E-06	8.76E-08	2.97E-03	2.99E-03
Lloyd Lake	1.72E-06	1.96E-06	1.01E-06	6.09E-06	8.76E-08	2.97E-03	2.98E-03

Table C.21: Maximum Radiation Dose to Ecological Receptors for the RFD Scenario - Project Lifespan and Far-Future

		Mallard T	otal Dose RFD Ca	ise (mGy/d)			
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose
			Application Case	e			
Reference (Broach Lake)	1.40E-05	1.59E-05	9.79E-06	7.84E-05	1.23E-06	2.55E-02	2.56E-02
Patterson Lake North Arm - West Basin	1.27E-04	1.45E-04	5.46E-05	1.27E-04	1.73E-06	3.49E-02	3.53E-02
Patterson Lake South Arm	8.62E-05	9.84E-05	4.42E-05	1.13E-04	1.60E-06	3.13E-02	3.16E-02
Beet Lake	4.04E-05	4.61E-05	2.76E-05	9.28E-05	1.28E-06	2.63E-02	2.65E-02
Lloyd Lake	1.64E-05	1.87E-05	1.16E-05	7.98E-05	1.23E-06	2.55E-02	2.56E-02
		U _l	oper Bound Scen	ario			
Reference (Broach Lake)	1.40E-05	1.59E-05	9.79E-06	7.84E-05	1.23E-06	2.55E-02	2.56E-02
Patterson Lake North Arm - West Basin	4.53E-04	5.17E-04	1.05E-05	1.01E-04	1.29E-06	2.66E-02	2.77E-02
Patterson Lake South Arm	1.35E-04	1.55E-04	1.02E-05	8.68E-05	1.23E-06	2.55E-02	2.59E-02
Beet Lake	5.91E-05	6.75E-05	1.00E-05	8.20E-05	1.23E-06	2.55E-02	2.57E-02
Lloyd Lake	1.81E-05	2.06E-05	9.82E-06	7.88E-05	1.23E-06	2.55E-02	2.56E-02

		Rusty Blackbi	rd Total Dose RF	D Case (mGy/d))		
Location	Uranium-238	Uranium-234	Thorium-230	Radium-226	Lead-210	Polonium-210	Total Dose
			Application Case	e			
Reference (Broach Lake)	1.01E-04	1.15E-04	2.74E-05	8.28E-04	3.65E-06	6.96E-02	7.07E-02
Patterson Lake North Arm - West Basin	5.12E-04	5.83E-04	1.39E-04	1.39E-03	4.71E-06	9.59E-02	9.85E-02
Patterson Lake South Arm	2.18E-04	2.48E-04	1.07E-04	9.64E-04	4.11E-06	8.54E-02	8.69E-02
Beet Lake	1.55E-04	1.76E-04	6.98E-05	8.98E-04	3.75E-06	7.20E-02	7.33E-02
Lloyd Lake	1.04E-04	1.19E-04	3.18E-05	8.32E-04	3.66E-06	6.98E-02	7.09E-02
		Up	pper Bound Scen	ario			
Reference (Broach Lake)	1.01E-04	1.15E-04	2.74E-05	8.28E-04	3.65E-06	6.96E-02	7.07E-02
Patterson Lake North Arm - West Basin	5.16E-04	5.89E-04	3.02E-05	1.18E-03	3.73E-06	7.28E-02	7.52E-02
Patterson Lake South Arm	2.13E-04	2.43E-04	2.87E-05	9.02E-04	3.65E-06	6.99E-02	7.12E-02
Beet Lake	1.44E-04	1.64E-04	2.81E-05	8.67E-04	3.65E-06	6.97E-02	7.09E-02
Lloyd Lake	1.04E-04	1.19E-04	2.75E-05	8.30E-04	3.65E-06	6.96E-02	7.07E-02

Table C.22: Sample Calculation - Mallard (Patterson Lake South) Dose and Risk Calculations for Copper

				Copper	
Parameter	Symbol	Calculation	Value	Unit	Source
Environmental Media Concentration					
Water Concentration	Cw	-	9.23E-04	mg/L	Appendix C
Sediment Concentration (dry weight)	$C_{sed(dw)}$	-	2.73E+00	mg/kg dw	Appendix C
Outdoor Air Concentration	Cair	-	3.00E-08	mg/m³	Appendix C
Aquatic Plant Concentration				·	
Bioaccumulation Factor (BAF)	BAF _{aquatic plant}	-	2.17E+02	L/kg fw	Table 3-8, IMPACT Model Report
Aquatic Plant Tissue Concentration	C _{aquatic} plant	$C_{\text{aquatic plant}} = C_{\text{w}} * BAF_{\text{aquatic plant}}$	2.00E-01	mg/kg fw	Calculated
Benthic Invertebrate Concentration				·	
Water-to-Sediment Partitioning Coefficient	k _d	-	3.00E+03	L/kg	Table 3-6, IMPACT Model Report
Dry Bulk Density	ρ	-	1.10E-01	kg/L	Assumed
Water Content (Volume Basis)	θ	-	9.60E-01	unitless	Assumed
Fraction in Porewater Phase	C _{pw (fraction)}	$C_{pw (fraction)} = 1 - (k_d * \rho / \theta) / (1 + k_d * \rho / \theta)$	2.90E-03	unitless	Calculated
Concentration Ratio – Porewater to Sediment	Ratio _{pw-sed}	$Ratio_{pw-sed} = C_{pw (fraction)} * \rho / \theta$	3.32E-04	kg/L	Calculated
Sediment Porewater Concentration	C _{pw} (sed)	$C_{pw (sed)} = C_{sed(dw)} * Ratio_{pw-sed}$	9.06E-04	mg/L	Calculated
Bioaccumulation Factor (BAF)	BAF _{benthic inv}	-	7.80E+03	L/kg fw	Table 3-8, IMPACT Model Report
Sediment Occupancy Factor	OF _{sed}	-	1.00E+02	%	Assumed
Benthic Invertebrate Tissue Concentration	C _{benthic} inv	$C_{benthic\ inv} = (1 - (OF_{sed}/100) * C_{sed(dw)} + (OF_{sed}/100) * C_{pw\ (sed)}) * BAF_{benthic\ inv}$	7.07E+00	Bq/kg fw	Calculated
Mallard Exposure Factors					
Water Intake	IR _w	-	6.40E-02	L/d	Table 6-8
Sediment Intake	IRs	-	2.00E-03	kg dw/d	Table 6-8
Air Intake	IR _{air}	-	2.00E-02	m³/d	Table 6-8
Aquatic Plant Intake	IR _{aquatic plant}	-	6.00E-02	kg fw/d	Table 6-8
Benthic Invertebrate Intake	IR _{benthic inv}	-	1.80E-01	kg fw/d	Table 6-8
Body Weight	BW	-	1.13	kg fw	Table 6-8
Toxicological Benchmark	TRV	-	7.52E+01	mg/kg d	Table 6-16
Mallard Dose and HQ					
Total Dose	D _{total}	$D_{total} = (C_w*IR_w + C_{s(dw)}*IR_s + C_{air}*IR_{air} + C_{aquatic\ plant}*IR_{aquatic\ plant} + C_{benthic\ inv}*IR_{benthic\ inv})/BW$	1.14E+00	mg/kg d	Calculated
Hazard Quotient	HQ	HQ = D _{total} /TRV	1.52E-02	unitless	Calculated

Table C.23: Sample Calculation - Adult Subsistence Harvester (Patterson Lake South) Dose and Risk Calculations for Copper

D	Considerati	Calmilation		Сор	pper
Parameter	Symbol	Calculation	Value	Unit	Source
Water Ingestion Dose					
Water Concentration (PLS)	C _{w (PLS)}	-	9.23E-04	mg/L	Appendix C
Water Concentration (Reference)	C _{w (Ref)}	-	8.43E-04	mg/L	Appendix C
Local Intake Fraction of Water (PLS)	IF _{w (PLS)}	-	5.00E-01	Unitless	Assumed
Local Intake Fraction of Water (Reference)	IF _{w (Ref)}	-	5.00E-01	Unitless	Assumed
Water Intake	IR _w	-	1.04E+00	L/d	CSA N288.1-20 (Table 21)
Human Adult Body Mass	BW	-	7.07E+01	kg	Health Canada, 2021
Ingestion Dose (Water)	D _w	$D_{w} = ((C_{w (PLS)} * IF_{w (PLS)} + C_{w (Ref)} * IF_{w (Ref)}) * IR_{w}) / BW$	1.30E-05	mg/kg/d fw	Calculated
Soil Ingestion Dose				•	
Soil Concentration (PLS)	C _{s (PLS)}	-	6.03E-01	mg/kg dw	Appendix C
Soil Concentration (Reference)	C _{s (Ref)}	-	6.01E-01	mg/kg dw	Appendix C
Local Intake Fraction of Soil (PLS)	IF _{s (PLS)}	-	5.00E-01	Unitless	Table 2-10, IMPACT Model Report
Local Intake Fraction of Soil (Reference)	IF _{s (Ref)}	-	5.00E-01	Unitless	Table 2-10, IMPACT Model Report
Soil Intake	IRs	-	4.00E-06	kg dw/d	Table 2-7, IMPACT Model Report
Days Exposed to Soil per Year	Exp _{days}	-	1.35E+02	d/y	Assumed
Human Adult Body Mass	BW	-	7.07E+01	kg	Health Canada, 2021
Ingestion Dose (Soil)	Ds	$D_{s} = (((C_{s (PLS)} * IF_{s (PLS)} + C_{s (Ref)} * IF_{s (Ref)}) * IR_{s}) / BW) * (Exp_{days} / 365)$	1.26E-08	mg/kg/d fw	Calculated
Terrestrial Plant Ingestion Dose	1				
Labrador Tea Concentration (PLS)	C _{LT (PLS)}	-	8.09E-01	mg/kg fw	Appendix C
Blueberry Concentration (PLS)	C _{BB (PLS)}	-	4.01E-01	mg/kg fw	Appendix C
Labrador Tea Concentration (Reference)	C _{LT (Ref)}	-	7.98E-01	mg/kg fw	Appendix C
Blueberry Concentration (Reference)	C _{BB (Ref)}	-	3.99E-01	mg/kg fw	Appendix C
Food Intake (Terrestrial Plants)	IR _{TP}	-	1.24E+01	kg/y fw	Assumed, site-specific
Local Intake Fraction of Labrador Tea (Reference)	IF _{LT (Ref)}	-	9.88E-02	Unitless	Table 2-10, IMPACT Model Report
Local Intake Fraction of Blueberry (Reference)	IF _{BB (Ref)}	-	4.01E-01	Unitless	Table 2-10, IMPACT Model Report
Local Intake Fraction of Labrador Tea (PLS)	IF _{LT (PLS)}	-	9.88E-02	Unitless	Table 2-10, IMPACT Model Report
Local Intake Fraction of Blueberry (PLS)	IF _{BB (PLS)}	-	4.01E-01	Unitless	Table 2-10, IMPACT Model Report
Human Adult Body Mass	BW	-	7.07E+01	kg	Health Canada, 2021
Food Processing Factor	FPF	-	1.00E+00	Unitless	Assumed
Ingestion Dose (Terrestrial Plants)	D _{TP}	$D_{TP} = ((C_{LT (PLS)} * IF_{LT (PLS)} + C_{BB (PLS)} * IF_{BB (PLS)} * IF_{LT (Ref)} * IF_{LT (Ref)} + C_{BB (Ref)} * IF_{BB (Ref)}) * IR_{TP} * FPF) / (BW * 365)$	2.30E-04	mg/kg/d fw	Calculated
Aquatic Animal Ingestion Dose	•	· · · · · · · · · · · · · · · · · · ·			
Northern Pike Concentration (PLS)	C _{NP (PLS)}	-	4.61E-01	mg/kg fw	Appendix C

Table C.23: Sample Calculation - Adult Subsistence Harvester (Patterson Lake South) Dose and Risk Calculations for Copper

				Сор	pper
Parameter	Symbol	Calculation	Value	Unit	Source
Whitefish Concentration (PLS)	C _{WF (PLS)}	-	4.59E-02	mg/kg fw	Appendix C
Northern Pike Concentration (Reference)	C _{NP (Ref)}	-	4.21E-01	mg/kg fw	Appendix C
Whitefish Concentration (Reference)	C _{WF (Ref)}	-	4.21E-02	mg/kg fw	Appendix C
Food Intake (Aquatic Animals)	IR _{AA}	-	8.47E+01	kg/y fw	Assumed, site-specific
Local Intake Fraction of Northern Pike (Reference)	IF _{NP (Ref)}	-	2.50E-01	Unitless	Table 2-10, IMPACT Model Report
Local Intake Fraction of Whitefish (Reference)	IF _{WF (Ref)}	-	2.50E-01	Unitless	Table 2-10, IMPACT Model Report
Local Intake Fraction of Northern Pike (PLS)	IF _{NP (PLS)}	-	2.50E-01	Unitless	Table 2-10, IMPACT Model Report
Local Intake Fraction of Whitefish (PLS)	IF _{WF (PLS)}	-	2.50E-01	Unitless	Table 2-10, IMPACT Model Report
Human Adult Body Mass	BW	-	7.07E+01	kg	Health Canada, 2021
Food Processing Factor	FPF	-	1.00E+00	Unitless	Assumed
Ingestion Dose (Aquatic Animals)	D _{AA}	$D_{AA} = ((C_{NP (PLS)} * IF_{NP (PLS)} + C_{WF (PLS)} * IF_{WF (PLS)} + C_{NP (Ref)} * IF_{NP (Ref)} * IF_{WF (Ref)} * IF_{$	7.97E-04	mg/kg/d fw	Calculated
Terrestrial Animal Ingestion Dose				•	
Moose Concentration (PLS)	C _{M (PLS)}	-	3.65E-01	mg/kg fw	Appendix C
Moose Organs Concentration (PLS)	C _{MO (PLS)}	-	1.46E+01	mg/kg fw	Appendix C
Beaver Concentration (PLS)	C _{B (PLS)}	-	3.81E-01	mg/kg fw	Appendix C
Mallard Concentration (PLS)	C _{Mal (PLS)}	-	1.23E+00	mg/kg fw	Appendix C
Grouse Concentration (PLS)	C _G (PLS)	-	2.61E-01	mg/kg fw	Appendix C
Store Food Concentration (PLS)	C _{SF (PLS)}	-	5.70E-01	mg/kg fw	Appendix C
Moose Concentration (Reference)	C _{M (Ref)}	-	3.51E-01	mg/kg fw	Appendix C
Moose Organs Concentration (Reference)	C _{MO (Ref)}	-	1.40E+01	mg/kg fw	Appendix C
Beaver Concentration (Reference)	C _{B (Ref)}	-	3.78E-01	mg/kg fw	Appendix C
Mallard Concentration (Reference)	C _{Mal (Ref)}	-	1.13E+00	mg/kg fw	Appendix C
Grouse Concentration (Reference)	C _{G (Ref)}	-	2.59E-01	mg/kg fw	Appendix C
Store Food Concentration (Reference)	C _{SF (Ref)}	-	5.70E-01	mg/kg fw	Appendix C
Food Intake (Terrestrial Animals)	IR _{TA}	-	6.09E+02	kg/y fw	Assumed, site-specific
Local Intake Fraction of Moose (PLS)	IF _{M (PLS)}	-	1.69E-02	Unitless	Table 2-10, IMPACT Model Report
Local Intake Fraction of Moose Organs (PLS)	IF _{MO (PLS)}	-	7.52E-03	Unitless	Table 2-10, IMPACT Model Report
Local Intake Fraction of Beaver (PLS)	IF _{B (PLS)}	-	4.68E-02	Unitless	Table 2-10, IMPACT Model Report
Local Intake Fraction of Mallard (PLS)	IF _{Mal (PLS)}	-	2.84E-03	Unitless	Table 2-10, IMPACT Model Report
Local Intake Fraction of Grouse (PLS)	IF _{G (PLS)}	-	1.01E-02	Unitless	Table 2-10, IMPACT Model Report
Local Intake Fraction of Store Food (PLS)	IF _{SF (PLS)}		4.16E-01	Unitless	Table 2-10, IMPACT Model Report
Local Intake Fraction of Moose (Reference)	IF _{M (Ref)}		1.69E-02	Unitless	Table 2-10, IMPACT Model Report
Local Intake Fraction of Moose Organs (Reference)	IF _{MO (Ref)}	-	7.52E-03	Unitless	Table 2-10, IMPACT Model Report

Table C.23: Sample Calculation - Adult Subsistence Harvester (Patterson Lake South) Dose and Risk Calculations for Copper

Parameter	Console al	Calculation	Copper				
	Symbol	Calculation	Value	Unitless Unitless Unitless Unitless Unitless Table 2-10, IMP, Unitless Table 2-10, IMP, Unitless Table 2-10, IMP, Unitless Table 2-10, IMP, Ag Health Ca	Source		
Local Intake Fraction of Beaver (Reference)	IF _{B (Ref)}	-	4.68E-02	Unitless	Table 2-10, IMPACT Model Report		
Local Intake Fraction of Mallard (Reference)	IF _{Mal (Ref)}	-	2.84E-03	Unitless	Table 2-10, IMPACT Model Report		
Local Intake Fraction of Grouse (Reference)	IF _{G (Ref)}	-	1.01E-02	Unitless	Table 2-10, IMPACT Model Report		
Local Intake Fraction of Store Food (Reference)	IF _{SF (Ref)}	-	4.16E-01	Unitless	Table 2-10, IMPACT Model Report		
Human Adult Body Mass	BW	-	7.07E+01	kg	Health Canada, 2021		
Food Processing Factor	FPF	-	1.00E+00	Unitless	Assumed		
Ingestion Dose (Terrestrial Animals)	D_{TA}	$D_{AA} = ((C_{M (PLS)} * IF_{M (PLS)} + C_{MO (PLS)} * IF_{MO (PLS)} * IF_{B (PLS)} * IF_{B (PLS)} * IF_{Mal (PLS)} * IF_{Mal (PLS)} * IF_{Mal (PLS)} + C_{G (PLS)} * IF_{G (PLS)} * IF_{G (PLS)} * IF_{G (Ref)} * IF_{MO (Ref)} * IF_{MO (Ref)} * IF_{MO (Ref)} * IF_{G (Ref)} * IF$	1.77E-02	mg/kg/d fw	Calculated		
Adult Subsistence Harvester Total Dose and HQ							
Total Ingestion Dose	D_{total}	$D_{total} = D_w + D_s + D_{TP} + D_{AA} + D_{TA}$	1.87E-02	mg/kg/d fw	Calculated		
Toxicological Benchmark	TRV	-	4.26E-01	mg/kg d	Table 5-17		
Hazard Quotient	HQ	$HQ = D_{total} / TRV$	4.39E-02	Unitless	Calculated		

Table C.24: Sample Calculation - Mallard (Patterson Lake South) Radiological Dose for Polonium-210

Parameter				Polonium-210			
	Symbol	Calculation	Value	Unit	Source		
Environmental Media Concentrations		·		·			
Water Concentration	Cw	-	5.31E-03	Bq/L	Appendix C		
Sediment Concentration (dry weight)	$C_{s(dw)}$	-	3.74E+02	Bq/kg dw	Appendix C		
Outdoor Air Concentration	Cair	-	1.51E-05	Bq/m³	Appendix C		
Sediment Dry Bulk Density	ρ _s	-	1.10E-01	kg dw/ L	CSA N288.1-20 clause 6.6.2.2		
Mixing Depth	d	-	3.00E-02	m	Assumed		
Sediment Surface Concentration (dry weight)	C _{s(dw)} '	$C_{s(dw)}' = C_{s(dw)} * \rho_s * d * 1000 L/m^3$	1.24E+03	Bq dw/ m ²	Calculated		
Aquatic Plant Concentration							
Bioaccumulation Factor - Aquatic Plant	BAF _{aquatic plant}	-	2.93E+02	L/kg fw	Assumed		
Aquatic Plant Concentration (fresh weight)	Caquatic plant	$C_{aquatic\ plant} = C_w * BAF_{aquatic\ plant}$	1.55E+00	Bq/kg fw	Calculated		
Benthic Invertebrate Concentration							
Water-to-Sediment Partitioning Coefficient	k _d	-	1.20E+05	L/kg	Assumed		
Dry Bulk Density	ρ	-	1.10E-01	kg/L	Assumed		
Water Content (Volume Basis)	θ	-	9.60E-01	unitless	Assumed		
Fraction in Porewater Phase	C _{pw (fraction)}	$C_{pw (fraction)} = 1 - (k_d * \rho / \theta) / (1 + k_d * \rho / \theta)$	7.27E-05	unitless	Calculated		
Concentration Ratio – Porewater to Sediment	Ratio _{pw-sed}	Ratio _{pw-sed} = $C_{pw (fraction)} * \rho / \theta$	8.33E-06	kg/L	Calculated		
Sediment Porewater Concentration	C _{pw (sed)}	$C_{pw (sed)} = C_{sed(dw)} * Ratio_{pw-sed}$	3.12E-03	Bq/L	Calculated		
Bioaccumulation Factor (BAF)	BAF _{benthic inv}	-	1.58E+04	L/kg fw	Assumed		
Sediment Occupancy Factor	OF _{sed}	-	1.00E+02	%	Assumed		
Benthic Invertebrate Tissue Concentration	C _{benthic} inv	$C_{benthic inv} = (1 - (OF_{sed}/100) * C_{sed(dw)} + (OF_{sed}/100) * C_{pw (sed)}) * BAF_{benthic inv}$	4.92E+01	Bq/kg fw	Calculated		
Mallard Exposure Factors							
Water Intake	IR _w	-	6.40E-02	L/d	Table 6-8		
Sediment Intake	IR _s	-	2.00E-03	kg dw/d	Table 6-8		
Air Intake	IR _{air}	-	2.00E-02	m³/d	Table 6-8		
Aquatic Plant Intake	IR _{aquatic plant}	-	6.00E-02	kg fw/d	Table 6-8		
Benthic Invertebrate Intake	IR _{benthic inv}	-	1.80E-01	kg fw/d	Table 6-8		
Fraction of Time Spent on Site	f_0	-	1.00E+00	unitless	Assumed		
Mallard Internal Dose (Radiological)				·			
Ingestion Transfer Factor	TF _{ing}	-	3.68E+00	d/kg fw	NCRP 123 (1996)		
Mallard Tissue Concentration	Ct	$C_t = f_0 * TF_{ing} * (C_w*IR_w + C_{s(dw)} * IR_s + C_{aquatic \ plant} * IR_{aquatic \ plant} + C_{benthic \ inv} * IR_{benthic \ inv})$	3.57E+01	Bq/kg fw	Calculated		
Dose Conversion Factor (Internal) - Duck	DCint	-	3.04E-02	(µGy/hr)/(Bq/kg fw)	ICRP 108 (2008) (Table C.4)		
Internal Dose	D _{int}	$D_{int} = C_t * DC_{int}$	1.09E+00	μGy/hr	Calculated		
Internal Dose (converted units)	D _{int} '	D _{int} ' = D _{int} * 24 h/d / 1000 μGy/mGy	2.61E-02	mGy/d	Calculated		

Table C.24: Sample Calculation - Mallard (Patterson Lake South) Radiological Dose for Polonium-210

Parameter	Counch of	Calculation		Polonium-210		
	Symbol	Calculation	Value	Unit	Source	
Mallard External Dose (Radiological)						
Occupancy Factor, Sediment - Riparian Birds	OFs	-	0.00E+00	unitless	Assumed	
Occupancy Factor, Sediment Surface - Riparian Birds	OF_{ss}	-	5.00E-01	unitless	Assumed	
Dose Conversion Factor (External) - Duck	DC _{ext (on soil)}	-	2.63E-11	(µGy/hr)/(Bq/m²)	Table 4.23	
External Dose	D_ext	$D_{ext} = f_0 * (C_{s(dw)}' * OF_{ss} * DC_{ext (on soil)})$	4.91E-09	μGy/hr	Calculated (Section 4.2.3.1)	
External Dose (converted units)	D _{ext} '	$D_{ext}' = D_{ext} * 24 \text{ h/d} / 1000 \mu \text{Gy/mGy}$	1.18E-10	mGy/d	Calculated	
Mallard Total Dose (Radiological)						
Total Dose	D _{total}	$D_{total} = D_{int} + D_{ext}$	1.09E+00	μGy/hr	Calculated	
Total Dose (converted units)	D _{total} '	$D_{total}' = D_{int}' + D_{ext}'$	2.61E-02	mGy/d	Calculated	

Table C.25: Sample Calculation - Adult Harvester (Patterson Lake South) Radiological Dose for Polonium-210

				Polonium	1-210
Parameter	Symbol	Calculation	Value	Unit	Source
Environmental Media Concentrations			·		
Water Concentration	C _w	-	5.31E-03	Bq/L	Appendix C
Water Concentration reference	C _{w (Ref)}	-	5.15E-03	Bq/L	Appendix C
Water Concentration incremental	C _w '	$C_w' = C_w - C_{w (Ref)}$	1.55E-04	Bq/L	Calculated
Soil Concentration (dry weight)	C _{s(dw)}	-	1.25E+01	Bq/kg dw	Appendix C
Soil Concentration reference (dry weight)	C _{s(dw) (Ref)}	-	1.12E+01	Bq/kg dw	Appendix C
Soil Concentration incremental (dry weight)	C _{s(dw)} '	$C_{s(dw)}' = C_{s(dw)} - C_{s(dw)}$ (Ref)	1.25E+00	Bq/kg dw	Calculated
Outdoor Air Concentration	Cair	-	1.51E-05	Bq/m³	Appendix C
Soil Dry Bulk Density (Sand)	ρ_{s}	-	1.50E+00	kg(dw)/ L	CSA N288.1-20 (cl. 6.3.2.2)
Mixing Depth	d	-	2.00E-01	m	CSA N288.1-20 (cl. 6.3.1.1)
Soil Surface Concentration	C _{ss}	$C_{ss} = C_{s(dw)}' * \rho_s * d * 1000 L/m^3$	3.76E+02	Bq(dw)/m ²	Calculated
Polygon Fraction Outdoor Air	PF _{air}	-	5.00E-01	Unitless	Assumed
Polygon Fraction Soil	PF _s	-	5.00E-01	Unitless	Assumed
Polygon Fraction Water	PFw	-	5.00E-01	Unitless	Assumed
Air Immersion Dose					
Fraction of Time at Location	fo	-	1.00E+00	Unitless	CSA N288.1-20 (cl. 6.14.3)
Outdoor Occupancy Factor	f _u	-	2.00E-01	Unitless	CSA N288.1-20 (cl. 6.14.3)
Building Shielding Factor	S _b	-	5.00E-01	Unitless	CSA N288.1-20 (cl. 6.2.5)
Dose Conversion Factor (Immersion)	DCF _a	-	1.23E-11	(Sv/a)/(Bq/m³)	Assumed
Air Immersion Dose	D _{air}	$D_{air} = C_{air} * f_o * (f_u + (1 - f_u) * S_b) * DCF_a * PF_{air}$	5.57E-17	Sv/a	Calculated
Unit conversion	D _{air} '	$D_{air}' = D_{air} * 1000 \text{ mSv} / \text{Sv}$	5.57E-14	mSv/a	Calculated
Air Inhalation Dose					
Inhalation Rate	IR _{air}	-	5.95E+03	m³/a	CSA N288.1-20 (Table 19)
Dose Conversion Factor (Inhalation)	DCF _{inh}	-	3.30E-06	Sv/Bq	Assumed
Air Inhalation Dose	D _{inh}	$D_{inh} = C_{air} * IR_{air} * DCF_{inh} * PF_{air}$	1.48E-07	Sv/a	Calculated
Unit conversion	D _{inh} '	$D_{inh}' = D_{inh} * 1000 \text{ mSv} / \text{Sv}$	1.48E-04	mSv/a	Calculated
Water Ingestion Dose					
Ingestion Rate	IR _w	-	3.80E+02	L/a	CSA N288.1-20 (Table 21)
Dose Conversion Factor (Food)	DCF _f	-	1.20E-06	Sv/Bq	Assumed
Water Ingestion Dose	D _w	$D_w = C_w' * IR_w * DCF_f * PF_w$	3.54E-08	Sv/a	Calculated
Unit Conversion	D _w '	$D_{w}' = D_{w} * 1000 \text{ mSv / Sv}$	3.54E-05	mSv/a	Calculated
Water Immersion Dose					
Bathtub Correction Factor	D _c	-	7.00E-01	Unitless	CSA N288.1-20 (cl. 6.16.12)
Occupancy Factor - Swimming Surface Water	OF ^s _w	-	1.04E-02	Unitless	CSA N288.1-20 (cl. 6.16.1.3)

Table C.25: Sample Calculation - Adult Harvester (Patterson Lake South) Radiological Dose for Polonium-210

	Combal	Calminsting		Poloniu	m-210
Parameter	Symbol	Calculation	Value	Unit	Source
Occupancy Factor - Pool Water	OF ^p _w	-	3.13E-02	Unitless	CSA N288.1-20 (cl. 7.6.1.2)
Occupancy Factor - Bath Water	OF ^b _w	-	1.40E-02	Unitless	CSA N288.1-20 (cl. 6.16.1.3)
Removal factor (sedimentation)	ρ	-	1.00E+00	Unitless	CSA N288.1-20 (cl. 7.10.4)
Dose Conversion Factor (Water)	DCF _w	-	2.66E-11	Sv/Bq	Assumed
Water Immersion Dose	D _{imm}	$D_{imm} = C_w' * (k_w^s * PF_w + D_c * PF_w * OF_w^b + PF_w * OF_w^p) * DCF_w$	1.06E-16	Sv/a	Calculated
Unit Conversion	D _{imm} '	$D_{imm}' = D_{imm} * 1000 \text{ mSv} / \text{Sv}$	1.06E-13	mSv/a	Calculated
External Soil Dose					
Dose reduction factor from surface roughness and terrain irregularities	f _r	-	6.80E-01	Unitless	CSA N288.1-20 (cl. 6.4.6.3)
Groundshine Shielding Factor	S _g	-	2.00E-01	Unitless	CSA N288.1-20 (cl. 6.14.3)
Dose Conversion Factor (Soil)	DCF _g	-	2.61E-13	(Sv/a)/(Bq/m²)	Assumed
Soil External Dose	Dg	$D_g = C_{ss} * f_o * f_r * (f_u + (1 - f_u) * S_g) * DCF_g * PF_s$	1.20E-11	Sv/a	Calculated
Unit conversion	D _g '	D _g ' = D _g * 1000 mSv / Sv	1.20E-08	mSv/a	Calculated
Internal Soil Dose			·		
Soil Intake Rate	IR _s	-	4.00E-06	kg(dw)/d	CSA N288.1-20 (Table 20)
Soil Exposure Frequency	t _d	-	1.35E+02	d/y	Assumed
Dose Conversion Factor (Food)	DCF _f	-	1.20E-06	Sv/Bq	Assumed
Internal Soil Dose	Ds	$D_s = C_{s(dw)}' * IR_s * t_d * DCF_f * PF_s$	4.07E-10	Sv/a	Calculated
Unit conversion	D _s '	$D_s' = D_s * 1000 \text{ mSv} / \text{Sv}$	4.07E-07	mSv/a	Calculated
Terrestrial Plant Ingestion Dose					
Labrador Tea Concentration (PLS)	C _{LT (PLS)}	-	4.32E+00	Bq/kg fw	Appendix C
Blueberry Concentration (PLS)	C _{BB (PLS)}	-	6.00E-01	Bq/kg fw	Appendix C
Labrador Tea Concentration (Reference)	C _{LT (Ref)}	-	8.67E-01	Bq/kg fw	Appendix C
Blueberry Concentration (Reference)	C _{BB (Ref)}	-	4.33E-01	Bq/kg fw	Appendix C
Labrador Tea Concentration (Incremental)	C _{LT (PLS)} '	$C_{LT (PLS)}' = C_{LT (PLS)} - C_{LT (Ref)}$	3.46E+00	Bq/kg fw	Calculated
Blueberry Concentration (Incremental)	C _{BB} (PLS)	$C_{BB (PLS)}' = C_{BB (PLS)} - C_{BB (Ref)}$	1.66E-01	Bq/kg fw	Calculated
Food Intake (Terrestrial Plants)	IR _{TP}	-	1.24E+01	kg/a fw	Assumed, site-specific
Local Intake Fraction of Labrador Tea (PLS)	IF _{LT (PLS)}	-	9.88E-02	Unitless	Table 2-10, IMPACT Model Report
Local Intake Fraction of Blueberry (PLS)	IF _{BB (PLS)}	-	4.01E-01	Unitless	Table 2-10, IMPACT Model Report
Dose Conversion Factor (Food)	DCF _f	-	1.20E-06	Sv/Bq	Assumed
Ingestion Dose (Terrestrial Plants)	D _{TP}	$D_{TP} = (C_{LT (PLS)}' * IF_{LT (PLS)} + C_{BB (PLS)}' * IF_{BB (PLS)}) * IR_{TP} * DCF_f$	6.07E-06	Sv/a	Calculated
Unit conversion	D _{TP} '	$D_{TP}' = D_{TP} * 1000 \text{ mSv / Sv}$	6.07E-03	mSv/a	Calculated
Aquatic Animal Ingestion Dose					
Northern Pike Concentration (PLS)	C _{NP (PLS)}	- -	7.96E-01	Bq/kg fw	Appendix C

Table C.25: Sample Calculation - Adult Harvester (Patterson Lake South) Radiological Dose for Polonium-210

				Poloniı	ım-210
Parameter	Symbol	Calculation	Value	Unit	Source
Whitefish Concentration (PLS)	C _{WF (PLS)}	-	7.14E-01	Bq/kg fw	Appendix C
Northern Pike Concentration (Reference)	C _{NP (Ref)}	-	7.73E-01	Bq/kg fw	Appendix C
Whitefish Concentration (Reference)	C _{WF (Ref)}	-	6.94E-01	Bq/kg fw	Appendix C
Northern Pike Concentration (Incremental)	C _{NP (PLS)} '	$C_{NP (PLS)}' = C_{NP (PLS)} - C_{NP (Ref)}$	2.33E-02	Bq/kg fw	Calculated
Whitefish Concentration (Incremental)	C _{WF (PLS)} '	$C_{WF (PLS)}' = C_{WF (PLS)} - C_{WF (Ref)}$	1.99E-02	Bq/kg fw	Calculated
Food Intake (Aquatic Animals)	IR _{AA}	-	8.47E+01	kg/a fw	Assumed, site-specific
Local Intake Fraction of Northern Pike (PLS)	IF _{NP (PLS)}	-	2.50E-01	Unitless	Table 2-10, IMPACT Model Report
Local Intake Fraction of Whitefish (PLS)	IF _{WF (PLS)}	-	2.50E-01	Unitless	Table 2-10, IMPACT Model Report
Dose Conversion Factor (Food)	DCF _f	-	1.20E-06	Sv/Bq	Assumed
Ingestion Dose (Aquatic Animals)	D _{AA}	$D_{AA} = ((C_{NP (PLS)}' * IF_{NP (PLS)} + C_{WF (PLS)}' * IF_{WF (PLS)}) * IR_{AA} * DCF_f$	1.10E-06	Sv/a	Calculated
Unit conversion	D _{AA} '	$D_{AA}' = D_{AA} * 1000 \text{ mSv} / \text{Sv}$	1.10E-03	mSv/a	Calculated
Terrestrial Animal Ingestion Dose					
Moose Concentration (PLS)	C _{M (PLS)}	-	1.04E+00	Bq/kg fw	Appendix C
Moose Organs Concentration (PLS)	C _{MO (PLS)}	-	1.04E-02	Bq/kg fw	Appendix C
Beaver Concentration (PLS)	C _{B (PLS)}	-	4.08E-01	Bq/kg fw	Appendix C
Mallard Concentration (PLS)	C _{Mal (PLS)}	-	3.57E+01	Bq/kg fw	Appendix C
Grouse Concentration (PLS)	C _{G (PLS)}	-	1.88E+00	Bq/kg fw	Appendix C
Store Food Concentration (PLS)	C _{SF (PLS)}	-	4.80E-02	Bq/kg fw	Appendix C
Moose Concentration (Reference)	C _{M (Ref)}	-	3.65E-01	Bq/kg fw	Appendix C
Moose Organs Concentration (Reference)	C _{MO (Ref)}	-	3.65E-03	Bq/kg fw	Appendix C
Beaver Concentration (Reference)	C _{B (Ref)}	-	2.94E-01	Bq/kg fw	Appendix C
Mallard Concentration (Reference)	C _{Mal} (Ref)	-	3.49E+01	Bq/kg fw	Appendix C
Grouse Concentration (Reference)	C _{G (Ref)}	-	1.30E+00	Bq/kg fw	Appendix C
Store Food Concentration (Reference)	C _{SF (Ref)}	-	4.80E-02	Bq/kg fw	Appendix C
Moose Concentration (Incremental)	C _{M (PLS)} '	$C_{M (PLS)}' = C_{M (PLS)} - C_{M (Ref)}$	6.78E-01	Bq/kg fw	Calculated
Moose Organs Concentration (Incremental)	C _{MO (PLS)} '	$C_{MO (PLS)}' = C_{MO (PLS)} - C_{MO (Ref)}$	6.78E-03	Bq/kg fw	Calculated
Beaver Concentration (Incremental)	C _{B (PLS)} '	$C_{B (PLS)}' = C_{B (PLS)} - C_{B (Ref)}$	1.14E-01	Bq/kg fw	Calculated
Mallard Concentration (Incremental)	C _{Mal} (PLS)	$C_{Mal (PLS)}' = C_{Mal (PLS)} - C_{Mal (Ref)}$	8.64E-01	Bq/kg fw	Calculated
Grouse Concentration (Incremental)	C _{G (PLS)} '	$C_{G (PLS)}' = C_{G (PLS)} - C_{G (Ref)}$	5.78E-01	Bq/kg fw	Calculated
Store Food Concentration (Incremental)	C _{SF (PLS)} '	$C_{SF (PLS)}' = C_{SF (PLS)} - C_{SF (Ref)}$	0.00E+00	Bq/kg fw	Calculated
Local Intake Fraction of Moose (PLS)	IF _{M (PLS)}	-	1.69E-02	Unitless	Table 2-10, IMPACT Model Report
Local Intake Fraction of Moose Organs (PLS)	IF _{MO (PLS)}	-	7.50E-03	Unitless	Table 2-10, IMPACT Model Report
Local Intake Fraction of Beaver (PLS)	IF _{B (PLS)}	-	4.68E-02	Unitless	Table 2-10, IMPACT Model Report
Local Intake Fraction of Mallard (PLS)	IF _{Mal (PLS)}	-	2.85E-03	Unitless	Table 2-10, IMPACT Model Report

Table C.25: Sample Calculation - Adult Harvester (Patterson Lake South) Radiological Dose for Polonium-210

Dayamatay	Councile of	Calculation	Polonium-210		
Parameter	Symbol	Calculation	Value	Unit Source Unitless Table 2-10, IMPACT Model Report	
Local Intake Fraction of Grouse (PLS)	IF _{G (PLS)}	-	1.01E-02	Unitless	Table 2-10, IMPACT Model Report
Local Intake Fraction of Store Food (PLS)	IF _{SF (PLS)}	-	1.68E-01	Unitless	Table 2-10, IMPACT Model Report
Food Intake (Terrestrial Animals)	IR _{TA}	-	6.09E+02	kg/a fw	Assumed, site-specific
Dose Conversion Factor (Food)	DCF _f	-	1.20E-06	Sv/Bq	Assumed
Ingestion Dose (Terrestrial Animals)	D _{TA}	$D_{TA} = ((C_{M (PLS)}' * IF_{M (PLS)} + C_{MO (PLS)}' * IF_{MO (PLS)} + C_{B (PLS)}' * IF_{B (PLS)} + C_{Mal (PLS)}' * IF_{Mal (PLS)} + C_{G (PLS)}' * IF_{G (PLS)} + C_{SF (PLS)}' * IF_{SF (PLS)}) * IR_{TA} * DCF_{f}$	1.83E-05	Sv/a	Calculated
Unit conversion	D _{TA} '	$D_{TA}' = D_{TA} * 1000 \text{ mSv} / \text{Sv}$	1.83E-02	mSv/a	Calculated
Harvester Total Dose					
Total Dose	D _{total}	$D_{total} = D_{air}' + D_{inh}' + D_{w}' + D_{imm}' + D_{g}' + D_{s}' + D_{TP}' + D_{AA}' + D_{TA}'$	2.57E-02	mSv/a	Calculated